

Wnt5a Exhibits Layer-Specific Expression in Adult Skin, Is Upregulated in Psoriasis, and Synergizes with Type 1 Interferon

Malgorzata Romanowska¹, Alan Evans¹, David Kellock², Susan E. Bray², Kathleen McLean², Susanne Donandt³, John Foerster^{1,4*}

1 Division of Experimental Medicine, University of Dundee, Dundee, Scotland, United Kingdom, **2** Tayside Tissue Bank, University of Dundee, Dundee, Scotland, United Kingdom, **3** Department of Dermatology, University of Dundee, Dundee, Scotland, United Kingdom, **4** Charité Medical University, Berlin, Germany

Abstract

Background: Wnt5a is a member of the wingless-type patterning regulators important in pre-natal development. The expression and distribution of Wnt5a and its receptors frizzled (fzd) 3 and fzd 5 in adult human skin have not been comprehensively studied to date.

Methodology/Principal Findings: We here show that Wnt5a, fzd3, fzd5, as well as fzd6 are restricted to specific layers in normal epidermis, analogous to their zonal distribution in hair follicles, suggesting a role in adult skin differentiation. In line, Wnt5a and fzd5 are both overexpressed and re-distributed in the epidermis of psoriasis which involves disturbed keratinocyte differentiation. Functionally, Wnt5a lowers the concentration of IFN required to induce target genes, and increases the magnitude of IFN target gene induction, suggesting a molecular mechanism underlying IFN hypersensitivity in psoriasis. Finally, we identify nedd8 and the amyloid precursor APP, previously shown to be upregulated in psoriasis, as targets of synergistic IFN α /Wnt5a induction.

Conclusions/Significance: The present data (i) suggest that Wnt5a regulates epidermal differentiation even in adult skin and (ii) identify synergistic induction of type 1 IFN target genes as a novel mode of Wnt5a action. Targeting Wnt5a in the skin may reduce IFN hypersensitivity and be of therapeutical value.

Citation: Romanowska M, Evans A, Kellock D, Bray SE, McLean K, et al. (2009) Wnt5a Exhibits Layer-Specific Expression in Adult Skin, Is Upregulated in Psoriasis, and Synergizes with Type 1 Interferon. PLoS ONE 4(4): e5354. doi:10.1371/journal.pone.0005354

Editor: Elizabeth Didier, Tulane National Primate Research Center, United States of America

Received: November 12, 2008; **Accepted:** March 17, 2009; **Published:** April 28, 2009

Copyright: © 2009 Romanowska et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: J.F. is supported by grants of Tenovus Scotland and Cancer Research UK. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: j.foerster@dundee.ac.uk

Introduction

Wnt proteins regulate diverse biological processes. Wnt5a represents the prototypical so-called non-canonical Wnt family members which signal not by inducing, or may even inhibit, the transcriptional activation of β -catenin. Apart from its complex role in development, Wnt5a is expressed in various adult tissues. Among its diverse biological activities, Wnt5a promotes proliferation in endothelial cells and glioblastoma cells [1–3], as well as adhesion in fibroblasts and breast cancer cells [4,5]. Most recently, evidence has accumulated for the involvement of Wnt5a in inflammatory responses and innate immunity. Thus, Wnt5a is induced by Toll-like receptors and elicits secretion of IL12 and TNF α in response to mycobacterial infection [6]. Furthermore, Wnt5a is drastically upregulated in macrophages by LPS and IFN γ and induces the production of the pro-inflammatory cytokines IL1, IL6, and IL8 [7]. Wnt5a is also upregulated in synovial fibroblasts of inflammatory joints in rheumatoid arthritis where it induces IL6, IL8, and IL15 [8].

The signal transduction of Wnt5a appears to be context-dependent [9]. In all of the inflammatory responses cited above, as

well as in endothelial cell activation, the active receptor appears to be Frizzled (Fzd) 5 [10]. However, signalling through Fzd3, and Ror2 have been described [4,9].

The epidermis represents a dynamic system of highly regulated keratinocyte stem-cell renewal, proliferation, differentiation, and migration. At the same time, the epidermis acts as first-line barrier in innate immunity and, as such, maintains complex communication with resident dendritic cells and sub-epidermal endothelial cells and fibroblasts. Although Wnt5a is expressed in all of these cell types and its role in the development of skin appendages including hair follicles [11] and mammary glands [12] has been examined, surprisingly little is known about the expression, distribution, and function of Wnt5a in post-natal human epidermis.

Psoriasis is a very common chronic inflammatory skin disease with a prevalence of 2–4% in Caucasian populations. In psoriasis, keratinocyte proliferation as well as differentiation are perturbed, leading to a grossly thickened epidermis with an altered composition of normal epidermal layers. Apart from keratinocyte turnover, psoriasis involves a hyperactive innate immune response. Thus, iatrogenic administration of type 1 IFN (IFN α or IFN β)

elicits disease flares in susceptible individuals, as does endogenous IFN release in the course of viral disease [13,14]. Moreover, keratinocytes from psoriasis patients are hypersensitive to type 1 IFN [15] and epidermal dendritic cells produce IFN α leading to massive upregulation of type 1 IFN target genes [16]. Indeed, hypersensitivity towards type 1 IFN induced by deletion of the IFN repressor IRF2 is sufficient to cause a psoriasis-like skin disease in mice [17].

We and others have previously found Wnt5a to be upregulated on the mRNA level in psoriasis [16,18]. Here, we show that Wnt5a and its putative receptors are localized to specific compartments in normal skin and that Wnt5a is both highly upregulated and re-distributed in the hyperproliferative epidermis in psoriasis. Functionally, we show that Wnt5a sensitizes keratinocytes towards type 1 IFN. Finally, we identify nedd8 and APP as novel targets of synergistic IFN/Wnt5a activation.

Materials and Methods

Immunohistochemistry

Paraffin-embedded samples were obtained from the Tayside Tissue bank. Prior to biopsy, patients gave written consent to storage and analysis of biopsy samples. Storage and use of all tissues included in the work presented here was approved by the Tayside Committee on Medical Research Ethics B (REC ref. Nr. 07/S1402/90). Antibodies used were anti wnt5a (R&D, order nr. AF645, final dilution 1:400), anti-frizzled 3 (Insight Biotech, ordered through Acris Antibodies, Germany, order nr. SP4568P, 1:200), anti-frizzled 5 (Cambridge Bioscience, ARP41245_P050, 1:800), anti-frizzled 6 (GenWay, 18-461-10538, ordered through Acris Antibodies, Germany; 1:400). A detailed protocol is given in Supporting Methods S1.

Keratinocyte primary culture and western blot

All studies on human samples were done with prior approval of the local institutional ethics committee. Processing of biopsy samples and primary keratinocyte culture was performed as described [16]. Cells were fractionated using the NE-PER kit (Pierce, order-nr 78833) and 20 μ g of protein loaded per lane onto 15% acrylamide-SDS gels. Blotting was performed on a semi-dry blotter onto nitro-cellulose membranes (Schleicher & Schell, 10 402 594). Blots were blocked in Tris-buffered saline containing 0.1% Tween-20 (TBST) and incubated with primary antibodies in TBST containing 4% dry milk as follows: rabbit anti-human nedd8 (Cell Signaling, #2745) and rabbit anti-human Amyloid β precursor protein (Biozol, ab12269) at 1:1000, respectively, overnight at 4°C. Secondary antibody was anti-rabbit-HRP (Amersham Biosciences, NA934) at 1:2000 for 1 h at RT. Blots were developed with ECL Plus (Amersham #RPN2 132) and scanned with a CCD camera.

Stimulation with Wnt5a-mimic peptide and IFN α

Formylated hexapeptide formyl-MDGCEL [5] was synthesized by Pepscan Systems, reconstituted to 10 mM in PBS, aliquoted and stored at -80°C. Primary keratinocytes at 50% confluence were stimulated with 100 μ M final concentration of Wnt5a-mimic peptide for 18 h. Stimulation with recombinant human IFN α (PBL Biomedical Labs, order-nr 3410) was carried out analogously with 20 ng/ml final concentration. For the concentration response, final concentrations were 3, 10, 30, 100 und 300 ng/ml, respectively.

Expression profiling

RNA was isolated and reverse transcribed to cDNA as described [16]. cDNA was fluorescence-labelled, hybridized to SkinPatho microarrays (Miltenyi Biotech), and scanned by the in-house microarray facility of the Charité Medical University, Berlin, Germany. Raw data were adjusted for red/green bias. Excel-based auto-filtering was used to exclude all transcripts lower than ten fluorescence units above the detection threshold in both the red- and green-labelled samples. P-values were determined using a standard t-test as indicated in the respective tables.

Wnt5a transfection

Full-length human wnt5a cDNA was purchased from Origene. Five $\times 10^5$ HaCat keratinocytes were cultivated in RPMI1640 medium and transfected in Advanced Medium (Gibco, order-nr 12491015) containing 2% FCS with 40 μ g Wnt5a-pCMV-XL4 or control vector (pCDNA6.1) using CaCl₂ precipitation as previously described for transfection of lentiviral vectors [19].

RT-PCR

RNA was extracted using the NucleoSpin II extraction kit (Machery-Nagel, #740 955) and cDNA prepared using SuperScript II RT, RnaseOut, Oligo(dT)₁₂₋₁₈ from Invitrogen. For PCR reactions, annealing temperature was uniformly 55°C. All genes were amplified for 25 cycles, except for wnt5a and IFI27 (30 cycles). Primers and conditions are detailed in Supporting Methods S1.

Results

Dysregulation of Wnt-signalling in hyperproliferative epidermis

Psoriasis is a skin disease characterized by hyperproliferation, as well as aberrant differentiation of epidermal keratinocytes. In the course of extensive expression profiling we previously noted that the non-canonical Wnt family member Wnt5a was greatly upregulated in psoriatic epidermis [16]. Accordingly, we carried out a more comprehensive expression analysis of Wnt-signalling molecules in psoriatic skin. As shown in table 1, only Wnt5a and Fzd6 were significantly upregulated whereas the β -catenin inhibitor ICAT was downregulated (shown in bold print), in confirmation of previous findings [18,20].

Expression of Wnt5a and Fzd receptors in adult human epidermis

Before analyzing Wnt5a expression in psoriasis on the protein level, we first performed a detailed immunohistochemical expression analysis in normal adult human skin which has not been published to date. We included Fzd3 and Fzd5 in this analysis, since these have been reported to bind Wnt5a, as well as Fzd6, as it was found upregulated in psoriasis (table 1) and squamous cell carcinoma [18]. The epidermal expression profiles are shown in figure 1. Wnt5a is mainly expressed in the basal layer and few cells in the upper spinous and granular layers. Fzd3 was found to be strictly confined to the spinous layer in the epidermis and was the only receptor exhibiting a non-homogeneous patchy expression pattern such that strongly positive cells are interspersed with Fzd3-negative cells in a mosaic-like pattern. Fzd5 expression is strictly limited to the granular layer (magnified inset) whereas Fzd6, similar to Wnt5a, is mainly found in the basal layer, corresponding to its distribution in murine skin [21] (fig. 1e therein). In the hair follicle, Fzd 3 is mainly expressed in the inner root sheath, Fzd 5 in the precortex, and Wnt5a in the papilla

(fig. 2). These findings are consistent with previous data obtained in postnatal murine skin [11,22], thereby also confirming the specificity of the antibodies employed here. In addition, we also observe Fzd6 expression in the cortex. Taken together, the highly compartmentalized distribution of Wnt5a and Fzd isoforms in adult human epidermis is reminiscent of their zonal restriction in hair follicles (see fig. 2, lower right), strongly suggesting that Wnt signalling regulates differentiation not only in hair follicles, but also in the epidermis.

Expression of Wnt5a and receptors in psoriasis

We next characterized the expression of Wnt5a, Fzd3, Fzd5, Fzd6 in psoriasis lesions which are characterized by hyperproliferation and dysregulated differentiation. As shown in figure 3, Wnt5a was found to be overexpressed throughout the entire spinous layer. Moreover, as shown in the top panel on the right, activated endothelial cells and dermal fibroblasts are strongly positive for Wnt5a. Finally, the neutrophil aggregates in the stratum corneum, a hallmark of psoriasis, also express Wnt5a (arrow, top left panel). While Fzd3 shows an expression pattern qualitatively unchanged to normal epidermis, Fzd5 is shifted from its exclusive expression in the granular layer towards overexpression in all epidermal layers as well as expression in the dermal endothelia (high power image on right). Fzd6 is also heavily overexpressed as well as in the endothelia but retains the same relative expression level as in normal epidermis (strongest staining in the basal layer). Taken together, Wnt5a, Fzd5, and Fzd6 are strongly upregulated in psoriasis. In addition, Wnt5a and Fzd5 exhibit an altered qualitative distribution.

Overexpression of Wnt5a in keratinocytes from psoriasis patients in vitro

We next sought to clarify whether any of the observed changes represent a cell-autonomous feature of the hyperproliferative keratinocytes in psoriasis patients, i.e. whether they can be detected in vitro in the absence of the inflammatory micro-environment. To this end, we expanded keratinocytes from non-lesional skin of psoriasis patients, or healthy control patients, *in vitro* and performed microarray-based expression profiling. As shown in table 2, Wnt5a was found significantly upregulated on the mRNA level in keratinocytes derived from psoriasis patients. This was not the case for Fzd3 and Fzd5 (Fzd6 was not represented on the Skin-Patho microarray used for this experiment). By western blot, we were able to confirm that Wnt5a protein is indeed overexpressed in primary human keratinocytes expanded from non-lesional skin of psoriasis patients (fig. 4). Although the low sample number precludes a definite statement, these data suggest that upregulation of Wnt5a may constitute of cell-autonomous feature of psoriatic keratinocytes. Regardless of this notion, the clear difference in Wnt5a expression between the psoriasis- and control-derived keratinocytes characterized here is suitable to study effects of endogenous Wnt5a overexpression (see below).

Wnt5a induces type I interferon target genes

In order to screen for cellular signalling pathways possibly affected by Wnt5a in keratinocytes, we stimulated keratinocytes in vitro with a formylated wnt5a-derived hexapeptide shown to mimic the effect of full-length wnt5a on adherence and motility in vivo and in vitro [5,23] and compared global gene expression between keratinocytes stimulated with Wnt5a-mimic peptide and control peptide. We performed this experiment both with keratinocytes from psoriasis patients and from control patients.

Table 1. Expression of Wnt-related genes in psoriasis.¹

Gene	Location	p-value	fold change
Wnt proteins			
WNT1	12q13	-	n.d.
WNT2	7q31	0.83	1.07
WNT2B	1p13	-	n.d.
WNT4	1p36.23-p35.1	0.78	-1.06
WNT5A	3p21-p14	0.0004	4.54
WNT6	2q35	0.017	1.79
WNT7A	3p25	-	n.d.
WNT8B	10q24	-	n.d.
WNT10B	12q13	0.24	-1.37
WNT11	11q13.5	0.14	-1.53
Frizzled-related proteins			
FZD1	7q21	0.18	-1.42
FZD2	17q21.1	-	n.d.
FZD3	17q21.1	0.39	-1.15
FZD4	11q14.2	0.75	1.15
FZD5	2q33-q34	0.18	1.34
FZD6	8q22.3-q23.1	0.027	1.99
FZD7	2q33	0.056	-1.58
FZD9	7q11.23	-	n.d.
FRZB	2qter	0.45	-1.41
SFRP1	8p12-p11.1	0.16	-1.49
SFRP4	7p14.1	0.10	-1.45
Other membrane receptors			
LRP5	11q13.4	0.79	-1.08
LRP6	12p11-p13	0.37	1.07
LRP1	12q13-q14	0.27	-1.30
ROR2	9q22	-	n.d.
RYK	3q22	0.22	-1.38
Wnt-inhibitors			
WIF1	12q13.13	0.24	-1.58
DKK1	10q11.2	0.39	-1.47
DKK4	8p11.2-11.1	0.37	1.12
Downstream effectors			
DVL3	3q27	0.99	-1.01
GSK3B	3q13.3	0.09	1.26
APC	5q21-q22	-	n.d.
CTNNB1	3p21	0.95	1.02
CTNNBIP1	1p36.22	0.006	-2.42

¹Expression of wnt-signalling related molecules was analyzed using the Affymetrix U95A microarray. Data shown represent fold-changes in psoriatic lesional vs. non-lesional skin of n = 5 independent donors, as detailed in [16]. **Bold**-print: p < 0.05 and fold-change > 1.5; n.d.: not detected. doi:10.1371/journal.pone.0005354.t001

Since there were no significant differences in the effect of the Wnt5a-mimic peptide on keratinocytes from psoriasis patients and controls (not shown), both groups were analyzed together. Of the 1133 transcripts represented on the Piquor SkinPatho microarray, only four exhibited a greater than 1.2-fold change: nedd8, MX1 (IFI-78), IFI-27, and APP (table 3). Both MX1 and IFI-27 are type




Figure 1. Expression of Wnt5a, Fzd3, Fzd5, and Fzd6 in adult human epidermis. Immunohistochemistry of paraffin-embedded skin samples was performed as detailed in Methods. Panels shown are at 200 \times magnification. doi:10.1371/journal.pone.0005354.g001

1 interferon target genes previously also found to be upregulated in psoriatic skin lesions [16]. Induction of all four target genes could subsequently be verified by RT-PCR of HaCat keratinocytes transfected either with full-length Wnt5a cDNA or control vector (fig. 5a). Importantly, type 1 IFN genes themselves (α or β) were not induced by Wnt5a (not shown).

Wnt5a causes hypersensitivity toward type 1 interferon

Since the magnitude of mRNA upregulation was rather modest for an IFN target gene, we hypothesized that a possible function of Wnt5a might be to "prime" cells for synergistic induction by type 1 IFN. To test this hypothesis, we transfected HaCat keratinocytes either with full-length Wnt5a cDNA or control vector. Transfected cells were then exposed to increasing concentrations of IFN α and expression of the IFN α target gene MX1 determined by PCR. As shown in figure 5b, concurrent overexpression of Wnt5a caused a left shift in the concentration response curve to all IFN α where the calculated apparent EC-50 shifted from 22 ng/ml to 6 ng/ml, suggesting that overexpression of Wnt5a indeed increases sensitivity toward IFN α . This observation predicts that low concentrations of IFN α should cause a greater magnitude of IFN target gene induction in cells with increased Wnt5a expression. To test this, we compared the response to type 1 IFN between primary epidermal keratinocytes from psoriasis patients harbouring increased endogenous Wnt5a protein levels (see above, fig. 4) to keratinocytes from control probands. We stimulated these cells with 20 ng/ml of IFN α , a concentration which, according to the data presented above, would be expected

to cause a higher magnitude of IFN α target gene induction in keratinocytes with endogenous Wnt5a overexpression. After 18 h of stimulation, global gene expression was analyzed using the Piquor SkinPatho microarray. Figure 5c shows all transcripts (41 of 1133) present on the array which were induced at least a two-fold by IFN α . The induction level is expressed as mean fold-change of all psoriasis keratinocytes ($n = 3$, along the y-axis) vs. mean fold-change of all control keratinocytes ($n = 4$, along the x-axis). Strikingly, all 41 IFN-responsive target genes are induced to a higher level in the keratinocytes harbouring higher Wnt5a levels, corresponding to a statistically highly significant 1.8 ± 1.1 -fold relative hyper-induction across all target genes ($p = 0.001$). These data confirm that hypersensitivity toward IFN α , i.e. increased IFN target gene induction at low IFN concentrations, is a general feature detectable for all target genes in primary cells with endogenous Wnt5a upregulation.

APP and nedd8 are synergistically induced by type 1 IFN and Wnt5a

Apart from MX1 and IFI-27, the only transcripts induced by Wnt5a in keratinocytes were APP and nedd8 (see above, table 3 and figure 5). Wnt5a alone showed only a modest induction of both genes in keratinocytes. Likewise, stimulation by IFN α alone only caused a slight upregulation of either gene (1.4 ± 0.6 fold and 1.6 ± 0.4 fold for APP and nedd8, respectively). Therefore, we hypothesized that these genes, too, might be subject to synergistic induction by Wnt5a and type 1 IFN. To test this hypothesis we transfected HaCat keratinocytes either with control vector or full-




Figure 2. Expression of Wnt5a and Fzd3, Fzd5, and Fzd6 in adult human hair follicles. Immunohistochemistry of paraffin-embedded skin samples was performed as detailed in Methods. Panels shown are at 200 \times magnification. The hair follicle schematic for the cartoon on lower right was adapted from [11].

doi:10.1371/journal.pone.0005354.g002

length Wnt5a, and cultivated the transfected cells either in the presence or absence of 20 ng/ml IFN α . As shown in fig. 6a, a specific band at appr. 70 kd, corresponding to soluble APP, was induced upon either Wnt5a or IFN α stimulation, but more strongly upon co-stimulation with both factors. Synergistic induction of soluble APP is also evident upon densitometric quantification of the blot, as shown on the bottom of the figure. Nedd8 is a ubiquitin-like peptide covalently linked to a number of target proteins. In nuclear extracts of HaCat cells, a nedd8-reactive protein migrating at appr. 50 kd was present in control cells, but more abundant after either Wnt5a-transfection or IFN α stimulation and clearly synergistically induced by both factors (fig. 6b), which is also seen upon densitometric quantification. In conjunction with the transcriptional induction of nedd8 by IFN α as well as Wnt5a seen on the mRNA level (figure 4a), these data strongly suggest that nedd8 is synergistically induced by Wnt5a and IFN α . Taken together, APP (previously implicated in psoriasis pathogenesis) and nedd8 represent novel interferon targets induced significantly by the concurrent action of Wnt5a and IFN α .

Discussion

We here show that Wnt5a and its putative receptor Fzd5 are overexpressed in psoriasis, that Wnt5a increases the sensitivity of keratinocytes towards type 1 interferon, and that nedd8 and APP are synergistically induced by Wnt5a and IFN α .

The unexpected finding that Wnt5a and fzd3, fzd5, and fzd6 proteins are highly compartmentalized in normal adult human epidermis, reminiscent of the distribution of wnt signalling

components in the hair follicle, raises the question of the role of this system in the dynamics of keratinocyte turnover in the skin. Of all relevant classical knock-out phenotypes, only Fzd6-deficient mice survive post term with intact epidermis except for abnormal hair whorls [21], suggesting that its physiological presence in the basal layer is dispensable. Therefore, inducible loss- or gain-of-function approaches will be required to delineate the functional role of wnt signalling in post-natal skin.

The marked overexpression of Wnt5a and Fzd5 in psoriasis suggest that this ligand-receptor pair may actively drive the chronic inflammatory and hyperproliferative nature of this phenotype. This idea receives further support from a large body of circumstantial evidence. First, WNT5a induces IL12 [6], which is central to psoriasis, as indicated by the therapeutic efficacy of antibodies targeted at IL12p40 and by the genetic association of IL12 with psoriasis [24,25]. Second, Wnt5a is itself strongly induced by STAT3 [26], thereby acting as a potential effector molecule for the psoriasis-like skin disease seen in STAT3-overexpressing transgenic mice [27]. Third, WNT5a induces epithelial differentiation and stimulates endothelial cell proliferation, both important elements of psoriasis. Fourth, Wnt5a is upregulated in wound healing [28] and psoriasis lesions can classically be triggered by skin wounding. Fifth, our data show that dermal fibroblasts in psoriasis are strongly positive for Wnt5a and activated fibroblasts from rheumatoid arthritis synthesize inflammatory cytokines, most notably IL15, subsequent to Wnt5a/Fzd5 signalling [29]. Sixth, Wnt5a derived from endothelia signals through Fzd5 to induce the translocation of NFAT and subsequent induction of IL-2 in T cells [10]. This effect may be highly relevant, since (i) continued T cell activation also represents a central element in psoriasis [30], and (ii) Wnt5a-




Figure 3. Expression of Wnt5a and Fzd proteins in psoriasis. (a) Immunohistochemistry of Wnt5a was performed as detailed in Methods. Panels on left are 200 \times , panels on right are at 400 \times magnification. Wnt5a staining was virtually identical in $n=12$ samples from independent patients.

doi:10.1371/journal.pone.0005354.g003

mediated T cell activation can be elicited by lithium which is one of the most potent clinical triggers of psoriasis flares [31]. Finally, and most importantly, WNT5a induces IFN hypersensitivity (see below), a central hallmark of psoriasis.

The relationship between Wnt5a and β -catenin signalling in psoriasis as well as physiological epidermal turn-over is unclear. Supra-basal accumulation of nuclear β -catenin in psoriasis has been reported [32]. Moreover, we here report down-regulation of two inhibitors of canonical wnt-signalling in psoriasis, ICAT (CTNNBIP1) and WIF (table 1). Furthermore, the psoriasis-inducing effect of lithium mentioned above may also involve GSK β -inhibition and β -catenin activation in vivo. Thus, Wnt5a and β -catenin activation may occur concurrently in psoriasis. Indeed, retroviral overexpression of Wnt5a after wounding in mice did not antagonize nuclear translocation of β -catenin [28].

Wnt5a signalling in psoriasis may also be related to the activation of the nuclear hormone receptor PPAR δ which we have previously identified as a central mediator in psoriasis. Thus, Wnt5a may synergize with IFN α to induce the expression of PPAR δ in activated T lymphocytes [29]. Moreover, PPAR δ stimulates keratinocyte proliferation and in several biological processes acts by antagonizing another PPAR isoform, PPAR γ [16]. Wnt5a also represses PPAR γ transactivation thereby synergizing with PPAR δ functionally [33]. Conversely, PPAR γ inhibits STAT3 [34] which, when overexpressed, induces Wnt5a and causes a psoriasis-like phenotype in vivo [27].

The molecular mechanism of Wnt5a/IFN α crosstalk is elusive as no data exist on the effect of frizzled-mediated signalling on downstream effectors of IFN signalling. Our results do suggest that Fzd5 may be the responsible molecule at the receptor level and

Table 2. Genes with altered expression in cultured psoriatic keratinocytes.¹

	FC in vitro ²	p-value ³	FC skin ⁴	p-value skin ⁴
Increased expression				
KYNURENINASE	5.5	0.003	3.4	0.06
COL3A1	3.2	0.006	2.9	0.018
COL5A2	3.1	0.006	n.s.	n.s.
COL6A1	3.0	0.000	n.s.	n.s.
WNT5A	2.9	0.003	4.5	0.0004
COL1A1	2.5	0.009	n.s.	n.s.
CXCL1 (GRO1)	2.0	0.008	n.s.	n.t.
MMP2	1.8	0.008	n.s.	n.s.
Decreased expression				
ARHE (Rho8)	0.6	0.002	3.3	0.005
PLK2 (SNK)	0.5	0.010	2.4	0.002

¹Keratinocytes from non-lesional skin of psoriasis patients (n = 3) or normal control skin (n = 4) were expanded in vitro and cultured for appr. 14 days, followed by expression profiling using the PIQUOR Skin-Patho microarray, as detailed in methods.

²Fold change (relative expression levels in keratinocytes from psoriasis patients vs. controls).

³Statistical significance, as determined by a non-paired two sided t-test.

⁴Overexpression in lesional vs. non-lesional psoriatic skin, as previously reported [16].

doi:10.1371/journal.pone.0005354.t002

antibody-mediated neutralization studies will be able to clarify this issue. Moreover, our data do not show a direct transcriptional upregulation of IFN genes or IFN receptors (data not shown) by Wnt5a activation. Down-stream of the membrane receptor, it is less clear at which point Fzd-mediated signals interact with IFN signalling relayers [35]. The present data should certainly prompt a more detailed analyses of the effect of Wnt5a on nuclear translocation and phosphorylation of factor involved in IFN signal transduction.

We here identify the APP amyloid precursor protein and the ubiquitin-like molecule nedd8 as bona fide targets of synergistic IFN α /Wnt5a induction. APP is upregulated and re-distributed in psoriasis lesions strikingly similar to Wnt5a and APP stimulates both proliferation and mobility of keratinocytes [36,37]. Furthermore, processed amyloid β induces transcription of kynureninase, which we found to be strongly upregulated in psoriasis in vivo, as well as in psoriasis-derived cells in vitro (table 2), and which can elicit an inflammatory skin reaction in vivo [38]. Therefore, the induction of APP by Wnt5a/IFN may contribute to increased keratinocyte turnover and migration in psoriasis. As for nedd8, further studies must clarify which target proteins are neddylated by concurrent Wnt5a/IFN stimulation and whether these are involved in epidermal dynamics. However, the results presented here clearly show that, in addition to established IFN target genes, another class of targets exists which are synergistically induced by Wnt5a/IFN α . Thus, screening for such targets in other cell types such as T cells, endothelial cells, or fibroblasts, would delineate the profile of innate immune activation elicited by Wnt5a.

In conclusion, we here demonstrate compartmentalized distribution of Wnt-signalling components in adult skin, up-regulation




Figure 4. Overexpression of Wnt5a in psoriatic keratinocytes in vitro. Keratinocytes from non-affected skin from psoriasis patients or healthy control skin were expanded in vitro for 14 days, fractionated in nuclear (n) or cytoplasmic/membrane fractions (c/m), and subjected to western blot, as detailed in Methods. 50 μ g of protein were loaded per lane. The bottom panel shows the Poinceau-S stain of the blot verifying even protein loading. * = non-specific band also visible in the Poinceau S stain.
doi:10.1371/journal.pone.0005354.g004

Table 3. Genes induced in keratinocytes by Wnt5a in vitro.¹

Gene ID	fold change			p-value ²
	Psoriasis	Controls	Total	
NEDD8	1.34	1.35	1.35	2×10^{-7}
IFI-78 (MX1)	1.53	1.16	1.35	0.14
IFI27	1.52	1.12	1.32	0.11
APP	1.41	1.14	1.28	0.002

¹Primary keratinocytes expanded from healthy control donors (n = 3) or psoriasis patients (n = 3) were cultivated in the presence or absence of Wnt5a mimic peptide [5] or control peptide for 48 h and gene expression subsequently analyzed as detailed in Methods. Total fold-changes shown represent the average across all six samples (wnt5a vs. control peptide).

²As determined by a two-sided paired t-test.

doi:10.1371/journal.pone.0005354.t003




Figure 5. Wnt5a induces type 1 IFN target genes and causes IFN hypersensitivity. (a) HaCat keratinocytes were transfected with full length Wnt5a cDNA or control vector (pCDNA6.1). 12 h after transfection, the expression of *nedd8*, *MX1*, *IFI-27*, and *APP* was determined by RT-PCR, as described in Methods. The expression of *Wnt5a* was also determined to verify overexpression. (b) HaCat keratinocytes transfected with full-length Wnt5a cDNA (open symbols) or control vector (closed symbols) and stimulated with the indicated concentrations of IFN α for 18 h. The expression of *IFI27* was then assessed using RT-PCR and quantified densitometrically. The data shown represent mean \pm s.d. of two independent experiments. (c) Keratinocytes from non-lesional psoriatic skin with elevated endogenous *Wnt5a* levels or from healthy control skin were expanded in vitro for 14 days and stimulated for 18 h with 20 ng/ml IFN α . Global gene expression was determined using the piquor skin-patho microarray. Plotted are the -fold change (IFN α vs. basal expression) of all genes (41 of 1133) exhibiting a ≥ 2 -fold induction by IFN α . The mean fold-changes of cells from $n = 3$ controls are plotted along the x-axis, the mean fold-change of cells from $n = 4$ cells are plotted along the y-axis. Dots represent mean values, horizontal lines represent s.d. among the controls, vertical lines represent s.d. among cells from psoriasis patients. The dashed line marks the theoretical equal magnitude of gene induction.

doi:10.1371/journal.pone.0005354.g005

of *Wnt5a* in hyperproliferative and differentiation-deficient epidermis, and identify sensitization for interferon as a novel function of *Wnt* signalling.

Supporting Information

Supporting Methods S1

Found at: doi:10.1371/journal.pone.0005354.s001 (0.06 MB DOC)

Acknowledgments

The authors wish to acknowledge the excellent technical support of Dr. Ute Ungethüm of the Charité Medical University in-house microarray facility.

Author Contributions

Conceived and designed the experiments: MR SB JF. Performed the experiments: MR AE DK SB KM SD JF. Analyzed the data: MR AE SD JF.




Figure 6. Synergistic induction of APP and nedd8 by Wnt5a and IFN α . (a) HaCat cells were transfected either with control vector or Wnt5a, followed by incubation in the absence or presence of 20 ng/ml IFN α for 18 h. Whole cell lysates were subjected to SDS-PAGE and blotted with anti-APP. The visible band migrating at appr. 70 kd is consistent with β -secretase-cleaved sAPP. Poinceau-5 stain of the membrane is shown as loading control. (b) HaCat cells were transfected and stimulated with IFN α as in (a), followed by preparation of nuclear extracts, SDS-PAGE, and western blot using anti-nedd8.

doi:10.1371/journal.pone.0005354.g006

References

- Cheng CW, Yeh JC, Fan TP, Smith SK, Charnock-Jones DS (2008) Wnt5a-mediated non-canonical Wnt signalling regulates human endothelial cell proliferation and migration. *Biochem Biophys Res Commun* 365: 285–290.
- Masckauchan TN, Agalliu D, Vorontchikhina M, Ahn A, Parmalee NL, et al. (2006) Wnt5a signaling induces proliferation and survival of endothelial cells in vitro and expression of MMP-1 and Tie-2. *Mol Biol Cell* 17: 5163–5172.
- Yu JM, Jun ES, Jung JS, Suh SY, Han JY, et al. (2007) Role of Wnt5a in the proliferation of human glioblastoma cells. *Cancer Lett.*
- Kawasaki A, Torii K, Yamashita Y, Nishizawa K, Kanekura K, et al. (2007) Wnt5a promotes adhesion of human dermal fibroblasts by triggering a phosphatidylinositol-3 kinase/Akt signal. *Cell Signal.*
- Safholm A, Leandersson K, Dejmeck J, Nielsen CK, Villoutreix BO, et al. (2006) A formylated hexapeptide ligand mimics the ability of Wnt-5a to impair migration of human breast epithelial cells. *J Biol Chem* 281: 2740–2749.
- Blumenthal A, Ehlers S, Lauber J, Buer J, Lange C, et al. (2006) The Wingless homolog WNT5A and its receptor Frizzled-5 regulate inflammatory responses of human mononuclear cells induced by microbial stimulation. *Blood* 108: 965–973.
- Pereira C, Schaer DJ, Bachli EB, Kurrer MO, Schoedon G (2008) Wnt5A/CaMKII signaling contributes to the inflammatory response of macrophages and is a target for the antiinflammatory action of activated protein C and interleukin-10. *Arterioscler Thromb Vasc Biol* 28: 504–510.
- Sen M, Lauterbach K, El-Gabalawy H, Firestein GS, Corr M, et al. (2000) Expression and function of wingless and frizzled homologs in rheumatoid arthritis. *Proc Natl Acad Sci U S A* 97: 2791–2796.
- Mikels AJ, Nusse R (2006) Purified Wnt5a protein activates or inhibits beta-catenin-TCF signaling depending on receptor context. *PLoS Biol* 4: e115.
- Murphy LL, Hughes CC (2002) Endothelial cells stimulate T cell NFAT nuclear translocation in the presence of cyclosporin A: involvement of the wnt/glycogen synthase kinase-3 beta pathway. *J Immunol* 169: 3717–3725.
- Reddy ST, Andl T, Lu MM, Morrissy EE, Millar SE (2004) Expression of Frizzled genes in developing and postnatal hair follicles. *J Invest Dermatol* 123: 275–282.
- Roarty K, Serra R (2007) Wnt5a is required for proper mammary gland development and TGF-beta-mediated inhibition of ductal growth. *Development* 134: 3929–3939.
- Downs AM, Dunnill MG (2000) Exacerbation of psoriasis by interferon-alpha therapy for hepatitis C. *Clin Exp Dermatol* 25: 351–352.
- Kowalick L (1997) Psoriasis flare caused by recombinant interferon beta injections. *J Am Acad Dermatol* 36: 501.
- Eriksen KW, Lovato P, Skov L, Krejsgaard T, Kalfot K, et al. (2005) Increased sensitivity to interferon-alpha in psoriatic T cells. *J Invest Dermatol* 125: 936–944.
- Romanowska M, al Yacoub N, Seidel H, Donandt S, Gerken H, et al. (2008) PPARdelta enhances keratinocyte proliferation in psoriasis and induces heparin-binding EGF-like growth factor. *J Invest Dermatol* 128: 110–124.
- Hida S, Ogasawara K, Sato K, Abe M, Takayanagi H, et al. (2000) CD8(+) T cell-mediated skin disease in mice lacking IRF-2, the transcriptional attenuator of interferon-alpha/beta signaling. *Immunity* 13: 643–655.
- Reischl J, Schwenke S, Beckman JM, Mrowietz U, Sturzebecher S, et al. (2007) Increased expression of Wnt5a in psoriatic plaques. *J Invest Dermatol* 127: 163–169.
- al Yacoub N, Romanowska M, Haritonova N, Foerster J (2007) Optimized production and concentration of lentiviral vectors containing large inserts. *J Gene Med* 9: 579–584.
- Haider AS, Peters SB, Kaporis H, Cardinale I, Fei J, et al. (2006) Genomic analysis defines a cancer-specific gene expression signature for human squamous cell carcinoma and distinguishes malignant hyperproliferation from benign hyperplasia. *J Invest Dermatol* 126: 869–881.
- Guo N, Hawkins C, Nathans J (2004) Frizzled6 controls hair patterning in mice. *Proc Natl Acad Sci U S A* 101: 9277–9281.
- Hung BS, Wang XQ, Cam GR, Rothnagel JA (2001) Characterization of mouse Frizzled-3 expression in hair follicle development and identification of the human homolog in keratinocytes. *J Invest Dermatol* 116: 940–946.
- Safholm A, Tuomela J, Rosenkvist J, Dejmeck J, Harkonen P, et al. (2008) The Wnt-5a-derived hexapeptide Foxy-5 inhibits breast cancer metastasis in vivo by targeting cell motility. *Clin Cancer Res* 14: 6556–6563.
- Cargill M, Schrodi SJ, Chang M, Garcia VE, Brandon R, et al. (2007) A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes. *Am J Hum Genet* 80: 273–290.
- Krueger GG, Langley RG, Leonardi C, Yehilding N, Guzzo C, et al. (2007) A human interleukin-12/23 monoclonal antibody for the treatment of psoriasis. *N Engl J Med* 356: 580–592.
- Katoh M, Katoh M (2007) STAT3-induced WNT5A signaling loop in embryonic stem cells, adult normal tissues, chronic persistent inflammation, rheumatoid arthritis and cancer (Review). *Int J Mol Med* 19: 273–278.

27. Sano S, Chan KS, Carbajal S, Clifford J, Peavey M, et al. (2005) Stat3 links activated keratinocytes and immunocytes required for development of psoriasis in a novel transgenic mouse model. *Nat Med* 11: 43–49.
28. Fathke C, Wilson L, Shah K, Kim B, Hocking A, et al. (2006) Wnt signaling induces epithelial differentiation during cutaneous wound healing. *BMC Cell Biol* 7: 4.
29. Sen M, Chamorro M, Reifert J, Corr M, Carson DA (2001) Blockade of Wnt-5A/frizzled 5 signaling inhibits rheumatoid synoviocyte activation. *Arthritis Rheum* 44: 772–781.
30. Boyman O, Hefti HP, Conrad C, Nickoloff BJ, Suter M, et al. (2004) Spontaneous development of psoriasis in a new animal model shows an essential role for resident T cells and tumor necrosis factor-alpha. *J Exp Med* 199: 731–736.
31. Wolf R, Ruocco V (1999) Triggered psoriasis. *Adv Exp Med Biol* 455: 221–225.
32. Hampton PJ, Ross OK, Reynolds NJ (2007) Increased nuclear beta-catenin in suprabasal involved psoriatic epidermis. *Br J Dermatol* 157: 1168–1177.
33. Takada I, Mihara M, Suzawa M, Ohtake F, Kobayashi S, et al. (2007) A histone lysine methyltransferase activated by non-canonical Wnt signalling suppresses PPAR-gamma transactivation. *Nat Cell Biol* 9: 1273–1285.
34. Wang LH, Yang XY, Zhang X, Huang J, Hou J, et al. (2004) Transcriptional inactivation of STAT3 by PPARgamma suppresses IL-6-responsive multiple myeloma cells. *Immunity* 20: 205–218.
35. Platanias LC (2005) Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nat Rev Immunol* 5: 375–386.
36. Siemes C, Quast T, Klein E, Bieber T, Hooper NM, et al. (2004) Normalized proliferation of normal and psoriatic keratinocytes by suppression of sAPPalpha-release. *J Invest Dermatol* 123: 556–563.
37. Herzog V, Kirfel G, Siemes C, Schmitz A (2004) Biological roles of APP in the epidermis. *Eur J Cell Biol* 83: 613–624.
38. Ito M, Ogawa K, Takeuchi K, Nakada A, Heishi M, et al. (2004) Gene expression of enzymes for tryptophan degradation pathway is upregulated in the skin lesions of patients with atopic dermatitis or psoriasis. *J Dermatol Sci* 36: 157–164.