



OPEN ACCESS

CASE REPORT

Cold agglutinins in a patient undergoing normothermic cardiac operation with warm cardioplegia

Tatsuhiko Ogawa

Department of Anaesthesiology,
Kochi Health Sciences Center,
Kochi, Japan

Correspondence to

Dr Tatsuhiko Ogawa,
tatsuhiko777@hotmail.com

Accepted 29 September 2017

SUMMARY

Cold agglutinins are autoantibodies that agglutinate red blood cells at low temperatures, leading to haemagglutination and haemolysis. They are generally of no clinical significance. However, when people with cold agglutinins undergo cardiac operation with hypothermia and cold cardioplegia, they can experience complications. Thus, different perioperative management is required for such patients. We describe a 74-year-old man with cold agglutinins incidentally detected on the preoperative screening test. He had never experienced any complications or developed a haematological disease. Since cold agglutinins were incidentally detected on the preoperative test, a special strategy was used to manage the temperature of cardiopulmonary bypass (CPB) and cardioplegia. He successfully underwent

normothermic cardiac operation with warm cardioplegia. A continuous retrograde hyperkalaemic infusion and intermittent antegrade infusion of warm cardioplegia with normothermic CPB is one of the best methods to avoid hypothermia and excessive activity and metabolism of the heart, and to provide a suitable operative field.

BACKGROUND

Cold agglutinins are autoantibodies that agglutinate red blood cells at low temperatures, leading to haemagglutination and haemolysis.¹⁻¹¹ They are generally of no clinical significance.^{1-3 5 6 8 10 11} However, when people who have cold agglutinins in their blood undergo cardiac operation under




Figure 1 Recommendations for preoperative testing and haematology consultation in patients with benign cold agglutinins and cold haemagglutinin disease. Reproduced from Barbara *et al.*¹ CA, cold agglutinins; CHAD, cold haemagglutinin disease; CPB, cardiopulmonary bypass.



CrossMark

To cite: Ogawa T. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-221888




Figure 2 Guidelines for managing patients with cold antibodies who undergo cardiovascular operation. During the preoperative visit, the patient must be evaluated for a history suggestive of pathogenic cold antibodies. Reproduced from Sapatnekar and Figueroa.² AHG XM, antihuman globulin crossmatch; RBC, red blood cell.

hypothermia and cold cardioplegia, they can experience complications.^{1 2 4 6 7 10 11} The reported incidence of cold agglutinins among screened cardiac surgical patients is low but not too rare (approximately 0.8%–4%^{4 6 7}), and different perioperative management is required for such patients. We describe a patient with incidentally detected cold agglutinins who successfully underwent normothermic cardiac operation with warm cardioplegia.

CASE PRESENTATION

A 74-year-old man was scheduled for cardiac operation because of severe mitral regurgitation and coronary stenosis. During preoperative screening, all initial preoperative routine test results were normal, except for the presence of the non-specific cold antibodies confirmed by the saline method. The direct antiglobulin test was negative for anti-immunoglobulin G, but we did not perform further testing of anti-C3. These findings had not been assessed until the preoperative evaluation by anaesthesiologists. On the basis of these indications, we postponed the operation and performed further tests, which resulted in the detection of cold agglutinins by the haemagglutination assay. The titre of the cold agglutinin was 1:512 at 4°C, and the thermal amplitude, defined as the highest temperature at which haemagglutination occurs, was 32°C. The patient had no history of anaemia, prior blood transfusion or any disease other than prostate cancer, for which he had undergone operation without complications.

Haematologists diagnosed him as having idiopathic cold agglutinin disease because of a lack of prior infection or haematological malignancy. There is no consensus on the clinical significance of the cold antibody titre and thermal amplitude that warrant appropriate precautions.¹ However,

he had no clinical symptoms, and his agglutinin titre was not extremely high compared with those of previously reported cases^{2 3}; there was no indication for preoperative plasmapheresis or the administration of immunoglobulin,⁴ corticosteroids and alkylating agents³ and rituximab therapy.⁵ However, thermal amplitude, which is more important than the titre in predicting the likelihood of complications,^{1 6} was 32°C; therefore, management using hypothermic cardiopulmonary bypass (CPB) with cold cardioplegia would have caused the temperatures of his core, peripheral and coronary arteries to reach unsafe levels.

Therefore, we performed normothermic cardiac operation with warm cardioplegia. We usually use a single infusion of cold blood cardioplegia in an antegrade and retrograde delivery system every 30 min, which was modified from the one-dose method.¹²

To avoid inadequate protection of the heart due to warm conditions and excessive cardiac activity and oxygen consumption, we added a continuous retrograde hyperkalaemic infusion to the intermittent antegrade infusion of cardioplegia. The patient was protected from exposure to hypothermia using warming blankets and a fluid warming system.

Retrograde cardioplegia (36°C) was performed continuously, except when antegrade cardioplegia was administered. We used blood cardioplegia made of the patient’s blood and potassium chloride (KCl, 8–20 mEq/L). We maintained cardiac arrest during almost the entire CPB period while keeping the patient’s temperature >35°C.

The patient had uneventful intraoperative and postoperative courses, with no clinically significant haemagglutination or haemolysis.

Written informed consent was obtained from the patient for publication of this case report.

TREATMENT

We changed the routine CPB plan (hypothermic CPB with cold cardioplegia) to normothermic CPB with warm cardioplegia. We administered a continuous retrograde hyperkalaemic infusion and intermittent antegrade infusion of warm cardioplegia with normothermic CPB, and we tried to keep the patient warm throughout the perioperative period.

OUTCOME AND FOLLOW-UP

The operation was successfully completed without complications, and the patient has been well without any symptoms 1 year postoperatively.

DISCUSSION

The optimal guidelines of patients with cold agglutinins undergoing CPB operation remain controversial because of the limited number of case reports. Some physicians believe that patients with low titres and low thermal amplitude antibodies may undergo CPB operation without any change in the routine management plan.^{1,2}

Recently, a new algorithm¹ was proposed to guide preoperative testing in patients with cold agglutinins (figure 1). The authors claimed that patients with non-symptomatic cold agglutinins can safely undergo normothermic cardiac operation with warm cardioplegia at 37°C, and they may not require additional testing. The determination of titres, thermal amplitude and haematology consultation are recommended in patients when systemic hypothermia or cold cardioplegia is required in benign cold agglutinins.

Furthermore, other authors developed guidelines² (figure 2) for pretransfusion testing and perioperative management of patients with cold agglutinins. They claimed that if cold agglutinins are incidentally detected and are not symptomatic, standard pretransfusion testing should be performed to determine whether antigen-negative red blood cells should be used. Considering the rarity of haemagglutination among patients with cold agglutinins, they concluded that special preoperative testing protocols are neither necessary nor justified in the absence of pathogenic cold agglutinins or symptoms. Physicians now tend to exclude unnecessary examinations related to cold agglutinins preoperatively given their cost–benefit relationships.

Regarding temperature management, protection of the myocardium and other organs may be jeopardised under normothermic surgical conditions and warm cardioplegia. It is difficult to accurately assess adverse effects of temperature management during CPB on myocardial oxygen consumption and energy metabolism at the cellular levels. Physicians have tried to determine the best CPB temperature management, and contraindicated outcomes between normothermic and hypothermic CPB operations have been reported.^{13–15} According to the Clinical Practice Guidelines for Temperature Management during CPB,¹⁶ surgical teams should limit the CPB temperature to <37°C to avoid cerebral hyperthermia (class I, level C), and temperature gradients between the arterial outlet and venous inflow should not exceed <10°C to avoid generation of gaseous emboli (class I, level C). No unified agreements have been reached on this difficult topic in the normal population. The presence of cold agglutinins presents more complicated

Patient's perspective

I considered that by avoiding cold conditions, I would be well without complications, such as haemagglutinations and a haemolytic crisis.

Learning points

- ▶ Cold agglutinins of low antibody titres can cause haemagglutination and a haemolytic crisis even in asymptomatic patients with an increased risk in cardiopulmonary bypass (CPB) operation.
- ▶ Complete preoperative screening for the cold antibody and direct communication with haematology specialists are required before additional tests are necessary and to avoid unexpected complications, especially if the patient is scheduled to undergo cardiac operation with CPB and could be exposed to cold conditions.
- ▶ Patients with a history or symptoms/signs suggesting cold agglutinin disease or the presence of any abnormal cold antibody should be referred to a haematology specialist, who would then determine the appropriate test, especially if the patient is scheduled to undergo cardiac operation with CPB and can be exposed to cold conditions.
- ▶ Continuous retrograde infusion of hyperkalaemic warm cardioplegia with normothermic CPB in patients with cold agglutinins is one of the best methods to avoid hypothermia and excessive activity and metabolism of the heart, and to provide a suitable operative field.
- ▶ This aforementioned method can be used by simply adding it to the intermittent routine administration of cardioplegia, and it is useful for physicians without much experience with this disease.

problems in optimal temperature management during CPB. According to the experts' opinion,¹ normothermic CPB operation with warm cardioplegia can be safely performed without any complications in patients with cold agglutinins. However, this disease has a special pathophysiology that is often ignored. Haemagglutination observed in patients with pathogenic cold agglutinins is not reliably correlated with the antibody titre.^{5,8} This phenomenon may occur in patients with titres as low as 1:32. Haemagglutination and the subsequent haemolysis caused by activated cold agglutinins have been observed in patients with low titres.^{9,10} This may be more hazardous during CPB under low-temperature conditions and, indeed, some lethal, irremediable cases have been reported.^{10,11}

On the basis of past reports, we presume that patients with cold agglutinins may exhibit haemagglutination and a haemolytic crisis, even with low antibody titres and examination results that are not particularly abnormal. Complete preoperative screening for the cold antibody and direct communication with haematology specialists are required before additional tests are necessary and to avoid unexpected complications, especially if the patient is scheduled to undergo cardiac operation with CPB and could be exposed to cold conditions. Patients with a history or symptoms/signs suggesting cold agglutinin disease or the presence of any abnormal cold antibody should be referred to a haematology specialist, who would then determine the appropriate

test, especially if the patient is scheduled to undergo cardiac operation with CPB and can be exposed to cold conditions.

If physicians do not know the patient's background characteristics and do not have much experience with this disease, as in the present case, they should take the most cautious measures, for example, preoperative testing of antibody titres and thermal amplitudes, after consulting with other medical professionals. A continuous retrograde hyperkalaemic infusion and intermittent antegrade infusion of warm cardioplegia with normothermic CPB is one of the best measures for patients with unknown background characteristics and for physicians without much experience in treating this condition.

We consider that this technique can be applied as a general replacement for cold cardioplegia without risk of agglutinations and other complications, if the systemic potassium concentration and myocardial energy metabolism are controlled. Unfortunately, we could not perform comparative experiments or draw any definitive conclusions; however, we believe further case reports and discussions can lead to better management of this disease.

Acknowledgements I would like to thank my colleagues, operating room staffs, and clinical engineers for their generous support, which enabled me to write this paper.

Contributors TO is the sole author of this paper and wrote the whole manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Barbara DW, Mauermann WJ, Neal JR, *et al*. Cold agglutinins in patients undergoing cardiac surgery requiring cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 2013;146:668–80.
- 2 Sapatnekar S, Figueroa PI. Cold antibodies in cardiovascular surgery: is preoperative screening necessary? *Am J Clin Pathol* 2016;145:789–95.
- 3 Berentsen S, Ulvestad E, Langholm R, *et al*. Primary chronic cold agglutinin disease: a population based clinical study of 86 patients. *Haematologica* 2006;91:460–6.
- 4 Kanemitsu S, Onoda K, Yamamoto K, *et al*. Simple preoperative management for cold agglutinins before cardiac surgery. *J Thorac Cardiovasc Surg* 2010;140:e73–4.
- 5 Swiecicki PL, Hegerova LT, Gertz MA, *et al*. Cold agglutinin disease. *Blood* 2013;122:1114–21.
- 6 Agarwal SK, Ghosh PK, Gupta D. Cardiac surgery and cold-reactive proteins. *Ann Thorac Surg* 1995;60:1143–50.
- 7 Osada H, Nakajima H, Shimizu A, *et al*. Type A aortic dissection with cold agglutinin disease. *Ann Thorac Surg* 2011;92:722–3.
- 8 Japan Intractable Diseases Information Center. Autoimmune hemolytic anemia (AIHA) (designated intractable disease 61). Cited 25 Jun 2017. <http://www.nanbyou.or.jp/entry/269>
- 9 Rousey SR, Smith RE. A fatal case of low titer anti-PR cold agglutinin disease. *Am J Hematol* 1990;35:286–7.
- 10 Madershahian N, Franke U, Jütte H, *et al*. Cold agglutinins in on-pump cardiac surgery: a rare but potentially lethal problem. *Int J Perfusionists* 2004;1:1–4.
- 11 Patel PA, Ghadimi K, Coetzee E, *et al*. Incidental cold agglutinins in cardiac surgery: intraoperative surprises and team-based problem-solving strategies during cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 2017;31:1109–18.
- 12 Buckberg GD, Athanasuleas CL. Cardioplegia: solutions or strategies? *Eur J Cardiothorac Surg* 2016;50:787–91.
- 13 Ho KM, Tan JA. Benefits and risks of maintaining normothermia during cardiopulmonary bypass in adult cardiac surgery: a systematic review. *Cardiovasc Ther* 2011;29:260–79.
- 14 Cakir H, Gur O, Ege T, *et al*. Comparison of the efficacy of the cardiac hypothermia and normothermia to myocardial damage in coronary artery bypass graft surgery with systemic normothermic cardiopulmonary bypass. *J Cardiovasc Surg* 2013;54:397–401.
- 15 Lomivorotov VV, Shmirev VA, Efremov SM, *et al*. Hypothermic versus normothermic cardiopulmonary bypass in patients with valvular heart disease. *J Cardiothorac Vasc Anesth* 2014;28:295–300.
- 16 Engelman R, Baker RA, Likosky DS, *et al*. The society of thoracic surgeons, the society of cardiovascular anesthesiologists, and the american society of extraCorporeal technology: clinical practice guidelines for cardiopulmonary bypass-temperature management during cardiopulmonary bypass. *Ann Thorac Surg* 2015;100:748–57.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow