



# Efficient Multiplex Genome Editing in *Streptomyces* via Engineered CRISPR-Cas12a Systems

Jun Zhang<sup>1†</sup>, Dan Zhang<sup>1,2†</sup>, Jie Zhu<sup>1</sup>, Huayi Liu<sup>1</sup>, Shufang Liang<sup>1,2\*</sup> and Yunzi Luo<sup>1,3\*</sup>

<sup>1</sup> Department of Gastroenterology, State Key Laboratory of Biotherapy, West China Hospital, Sichuan University and Collaborative Innovation Center of Biotherapy, Chengdu, China, <sup>2</sup> State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University and Collaborative Innovation Center of Biotherapy, Chengdu, China, <sup>3</sup> Key Laboratory of Systems Bioengineering (Ministry of Education), Frontiers Science Center of Synthetic Biology, School of Chemical Engineering and Technology, Tianjin University, Tianjin, China

## OPEN ACCESS

### Edited by:

Yuan Lu,  
Tsinghua University, China

### Reviewed by:

Yiping Qi,  
University of Maryland, United States  
Dae-Hee Lee,  
Korea Research Institute  
of Bioscience & Biotechnology  
(KRIBB), South Korea  
Steven Lin,  
Institute of Biological Chemistry,  
Academia Sinica, Taiwan

### \*Correspondence:

Shufang Liang  
zizi2006@scu.edu.cn  
Yunzi Luo  
yunzi.luo@tju.edu.cn;  
luoyunzi827@aliyun.com

† These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Synthetic Biology,  
a section of the journal  
Frontiers in Bioengineering and  
Biotechnology

Received: 07 January 2020

Accepted: 09 June 2020

Published: 30 June 2020

### Citation:

Zhang J, Zhang D, Zhu J, Liu H,  
Liang S and Luo Y (2020) Efficient  
Multiplex Genome Editing  
in *Streptomyces* via Engineered  
CRISPR-Cas12a Systems.  
Front. Bioeng. Biotechnol. 8:726.  
doi: 10.3389/fbioe.2020.00726

*Streptomyces* strains produce a great number of valuable natural products. With the development of genome sequencing, a vast number of biosynthetic gene clusters with high potential for use in the discovery of valuable clinical drugs have been revealed. Therefore, emerging needs for tools to manipulate these biosynthetic pathways are presented. Although the clustered regularly interspaced short palindromic repeats/CRISPR-associated protein 9 (CRISPR/Cas 9) system has exhibited great capabilities for gene editing in multiple *Streptomyces* strains, it has failed to work in some newly discovered strains and some important industrial strains. Additionally, the protospacer adjacent motif (PAM) recognition scope of this system sometimes limits its applications for generating precise site mutations and insertions. Here, we developed three efficient CRISPR-*F*nCas12a systems for multiplex genome editing in several *Streptomyces* strains. Each system exhibited advantages for different applications. The CRISPR-*F*nCas12a1 system was efficiently applied in the industrial strain *Streptomyces hygrosopicus*, in which *Sp*Cas9 does not work well. The CRISPR-*F*nCas12a2 system was used to delete large fragments ranging from 21.4 to 128 kb. Additionally, the CRISPR-*F*nCas12a3 system employing the engineered *F*nCas12a mutant EP16, which recognizes a broad spectrum of PAM sequences, was used to precisely perform site mutations and insertions. The CRISPR-*F*nCas12a3 system addressed the limitation of TTN PAM recognition in *Streptomyces* strains with high GC contents. In summary, all the CRISPR-*F*nCas12a systems developed in this study are powerful tools for precise and multiplex genome editing in *Streptomyces* strains.

**Keywords:** CRISPR, *F*nCas12a, *Streptomyces*, genome editing, PAM recognition

## INTRODUCTION

*Streptomyces*, the largest genus of actinobacteria, has been well studied, as it contains the most prolific producers of a vast array of bioactive natural products, including antibiotics, antifungals, and anticancer agents (Baltz, 2008; Zhu et al., 2011; Cho et al., 2017; Frattaruolo et al., 2017). Over the past decades, large-scale genome sequencing efforts have revealed that great potential still remains for the discovery of new natural products produced by members of this genus. Thus,

increasing numbers of genetic engineering tools have been developed to explore these products (Luo et al., 2015b, 2016; Zhao et al., 2019). In particular, with the rapid development of the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein (Cas) system (Jiang et al., 2013; Mali et al., 2013; Yan et al., 2018), effective genome editing has become increasingly easy and convenient, paving the way for us to assemble or activate uncharacterized gene clusters. However, CRISPR/Cas9 has been reported to not work in some *Streptomyces* strains, such as *Streptomyces* sp. KY 40-1 (Salem et al., 2017), *Streptomyces* sp. NRRL S-244 (Yeo et al., 2019) and *Streptomyces hygroscopicus* SIPI-KF (Li et al., 2018), because of its toxicity to the hosts.

Cas12a, a Class 2 CRISPR effector, is a single RNA-guided endonuclease (Zetsche et al., 2015). Due to their simplicity, AsCas12a from *Acidaminococcus* sp. BV3L6 and LbCas12a from *Lachnospiraceae bacterium* ND2006 have generally been applied to mammalian cells (Toth et al., 2016) and plants (Tang et al., 2017), while FnCas12a from *Francisella novicida* U112 has successfully been applied to yeast (Swiat et al., 2017), *Corynebacterium glutamicum* (Jiang et al., 2017) and other bacteria (Ungerer and Pakrasi, 2016). Wild-type (WT) FnCas12a has been used in *Streptomyces*, particularly in some strains in which the CRISPR/Cas9 system does not function properly, such as *S. hygroscopicus* SIPI-KF (Li et al., 2018) and *Streptomyces* sp. NRRL S-244 (Yeo et al., 2019).

However, the TTN protospacer adjacent motif (PAM) requirement restricts the application of FnCas12a in *Streptomyces* strains, which are GC rich. In particular, target site selection is limited for site mutations or insertions in the genome. Thus, there remains a demand for highly efficient and versatile genome editing tools for *Streptomyces*. Recently, we constructed an engineered FnCas12a variant, EP16, which shows broad PAM recognition abilities *in vitro*, including YN (Y = C or T), TAC and CAA (Wang et al., 2019). Therefore, EP16 provides additional opportunities to precisely edit the high-GC-contents genomes of *Streptomyces* strains.

In this study, three optimized CRISPR-FnCas12a systems, CRISPR-FnCas12a1, CRISPR-FnCas12a2 and CRISPR-FnCas12a3 (Figure 1 and Supplementary Table 2), were developed to edit the genomes of different *Streptomyces* strains based on homology-directed repair (HDR), as non-homologous end joining (NHEJ) does not occur in most streptomycetes. (i) The CRISPR-FnCas12a1 system exhibited a higher transformation efficiency than the CRISPR-FnCas12a2 system, as it contains a *traJ* gene encoding an activator of the transfer operon (Will and Frost, 2006). (ii) A higher editing efficiency was observed for the CRISPR-FnCas12a2 system than the CRISPR-FnCas12a1 system when FnCas12a was driven by the constitutive promoters *kasOp\** and *rpsLp(XC)*. (iii) The FnCas12a3 system containing the FnCas12a mutant EP16 (N607R/K613V/N617R/K180S/K660R/D616N) with expanded PAM recognition ability was used to generate site mutation, insertion and subsequent biosynthetic gene cluster activation (Figure 1). Overall, our findings describe powerful tools for precise genome editing and subsequently for discovering and activating valuable natural products in *Streptomyces* strains.

## MATERIALS AND METHODS


### Strains, Cultivation and Reagents

*Streptomyces coelicolor* A3(2), *Streptomyces griseus*, *Streptomyces lividans*, *E. coli* DH5 $\alpha$  and *E. coli* ET12567/pUZ8002 were gifts from Professor Huimin Zhao at the University of Illinois at Urbana-Champaign. *S. hygroscopicus* NRRL5491 and *Streptomyces roseosporus* NRRL 11379 were purchased from the American Type Culture Collection (ATCC, Manassas, VA, United States). All *Streptomyces* strains were grown in MYG liquid medium (10 g/L malt extract broth, 4 g/L yeast extract, and 4 g/L glucose) or on M-ISP4 agar medium (1 g/L yeast extract, 2 g/L tryptone, 5 g/L soluble starch, 5 g/L D-mannitol, 5 g/L soya flour, 1 g/L NaCl, 2 g/L (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1 g/L K<sub>2</sub>HPO<sub>3</sub>, 2 g/L CaCO<sub>3</sub>, 20 g/L agar, 1 g/L FeSO<sub>4</sub>, 1 g/L MnCl<sub>2</sub>, and 1 g/L ZnSO<sub>4</sub>, pH 7.0) at 30°C. Next, 2 × YT broth (1% yeast extract, 1% tryptone, and 0.5% NaCl, pH 7.0) was used for strain washing and spore germination before conjugation. M-ISP4 agar medium supplemented with 25 mM MgCl<sub>2</sub> or 20 μg/mL apramycin and 40 μg/mL nalidixic acid was used for conjugation from *E. coli* ET12567/pUZ8002 to *Streptomyces* (Kieser et al., 2000) or for plasmid selection in *Streptomyces*. R2YE medium supplemented with 20 μg/mL apramycin and 40 μg/mL nalidixic acid was used to screen potentially edited *Streptomyces*. *E. coli* DH5 $\alpha$  was grown in Luria-Bertani (LB) broth with 50 μg/mL apramycin and used for plasmid cloning and maintenance. *E. coli* ET12567/pUZ8002 was grown in LB broth supplemented with 25 μg/mL apramycin, 12.5 μg/mL chloramphenicol and 25 μg/mL kanamycin.

D-Mannitol was obtained from Sigma-Aldrich (St. Louis, MO, United States). All media components of LB broth were purchased from Oxoid (Basingstoke, Hampshire, United Kingdom), and other reagents added to media were purchased from Sangon Biotech Co., Ltd. (Shanghai, China). PCR was performed with Q5 DNA polymerase (New England Biolabs, Ipswich, MA, United States) or T5 mix (Tsingke, Beijing, China). All PCR products were purified using a Wizard Genomic DNA Purification Kit (Promega, Madison, WI, United States). The restriction endonucleases *Xba*I, *Bbs*I, and *CIP* were purchased from New England Biolabs. A one-step cloning kit for two or more fragment assemblies was purchased from Vazyme Biotech Co., Ltd. (Nanjing, Jiangsu, China).

### Plasmid Construction

Three CRISPR-FnCas12a systems (CRISPR-FnCas12a1, CRISPR-FnCas12a2 and CRISPR-FnCas12a3) were constructed in this study. The CRISPR-FnCas12a1 system was constructed based on pWHU2653 (Zeng et al., 2015). The CRISPR-FnCas12a2 system and the CRISPR-FnCas12a3 system were constructed based on pCRISPomyces-2 (Cobb et al., 2015). The CRISPR-FnCas12a1 system consisted of a counterselection marker *codA(sm)*. The CRISPR-FnCas12a systems were constructed in several steps. The *kasOp\** promoter (Wang et al., 2013), *rpsLp(XC)* promoter (Shao et al., 2013), *ermEp\** promoter (Luo et al., 2015a), and *Potr\** system (Wang et al., 2016) were amplified by PCR from template plasmids. *Streptomyces* codon-optimized FnCas12a was



chemically synthesized by Genewiz (Suzhou, Jiangsu, China). Other elements, including the *lacZ* cassette, unique *BbsI* and *XbaI* restriction sites, temperature-sensitive pSG5 *rep* region, *aac(3)IV* coding sequence, *colE1* origin, and transfer *oriT* region, were amplified from the pWHU2653 or pCRISPOmyces-2 plasmid. The yeast helper fragment was amplified from pRS416, which was a gift from Professor Huimin Zhao. All fragments were assembled into a plasmid with a one-step assembly kit or DNA assembler (Shao et al., 2009). A specific 23-nt spacer with a TTN PAM sequence located at the 5' end of the coding strand was chosen. The PAM sequence together with the nearby 12-nt spacer sequence was analyzed using BLAST to confirm its specificity. The 19-nt or 36-nt direct repeat (DR) sequences and the 23-nt spacer targeting genes and gene clusters were synthesized and inserted into pYL-*FnCas12a* plasmids through Golden Gate assembly. Next, two 1- or 2-kb homologous arms corresponding to the upstream and downstream regions of the target genes or gene clusters were amplified from purified genomic DNA and fused into the *XbaI* site of the desired plasmid by one-step assembly. The presence of the correct plasmids was confirmed

by sequencing (Tsingke, Beijing, China). All primers used in this study are listed in **Supplementary Table 1**.

## Transformation

DNA was transformed into *E. coli* DH5 $\alpha$  or ET12567/pUZ8002 using the heat shock method. Yeast transformation was performed by electroporation following a protocol described in a previous study (Shao et al., 2009). The conjugation of plasmids from *E. coli* ET12567/pUZ8002 to *Streptomyces* strains was performed following a previously described protocol (Kieser et al., 2000).

## Screening of Potentially Edited *Streptomyces* Strains

Seven days after conjugation, seven single exconjugants were randomly picked, restreaked on R2YE or M-ISP4 agar plates supplemented with 20  $\mu$ g/mL apramycin and 40  $\mu$ g/mL nalidixic acid, and grown at 30°C for 7 days. Then, mycelia were collected from the plates for genomic DNA isolation using a bacterial

genomic DNA extraction kit (Tiangen, Beijing, China). Deletions were identified using PCR amplification of the genomic DNA or spores with primers located upstream and downstream of the target genes or primers annealing within or outside of the target gene clusters. Then, sequencing of the PCR products was performed to confirm the sequences (Tsingke, Beijing, China). For CRISPR-*FnCas12a1* system clearance, M-ISP4 plates containing 800 µg/mL 5FC were used to cultivate the strains in the dark at 28°C for 3 or 4 days. For the CRISPR-*FnCas12a2* system, plasmid clearance was carried out through high-temperature cultivation at 37°C for 2–3 days after normal cultivation at 30°C for 1 day.

### Transcription Analysis of *FnCas12a* by Real-Time PCR (RT-qPCR)

Spores from every *Streptomyces* strain were separately inoculated into MYG medium for mycelium growth. After cultivation for 72 h, total RNA was extracted using TRIzol (Thermo Fisher, Waltham, MA, United States). Reverse transcription was completed using a first-strand cDNA synthesis kit (Bio-Rad, Carlsbad, CA, United States). SYBR Green PCR Master Mix (Bio-Rad) was used for RT-qPCR. Primers were designed with an online tool.<sup>1</sup> The reaction mixtures for RT-qPCR included 1 µL of cDNA templates, 1 µL of primers, 3 µL of ddH<sub>2</sub>O and 5 µL of SYBR Green PCR Master Mix and were assayed using the following program: 2 min at 50°C and 3 min at 95°C for one cycle; 15 s at 95°C, 30 s at 62°C and 30 s at 72°C for 30 cycles; and 10 min at 72°C for a final cycle. The endogenous *hrdB* gene was used as an internal control. The transcription levels of the other genes were normalized to the control.

### Fermentation and HPLC Analysis

After plasmid clearance, the culture of vegetative mycelium was spread over M-ISP4 plates and cultivated at 30°C for 4–7 days. The spores were then scraped and counted by spreading appropriate dilutions on plates and counting single colonies that had grown for 3 days. For daptomycin fermentation,  $1 \times 10^9$  *Streptomyces* spores were inoculated into 15 mL of MYG medium and grown at 30°C for 48 h as the seed culture. Then, the seed culture was inoculated in 500 mL of fermentation medium (40 g/L dextrin, 5 g/L casein, 80 g/L glucose, and 5 g/L MgSO<sub>4</sub>, pH 7.5) at 30°C for 9 days. After 48 h of fermentation, decanoate (final concentration 0.05%, w/v) was added every 12 h until the end of the fermentation period. After fermentation, the broth was centrifuged at 10,000 rpm for 30 min. The supernatant was then mixed with an equal volume of ethyl acetate to extract the metabolites. Next, the organic phase was dried under a vacuum. Finally, the metabolites were redissolved in 4 mL of methanol and filtered through a 0.22-µm membrane before the HPLC analysis.

The metabolites were analyzed on an HPLC system (Agilent Technologies Inc., Carpinteria, CA, United States) with an Agilent C18 reverse-phase column (internal diameter 4.6 mm × 250 mm, 5-µm particle size, Agilent Technologies, Inc.) at room temperature. The flow rate was 1 mL/min, and the

absorbance of the eluate was monitored at 223 nm. The mobile phase, which was buffered with 0.01% formic acid, was initially maintained at a 75:25 water/acetonitrile for 5 min, followed by a linear gradient of 100% acetonitrile for 20 min.

### Statistical Analysis

The transformation frequency was calculated based on the ratio of the number of exconjugants to the number of spores used for conjugation. The experiments were performed in triplicate. The editing efficiency was calculated with the equation for editing efficiency (EF) = number of edited colonies evaluated by PCR/total number of evaluated colonies × 100%. All the colonies that showed double bands of WT and mutant products were calculated as wild type, and they were not included in the edited colonies. The editing efficiencies were calculated based on three replicates. Significant differences between the two groups were analyzed using *t* test. *p* < 0.05 was considered to indicate statistical significance.


## RESULTS

### Design and Construction of Three CRISPR-*FnCas12a* Systems

In previous studies, the pCRISPomyces-2 system (Cobb et al., 2015) and the pWHU2653 system (Zeng et al., 2015) have exhibited high editing efficiencies in model *Streptomyces* strains. To develop efficient and versatile genome editing tools in multiple *Streptomyces* strains, two *FnCas12a* systems were constructed: one based on the pWHU2653 system (Zeng et al., 2015) (CRISPR-*FnCas12a1*) and the other based on the pCRISPomyces-2 system (Cobb et al., 2015) (CRISPR-*FnCas12a2*). The *FnCas12a1* system carries the selection markers *codA(sm)* (Dubeau et al., 2009) and *aac(3)IV*, the *rep(pIJ101)* replicon, and the B1006 terminator that terminates the transcription of the crRNA cassette, while the *FnCas12a2* system carries only the selection marker *aac(3)IV*, the *pSG5* origin of replication, and the *fd* terminator that terminates the transcription of the crRNA cassette (**Supplementary Table 2**). Three constitutive promoters, *kasOp\** (Wang et al., 2013), *rpsLp(XC)* (Shao et al., 2013), and *ermEp\** (Luo et al., 2015a), and an inducible *Potr\** system were selected to control the expression of *FnCas12a* in each system. Overall, 8 plasmids were constructed: pYL-*kasOp\**-*FnCas12a1*, pYL-*rpsLp(XC)*-*FnCas12a1*, pYL-*ermEp\**-*FnCas12a1*, pYL-*Potr\**-*FnCas12a1*, pYL-*kasOp\**-*FnCas12a2*, pYL-*rpsLp(XC)*-*FnCas12a2*, pYL-*ermEp\**-*FnCas12a2*, and pYL-*Potr\**-*FnCas12a2* (**Figures 2A,D**, **Supplementary Figure 1**, and **Table 3**). The CRISPR-*FnCas12a3* system was constructed based on the *FnCas12a2* system by replacing the WT *FnCas12a* protein with the *FnCas12a* variant EP16.

All the above plasmids were transformed into different *Streptomyces* strains. Over 10,000 exconjugants were generated by the transformation of *S. coelicolor* with pYL-*Potr\**-*FnCas12a1* with or without addition of oxytetracycline (OTC), a value that was significantly greater than the number of exconjugants generated by plasmids pYL-*kasOp\**-*FnCas12a1*,

<sup>1</sup><https://www.idtdna.com/scitools/Applications/RealTimePCR/>



*pYL-rpsLp(XC)-Fncas12a1*, and *pYL-ermEp\*-Fncas12a1* (**Figure 2B**). Meanwhile, 0.6  $\mu\text{M}$  OTC in M-ISP4 medium induced the greatest number of exconjugants ( $12453 \pm 4582$ ). Higher OTC concentrations resulted in fewer exconjugants. Similarly, introducing *pYL-Potr\*-Fncas12a1* into *S. griseus* generated  $308 \pm 13$  exconjugants. Other *Fncas12a1* plasmids were also transformed into *S. griseus* and generated  $95 \pm 39$  (*pYL-rpsLp(XC)-Fncas12a1*),  $62 \pm 4$  (*pYL-kasOp\*-Fncas12a1*), and  $143 \pm 23$  (*pYL-ermEp\*-Fncas12a1*) exconjugants, respectively (**Figure 2C**).

A small number of *S. coelicolor* exconjugants ( $178 \pm 27$ ) were generated after introducing *pYL-kasOp\*-Fncas12a2*, while more exconjugants ( $303 \pm 83$  and  $391 \pm 21$ , respectively) were generated after introducing *pYL-rpsLp(XC)-Fncas12a2*, *pYL-ermEp\*-Fncas12a2*, and *pYL-Potr\*-Fncas12a2*. The transformation of the plasmid *pYL-Potr\*-Fncas12a2* in the absence of OTC resulted in more exconjugants ( $546 \pm 66$ ) than the transformation of other *Fncas12a2* system plasmids with constitutive promoters. Moreover, in the presence of OTC at a final concentration of 0.6  $\mu\text{M}$ , the transformation

of *pYL-Potr\*-Fncas12a2* generated the greatest number of exconjugants ( $892 \pm 166$ ) (**Figure 2E**). Similarly, the introduction of *pYL-Potr\*-Fncas12a2* into *S. griseus* generated the most exconjugants in the absence of OTC (**Figure 2F**). A comparison of editing efficiencies between two systems showed that all *Fncas12a1* system plasmids exhibited significantly higher transformation frequencies than *Fncas12a2* system plasmids. The transformation frequencies of the *Fncas12a1* system plasmids were higher than those of the *Fncas12a2* system plasmids in *S. griseus*. However, there were no statistically significant differences in *S. griseus* unlike *S. coelicolor* (**Supplementary Figure 2**).

Additionally, *Fncas12a* systems controlled by the promoters *ermEp\** and *Potr\** showed significantly higher transformation frequencies than those controlled by *kasOp\** (**Figures 2B,C,E,F**). These results are consistent with those of a study by Ungerer and Pakrasi, which revealed that a vector carrying *Fncas12a* yielded fewer colonies than an empty vector. *Fncas12a* is toxic to the host, and host toxicity increases with increasing *Fncas12a* promoter strength. Therefore, we chose the *Fncas12a1* system

driven by the *Potr\** or *ermEp\** promoter to perform genome editing in *Streptomyces* strains that are difficult to be transformed.


## Optimization of the CRISPR-*FnCas12a* Systems

To optimize the CRISPR-*FnCas12a* systems, we assessed the impacts of the DR length and the *FnCas12a* transcription level on genome editing efficiency. In some hosts, a shorter DR length resulted in a higher editing efficiency, as observed for *FnCas12a* in *Saccharomyces cerevisiae* (Swiat et al., 2017) and for *AsCas12a* in mammalian cells (Zetsche et al., 2017). In some other hosts, such as *Synechococcus* UTEX 2973 (Ungerer and Pakrasi, 2016), *Streptomyces* sp. NRRL S-244 (Yeo et al., 2019), and *Streptomyces albus* J1074 (Yeo et al., 2019), 36-nt DRs have shown high editing efficiency. However, there have been no reports focusing on the impact of the DR length on the editing efficiency in *Streptomyces* strains.

To confirm the impact of DR length on *FnCas12a*-mediated genome editing efficiency in *Streptomyces* strains, we chose two crRNA arrays with different DR lengths to guide *actII-orf4* gene editing in *S. coelicolor*. Array 1 contained a 19-nt DR and a

23-nt spacer, while array 2 contained a 36-nt DR and a 23-nt spacer (Figure 3A). CRISPR array 1 or 2 and an editing template were introduced into every pYL-*FnCas12a* plasmid to obtain a series of *actII-orf4* deletion plasmids (Figure 3B and Supplementary Table 3). When controlled by *kasOp\**, *FnCas12a* pairing with array 2 showed a significantly higher deletion efficiency than that pairing with array 1 for both *FnCas12a1* and *FnCas12a2* systems in *S. coelicolor* (Table 1 and Supplementary Figures 3A,B). Specifically, plasmid pYL-*kasOp\**-*FnCas12a2-actII-orf4*-DR36 led to a loss of blue pigment and complete *actII-orf4* deletion with the highest editing efficiency of 100% (Figure 3C and Table 1). In addition, when driven by *kasOp\** or *rpsLp(XC)*, the *FnCas12a2* system showed significantly higher editing efficiencies than *FnCas12a1* system for both 19-nt and 36-nt DRs (Supplementary Figure 4). In summary, pYL-*kasOp\**-*FnCas12a2* carrying the 36-nt DR showed the highest and most stable editing efficiency among all *FnCas12a* plasmid combinations (Table 1 and Supplementary Figure 4). Therefore, pYL-*kasOp\**-*FnCas12a2* with a 36-nt DR was deemed the most suitable system for genome editing.

To explore the effects of *FnCas12a* expression level on genome editing efficiency, we measured the transcription levels of *FnCas12a* in *S. coelicolor*. We found that the transcription



**TABLE 1** | Deletion efficiencies in the *actII-orf4* gene deletion experiments by using two *FnCas12a* systems.

Plasmid	Editing efficiency
pYL- <i>kasOp</i> *- <i>FnCas12a1-actII-orf4</i> -DR19	28.6% ± 11.7%
pYL- <i>kasOp</i> *- <i>FnCas12a1-actII-orf4</i> -DR36	76.2% ± 6.7%
pYL-rpsLp(XC)- <i>FnCas12a1-actII-orf4</i> -DR19	33.3% ± 6.7%
pYL-rpsLp(XC)- <i>FnCas12a1-actII-orf4</i> -DR36	71.4% ± 11.7%
pYL- <i>ermEp</i> *- <i>FnCas12a1-actII-orf4</i> -DR19	14.3% ± 11.7%
pYL- <i>ermEp</i> *- <i>FnCas12a1-actII-orf4</i> -DR36	23.8% ± 6.7%
pYL-Potr*- <i>FnCas12a1-actII-orf4</i> -DR19	9.5% ± 6.7%
pYL-Potr*- <i>FnCas12a1-actII-orf4</i> -DR36	28.6% ± 14.3%
pYL- <i>kasOp</i> *- <i>FnCas12a2-actII-orf4</i> -DR19	80.2% ± 6.3%
pYL- <i>kasOp</i> *- <i>FnCas12a2-actII-orf4</i> -DR36	100.0% ± 0.0%
pYL-rpsLp(XC)- <i>FnCas12a2-actII-orf4</i> -DR19	81.0% ± 17.8%
pYL-rpsLp(XC)- <i>FnCas12a2-actII-orf4</i> -DR36	94.4% ± 7.9%
pYL- <i>ermEp</i> *- <i>FnCas12a2-actII-orf4</i> -DR19	14.3% ± 0.0%
pYL- <i>ermEp</i> *- <i>FnCas12a2-actII-orf4</i> -DR36	28.6% ± 11.7%
pYL-Potr*- <i>FnCas12a2-actII-orf4</i> -DR19	23.8% ± 6.7%
pYL-Potr*- <i>FnCas12a2-actII-orf4</i> -DR36	21.4% ± 7.1%

levels of *FnCas12a* were significantly higher under the control of *kasOp*\* than under the control of other promoters or the inducible system for both *FnCas12a1* and *FnCas12a2* systems (Supplementary Figure 5). With regard to editing efficiency, the plasmids carrying the 36-nt crRNA and *FnCas12a* driven by the strong promoter *kasOp*\* resulted in higher editing efficiencies than those driven by a weak constitutive promoter (*ermEp*\*) or an inducible system (Potr\*) (Table 1, Supplementary Figure 6, and Tables 3, 4). Notably, the inducible system has been reported to be fully induced by OTC at a final concentration of 3  $\mu$ M in *S. coelicolor* M1146 (Wang et al., 2016). Thus, we performed transformations on M-ISP4 plates supplemented with OTC at final concentrations of 0, 0.6, 1.2, and 3.0  $\mu$ M to maintain the Potr\* system at low, medium and high activity levels. Among the four OTC concentrations, 3.0 or 1.2  $\mu$ M OTC was sufficient to induce a high editing efficiency of pYL-Potr\*-*FnCas12a1-actII-orf4*-DR19 (23.8% ± 13.5%) or pYL-Potr\*-*FnCas12a1-actII-orf4*-DR36 (28.6% ± 11.7%). In the *FnCas12a2* system, 1.2  $\mu$ M OTC supplementation was sufficient to induce a high editing efficiency of pYL-Potr\*-*FnCas12a2-actII-orf4*-DR19 (35.7% ± 7.1%) or pYL-Potr\*-*FnCas12a2-actII-orf4*-DR36 (69.1% ± 2.4%) (Supplementary Table 4). Then, we measured the transcription levels of *FnCas12a* driven by the inducible Potr\* system in the presence of 0, 0.6, 1.2 and 3.0  $\mu$ M OTC. Significantly higher transcription levels of the *FnCas12a* were produced by both the *FnCas12a1* and *FnCas12a2* systems in the presence of 0.6, 1.2 and 3.0  $\mu$ M OTC than in the absence of OTC (Supplementary Figures 5A,B). Pearson's correlation coefficients were calculated to explore the correlations between *FnCas12a* transcription levels and editing efficiencies. The analysis revealed that *FnCas12a* expression and the editing efficiency were positively correlated with the *FnCas12a1* (correlation coefficient:  $p = 0.050$ ,  $r = 0.750$ ) and *FnCas12a2* systems (correlation coefficient:  $p = 0.036$ ,  $r = 0.790$ ) when pairing with the 36-nt DR-containing crRNA (Supplementary


Figures 5C,D). Briefly, elevated *FnCas12a* expression leads to increased genome editing efficiency of *FnCas12a* pairing with the 36-nt DR-containing crRNA.

## The Ability of *FnCas12a* to Delete Large Chromosomal Fragments

pYL-*kasOp*\*-*FnCas12a2* carrying a 36-nt DR has been proven to be efficient for genome editing in *Streptomyces* strains. Thus, the ability of *FnCas12a* to delete large DNA fragments was evaluated. The actinorhodin biosynthesis gene cluster (ACT, 21.4 kb) and  $Ca^{2+}$ -dependent antibiotic biosynthesis gene cluster (CDA, 82.8 kb) in *S. coelicolor* and the daptomycin biosynthesis gene cluster (DAP, 127.6 kb) in *S. roseosporus* were selected. For single cuts, one to three spacers were selected for each cluster. The ACT-deletion strains produced red pigment on R2YE plates, while the WT strains produced blue pigment (Figure 4A). Subsequently, the PCR results showed that the 577-bp bands were amplified from the genomic DNA of WT strains but not from that of the edited strains. In addition, the 1,176-bp bands were amplified from the genomic DNA of successfully edited strains but were not amplified from the genomic DNA of WT strains (Figure 4A). The sequencing of the 1,176-bp fragments indicated that the ACT gene cluster was completely deleted from the genome of *S. coelicolor*. The deletion efficiency of the 21.4-kb gene fragment was 92.9% ± 7.2% (Supplementary Table 5). The efficiencies decreased with increasing deletion fragment sizes. The deletion efficiencies of the CDA gene cluster were 55.6% ± 7.9% (sp1), 18.8% ± 6.3% (sp2), and 25.0% ± 0% (sp3) (Supplementary Table 5 and Figure 4B). The deletion efficiencies of the DAP gene cluster were 25.0% ± 0% (sp1) and 0% (sp2) (Supplementary Table 5 and Figure 4C). To increase editing efficiency, we also attempted to introduce double cuts in the CDA gene cluster using an *FnCas12a2* system carrying two crRNA cassettes (36-nt DR+2sp-1+36-nt DR+2sp-2). Two spacers (2sp-1 and 2sp-2) flanking the edges of the CDA gene cluster were selected to cut the two loci in the cluster. Two 2-kb arms homologous to the corresponding upstream and downstream sequences of the target gene clusters were introduced into the plasmids to repair the double-strand breaks at the edges of the CDA gene cluster. However, no increase in the editing efficiency was observed (25.0% ± 0%) (2sp) (Supplementary Table 5 and Figure 4B).

## Application of the CRISPR-*FnCas12a* System for Multiplex Genome Editing in *Streptomyces* Species

Editing multiple genes step by step in *Streptomyces* is labor-intensive and time-consuming since the *Streptomyces* growth cycle is relatively long. Thus, an efficient tool is needed to perform multigene editing. pYL-*kasOp*\*-*FnCas12a2* was used to assess the possibility of CRISPR/*FnCas12a*-mediated multiplex gene deletion. We constructed an *actII-orf4/redD* double-deletion mutant and an *actI-orf1/redX* double-deletion mutant based on the pYL-*kasOp*\*-*FnCas12a2* system. Both double-deletion constructs contained two arrays. Array 2-1 consisted of the *gapdhp*(EL) promoter, DR-spacer1 and the T7 terminator. Array 2-2 consisted of the rpsLp(XC) promoter, DR-spacer2 and oop



**FIGURE 4 |** Evaluation of a large-fragment deletion via the *FnCas12a2* system. **(A)** Identification of the ACT gene cluster deletion. Red pigment was produced by the ACT-deletion strains but not by the WT strains. The primers P3 and P4 produced a 577-bp amplicon for the WT strains and no products for the ACT-deletion strains. The primers P3 and P5 generated an 1,176-bp amplicon for the ACT-deletion strains and no products for the WT strains. The sequencing data for the 1,176-bp amplicon showed the complete deletion of the ACT gene cluster. **(B)** Evaluation of CDA gene cluster deletion. For single cuts, three single spacers, sp1, sp2, and sp3, were selected. For double cuts, the paired spacers 2sp-1 and 2sp-2 were selected. The primers P6 and P7 produced a 2,438-bp amplicon for the WT strains and no products for the CDA-deletion strains. The primers P6 and P8 generated a 4,483-bp amplicon for the WT strains, but no products for the WT strains. **(C)** Identification of DAP gene cluster deletion. The primers P9 and P10 generated a 439-bp amplicon for the WT strains, but no products for the DAP-deletion strains. The primers P9 and P11 generated a 1,233-bp amplicon for the DAP-deletion strains but no products for the WT strains. The sequencing data for the 1,233-bp amplicon showed the complete deletion of the DAP gene cluster. The experiments were performed in triplicate. Red arrows indicate the successfully deleted colonies.


terminator (**Figure 5A**). *ActII-orf4* and *redD* are pathway-specific regulatory genes, and *actI-orf1* and *redX* are beta-ketoacyl synthase genes. *ActII-orf4* and *actII-orf1* are responsible for actinorhodin biosynthesis, while *redD* and *redX* contribute to undecylprodigiosin biosynthesis in *S. coelicolor*. *S. coelicolor* lacking *actII-orf4* or *actI-orf1* failed to synthesize actinorhodin and produced a red pigment on R2YE medium. *RedD* or *redX* deletion abolished undecylprodigiosin synthesis and resulted in a blue color on R2YE medium (**Figures 5B,C**). The success of double deletion of *actII-orf4* and *redD* was evaluated using PCR. For the double deletions of *actII-orf4* and *redD*, the generation of an 1,157-bp band indicated the deletion of *actII-orf4*, and the amplification of an 1,102-bp band indicated the deletion of *redD*. Among the seven randomly selected exconjugants, exconjugant 4 harbored the correct double deletion (**Figure 5B**). For the double deletion of *actI-orf1* and *redX*, the amplification of an 1,129-bp band indicated the deletion of *actI-orf1*, and the

amplification of a 2,283-bp band indicated the deletion of *redX*. Among the seven randomly selected exconjugants, exconjugant 1 harbored the correct double deletion, and colonies 3, 4, 6, and 7 showed mixtures of the *redX*-deletion mutant and the WT strain (**Figure 5C**). This phenomenon is common in microbes and has been called “incomplete genome editing” (Huang et al., 2015b; Ungerer and Pakrasi, 2016). The colonies that showed double bands with WT and mutant sizes were not classified as mutant colonies. The double-deletion efficiencies of *actII-orf4/redD* and *actI-orf1/redX* were  $14.3\% \pm 0\%$  and  $14.3\% \pm 0\%$ , respectively (**Supplementary Table 5** and **Supplementary Figure 7**).

## Precise Genome Editing With the CRISPR-*FnCas12a* Systems

pYL-*ermEp\**-*FnCas12a1* and pYL-Potr\*-*FnCas12a1* have been indicated to have increased transformation frequencies in






*Streptomyces*. Thus, the genome editing efficiencies of these constructs were tested in *S. hygroscopicus* NRRL5491, in which it is difficult to perform genetic editing. Rapamycin is a macrolide immunosuppressant produced by *S. hygroscopicus* NRRL5491 and *Actinoplanes* sp. N902-109 (Huang et al., 2015a), and it has been approved as a treatment for select conditions by the FDA (Arriola Apelo and Lamming, 2016). In addition, rapamycin exhibits antifungal and anticancer activities. According to a previous study, *Actinoplanes* sp. N902-109 carries an additional *rapTH* gene and exhibits higher production of rapamycin than *S. hygroscopicus* NRRL5491. *RapTH* encodes a homolog of type II thioesterase (Huang et al., 2015a). The *rapTH* gene has been proposed to play an important role in rapamycin generation. To introduce *rapTH* into *S. hygroscopicus* NRRL5491, we inserted the *rpsLp*(CF) promoter in front of *rapTH* and inserted the *ermEp\** promoter in front of *rapQ* in constructs

based on *pYL-ermEp\*-FnCas12a1* and *pYL-Potr\*-FnCas12a1* (Supplementary Figure 8A). The insertion fragment contained the *ermEp\** promoter (282 bp), the *rapTH* gene (756 bp) and the *rpsLp*(CF) promoter (302 bp) (a total of 1,340 bp) (Supplementary Figure 8A). Eight to twelve single colonies were selected and evaluated. A 2,277-bp band was amplified from the genomic DNA of WT strains, while a 3,668-bp band was amplified from genomic DNA of successfully edited strains (Supplementary Figure 8B). The maximum insertion efficiency of the two promoters and the *rapTH* gene was 25.0% when *pYL-Potr\*-FnCas12a1* was used in the presence of 1.2  $\mu$ M OTC (Supplementary Table 5).

WT *FnCas12a* requires a TTN PAM sequence, which may limit its application in *Streptomyces* strains with high-GC-content genomes. For gene deletion, a TTN or NGG PAM can easily be selected in the whole gene reading frame. Thus,



there are no significant restrictions in the use of CRISPR-Cas9 or CRISPR-Cas12a. However, the selection region is sometimes limited for insertions or site mutations. Therefore, the use of an NGG or TTN PAM may be restricted. Daptomycin, a cyclic lipopeptide produced in *S. roseosporus*, shows significant activity against Gram-positive pathogens, such as methicillin-resistant *Staphylococcus aureus* (MRSA) (Vilhena and Bettencourt, 2012). The daptomycin biosynthetic pathway contains three nonribosomal peptide synthetase (NRPS) genes, *dptA*, *dptBC* and *dptD* (Miao et al., 2005). To increase the production of daptomycin in *S. roseosporus*, we planned to introduce *kasOp\** in front of the *dptA* gene, which encodes the first subunit of NRPS (Figure 6A). However, no TTN PAM sequences suitable for *FnCas12a* recognition were located in front of the NRPS genes of daptomycin. Therefore, it was impossible for us to introduce a strong promoter in front of the NRPS genes via the *FnCas12a* system with

WT *FnCas12a*. To overcome this limitation, we applied a CRISPR-Cas12a3 system containing the *FnCas12a* mutant EP16, which was previously generated by our laboratory (Wang et al., 2019). Among the tested variants, the *FnCas12a* mutant EP16 (N607R/K613V/N617R/K180S/K660R/D616N) exhibited the best recognition capabilities *in vitro*, as it could recognize YN (Y = C or T), TAC and CAA PAMs. We selected three spacers adjacent to CCG, CCA and ATC PAMs with which the *FnCas12a* variant EP16, but not WT *FnCas12a*, worked well *in vitro*. The *kasOp\** promoter insertion efficiencies of CRISPR-*FnCas12a3* with CCG, CCA and ATC PAMs were  $50.0\% \pm 12.5\%$ ,  $40.0\% \pm 17.4\%$ , and  $23.6\% \pm 8.6\%$ , respectively (Supplementary Table 5). In summary, the CRISPR-*FnCas12a3* system provides opportunities to select suitable PAMs for precise genome editing in GC-rich organisms.

After we introduced *kasOp\** in front of the daptomycin biosynthesis genes, we performed plasmid clearance to generate

the strain *S. roseosporus*/PkasA (Figure 9A). Introduction of *kasOp\** increased the transcription levels of the *dptA*, *dptBC* and *dptD* genes by 1.83-, 1.12- and 2.04-fold, respectively (Supplementary Figure 9A). Compared with WT *S. roseosporus*, *S. roseosporus*/PkasA showed increased production of daptomycin (Supplementary Figures 9B,C).

In addition to insertions, CRISPR-*FnCas12a3* was also applied to generate precise site mutations. *RpsL*(SCO4659) encodes the ribosomal protein S12 in *S. coelicolor* A3(2), and its specific site mutation (K88E) confers resistance to high concentrations of streptomycin (100 µg/mL) (Shima et al., 1996). Here, we introduced a GAA mutation (K88E) to ensure the resistance of *S. coelicolor* A3(2) to streptomycin (Figure 6B). Moreover, a specific C-to-T mutation was introduced at the 267th nucleotide of *rpsL* to introduce a *BglII* restriction site for rapid identification of the correct exconjugants (Figure 6B). To avoid undesirable re-editing, the double-strand break (DSB) was repaired using a template containing altered nucleotides at the 5' end of the spacer sequence (Figure 6B). PCR amplicons of 1,726 bp were amplified from the genomes of eight exconjugants and digested separately with the *BglII* restriction enzyme. One 1,726-bp PCR amplicon was successfully digested into 1,113-bp and 613-bp fragments. Subsequent sequencing of the 1,726-bp PCR amplicon from exconjugant 3 indicated that the site mutations were successfully created (Figure 6C). The restriction digestion and DNA sequencing results showed a site mutation efficiency  $12.5\% \pm 0\%$  (Figure 6C and Supplementary Table 5). However, phenotype screening indicated that 33.3% of the *S. coelicolor* A3(2) colonies harboring pYL-*kasOp\**-*FnCas12a3*-*rpsL* were resistant to streptomycin (Figure 6D). These results suggest that some strains failed to be mutated.

## DISCUSSION

In the present study, we developed three useful genome editing tools with different advantages and applied them to examine their transformation frequencies and genome editing efficiencies in several *Streptomyces* species. The successful application of the CRISPR-*FnCas12a3* system with expanded PAM recognition ability for precise insertions and site mutations overcomes the restricted applications of the TTN PAM in organisms with high GC contents. Its flexibility in PAM selection will promote the application of *FnCas12a* in *Streptomyces* species. The *FnCas12a1* system worked well in *S. hygroscopicus*, in which CRISPR-Cas9 is ineffective. The *FnCas12a2* system efficiently deleted large chromosomal fragments (~128 kb) and was useful for deleting multiple genes. Altogether, these three systems have different advantages and are complementary to each other.

Due to the high GC content of the *Streptomyces* genome, it will not be easy for us to select a TTN PAM with high prediction scores. In particular, the number of suitable sequence region that can be targeted for insertions and site mutations is very limited. Thus, it is inconvenient to select a TTN PAM for large-scale genome engineering in *Streptomyces*. The *FnCas12a* mutant EP16 has a wide range of PAM recognition sites (60/64 sites), including YN (Y = C or T), TAC and CAA sites. EP16 cleaves

target DNA by the PAMs CCG, CCA and ATC *in vitro* with high efficiencies of 97, 94 and 96%, respectively. However, the efficiencies of CRISPR-*FnCas12a3* harboring EP16 with CCG, CCA and ATC recognition sites were only  $50.0\% \pm 12.5\%$ ,  $40.0\% \pm 17.4\%$ , and  $23.6\% \pm 8.6\%$ , respectively, *in vivo*. This phenomenon is common. As shown in the studies by Zetsche et al. and Tu et al., *FnCas12a* recognizes TTN sites *in vitro* but frequently fails to recognize TTN sites in human cells (Zetsche et al., 2015; Tu et al., 2017). *FnCas12a* has been reported to prefer KYTV in human cells (Sun et al., 2018), but prefer TTTV PAMs in rice (Zhong et al., 2018). Moreover, our unpublished data from human HEK293T cells also show a similar phenomenon. This phenomenon may be attributable to the complex microenvironments in living organisms. For instance, post-translational modifications, such as acetylation (Ishigaki et al., 2017) and methylation (Huang et al., 2018), are common in *Streptomyces*. *Moraxella bovoculi* (Mb) Cas12a has been demonstrated to lose its cleavage functions upon the acetylation of the critical PAM recognition residue Lys635 (Dong et al., 2019) *in vitro*. Taken together, results indicate that the *FnCas12a3* system overcomes the restricted applications of WT *FnCas12a*, which requires a TTN PAM for editing in *Streptomyces*. Although the editing efficiencies of the *FnCas12a3* system were low, the identification of a new Cas12a variant (EP16) that works on a broad range of PAMs in *Streptomyces* is a very important step forward. This powerful tool will enable researchers to generate desired insertions and precise site mutations and will also enable the activation of biosynthetic pathways to generate valuable natural products.

In a previous study, conservation was discovered at the 3' end of the DR sequence among all *FnCas12a* family proteins (Zetsche et al., 2015). In addition, 19-nt DR-containing crRNA cassettes have been proven to exhibit good editing efficiencies *in vitro* (Wang et al., 2019) and in some host cells, such as human HEK293T cells (Zetsche et al., 2017) and *S. cerevisiae* (Swiat et al., 2017). In another study, a 36-nt DR was applied for markerless editing in *Cyanobacteria* species (Ungerer and Pakrasi, 2016). Although *FnCas12a* has been used in many hosts, to the best of our knowledge, only one report has focused on the impacts of the DR length on the editing efficiency in *Saccharomyces cerevisiae* (Swiat et al., 2017). In the current study, a crRNA containing a 19-nt DR led to much higher editing efficiency than that containing a 36-nt DR in *Saccharomyces cerevisiae*. In *Streptomyces* species, when *FnCas12a* was controlled by strong constitutive promoters, the crRNA containing a 36-nt DR led to high editing efficiency. However, when *FnCas12a* was controlled by a weak promoter or an inducible system, significant differences in the editing efficiencies resulting from 36-nt DR-containing crRNA and 19-nt DR-containing crRNA were not observed (Table 1 and Supplementary Figure 3). Therefore, the impact of the DR length is different for different plasmid systems or hosts. Thus, the impact of the DR length should be evaluated whenever Cas12a-mediated genome editing is performed for the first time in a specific organism.

The transformation frequencies and editing efficiencies of the *FnCas12a1* and *FnCas12a2* systems were significantly different. The plasmids constructed from the *FnCas12a1* system

induced markedly higher transformation frequencies than those constructed from the *FnCas12a2* system. A comparison of the elements related to transformation revealed that the *FnCas12a1* system carries a *traJ* gene. This gene encodes an activator of the transfer (*tra*) operon, which is crucial for the transfer region of the fertility factor (Will and Frost, 2006). After the *traJ* gene was deleted from the *FnCas12a1* system, the transformation frequency was significantly decreased (**Supplementary Figure 10**). In this case, the *traJ* gene in *FnCas12a1* system was required for its high transformation frequency. Moreover, the *FnCas12a1* system contains the counterselection marker *codA(sm)*, which saves a substantial amount of time during plasmid elimination (Zeng et al., 2015). Thus, the *FnCas12a1* system is more suitable than the *FnCas12a2* system for applications in some strains with low transformation frequencies. On the other hand, Wei et al. (2017) found that strong terminators are responsible for high gene expression. To explore whether the different terminators of crRNA arrays contribute to the different editing efficiencies between *FnCas12a1* system and *FnCas12a2* system, we exchanged the terminators of crRNA arrays of the two systems to construct the plasmids pYL-*kasOp*\*-*FnCas12a1-actII-orf4-DR36-ooop* and pYL-*kasOp*\*-*FnCas12a2-actII-orf4-DR36-B1006*. The pYL-*kasOp*\*-*FnCas12a1-actII-orf4-DR36-ooop* plasmid led to significantly higher editing efficiency ( $95.2\% \pm 6.7\%$ ) than the pYL-*kasOp*\*-*FnCas12a1-actII-orf4-DR36* plasmid ( $76.2\% \pm 6.7\%$ ), and pYL-*kasOp*\*-*FnCas12a2-actII-orf4-DR36-B1006* led to a significantly lower efficiency ( $47.6\% \pm 17.8\%$ ) than pYL-*kasOp*\*-*FnCas12a2-actII-orf4-DR36* (100%) (**Supplementary Figure 11**). Thus, the oop terminator of the crRNA array was responsible for the higher editing efficiency of the *FnCas12a* system. In conclusion, the two systems described in the present study provide researchers with additional choices for manipulation of different *Streptomyces* strains.

Genome editing and transcriptional repression in *Streptomyces* with the CRISPR-Cas12a system have also recently been reported by Li et al. (2018). Those researchers obtained an editing efficiency for single-gene deletion with a CRISPR-Cas12a system of 95%, which was lower than that obtained with our *FnCas12a2* system (100%). However, the editing efficiencies of pYL-*ermEp*\*-*FnCas12a1-actII-orf4-DR19* and pYL-*ermEp*\*-*FnCas12a2-actII-orf4-DR19* were lower than those of their CRISPR-*FnCas12a* system carrying Cas12a controlled by the same promoter, *ermEp*\*. Several factors may explain this discrepancy. First, the promoter controlling the crRNA cassette in the CRISPR-Cas12a system described by Lei Li et al. was *kasOp*\*, which is stronger than the *gapdh*(EL) (Shao et al., 2013; Myronovskyi and Luzhetskyy, 2016) promoter used in our study. The high crRNA expression level may have increased the editing efficiency. Second, the strains were different. The authors of the previous study used *S. hygroscopicus* SIPI-KF and *E. coli* S17-1. *E. coli* S17-1 contains a chromosomally integrated derivative of RP4, which stimulates the integration of DNA from the donor strain into the recipient genome (Simon et al., 1983; Voeykova et al., 1998; Kieser et al., 2000). Thus, the status of the *FnCas12a* system in *Streptomyces* may be influenced by different donor strains. Importantly, *FnCas12a* was successfully applied

in the current study to accurately delete large chromosomal DNA fragments ranging from 24 to 128 kb with efficiencies ranging from 25% to  $92.9\% \pm 7.2\%$  based on HDR, and the editing efficiency obtained using HDR was much higher than that obtained using NHEJ repair (10%, 27.6 kb). Thus, HDR was more suitable than NHEJ for deleting large DNA fragments. Moreover, the *FnCas12a3* system has an expanded PAM recognition ability, which might provide increased opportunities for researchers to conduct precise genome editing. This system will be particularly useful for generating insertions and site mutations, as it facilitates selection of PAMs in the limited DNA sequence regions in *Streptomyces* strains with a high GC content. Therefore, the three *FnCas12a* systems we developed are versatile tools for precise genome editing in different *Streptomyces* strains.

In summary, the engineering tools developed in the present study are applicable for biosynthetic pathway reconstruction, metabolic engineering, and chassis cell construction in different *Streptomyces* strains. These tools will also be beneficial for natural product discovery and overproduction.

## DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/**Supplementary Material**.

## AUTHOR CONTRIBUTIONS

JZha, HL, SL, and YL designed the experiments. JZha, DZ, and JZhu performed the experiments. DZ, JZha, and YL wrote the manuscript. All authors contributed to the article and approved the submitted version.

## FUNDING

This study was supported by the Natural Science Foundation of Tianjin City (19JCYBJC24200), the National Key R&D Program of China (2018YFA0903300), the National Natural Science Foundation of China (81502966) the Projects of International Cooperation and Exchanges of NSFC (31961143005), and the International Cooperation of Sichuan Province Scientific Committee (20GJHZ0190).

## ACKNOWLEDGMENTS

We are grateful to Professor Huimin Zhao from the University of Illinois at Urbana-Champaign for providing the plasmid pCRISPomyces-2 and Professor Yuhui Sun from Wuhan University for kindly sharing the plasmid pWHU2653.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fbioe.2020.00726/full#supplementary-material>

## REFERENCES

- Arriola Apelo, S. I., and Lamming, D. W. (2016). Rapamycin: an inhibitor of aging emerges from the soil of Easter Island. *J. Gerontol. A. Biol. Sci. Med. Sci.* 71, 841–849. doi: 10.1093/gerona/glw090
- Baltz, R. H. (2008). Renaissance in antibacterial discovery from actinomycetes. *Curr. Opin. Pharmacol.* 8, 557–563. doi: 10.1016/j.coph.2008.04.008
- Cho, G., Kim, J., Park, C. G., Nislow, C., Weller, D. M., and Kwak, Y. S. (2017). Caryolan-1-ol, an antifungal volatile produced by *Streptomyces* spp., inhibits the endomembrane system of fungi. *Open Biol.* 7:170075. doi: 10.1098/rsob.170075
- Cobb, R. E., Wang, Y., and Zhao, H. (2015). High-efficiency multiplex genome editing of *Streptomyces* species using an engineered CRISPR/Cas system. *ACS Synth. Biol.* 4, 723–728. doi: 10.1021/sb500351f
- Dong, L., Guan, X., Li, N., Zhang, F., Zhu, Y., Ren, K., et al. (2019). An anti-CRISPR protein disables type V Cas12a by acetylation. *Nat. Struct. Mol. Biol.* 26, 308–314. doi: 10.1038/s41594-019-0206-1
- Dubeau, M. P., Ghinet, M. G., Jacques, P. E., Clermont, N., Beaulieu, C., and Brzezinski, R. (2009). Cytosine deaminase as a negative selection marker for gene disruption and replacement in the genus *Streptomyces* and other actinobacteria. *Appl. Environ. Microbiol.* 75, 1211–1214. doi: 10.1128/AEM.02139-08
- Frattaruolo, L., Lacret, R., Cappello, A. R., and Truman, A. W. (2017). A genomics-based approach identifies a thioviridamide-like compound with selective anticancer activity. *ACS Chem. Biol.* 12, 2815–2822. doi: 10.1021/acscchembio.7b00677
- Huang, H., Ren, S. X., Yang, S., and Hu, H. F. (2015a). Comparative analysis of rapamycin biosynthesis clusters between *Actinoplanes* sp. N902-109 and *Streptomyces hygroscopicus* ATCC29253. *Chin. J. Nat. Med.* 13, 90–98. doi: 10.1016/S1875-5364(15)60012-7
- Huang, H., Zheng, G., Jiang, W., Hu, H., and Lu, Y. (2015b). One-step high-efficiency CRISPR/Cas9-mediated genome editing in *Streptomyces*. *Acta Biochim. Biophys. Sin.* 47, 231–243. doi: 10.1093/abbs/gmv007
- Huang, T., Duan, Y., Zou, Y., Deng, Z., and Lin, S. (2018). NRPS protein MarQ catalyzes flexible adenylation and specific S-methylation. *ACS Chem. Biol.* 13, 2387–2391. doi: 10.1021/acscchembio.8b00364
- Ishigaki, Y., Akanuma, G., Yoshida, M., Horinouchi, S., Kosono, S., and Ohnishi, Y. (2017). Protein acetylation involved in streptomycin biosynthesis in *Streptomyces griseus*. *J. Proteomics* 155, 63–72. doi: 10.1016/j.jprot.2016.12.006
- Jiang, W., Bikard, D., Cox, D., Zhang, F., and Marraffini, L. A. (2013). RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat. Biotechnol.* 31, 233–239. doi: 10.1038/nbt.2508
- Jiang, Y., Qian, F., Yang, J., Liu, Y., Dong, F., Xu, C., et al. (2017). CRISPR-Cpf1 assisted genome editing of *Corynebacterium glutamicum*. *Nat. Commun.* 8:15179. doi: 10.1038/ncomms15179
- Kieser, T., Bibb, M. J., Buttner, M. J., Chater, K. F., and Hopwood, D. A. (2000). *Practical Streptomyces Genetics*. Norwich: John Innes Foundation.
- Li, L., Wei, K., Zheng, G., Liu, X., Chen, S., Jiang, W., et al. (2018). CRISPR-Cpf1-assisted multiplex genome editing and transcriptional repression in *Streptomyces*. *Appl. Environ. Microbiol.* 84:e00827-18. doi: 10.1128/AEM.00827-18
- Luo, Y., Enghiad, B., and Zhao, H. (2016). New tools for reconstruction and heterologous expression of natural product biosynthetic gene clusters. *Nat. Prod. Rep.* 33, 174–182. doi: 10.1039/c5np00085h
- Luo, Y., Zhang, L., Barton, K. W., and Zhao, H. (2015a). Systematic identification of a panel of strong constitutive promoters from *Streptomyces albus*. *ACS Synth. Biol.* 4, 1001–1010. doi: 10.1021/acssynbio.5b00016
- Luo, Y., Li, B. Z., Liu, D., Zhang, L., Chen, Y., Jia, B., et al. (2015b). Engineered biosynthesis of natural products in heterologous hosts. *Chem. Soc. Rev.* 44, 5265–5290. doi: 10.1039/c5cs00025d
- Mali, P., Yang, L., Esvelt, K. M., Aach, J., Guell, M., DiCarlo, J. E., et al. (2013). RNA-guided human genome engineering via Cas9. *Science* 339, 823–826. doi: 10.1126/science.1232033
- Miao, V., Coeffet-Legal, M. F., Brian, P., Brost, R., Penn, J., Whiting, A., et al. (2005). Daptomycin biosynthesis in *Streptomyces roseosporus*: cloning and analysis of the gene cluster and revision of peptide stereochemistry. *Microbiology* 151(Pt 5), 1507–1523. doi: 10.1099/mic.0.27757-0
- Myronovskiy, M., and Luzhetskyy, A. (2016). Native and engineered promoters in natural product discovery. *Nat. Prod. Rep.* 33, 1006–1019. doi: 10.1039/c6np00002a
- Salem, S. M., Weidenbach, S., and Rohr, J. (2017). Two cooperative glycosyltransferases are responsible for the sugar diversity of saquayamycins isolated from *Streptomyces* sp. KY 40-1. *ACS Chem. Biol.* 12, 2529–2534. doi: 10.1021/acscchembio.7b00453
- Shao, Z., Rao, G., Li, C., Abil, Z., Luo, Y., and Zhao, H. (2013). Refactoring the silent spectinabilin gene cluster using a plug-and-play scaffold. *ACS Synth. Biol.* 2, 662–669. doi: 10.1021/sb400058n
- Shao, Z., Zhao, H., and Zhao, H. (2009). DNA assembler, an in vivo genetic method for rapid construction of biochemical pathways. *Nucleic Acids Res.* 37:e16. doi: 10.1093/nar/gkn991
- Shima, J., Hesketh, A., Okamoto, S., Kawamoto, S., and Ochi, K. (1996). Induction of actinorhodin production by rpsL (encoding ribosomal protein S12) mutations that confer streptomycin resistance in *Streptomyces lividans* and *Streptomyces coelicolor* A3(2). *J. Bacteriol.* 178, 7276–7284. doi: 10.1128/jb.178.24.7276-7284.1996
- Simon, R., Priefer, U., and Pühler, A. (1983). A broad host range mobilization system for *in vivo* genetic engineering transposon mutagenesis in Gram negative bacteria. *Nat. Biotechnol.* 1, 784–791. doi: 10.1038/nbt1183-784
- Sun, H. H., Li, F. F., Liu, J., Yang, F. Y., Zeng, Z. H., Lv, X. J., et al. (2018). A single multiplex crRNA array for FnCpf1-mediated human genome editing. *Mol. Ther.* 26, 2070–2076. doi: 10.1016/j.yth.2018.05.021
- Swiat, M. A., Dashko, S., den Ridder, M., Wijsman, M., van der Oost, J., Daran, J. M., et al. (2017). FnCpf1: a novel and efficient genome editing tool for *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 45, 12585–12598. doi: 10.1093/nar/gkx1007
- Tang, X., Lowder, L. G., Zhang, T., Malzahn, A. A., Zheng, X., Voytas, D. F., et al. (2017). A CRISPR-Cpf1 system for efficient genome editing and transcriptional repression in plants. *Nat. Plants* 3:17018. doi: 10.1038/nplants.2017.18
- Toth, E., Weinhardt, N., Bencsura, P., Huszar, K., Kulcsar, P. I., Talas, A., et al. (2016). Cpf1 nucleases demonstrate robust activity to induce DNA modification by exploiting homology directed repair pathways in mammalian cells. *Biol. Direct.* 11:46. doi: 10.1186/s13062-016-0147-0
- Tu, M. J., Lin, L., Cheng, Y. L., He, X. B., Sun, H. H., Xie, H. H., et al. (2017). A 'new lease of life': FnCpf1 possesses DNA cleavage activity for genome editing in human cells. *Nucleic Acids Res.* 45, 11295–11304. doi: 10.1093/nar/gkx783
- Ungerer, J., and Pakrasi, H. B. (2016). Cpf1 is a versatile tool for CRISPR genome editing across diverse species of Cyanobacteria. *Sci. Rep.* 6:39681. doi: 10.1038/srep39681
- Vilhena, C., and Bettencourt, A. (2012). Daptomycin: a review of properties, clinical use, drug delivery and resistance. *Mini Rev. Med. Chem.* 12, 202–209. doi: 10.2174/1389557511209030202
- Voeykova, T., Emeljanova, L., Tabakov, V., and Mkrtumyan, N. (1998). Transfer of plasmid pTO1 from *Escherichia coli* to various representatives of the order Actinomycetales by intergeneric conjugation. *FEMS Microbiol. Lett.* 162, 47–52. doi: 10.1016/s0378-1097(98)00100-1
- Wang, L., Wang, H., Liu, H., Zhao, Q., Liu, B., Wang, L., et al. (2019). Improved CRISPR-Cas12a-assisted one-pot DNA editing method enables seamless DNA editing. *Biotechnol. Bioeng.* 116, 1463–1474. doi: 10.1002/bit.26938
- Wang, W., Yang, T., Li, Y., Li, S., Yin, S., Styles, K., et al. (2016). Development of a synthetic oxytetracycline-inducible expression system for *Streptomyces* using de novo characterized genetic parts. *ACS Synth. Biol.* 5, 765–773. doi: 10.1021/acssynbio.6b00087
- Wang, W. L., Wang, J., Xiang, S., Feng, X., and Yang, K. (2013). An engineered strong promoter for *Streptomyces*. *Appl. Environ. Microbiol.* 79, 4484–4492. doi: 10.1128/aem.00985-13
- Wei, L. N., Wang, Z. X., Zhang, G. L., and Ye, B. E. (2017). Characterization of terminators in *Saccharomyces cerevisiae* and an exploration of factors affecting their strength. *Chembiochem* 18, 2422–2427. doi: 10.1002/cbic.201700516
- Will, W. R., and Frost, L. S. (2006). Characterization of the opposing roles of H-NS and TraJ in transcriptional regulation of the F-plasmid tra operon. *J. Bacteriol.* 188, 507–514. doi: 10.1128/JB.188.2.507-514.2006
- Yan, X. T., An, Z., Huangfu, Y., Zhang, Y. T., Li, C. H., Chen, X., et al. (2018). Polycyclic polyprenylated acylphloroglucinol and phenolic metabolites from the aerial parts of *Hypericum elatoides* and their neuroprotective and anti-neuroinflammatory activities. *Phytochemistry* 159, 65–74. doi: 10.1016/j.phytochem.2018.12.011
- Yeo, W. L., Heng, E., Tan, L. L., Lim, Y. W., Lim, Y. H., Hoon, S., et al. (2019). Characterization of Cas proteins for CRISPR-Cas editing in streptomycetes. *Biotechnol. Bioeng.* 116, 2330–2338. doi: 10.1002/bit.27021

- Zeng, H., Wen, S., Xu, W., He, Z., Zhai, G., Liu, Y., et al. (2015). Highly efficient editing of the actinorhodin polyketide chain length factor gene in *Streptomyces coelicolor* M145 using CRISPR/Cas9-CodA(sm) combined system. *Appl. Microbiol. Biotechnol.* 99, 10575–10585. doi: 10.1007/s00253-015-6931-4
- Zetsche, B., Gootenberg, J., Abudayyeh, O., Slaymaker, M., Makarova, K., Essletzbichler, P., et al. (2015). Cpf1 is a single RNA-guided endonuclease of a Class 2 CRISPR-Cas system. *Cell* 163, 759–771. doi: 10.1016/j.cell.2015.09.038
- Zetsche, B., Heidenreich, M., Mohanraju, P., Fedorova, I., Kneppers, J., DeGennaro, E. M., et al. (2017). Multiplex gene editing by CRISPR-Cpf1 using a single crRNA array. *Nat. Biotechnol.* 35, 31–34. doi: 10.1038/nbt.3737
- Zhao, Y., Wang, L. P., and Luo, Y. Z. (2019). Recent advances in natural products exploitation in *Streptomyces* via synthetic biology. *Eng. Life Sci.* 19, 452–462. doi: 10.1002/elsc.201800137
- Zhong, Z. H., Zhang, Y. X., You, Q., Tang, X., Ren, Q. R., Liu, S. S., et al. (2018). Plant genome editing using FnCpf1 and LbCpf1 nucleases at redefined and altered PAM sites. *Mol. Plant* 11, 999–1002. doi: 10.1016/j.molp.2018.03.008
- Zhu, F., Qin, C., Tao, L., Liu, X., Shi, Z., Ma, X., et al. (2011). Clustered patterns of species origins of nature-derived drugs and clues for future bioprospecting. *Proc. Natl. Acad. Sci. U.S.A.* 108, 12943–12948. doi: 10.1073/pnas.1107336108
- Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Copyright © 2020 Zhang, Zhang, Zhu, Liu, Liang and Luo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.