



Performance and safety of diagnostic procedures in superior vena cava syndrome

To the Editor:

Superior vena cava syndrome (SVCS) is an uncommon condition resulting from extrinsic compression or intraluminal blockade of the superior vena cava. The increased upper body venous pressure results in distended subcutaneous vessels and oedema of the head, neck and arms. SVCS can be a medical emergency if associated with laryngeal or cerebral oedema. The most common SVCS aetiologies are intrathoracic malignancies, accounting for 60 to 86% of cases [1–3].

Promptly obtaining a tissue diagnosis before performing therapeutic interventions is the preferred approach in most cases [3, 4] as the majority of patients present without a prior cancer diagnosis [5]. Invasive procedures, such as bronchoscopy and endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA), may be required to obtain a histological diagnosis. Rigid bronchoscopy may also be indicated in cases of concomitant malignant central airway obstruction [6]. Endoscopic procedures are generally performed in the recumbent position, increasing intracranial pressure [7, 8], and could potentially increase the risk of complications in a population presenting subclinical cerebral oedema. Impaired venous return may also cause vascular congestion of the airway leading to an increased bleeding risk [9–11]. Impaired venous return may be further decreased in the supine position [12] with a potential for haemodynamic consequences, especially in the context of the sedations administered [10, 13, 14].

The performance and safety of endoscopic procedures in SVCS patients has not been well studied [9, 15–20]. The objectives of the present study are to evaluate the safety and performance of different diagnostic and therapeutic procedures in SVCS, with a particular focus on endoscopic procedures.

We performed a retrospective chart review of all patients who received a diagnosis of SVCS in our institution from April 1, 2012 to March 31, 2017. We included all patients with a superior vena cava occlusion on contrast computed tomography (CT) scan, including those who were initially asymptomatic. Patients were included if they were older than 18 years old and had a diagnostic or a therapeutic procedure, including standard bronchoscopy, EBUS-TBNA, transthoracic needle biopsy (TNB) and rigid bronchoscopy.

SVCS severity was stratified according to the grading system described by Yu *et al.* [4]. Central airway lumen obstruction was measured on CT scan as the ratio of the most narrowed point of the involved airway (trachea or main bronchi) *versus* its estimated normal diameter, and was considered significant if >50%. The presence of an association between SVCS severity or central airway obstruction and procedural complications was explored.

Bronchoscopies and EBUS-TBNA were performed under conscious sedation using fentanyl and midazolam in the recumbent position, whereas rigid bronchoscopies were performed under general anaesthesia. Minor complications, comprising mainly minor haemorrhages and hypoxaemias, were



@ERSpublications

Standard bronchoscopy and EBUS-TBNA have good diagnostic yield and are relatively safe procedures in the setting of SVCS. However, complications may arise from the underlying malignancy and its proximity to central vital structures. <https://bit.ly/37HXFUY>

Cite this article as: Boily-Daoust C, Plante A, Adam C. Performance and safety of diagnostic procedures in superior vena cava syndrome. *ERJ Open Res* 2021; 7: 00392-2020 [<https://doi.org/10.1183/23120541.00392-2020>].

Copyright ©ERS 2021. This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.



respectively defined as bleedings requiring local treatments and hypoxaemia extending more than 1 hour after the procedure. Complications were considered as major if they led to an escalation of care.

Forty-three patients underwent 73 procedures. Diagnostic procedures included 16 bronchoscopies, 23 EBUS-TBNA and 9 TNB. Nineteen bronchoscopies were performed for nondiagnostic purposes, including anatomic evaluation before stent placement (n=3, 15.8%), stent revision (n=8, 42.1%) and bronchial cleaning (n=8, 42.1%). Six therapeutic bronchoscopies were performed, including five for stent placement and one for tumour debulking.

Mean age was 57.2 (± 16.0) years and the vast majority of patients had a malignant diagnosis (n=40, 93%), the most frequent being nonsmall cell lung cancer (n=23, 53.5%). Twelve (19%) endoscopic procedures were performed when the SVCS severity score was elevated ($\geq 3/4$) and 22 (40%) when the central airway lumen was obstructed $>50\%$. Mean procedural time for EBUS-TBNA, diagnostic and therapeutic bronchoscopy were 18.0 (± 4.9), 13.1 (± 6.8) and 52.3 min (± 48.9), respectively.

Diagnostic yield of bronchoscopy, EBUS-TBNA and TNB were 81.3%, 87.0 and 100.0%, respectively. Regarding EBUS-TBNA, 31 stations were sampled in 23 patients (average 1.3 stations ± 0.6 per patient). Either the primary lesion or the 4R station were sampled in 20 (87%) patients.

There were no complications following TNB. Minor complications were reported in eight (13%) endoscopic procedures, four (6%) haemorrhages requiring topical treatments and six (9%) transient hypoxaemias extending more than 1 hour after the procedure but not requiring escalation of care (table 1). Two procedures were associated with two complications. There were significantly more transient hypoxaemias following EBUS compared to standard bronchoscopies (5 (21.7%) *versus* 1 (2.9%), $p=0.03$).

Three patients suffered major complications, two of which occurred during therapeutic bronchoscopies. One patient with a tracheoesophageal fistula developed a tension pneumothorax during positive pressure ventilation. Another patient with a tracheal tumour could not be ventilated or intubated following general anaesthesia induction and needed an emergency cricotomy. The third patient developed haemodynamically unstable atrial fibrillation during a standard bronchoscopy, in the setting of a neoplastic pericardial effusion. No deaths related to the complications were recorded.

SVCS severity or presence of significant central airway obstruction were not associated with more frequent complications. There were three (25%) minor complications in procedures performed with a SVCS score $\geq 3/4$ compared to five (9.6%) amongst procedures with a SVCS score $\leq 2/4$ ($p=0.70$) and there were four (18.3%) minor complications in procedures performed with a central airway obstruction $>50\%$ compared to four (12.1%) in those with a lesser degree of airway obstruction ($p=0.16$).

Timely establishment of a histological diagnosis is required in order to provide optimal treatment for patients with SVCS. In our cohort, endoscopic procedures and TNB proved to have a good diagnostic yield in patients with SVCS, which is consistent with existing literature [16, 18, 19].

The mean procedural time for EBUS-TBNA (18.7 \pm 20.4 min) was consistent with the mean procedural time reported in the literature for EBUS-TBNA performed for various indications under conscious sedation [21]. No major complications were observed with EBUS-TBNA, even if 87% patients had punctures in the vicinity of their compressed superior vena cava. Our results suggest that EBUS-TBNA in the setting of SVCS is safe.

Minor complications during endoscopic procedures, consisting of bleeding requiring topical treatments and transient hypoxaemia were not uncommon, but did not alter patient trajectory.

TABLE 1 Procedural complications

	Endoscopic procedures (n=64)				Total (n=64)
	Diagnostic		Therapeutic (n=6)	Other [#] (n=19)	
	Diagnostic bronchoscopy (n=16)	EBUS (n=23)			
Minor complications	2 (12.5)	5 (21.7)	0 (0)	1 (5.3)	8 (13)
Major complications	1 (6.3)	0 (0)	2 (33.3)	0 (0)	3 (4.7)
Total	3 (18.8)	5 (21.7)	2 (33.3)	1 (5.3)	11 (17.2)

Data are presented as n (%). EBUS: endobronchial ultrasound. [#]: stent revision (n=8), anatomic evaluation before stent placement (n=3) and bronchial cleaning (n=8).

Minor adverse events occurred in 8.6% of flexible bronchoscopies, which is consistent with other series [22–25]. The EBUS-TBNA complications rate observed in our series (21.7%) may seem higher than in previous reports. The systematic review by VON BARTHELD *et al.* [26] looked at 16 181 patients and reported an overall complication rates (minor and serious adverse events) of 0.35% for EBUS-TBNA and endoscopic ultrasound, whereas the American College of Chest Physicians Quality Improvement Registry, Evaluation and Education (AQuIRE) reported an EBUS-TBNA complication rate of 1.44% [27]. This difference can be explained by the different definitions, mainly the definition of hypoxaemia, used for complications leading to the inclusion of events that would not have been included in previous studies.

Finally, we observed complications in two of six (33.3%) therapeutic procedures, which seems higher than what was reported in previous studies [28–33]). We previously reported in a multicentric retrospective study, a 6.7% rate of nonlethal complications and the AQuIRE registry reported a complication rate of 3.9% [30]. However, the limited samples of the subgroup analysis prevented any firm conclusions.

Major complications occurred in three endoscopic procedures (4.7%), including two therapeutic bronchoscopies. We feel that major complications in our cohort were not directly related to the SVCS itself, but rather to the underlying malignancy and its proximity to central vital structures. Similarly, SCHRAUFNAGEL and colleagues [5] reported no complication from the superior vena cava obstruction itself nor the procedures, but rather from other tumour-associated complications, such as cardiac tamponade or airway obstruction.

In conclusion, TNB, standard bronchoscopy and EBUS-TBNA have good diagnostic yield and are relatively safe procedures in the setting of SVCS. However, when selecting the best diagnostic procedure, clinicians should bear in mind the comorbid conditions associated with central lesions.

Catherine Boily-Daoust, Alexandre Plante , Cedrick Adam and Marc Fortin

Dept of Pulmonary Medicine, Institut Universitaire de Cardiologie et de Pneumologie de Québec, Université Laval, Québec, QC, Canada.

Correspondence: Catherine Boily-Daoust, Dept of Pulmonary Medicine, Institut Universitaire de Cardiologie et de Pneumologie de Québec, Université Laval, 2725 Chemin Ste-Foy, Québec City, G1V 4G5, Canada. E-mail: boilydaoust.c@gmail.com

Received: 16 June 2020 | Accepted after revision: 19 Oct 2020

Conflict of interest: None declared.

References

- 1 Wilson LD, Detterbeck FC, Yahalom J. Superior vena cava syndrome with malignant causes. *N Engl J Med* 2007; 356: 1862–1869.
- 2 Cheng S. Superior vena cava syndrome: a contemporary review of a historic disease. *Cardiol Rev* 2009; 17: 16–23.
- 3 Straka C, Ying J, Kong FM, *et al.* Review of evolving etiologies, implications and treatment strategies for the superior vena cava syndrome. *Springerplus* 2016; 5: 229.
- 4 Yu JB, Wilson LD, Detterbeck FC. Superior vena cava syndrome—a proposed classification system and algorithm for management. *J Thorac Oncol* 2008; 3: 811–814.
- 5 Schraufnagel DE, Hill R, Leech JA, *et al.* Superior vena caval obstruction. Is it a medical emergency? *Am J Med* 1981; 70: 1169–1174.
- 6 Jung B, Murgu S, Colt H. Rigid bronchoscopy for malignant central airway obstruction from small cell lung cancer complicated by SVC syndrome. *Ann Thorac Cardiovasc Surg* 2011; 17: 53–57.
- 7 Kerwin AJ, Croce MA, Timmons SD, *et al.* Effects of fiberoptic bronchoscopy on intracranial pressure in patients with brain injury: a prospective clinical study. *J Trauma* 2000; 48: 878–882.
- 8 Peerless JR, Snow N, Likavec MJ, *et al.* The effect of fiberoptic bronchoscopy on cerebral hemodynamics in patients with severe head injury. *Chest* 1995; 108: 962–965.
- 9 Carr IM, Koegelenberg CF, von Groote-Bidlingmaier F, *et al.* Blood loss during flexible bronchoscopy: a prospective observational study. *Respiration* 2012; 84: 312–318.
- 10 Chaudhary K, Gupta A, Wadhawan S, *et al.* Anesthetic management of superior vena cava syndrome due to anterior mediastinal mass. *J Anaesthesiol Clin Pharmacol* 2012; 28: 242–246.
- 11 Pullerits J, Holzman R. Anaesthesia for patients with mediastinal masses. *Can J Anaesth* 1989; 36: 681–688.
- 12 Mackie AM, Watson CB. Anaesthesia and mediastinal masses. A case report and review of the literature. *Anaesthesia* 1984; 39: 899–903.
- 13 Massullo D, Di Benedetto P, Pinto G. Intraoperative strategy in patients with extended involvement of mediastinal structures. *Thorac Surg Clin* 2009; 19: 113–120.
- 14 Narang S, Harte BH, Body SC. Anesthesia for patients with a mediastinal mass. *Anesthesiol Clin North Am* 2001; 18: 559–579.
- 15 Ahmann FR. A reassessment of the clinical implications of the superior vena caval syndrome. *J Clin Oncol* 1984; 2: 961–969.
- 16 Wong MK, Tam TC, Lam DC, *et al.* EBUS-TBNA in patients presented with superior vena cava syndrome. *Lung Cancer* 2012; 77: 277–280.

- 17 Lee D, Moon SM, Kim D, *et al.* Lung cancer with superior vena cava syndrome diagnosed by intravascular biopsy using EBUS-TBNA. *Respir Med Case Rep* 2016; 19: 177–180.
- 18 Zhou ZL, Zhao H, Li Y, *et al.* Diagnostic value of endobronchial ultrasound guided transbronchial needle aspiration in superior vena cava syndrome. *Chin Med J* 2013; 126: 4453–4456.
- 19 Selçuk ZT, Firat P. The diagnostic yield of transbronchial needle aspiration in superior vena cava syndrome. *Lung Cancer* 2003; 42: 183–188.
- 20 Brundyn K, Koegelenberg CF, Diacon AH, *et al.* Transbronchial fine needle aspiration biopsy and rapid on-site evaluation in the setting of superior vena cava syndrome. *Diagn Cytopathol* 2013; 41: 324–329.
- 21 Casal RF, Lazarus DR, Kuhl K, *et al.* Randomized trial of endobronchial ultrasound-guided transbronchial needle aspiration under general anesthesia versus moderate sedation. *Am J Respir Crit Care Med* 2015; 191: 796–803.
- 22 Leiten EO, Martinsen EM, Bakke PS, *et al.* Complications and discomfort of bronchoscopy: a systematic review. *Eur Clin Respir J* 2016; 3: 33324.
- 23 Bechara R, Beamis J, Simoff M, *et al.* Practice and complications of flexible bronchoscopy with biopsy procedures. *J Bronchol* 2005; 12: 139–142.
- 24 Grendelmeier P, Kurer G, Pflimlin E, *et al.* Feasibility and safety of propofol sedation in flexible bronchoscopy. *Swiss Med Wkly* 2011; 141: w13248.
- 25 Hehn BT, Haponik E, Rubin HR, *et al.* The relationship between age and process of care and patient tolerance of bronchoscopy. *J Am Geriatr Soc* 2003; 51: 917–922.
- 26 von Bartheld MB, van Breda A, Annema JT. Complication rate of endosonography (endobronchial and endoscopic ultrasound): a systematic review. *Respiration* 2014; 87: 343–351.
- 27 Eapen GA, Shah AM, Lei X, *et al.* Complications, consequences, and practice patterns of endobronchial ultrasound-guided transbronchial needle aspiration: results of the AQUiRE registry. *Chest* 2013; 143: 1044–1053.
- 28 Stahl DL, Richard KM, Papadimos TJ. Complications of bronchoscopy: a concise synopsis. *Int J Crit Illn Inj Sci* 2015; 5: 189–195.
- 29 Fortin M, Yarmus L, Rendina EA, *et al.* Multi-institutional retrospective analysis of adverse events following rigid tracheobronchoscopy. *Respirology* 2021; 26: 87–91.
- 30 Ost DE, Ernst A, Grosu HB, *et al.* Therapeutic bronchoscopy for malignant central airway obstruction: success rates and impact on dyspnea and quality of life. *Chest* 2015; 147: 1282–1298.
- 31 Drummond M, Magalhães A, Hespagnol V, *et al.* Rigid bronchoscopy complications in a university hospital. *J Bronchol* 2003; 10: 177–182.
- 32 Perrin G, Colt HG, Martin C, *et al.* Safety of interventional rigid bronchoscopy using intravenous anesthesia and spontaneous assisted ventilation. A prospective study. *Chest* 1992; 102: 1526–1530.
- 33 Manalan K, Ryan K, Basheer H. An audit review of safety and complication rates, of rigid bronchoscopy and large airways intervention, in a London tertiary centre. *Eur Respir J* 2018; 52: Suppl. 62, PA1748.