

We investigated the effect of transforming factor factor- β_1 (TGF- β_1) on thromboxane B₂ (TXB₂) and prostaglandin E₂ (PGE₂) production in *in vitro* silica dust-exposed rat alveolar macrophages (AM). In the presence of 5 μ g of anti-TGF- β_1 antibodies, TXB₂ production decreased, but PGE₂ production increased. Addition of 2 ng of TGF- β_1 to the culture medium potentiated TXB₂ production, but PGE₂ production apparently did not change. At 50 ng of TGF- β_1 , TXB₂ production decreased, and PGE₂ production varied. Our data suggest that in rat AM: (1) both endogenous and exogenous TGF- β_1 regulate TXB₂ production; and (2) in the absence of endogenous TGF- β_1 the liberation of PGE₂ increases; however, exogenous TGF- β_1 does not have a regulatory effect on PGE₂.

Key words: Rat alveolar macrophages, PGE₂, TGF- β_1 , TXB₂

Regulation of TXB₂ and PGE₂ production by TGF- β_1 in *in vitro* silica dust-exposed rat alveolar macrophage

Urszula Orlinska^{CA,*} and Douglas C. Kuhn

Department of Pathology, Hershey Medical Center, The Pennsylvania State University, P.O. Box 850, Hershey, PA 17033, USA

*Present address: Paradigm Biosciences, Inc., 2401 Foothill Dr., Salt Lake City, UT 84109, USA

^{CA}Corresponding Author

Introduction

Chronic airway inflammation and lung fibrosis are common features of prolonged exposure to mineral particles of silica in humans and in rodents.¹ Alveolar macrophages (AM) have an essential role in lung clearance and defence mechanisms against inhaled particles.² They fulfil these functions via inflammatory mediators produced upon macrophage activation. Among the many mediators are arachidonic acid metabolites such as thromboxane A₂ (TXA₂) and prostaglandin E₂ (PGE₂). Thromboxane A₂ (of which TXB₂ is the stable metabolite) is a very potent platelet aggregating factor and pulmonary vasoconstrictor. A recent study of the regulatory effect of TXB₂ on the proliferation of vascular smooth muscle cells from rats demonstrated a rapid build-up of cytoskeletal protein in these cells in hypertension, suggesting that TXB₂ may play a role in the remodelling of the vascular wall in hypertension.^{3,4}

Prostaglandin E₂ is one of the key inflammatory mediators with multiple functions. In the lung, it induces bronchodilation and causes an increase in epithelial cell chloride secretion.⁵ PGE₂ induces vasodilatation in unventilated foetal lung.⁶

Another mediator that is produced by alveolar macrophages is transforming growth factor- β_1 (TGF- β_1). TGF- β_1 is a multifunctional peptide. TGF- β_1 can influence cell proliferation or differentiation and other functions in various types of

cells, some of which are related to inflammation and tissue fibrosis. For example, TGF- β_1 increases the rate of fibronectin gene transcription,⁷ can down-regulate interleukin-1 β receptors,⁸ and inhibits interleukin-6 release from fibroblasts.⁹ The most profound effect of TGF- β_1 is its ability to enhance the formation of extracellular matrix. TGF- β_1 promotes the formation of collagen and fibronectin in fibroblasts of human, rat and mouse origin. From *in vivo* studies, it has been shown that TGF- β_1 can stimulate the formation of the typical granulation tissue found in tissue repair.

Although pulmonary macrophage-derived mediators have a pivotal role in lung inflammation and fibrosis, the interrelationship between these mediators is just beginning to be understood. We have shown the very early release (15 min) of TGF- β_1 from silica dust-exposed rat alveolar macrophages.¹⁰ The amount of TGF- β_1 released at this time was the greatest over 24 h kinetic studies, and preceded the production of TXB₂ and PGE₂. To further unravel the interrelationship between eicosanoids and TGF- β_1 produced by alveolar macrophage, the modification of TXB₂ and PGE₂ production by TGF- β_1 in *in vitro* silica-exposed rat alveolar macrophages was investigated. We observed decreased production of TXB₂ in the presence of anti-TGF- β_1 antibodies, or high concentration (50 ng) of TGF- β_1 . TXB₂ production increased in the presence of a low concentration of TGF- β_1 (2 ng) in the culture medium. PGE₂ production increased in

the presence of anti-TGF- β_1 antibodies. Exogenous TGF- β_1 at low or high concentration did not have any effect on PGE₂ production in rat alveolar macrophages.

Materials and Methods

Materials: Hanks' solution and Dulbecco's phosphate buffered saline (DPBS), and HEPES (4-(2-hydroxyethyl)-1-piperazine ethanesulfonic acid) buffer were prepared in our laboratory. Six-well tissue culture plates were obtained from Falcon Laboratory Products (Beckton and Dickinson Labware, Lincoln Park, NJ). DMSO (dimethylsulfoxide) was obtained from Calbiochem (San Diego, CA). Allopurinol, oxypurinol, allopurinol riboside, Penn/Strep, M199 medium (without phenol red), and human serum albumin were all obtained from Sigma (St. Louis, MO). Human recombinant and polyclonal anti-TGF- β_1 antibody were obtained from R&D Systems (Minneapolis, MN). RIA reagents for TXB₂ and PGE₂ assay were purchased from Advanced Magnetics, Inc. (Boston, MA).

Isolation and culture of rat alveolar macrophages: Sprague-Dawley rats weighing approximately 225 g were lavaged *ex vivo*. Rats were given an intramuscular injection of sodium pentobarbital (10 mg) and the rib cage was opened to expose the lungs. An incision halfway through the trachea was made and the lavage tube of the trachea cannula was inserted into the incision. The lungs were washed five times each with 10 ml of lavage fluid (phosphate-buffered saline containing 0.1% EDTA, at 37°C). The collected lavage fluid was filtered through sterile Nitex gauze and centrifuged at 250 × g for 10 min. The supernatant was discarded, and the cells resuspended in M199, 10 mM HEPES 0.3% human serum albumin, 100 U/ml penicillin and 100 µg/ml streptomycin (culture medium). Cells from all rats were pooled and centrifuged again at 250 × g for 10 min. Cells were enumerated in a haemocytometer. Viability was determined by the exclusion of trypan blue and cells were plated at 1 × 10⁶/ml of culture medium in 6-well culture plates. The cells were preincubated for 1 h at 37°C in a humidified atmosphere at 5% CO₂ in air. The nonadherent cells were removed by aspiration. Fresh culture medium (37°C) was placed in the wells and unlabelled arachidonic acid was added at a final concentration of 3 µM.¹⁰ This incubation with arachidonic acid was conducted in order to replenish endogenous arachidonic acid utilized during the exaggerated eicosanoid production which occurs during cell isolation and attachment.¹¹ The cells were incu-

bated at 37°C for 24 h. After incubation the medium containing arachidonic acid was removed and fresh culture medium was placed in the wells. Macrophages were activated with a microcrystalline form of silica dust (Min-U Sil, 2–7 µm diameter, provided by Dr Peter Bolsitis, MIT) at a final concentration of 200 µg/ml, at 37°C for 24 h, in the absence or presence of anti-TGF- β_1 antibodies, or TGF- β_1 . Nonspecific antibody as an isotype-matched control for anti-TGF- β_1 antibody was tested. Media were collected into silianized tubes for the determination of eicosanoids.

Radioimmunoassay for TXB₂ and PGE₂: Following incubation, the culture medium was removed and acidified to pH 3.0 with 1 N HCl. Eicosanoids were extracted twice with 2 ml ethyl acetate/cyclohexane (1:1). The extracts were stored at –20°C until analysis of eicosanoids by specific radioimmunoassay, as described previously.¹² The antibodies utilized in the assay of TXB₂ or PGE₂ showed less than 1% cross-reactivity with other eicosanoids.

Cell viability test: The lactate dehydrogenase (LDH) activity in rat culture conditions was measured using a Sigma kit for LDH. There was no detectable LDH enzyme activity in the culture medium from rat alveolar macrophages.

Data analysis: Results are presented as means ± S.E. of a number of independent experiments.

Results

The effect of TGF- β_1 on TXB₂ production in rat alveolar macrophages: Table 1 shows the effect of anti-TGF- β_1 antibodies, or low or high concentrations of TGF- β_1 , on TXB₂ production in silica-exposed rat alveolar macrophages. In the presence of 5 µg of anti-TGF- β_1 antibodies in the culture medium, during the process of macrophage activation with silica, the production of TXB₂ was inhibited. The TGF- β_1 , at 2 ng concentration in the culture medium, potentiated TXB₂ production. At 50 ng of TGF- β_1 in the culture medium, the production of TXB₂ was inhibited.

The effect of TGF- β_1 on PGE₂ production in rat alveolar macrophages: Table 2 shows the effect of anti-TGF- β_1 antibodies, or low or high concentrations of TGF- β_1 , on PGE₂ production in silica-exposed rat alveolar macrophage. In the presence of 5 µg of anti-TGF- β_1 antibodies in the culture medium, the production of PGE₂ was

Table 1. The effect of TGF-β₁ on TXB₂ production by rat alveolar macrophages exposed to silica dust

Treatment	Experiment 1		Experiment 2		Experiment 3	
	TXB ₂ (pg/ml)	% Control	TXB ₂ (pg/ml)	% Control	TXB ₂ (pg/ml)	% Control
Control	62 ± 2		179 ± 5		79 ± 3	
Silica	132 ± 2	213	369 ± 2	206	172 ± 9	218
Silica + 5 μg anti-TGF-β ₁	60 ± 5	100	313 ± 2	175	103 ± 7	130
Silica + 2 ng TGF-β ₁	ND	ND	491 ± 3	274	219 ± 2	277
Silica + 50 ng TGF-β ₁	47 ± 5	76	69 ± 0	39	158 ± 3	200

Alveolar macrophages at 1×10^6 cells were activated with 200 μg of silica dust in the absence or presence of anti-TGF-β₁ antibodies, or TGF-β₁ added at the time of cell activation, and incubated at 37°C for 24 h. TXB₂ was assayed by the RIA as described in Methods. $n=2$ in each experimental group. ND = not done.

Table 2. The effect of TGF-β₁ on PGE₂ production by rat alveolar macrophages exposed to silica dust

Treatment	Experiment 1		Experiment 2		Experiment 3	
	PGE ₂ (pg/ml)	% Control	PGE ₂ (pg/ml)	% Control	PGE ₂ (pg/ml)	% Control
Control	445 ± 6		207 ± 9		411 ± 10	
Silica	939 ± 3	211	241 ± 10	116	438 ± 10	107
Silica + 5 μg anti-TGF-β ₁	1137 ± 75	256	330 ± 9	159	783 ± 20	191
Silica + 2 ng TGF-β ₁	ND	ND	253 ± 3	122	457 ± 2	111
Silica + 50 ng TGF-β ₁	840 ± 91	189	227 ± 7	110	504 ± 20	123

Alveolar macrophages at 1×10^6 cells were activated with 200 μg of silica dust in the absence or presence of anti-TGF-β₁ antibodies, or TGF-β₁ added at the time of cell activation, and incubated at 37°C for 24 h. PGE₂ was assayed by the RIA as described in Methods. $n=2$ in each experimental group. ND = not done.

potentiated. The addition of 2 ng of TGF-β₁ to the culture medium had no apparent effect on PGE₂ production in either of two experiments performed. At 50 ng of TGF-β₁, the production of PGE₂ varied.

Discussion

This study was designed to evaluate the regulation of TXB₂ and PGE₂ production by TGF-β₁ in alveolar macrophages. The reason for these studies was to supplement the limited information in the literature on the interrelationship between the inflammatory mediators released by a specific inflammatory cell. TXB₂, PGE₂ and TGF-β₁ are the principal inflammatory mediators released by alveolar macrophages during the lung's defence process against pathogens or foreign matter. We have demonstrated in earlier studies¹⁰ that these three mediators are released from rat alveolar macrophages upon exposure to silica dust. In those studies we observed a striking difference between the kinetics of release of TGF-β₁ (maximum release at 15 min post-silica dust) and TXB₂ (plateau release at 6 h post-silica dust), or PGE₂ release (plateau release at 6 h post-silica dust). Because it is thought that TGF-β₁ has a central regulatory role in vascular physiology and pathology, we formulated a hypothesis that perhaps TGF-β₁ has an impact on the production of either TXB₂ or PGE₂, or both, by alveolar macrophage upon silica dust-exposure.

We observed a positive effect of TGF-β₁ on TXB₂ production in rat alveolar macrophages which had been exposed to silica dust. In the absence of TGF-β₁ (plus anti-TGF-β₁ antibodies), the quantity of TXB₂ released from rat alveolar macrophages to the culture medium in response to silica exposure was significantly smaller when compared to the response with silica alone. Exogenous TGF-β₁, at low concentration, potentiated the action of silica dust, and the amount of TXB₂ produced by rat alveolar macrophages was significantly greater than the amount of TXB₂ released by alveolar macrophages activated by silica alone. High concentration of TGF-β₁, however, inhibited TXB₂ production in rat alveolar macrophages; in response to silica, the amount of TXB₂ released was smaller than the amount of TXB₂ produced in rat alveolar macrophages activated with silica alone, or in the absence of TGF-β₁ (in the presence of anti-TGF-β₁ antibodies). This observation suggests that upon exposure to silica dust a homeostatic auto-crine loop for TGF-β₁, that regulates the production of TXB₂, appears to function in rat alveolar macrophages. Our data also suggest the role for concentration of TGF-β₁ in the regulation of TXB₂ production; depending on the concentration, this growth factor appears to be either stimulatory or inhibitory to rat alveolar macrophage, and to TXB₂ production.

At the same time, however, the effect of TGF-β₁ on the generation of another eicosanoid,

namely PGE₂, in silica-exposed rat alveolar macrophages appeared to be very different to that observed for TXB₂. In the absence of TGF-β₁ (plus anti-TGF-β₁ antibodies), the amount of PGE₂ produced in silica-exposed alveolar macrophages was greater than when compared to silica alone. When anti-TGF-β₁ antibodies were present in the culture medium during the exposure of macrophages to silica, we observed that (1) these macrophages liberated greater amount of PGE₂ compared to silica alone; and (2) there was a stimulation of PGE₂ production by alveolar macrophages which did not produce PGE₂ after exposure to silica. Furthermore, the presence of exogenous TGF-β₁, at low or high concentrations, had either no apparent effect, or a variable effect, respectively, on PGE₂ production in silica-exposed rat alveolar macrophages when compared to silica alone. Our data suggest, therefore, a possible distinct biological role for endogenous, but not exogenous TGF-β₁ to down-regulate PGE₂ production in rat alveolar macrophages after exposure to silica.

Data presented in this manuscript provide evidence of the regulatory role of TGF-β₁ in the liberation of inflammatory eicosanoids, TXB₂ and PGE₂, in rat alveolar macrophages, when the cells are exposed to silica dust. In support of our observation are the reports of Datta *et al.*, where the authors demonstrate that TGF-β₁ inhibits lipoxygenase and epoxygenase eicosanoid production by osteosarcoma cells,¹³ or eicosanoid metabolism in osteogenic osteosarcoma cells.¹⁴ The regulation of eicosanoid liberation by TGF-β₁, or the regulation of TNF-α production by TGF-β₁¹³ suggests that the mechanism of action of TGF-β₁ on tissues may not necessarily be direct, but rather it requires actions of other mediators. The interrelationship between TGF-β₁ itself and all the inflammatory mediators directed by TGF-β₁ in inflammation appears to be very intriguing.

In summary, our data suggest that in silica-exposed alveolar macrophages, both endogenous and exogenous TGF-β₁ regulate TXB₂ production. Our data suggest that endogenous, but not exogenous TGF-β₁ has a regulatory role in PGE₂ production. Because both TXB₂ and PGE₂ were measured in the same samples, and because there is a consistent increase in silica-stimulated

PGE₂ synthesis in the presence of anti-TGF-β₁ antibodies in the absence of an effect of authentic TGF-β₁, it appears that in silica-exposed alveolar macrophages the regulatory effects of TGF-β₁ are different between the different components of the arachidonic acid pathway.

References

1. Davis GS. Pathogenesis of silicosis: current concepts and hypotheses. *Lung* 1986; **164**: 139–154.
2. Tarling J, Lin HS, Hsu S. Self-renewal of pulmonary alveolar macrophages: evidence from radiation chimera studies. *J Leukocyte Biol* 1987; **42**: 443–446.
3. Ishimutsu T, Uehara Y, Ishii M, Ikeda T, Matsuoaka H, Sugimoto T. Thromboxane and vascular smooth muscle cell growth in genetically hypertensive rats. *Hypertension* 1988; **12**: 46–51.
4. Nagata T, Uehara Y, Numabe A, *et al*. Regulatory effect of thromboxane A₂ on proliferation of vascular smooth cells from rats. *Am J Physiol* 1992; **263**: H1331–H1338.
5. Sigal E. The molecular biology of mammalian arachidonic acid metabolism. *Am J Physiol* 1991; **260**: L13–L28.
6. Cassin S, Tod M, Philips J, Frisinger J, Jordan J, Gibbs C. Effects of prostaglandin D₂ on perinatal circulation. *Am J Physiol* 1981; **240**: H755–H760.
7. Dean DC, Newby RF, Burgois S. Regulation of fibronectin biosynthesis by dexamethasone, transforming growth factor β, and cAMP in human cell lines. *J Cell Biol* 1988; **106**: 2159–2170.
8. Dubois CM, Ruscetti FW, Palaszynski EW, Falk LA, Oppenheim JJ, Keller JR. Transforming growth factor β is a potent inhibitor of interleukin 1 (IL-1) receptor expression: proposed mechanism of inhibition of IL-1 action. *J Exp Med* 1990; **172**: 737–744.
9. Elias JA, Lentz V, Cummings PJ. Transforming growth factor-β regulation of IL-6 production by unstimulated and IL-1 stimulated human fibroblasts. *J Immunol* 1990; **146**: 3437–3443.
10. Orlinska U, Palmer C, Kuhn DC, Demers LM. Interrelationship between hydroxyl radical formation, and thromboxane B₂ and transforming growth factors β₁ production in *in vitro* silica dust-exposed rat alveolar macrophage. *Am Rev Res Dis* (accepted).
11. Kouzan S, Nolan RD, Fournier T, Bignon J, Eling TE, Brody AR. Stimulation of arachidonic acid metabolism by adherence of alveolar macrophages to a plastic substrate. *Am Rev Res Dis* 1988; **137**: 38–43.
12. Demers LM. Prostaglandins, thromboxane, and leukotrienes. In Race GJ, ed. *Laboratory Medicine*. Philadelphia: Harper and Row, 1983, pp. 1–21.
13. Datta H, Sullivan M, Ohri S, MacIntyre I. Transforming growth factor-β₁ inhibits lipoxygenase and epoxygenase eicosanoid production by osteosarcoma cells. *Biochem Soc Trans* 1990; **18**: 1250–1260.
14. Datta HK, Sullivan M, Rathod H, MacIntyre I. Transforming growth factor-β modulates eicosanoid metabolism in osteogenic osteosarcoma cells. *Biochem Biophys Res Comm* 1991; **178**: 940–945.

ACKNOWLEDGEMENTS. Work presented in this manuscript was supported by a grant from the US Dept. of Interior's Mineral Institute Program administered through the Generic Mineral Technology Center for Respirable Dust, grant number 1135142. We thank Stephen Bobar for his great help with lavages of rat lungs and for the excellent performance of radio-immunoassays.

Received 4 July 1995;
accepted in revised form 31 August 1995