

Current awareness on yeast

(4 weeks journals - Search completed at 10th. Nov. 2004)

CURRENT AWARENESS - USE THE LINKS ONLINE

View this bibliography online at



www.interscience.wiley.com
Your gateway to Wiley journals online

The articles use , the innovative multi-publisher reference linking. This enables readers to move seamlessly from the reference to the cited publication, typically located on a different server and published by a different publisher. Links to MEDLINE abstracts are also available for key journals in the biosciences.

27/45

I Books, reviews & symposia

- Barnett JA. (2004). Univ East Anglia, Sch Biol Sci, Norwich NR4 7TJ, England. A history of research on yeasts 8: Taxonomy (Review). *Yeast* **21**, (14) 1141
- Feldhaus MJ, Siegel RW. (2004). Pacific NW Natl Lab, MSIN K4-12, POB 999, 902 Battelle Blvd, Richland, Wa 99352, USA. Yeast display of antibody fragments: A discovery and characterization platform (Review). *J Immunol Methods* **290**, (1-2) 69
- Novak S, Zechner-Krpan V*, Maric V. (2004). *Univ Zagreb, Fac Food Science & Biotechnology, Pierottijeva 6, HR-10000 Zagreb, Croatia. Regulation of maltose transport and metabolism in *Saccharomyces cerevisiae* (Review). *Food Technol Biotechnol* **42**, (3) 213
- Ton VK, Rao R*. (2004). *Johns Hopkins Univ Sch Med, Department Physiol, 725 Nth Wolfe St, Baltimore, Md 21205, USA. Functional expression of heterologous proteins in yeast: Insights into Ca²⁺ signaling and Ca²⁺-transporting ATPases (Review). *Am J Physiol* **287**, (3) C580
- Valdivia RH. (2004). Duke Univ Med Ctr, Dept Mol Genet & Microbiol, Ctr Microbial Pathogenesis, Durham, NC 27710, USA. Modeling the function of bacterial virulence factors in *Saccharomyces cerevisiae* (Review). *Eukaryot Cell* **3**, (4) 827
- Watanabe R, Riezman H*. (2004). *Univ Geneva, Sci II, 30 Quai E Ansermet, CH-1211 Geneva 4, Switzerland. Differential ER exit in yeast and mammalian cells. *Curr Opin Cell Biol* **16**, (4) 350
- Weston AD, Baliga NS, Bonneau R, Hood L. (2003). Inst Syst Biol, Seattle, Wa 98103, USA. Systems approaches applied to the study of *Saccharomyces cerevisiae* and *Halobacterium* sp. *Cold Spring Harb Symp Quant Biol* (68) 345
- Bariloche, Rio Negro, Argentina. Constitutive and UV-inducible synthesis of photoprotective compounds (carotenoids and mycosporines) by freshwater yeasts. *Photochem Photobiol* **3**, (3) 281
- Lu HZ, Jia JH, Wang QM, Bai FY*. (2004). *Chinese Acad Sci, Institute Microbiol, Systemat Mycol & Lichenol Laboratory, CN-100080 Beijing, Peoples Rep China. *Candida asparagi* sp nov., *Candida diospyri* sp nov., and *Candida qinlingensis* sp nov., novel anamorphic, ascomycetous yeast species. *Int J Syst Evol Microbiol* **54**, (4) 1409
- Paraggio M. (2004). Univ Basilicata, Dept Biol, Viale Ateneo, Lucano 10, IT-85100 Potenza, Italy. Biodiversity of a natural population of *Saccharomyces cerevisiae* and *Hanseniaspora uvarum* from Aglianico del Vulture. *Food Technol Biotechnol* **42**, (3) 165
- Peter G, Dlačhy D, Vasdinyei R, Tornai-Lehoczki J, Deak T. (2004). Budapest Univ Econ Sci & Publ Adm, Natl Collect Agr & Ind Microorganisms, Somloi ut 14-16, HU-1118 Budapest, Hungary. *Candida galli* sp nov., a new yeast from poultry. *Antonie van Leeuwenhoek* **86**, (2) 105
- Sommaruga R, Libkind D, Van Broock M, Whitehead K. (2004). University Innsbruck, Institute Zool & Limnol, Laboratory Aquatic Photobiology & Plankton Ecology, Technikerstr 25, AT-6020 Innsbruck, Austria. Mycosporine- glutaminol-glucoside, a UV-absorbing compound of two *Rhodotorula* yeast species. *Yeast* **21**, (13) 1077

3 Biochemistry

2 General

- Abdrakhmanova A, Zickermann V, Bostina M, Radermacher M, Schagger H, Kerscher S, Brandt U*. (2004). *Univ Frankfurt, Inst Biochem I, Fachbereich Med, ZBC, Theodor Stern Kai 7, Haus 25B, DE-60590 Frankfurt, Germany. Subunit composition of mitochondrial complex I from the yeast *Yarrowia lipolytica*. *Biochim Biophys Acta* **1658**, (1-2) 148
- Atyaksheva LF, Poltorak OM, Chukhrai ES. (2004). Moscow MV Lomonosov State Univ, Fac Chem, Vorobeyevy gory, RU-119899 Moscow, Russia. Thermal stability of yeast β -galactosidase. *Russ J Phys Chem* **78**, (2) 287
- Bhatnagar A, Raghavendra PR, Kranthli BV, Rangarajan PN*. (2004). *Indian Inst Sci, Dept Biochem, IN-560012 Bangalore, India. Yeast cytochrome *c* is a sequence-specific DNA-binding protein. *Biochem Biophys Res Commun* **321**, (4) 900
- Ergin C, Ilkit M, Hilmioglu S, Kaleli I, Gulbaba AG, Demirci M, Kaya S. (2004). Pamukkale Univ, Fac Med, Dept Microbiol, Posta Kutusu 119, TR-20100 Denizli, Turkey. The first isolation of *Cryptococcus neoformans* from eucalyptus trees in South Aegean and Mediterranean Regions of Anatolia in Turkey despite Taurus Mountains alkalinity. *Mycopathologia* **158**, (1) 43
- Libkind D, Perez P, Sommaruga R, Dieguez MD, Ferraro M, Brizzio S, Zagarese H, Van Broock M. (2004). Univ Nacl Comahue, Ctr Reg Univ Bariloche, Unidad Postal Univ, AR-8400 San Carlos de

In order to keep subscribers up-to-date with the latest developments in their field, this current awareness service is provided by John Wiley & Sons and contains newly-published material on yeasts. Each bibliography is divided into 10 sections. 1 Books, Reviews & Symposia; 2 General; 3 Biochemistry; 4 Biotechnology; 5 Cell Biology; 6 Gene Expression; 7 Genetics & Genomics; 8 Physiology; 9 Medical Mycology; 10 Recombinant DNA Technology. Within each section, articles are listed in alphabetical order with respect to author. If, in the preceding period, no publications are located relevant to any one of these headings, that section will be omitted.

- Bowman EJ, Graham LA, Stevens TH, Bowman BJ*. (2004). *Univ Calif Santa Cruz, Dept Mol Cell & Dev Biol, Santa Cruz, Ca 95064, USA. The bafilomycin/concanamycin binding site in subunit *c* of the V-ATPases from *Neurospora crassa* and *Saccharomyces cerevisiae*. *J Biol Chem* **279**, (32) 33131
- Cetkauskaite A, Pessala P, Sodergren A. (2004). Vilnius State Univ, Dept Biochem & Biophys, Ciurlionio Str 21, LT-2009 Vilnius, Lithuania. Elemental sulfur: Toxicity *in vivo* and *in vitro* to bacterial luciferase, *in vitro* yeast alcohol dehydrogenase, and bovine liver catalase. *Environ Toxicol* **19**, (4) 372
- Ebert CE, Beattie DS*. (2004). *West Virginia Univ Sch Med, Dept Biochem & Mol Pharmacol, Morgantown, WV 26506, USA. A compensatory double mutation of the alanine-86 to leucine mutant located in the hinge region of the iron-sulfur protein of the yeast cytochrome *bc₁* complex. *Arch Biochem Biophys* **429**, (1) 16
- Hornig YC, Cobine PA, Maxfield AB, Carr HS, Winge DR*. (2004). *Univ Utah Hlth Sci Ctr, Salt Lake City, Ut 84132, USA. Specific copper transfer from the Cox17 metallochaperone to both Sco1 and Cox11 in the assembly of yeast cytochrome *c* oxidase. *J Biol Chem* **279**, (34) 35334
- Ichikawa N, Mizuno M. (2004). Osaka City Univ, Fac Human Life Sci, Dept Food & Nutr, 3-3-138 Sugimoto, Osaka 558 5855, Japan. Functional expression of hexahistidine-tagged β -subunit of yeast F₁-ATPase and isolation of the enzyme by immobilized metal affinity chromatography. *Protein Expr Purif* **37**, (1) 97
- Kersting MC, Choi HS, Carman GM*. (2004). *Rutgers State Univ, Dept Food Sci, 65 Dudley Rd, New Brunswick, NJ 08901, USA. Regulation of the yeast *EKII*-encoded ethanolamine kinase by inositol and choline. *J Biol Chem* **279**, (34) 35353
- Kumar DP, Tiwari A, Bhat R*. (2004). *Jawaharlal Nehru Univ, Ctr Biotechnol, IN-110067 New Delhi, India. Effect of pH on the stability and structure of yeast hexokinase A - Acidic amino acid residues in the cleft region are critical for the opening and the closing of the structure. *J Biol Chem* **279**, (31) 32093
- Liger D, Graillie M, Zhou CZ, Leulliot N, Quevillon-Cheruel S, Blondeau K, Janin J, Van Tilbeurgh H*. (2004). *University Paris 11, Inst Biochim & Biophys Mol & Cellulaire, CNRS UMR 8619, Batiment 430, FR-91405 Orsay, France. Crystal structure and functional characterization of yeast YLR011wp, an enzyme with NAD(P)H-FMN and ferric iron reductase activities. *J Biol Chem* **279**, (33) 34890
- Marina P, Martinez-Costa OH, Calderon IL, Aragon JJ*. (2004). *Univ Autonoma Madrid, Dept Bioquim, ES-28049 Madrid, Spain. Characterization of the aspartate kinase from *Saccharomyces cerevisiae* and of its interaction with threonine. *Biochem Biophys Res Commun* **321**, (3) 584
- McBryant SJ, Peersen OB*. (2004). *Colorado State Univ, Dept Biochem & Mol Biol, Fort Collins, Co 80523, USA. Self-association of the yeast nucleosome assembly protein 1. *Biochemistry* **43**, (32) 10592
- Nayar S, Brahma A, Barat B, Bhattacharyya D*. (2004). *Indian Institute Chem Biology, 4 Raja SC Mullick Rd, IN-700032 Calcutta, India. UDP-galactose 4-epimerase from *Kluyveromyces fragilis*: Analysis of its hysteretic behavior during catalysis. *Biochemistry* **43**, (31) 10212
- Ravanal MC, Flores M, Perez E, Aroca F, Cardemil E*. (2004). *Univ Santiago Chile, Fac Quim & Biol, Dept Ciencias Quim, Casilla 40, Santiago 33, Chile. *Saccharomyces cerevisiae* phosphoenolpyruvate carboxykinase: Relevance of arginine 70 for catalysis. *Biochimie* **86**, (6) 357
- Shinyashiki M, Pan CJG, Lopez BE, Fukuto JM*. (2004). *UCLA Sch Med, Dept Pharmacol, Hlth Sci Ctr, Los Angeles, Ca 90095, USA. Inhibition of the yeast metal reductase heme protein Fre1 by nitric oxide (NO): A model for inhibition of NADPH oxidase by NO. *Free Radic Biol Med* **37**, (5) 713
- Solovoyov A, Xiao RY, Gilbert HF*. (2004). *Baylor Coll Med, Verna & Marrs McLean Dept Biochem, 1 Baylor Plaza, Houston, Tx 77030, USA. Sulfhydryl oxidation, not disulfide isomerization, is the principal function of protein disulfide isomerase in yeast *Saccharomyces cerevisiae*. *J Biol Chem* **279**, (33) 34095
- Valencia E, Larroy C, Ochoa WF, Pares X, Fita I, Biosca JA*. (2004). *Univ Autonoma Barcelona, Dept Biochem & Mol Biol, ES-08193 Bellaterra, Barcelona, Spain. *Apo* and *Holo* structures of an NADP(H)-dependent cinnamyl alcohol dehydrogenase from *Saccharomyces cerevisiae*. *J Mol Biol* **341**, (4) 1049
- Weissman Z, Kornitzer D*. (2004). *Technion - Israel Inst Technol, Bruce Rappaport Fac Med, Dept Mol Microbiol, POB 9649, IL-31096 Haifa, Israel. A family of *Candida* cell surface haem-binding proteins involved in haemin and haemoglobin-iron utilization. *Mol Microbiol* **53**, (4) 1209
- Yamamoto K, Nakayama A, Yamamoto Y*, Tabata S. (2004). *Nara Med Univ, Dept Chem, Nara 634 8521, Japan. Val216 decides the substrate specificity of α -glucosidase in *Saccharomyces cerevisiae*. *Eur J Biochem* **271**, (16) 3414

4 Biotechnology

- Bednarski W, Adamczak M, Tomasik J, Plaszczyk M. (2004). Univ Warmia & Mazury Olsztyn, Chair Food Biotechnol, J Heweliusz St 1, PL-10718 Olsztyn, Poland. Application of oil refinery waste in the biosynthesis of glycolipids by yeast. *Bioresource Technol* **95**, (1) 15
- Campelo AF, Belo I*. (2004). *Univ Minho, Ctr Engn Biol, PT-4710-057 Braga, Portugal. Fermentative capacity of baker's yeast exposed to hyperbaric stress. *Biotechnol Lett* **26**, (15) 1237
- Cavazza A, Poznanski E, Trioli G. (2004). Ist Agrario San Michele All'Adige, Via E Mach 1, IT-38010 San Michele All'Adige, TN, Italy. Restart of fermentation of simulated stuck wines by direct inoculation of active dry yeasts. *Am J Enol Viticult* **55**, (2) 160
- Chtourou M, Ammar E*, Nasri M, Medhioub K. (2004). *Ecole Natl Ingn Sfax, LARSEN, UR Etude & Gestion Environm Urbains & Cotiers, BP W, TN-3038 Sfax, Tunisia. Isolation of a yeast, *Trichosporon cutaneum*, able to use low molecular weight phenolic compounds: Application to olive mill waste water treatment. *J Chem Technol Biotechnol* **79**, (8) 869
- Comitini F, Di Pietro N, Zacchi L, Mannazzu I, Ciani M*. (2004). *Univ Politecn Marche, Dipt Sci Alimenti, Via Breccia Bianche, IT-60131 Ancona, Italy. *Kluyveromyces phaffii* killer toxin active against wine spoilage yeasts: Purification and characterization. *Microbiology* **150**, (8) 2535
- De Maria PD, Xenakis A, Stamatis H, Sinisterra JV*. (2004). *Univ Complutense Madrid, Fac Pharm, Lab Biotransformat, ES-28040 Madrid, Spain. Lipase factor (LF) as a characterization parameter to explain the catalytic activity of crude lipases from *Candida rugosa*, free or immobilized in microemulsion-based organogels. *Enzyme Microb Technol* **35**, (4) 277
- Fernandez-Gonzalez M, Ubeda JF, Vasudevan TG, Otero RRC, Briones AI*. (2004). *Univ Castilla La Mancha, Fac Quim, Dept Quim Analit & Technol Alimentos, ES-13071 Ciudad Real, Spain. Evaluation of polygalacturonase activity in *Saccharomyces cerevisiae* wine strains. *FEMS Microbiol Lett* **237**, (2) 261
- An GH, Song JY, Chang KS, Lee BD, Chae HS, Jang BC. (2004). Chungnam Natl Univ, Dept Food Sci & Technol, 220 Gung-dong, Taejon 305 764, South Korea. Pigmentation and delayed oxidation of broiler chickens by the red carotenoid, astaxanthin, from chemical synthesis and the yeast, *Xanthophyllomyces dendrorhous*. *Asian Australas J Anim Sci* **17**, (9) 1309
- Gorgens J, Planas J, Van Zyl W, Knoetze J, Hahn-Hagerdal B*. (2004). *Lund Univ, Dept Appl Microbiol, Box 124, SE-22100 Lund, Sweden. Comparison of three expression systems for heterologous xylanase production by *S. cerevisiae* in defined medium. *Yeast* **21**, (14) 1205
- Gulbinienė G, Kondratienė L, Jokantaite T, Serviene E, Melvydas V, Petkuniene G*. (2004). *Inst Bot, Genet Lab, Zaliuju Ezeru 49, LT-08406 Vilnius, Lithuania. Occurrence of killer yeast strains in fruit and berry wine yeast populations. *Food Technol Biotechnol* **42**, (3) 159
- He H, Dong XP, Yang M, Yang QX, Duan SM, Yu YB, Han J, Zhang CB, Chen L, Yang X. (2004). Chinese Acad Sci, Res Ctr Eco-Environm Sci, POB 2871, 18 Shuangqing Rd, CN-100085 Beijing, Peoples Rep China. Catalytic inactivation of SARS coronavirus, *Escherichia coli* and yeast on solid surface. *Catal Commun* **5**, (3) 170

- Jin YS, Jeffries TW*. (2004). *USDA/US Forest Serv, Forest Prod Lab, 1 Gifford Pinchot Dr, Madison, WI 53705, USA. Stoichiometric network constraints on xylose metabolism by recombinant *Saccharomyces cerevisiae*. *Metab Eng* **6**, (3) 229
- Keating JD, Robinson J, Bothast RJ, Saddler JN, Mansfield SD*. (2004). *Univ British Columbia, Dept Wood Sci, 2424 Main Mall, Vancouver, Brit Columbia, Canada V6T 1Z4. Characterization of a unique ethanologenic yeast capable of fermenting galactose. *Enzyme Microb Technol* **35**, (2-3) 242
- Keating JD, Robinson J, Cotta MA, Saddler JN, Mansfield SD*. (2004). *Address as above. An ethanologenic yeast exhibiting unusual metabolism in the fermentation of lignocellulosic hexose sugars. *J Ind Microbiol Biotechnol* **31**, (5) 235
- Kim MD, Lee TH, Lim HK, Seo JH*. (2004). *Seoul Natl Univ, Dept Agr Biotechnol, Seoul 151 742, South Korea. Production of antithrombotic hirudin in *GALI*-disrupted *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol* **65**, (3) 259
- Kobayashi F, Nakamura Y*. (2004). *Kanazawa Univ, Fac Engr, Dept Chem & Chem Engr, 2-40-20 Kodatsuno, Ishikawa 920 866, Japan. Mathematical model of direct ethanol production from starch in immobilized recombinant yeast culture. *Biochem Eng J* **21**, (1) 93
- Lagos FM, Carballeira JD, Bermudez JL, Alvarez E, Sinisterra JV*. (2004). *Univ Complutense, Fac Pharm, Grp Biotransform, ES-28040 Madrid, Spain. Highly stereoselective reduction of haloketones using three new yeasts: Application to the synthesis of (*S*)-adrenergic β -blockers related to propranolol. *Tetrahedron Asymmetry* **15**, (5) 763
- Lotter J, Botes A, Van Dyk M, Breytenbach J. (2004). NW Univ, Dept Pharm Chem, PB X6001, ZA-2520 Potchefstroom, Rep Sth Africa. Correlation between the physicochemical properties of organic solvents and their biocompatibility toward epoxide hydrolase activity in whole-cells of a yeast, *Rhodotorula* sp. *Biotechnol Lett* **26**, (15) 1191
- Lotter J, Botes AL, Van Dyk MS, Breytenbach JC. (2004). NW Univ, Dept Pharmaceut Chem, Private Bag X6001, ZA-2520 Potchefstroom, Rep Sth Africa. Hydrolytic kinetic resolution of the enantiomers of the structural isomers *trans*-1-phenylpropene oxide and (2,3-epoxypropyl) benzene by yeast epoxide hydrolase. *Biotechnol Lett* **26**, (15) 1197
- Lou WY, Zong MH*, Zhang YY, Wu H. (2004). *Sth China Univ Technol, Dept Biotechnol, CN-510640 Guangzhou, Peoples Rep China. Efficient synthesis of optically active organosilyl alcohol via asymmetric reduction of acyl silane with immobilized yeast. *Enzyme Microb Technol* **35**, (2-3) 190
- Lozano P, Villora G, Gomez D, Gayo AB, Sanchez-Conesa JA, Rubio M, Iborra JL*. (2004). *Univ Murcia, Fac Chem, Dept Biochem & Mol Biol B & Immunol, POB 4021, Campus Espinardo, ES-30100 Murcia, Spain. Membrane reactor with immobilized *Candida antarctica* lipase B for ester synthesis in supercritical carbon dioxide. *J Supercrit Fluid* **29**, (1-2) 121
- Lu YN, Bellgrau D, Dwyer-Nield LD, Malkinson AM, Duke RC, Rodell TC, Franzusoff A*. (2004). *GlobeImmune Inc, 12635 East Montview Blvd, Aurora, Co 80010, USA. Mutation-selective tumor remission with Ras-targeted, whole yeast-based immunotherapy. *Cancer Res* **64**, (15) 5084
- Mateo C, Monti R, Pessela BCC, Fuentes M, Torres R, Guisan JM, Fernandez-Lafuente R*. (2004). *CSIC - Inst Catalis, Dept Biocatalis, Campus Univ Autonomia, ES-28049 Madrid, Spain. Immobilization of lactase from *Kluyveromyces lactis* greatly reduces the inhibition promoted by glucose. Full hydrolysis of lactose in milk. *Biotechnol Prog* **20**, (4) 1259
- McCabe RW, Taylor A*. (2004). Univ Cent Lancashire, Ctr Mat Sci, Preston PR1 2HE, England. Synthesis of novel polyurethane polyesters using the enzyme *Candida antarctica* lipase B. *Green Chem* **6**, (2) 151
- Molinari P, Pilosof AMR, Jagus RJ*. (2004). *Univ Buenos Aires, Fac Ingn, Dept Ingn Ind, Ciudad Univ, AR-1428 Buenos Aires, DF, Argentina. Effect of growth phase and inoculum size on the inactivation of *Saccharomyces cerevisiae* in fruit juices, by pulsed electric fields. *Food Res Int* **37**, (8) 793
- Moller K, Sharif MZ, Olsson L. (2004). Tech Univ Denmark, Ctr Microbial Biotechnol, Biocentrum DTU, Bldg 223, DK-2800 Lyngby, Denmark. Production of fungal α -amylase by *Saccharomyces kluyveri* in glucose-limited cultivations. *J Biotechnol* **111**, (3) 311
- Nishida O, Kuwazaki S, Suzuki C, Shima J*. (2004). *Natl Food Res Inst, 2-1-12 Kannondai, Tsukuba, Ibaraki 305 8642, Japan. Superior molasses assimilation, stress tolerance, and trehalose accumulation of baker's yeast isolated from dried sweet potatoes (hoshi-imo). *Biosci Biotechnol Biochem* **68**, (7) 1442
- Ovsejevi K, Grazu V, Cuadra K, Batista-Viera F. (2004). Fac Quim Montevideo, Gen Flores 2124, UY-CC1157 Montevideo, Uruguay. Enzyme reduction on solid phase as a tool for the reversible immobilization of yeast β -galactosidase onto a thiol-reactive support. *Enzyme Microb Technol* **35**, (2-3) 203
- Rahman MBA, Basri M, Hussein MZ, Rahman RNZA, Zainol DH, Salleh AB. (2004). Univ Pertanian Malaysia, Fac Sci & Environm Studies, Ctr Res Enzyme & Microbial Technol, MY-43400 Serdang, Selangor, Malaysia. Immobilization of lipase from *Candida rugosa* on layered double hydroxides for esterification reaction. *Appl Biochem Biotechnol* **118**, (1-3) 313
- Ramirez M, Vinagre A, Ambrona J, Molina F, Maqueda M, Robello JE. (2004). Univ Extremadura, Fac Ciencias, Dept Microbiol, Antiguo Rectorado, ES-06071 Badajoz, Spain. Genetic instability of heterozygous, hybrid, natural wine yeasts. *Appl Environ Microbiol* **70**, (8) 4686
- Reeleder RD. (2004). Agr & Agri-Food Canada, Sthn Crop Protect & Food Res Ctr, 1391 Sandford St, London, Ontario, Canada N5V 4T3. The use of yeasts for biological control of the plant pathogen *Sclerotinia sclerotiorum*. *Biocontrol* **49**, (5) 583
- Romero AM, Doval MM, Sturla MA, Judis MA*. (2004). *Univ Nacl Nordeste, Fac Agroind, Dept Tecnol, Comandante Fernandez 755, AR-3700 Chaco, Argentina. Antioxidant properties of polyphenol-containing extract from soybean fermented with *Saccharomyces cerevisiae*. *Eur J Lipid Sci Technol* **106**, (7) 424
- Shigechi H, Koh J, Fujita Y, Matsumoto T, Bito Y, Ueda M, Satoh E, Fukuda H, Kondo A*. (2004). *Kobe Univ, Dept Chem Sci & Engr, Fac Engr, 1-1 Rokkodaicho, Nada ku, Kobe, Hyogo 654 78501, Japan. Direct production of ethanol from raw corn starch via fermentation by use of a novel surface-engineered yeast strain codisplaying glucosylase and α -amylase. *Appl Environ Microbiol* **70**, (8) 5037
- Solaiman DKY, Ashby RD, Nunez A, Foglia TA. (2004). USDA/ARS, Fats Oils & Anim Coprod Res Unit, Eastern Reg Res Ctr, 600 East Mermaid Lane, Wyndmoor, Pa 19038, USA. Production of sophorolipids by *Candida bombicola* grown on soy molasses as substrate. *Biotechnol Lett* **26**, (15) 1241
- Teoh AL, Heard G, Cox J*. (2004). *Univ New Sth Wales, Sch Chem Sci, Sydney, NSW 2052, Australia. Yeast ecology of kombucha fermentation. *Int J Food Microbiol* **95**, (2) 119
- Tian SP, Qin GZ, Xu Y. (2004). Chinese Acad Sci, Inst Bot, Key Lab Photosynth & Environm Mol Physiol, CN-100093 Beijing, Peoples Rep China. Survival of antagonistic yeasts under field conditions and their biocontrol ability against postharvest diseases of sweet cherry. *Postharvest Biol Technol* **33**, (3) 327
- Toikkanen JH, Sundqvist L, Keranen S. (2004). VTT Biotechnol, PO Box 1500, Tietotie 2, FI-02044 VTT, Finland. *Kluyveromyces lactis* *SSO1* and *SEB1* genes are functional in *Saccharomyces cerevisiae* and enhance production of secreted proteins when overexpressed. *Yeast* **21**, (12) 1045
- Tsapatsaris S, Kotzekidou P*. (2004). *Aristotle Univ Thessaloniki, Fac Agr, Dept Food Sci & Technol, Lab Food Microbiol & Hyg, PB 250, GR-54124 Thessaloniki, Greece. Application of central composite design and response surface methodology to the fermentation of olive juice by *Lactobacillus plantarum* and *Debaryomyces hansenii*. *Int J Food Microbiol* **95**, (2) 157
- Tsukatani T, Ide S, Ukeda H, Matsumoto K. (2004). Fukuoka Ind Technol Ctr, Biotechnol & Food Res Inst, Kurume, Fukuoka 839 0861, Japan. Metabolism of benzoquinone by yeast cells and oxidative characteristics of corresponding hydroquinone: Application to highly sensitive measurement of yeast cell density by using benzoquinone and a chemiluminescent probe. *Biosci Biotechnol Biochem* **68**, (7) 1525
- Wongsumpanchai W, Anamart S, Laoteng K, Petsom A*. (2004). *Chulalongkorn Univ, Fac Sci, Program Biotechnol, TH-10330 Bangkok, Thailand. Elongation of C16:0 to C18:0 fatty acids in methylotrophic yeast *Hansenula polymorpha* CBS 1976 and fatty acid auxotrophic mutants. *FEMS Microbiol Lett* **237**, (2) 213

5 Cell biology

- Barale S, McCusker D, Arkowitz RA*. (2004). *Univ Nice, Fac Sci Parc Valrose, Inst Signaling Dev Biol & Canc, CNRS UMR 6543, FR-06108 Nice 2, France. The exchange factor Cdc24 is required for cell fusion during yeast mating. *Eukaryot Cell* **3**, (4) 1049
- De Groot PWJ, De Boer AD, Cunningham J, Dekker HL, De Jong L, Hellingwerf KJ, De Koster C, Klis FM. (2004). Univ Amsterdam, Swammerdam Inst Life Sci, Microbiol Lab, Nieuwe Achtergracht 166, NL-1018 WV Amsterdam, The Netherlands. Proteomic analysis of *Candida albicans* cell walls reveals covalently bound carbohydrate-active enzymes and adhesins. *Eukaryot Cell* **3**, (4) 955
- Gualtieri T, Ragni E, Mizzi L, Fascio U, Popolo L*. (2004). *Univ Milan, Dipt Sci Biomol & Biotechnol, Via Celoria 26, IT-20133 Milano, Italy. The cell wall sensor Wsc1p is involved in reorganization of actin cytoskeleton in response to hypo-osmotic shock in *Saccharomyces cerevisiae*. *Yeast* **21**, (13) 1107
- Guldener U, Koehler GJ, Haussmann C, Bacher A, Kricke J, Becher D, Hegemann JH*. (2004). *Univ Dusseldorf, DE-40225 Dusseldorf, Germany. Characterization of the *Saccharomyces cerevisiae* Foli1 protein: Starvation for C1 carrier induces pseudohyphal growth. *Mol Biol Cell* **15**, (8) 3811
- Harris RL, Silverman PA*. (2004). *Oklahoma Med Res Fdn, Program Mol Cell & Dev Biol, 825 NE 13th St, Oklahoma City, Ok 73104, USA. Tra proteins characteristic of F-like type IV secretion systems constitute an interaction group by yeast two-hybrid analysis. *J Bacteriol* **186**, (16) 5480
- Jeong HT, Oowatari Y, Abe M, Tanaka K, Matsuda H, Kawamukai M*. (2004). *Shimane Univ, Fac Life & Environm Sci, Dept Life Sci & Biotechnol, 1060 Nishikawatsu, Matsue, Shimane 690 8504, Japan. Interaction between a negative regulator (Msa2/Nrd1) and a positive regulator (Cpc2) of sexual differentiation in *Schizosaccharomyces pombe*. *Biosci Biotechnol Biochem* **68**, (7) 1621
- Kaneko A, Umeyama T, Hanaoka N, Monk BC, Uehara Y, Niimi M*. (2004). *National Institute Infect Dis, Dept Bioactive Mol, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162 8640, Japan. Tandem affinity purification of the *Candida albicans* septin protein complex. *Yeast* **21**, (12) 1025
- Kellermayer R, Szigeti R, Kellermayer M, Miseta A. (2004). Univ Pecs, Dept Med Genet & Child Dev Med, Szigeti 12, HU-7623 Pecs, Hungary. The intracellular dissipation of cytosolic calcium following glucose re-addition to carbohydrate depleted *Saccharomyces cerevisiae*. *FEBS Lett* **571**, (1-3) 55
- Kerscher S, Benit P, Abdrakhmanova A, Zwicker K, Rais I, Karas M, Rustin P, Brandt U. (2004). Univ Frankfurt, Fachbereich Med, Inst Biochem 1, DE-6000 Frankfurt, Germany. Processing of the 24 kDa subunit mitochondrial import signal is not required for assembly of functional complex I in *Yarrowia lipolytica*. *Eur J Biochem* **271**, (17) 3588
- Kettner K, Blomberg A, Rodel G*. (2004). *Dresden Univ Technol, Inst Genet, DE-01262 Dresden, Germany. *Schizosaccharomyces pombe* ER oxidoreductin-like proteins SpEro1a p and SpEro1b p. *Yeast* **21**, (12) 1035
- Kruppa M, Krom BP, Chauhan N, Bambach AV, Cihlar RL, Calderone RA*. (2004). *Georgetown Univ Med Ctr, Dept Microbiol & Immunol, Washington, DC 20057, USA. The two-component signal transduction protein Chk1p regulates quorum sensing in *Candida albicans*. *Eukaryot Cell* **3**, (4) 1062
- Lasorsa FM, Scarcia P, Erdmann R, Palmieri F, Rottensteiner H*, Palmieri L. (2004). *Ruhr Univ Bochum, Inst Physiol Chem, DE-44780 Bochum, Germany. The yeast peroxisomal adenine nucleotide transporter: Characterization of two transport modes and involvement in Δ pH formation across peroxisomal membranes. *Biochem J* **381**, (3) 581
- Leger-Silvestre I, Milkereit P, Ferreira-Cerca S, Saveanu C, Rousselle JC, Choismel V, Guinefoleau C, Gas N, Gleizes PE*. (2004). *CNRS, Lab Biol Mol Eucaryote, 118 Route Narbonne, FR-31062 Toulouse, France. The ribosomal protein Rps15p is required for nuclear exit of the 40S subunit precursors in yeast. *EMBO J* **23**, (12) 2336
- Li LT, Kaplan J*. (2004). *Univ Utah Sch Med, Dept Pathol, Div Cell Biol & Immunol, Salt Lake City, Ut 84132, USA. A mitochondrial-vacuolar signaling pathway in yeast that affects iron and copper metabolism. *J Biol Chem* **279**, (32) 33653
- Li WM, Wang ZX, Jia SR*. (2004). *Chinese Acad Agr Sci, Biotechnol Res Inst, CN-100081 Beijing, Peoples Rep China. Effect of *GbkTNI* from *Gossypium barbadense* on cell elongation of fission yeast (*Schizosaccharomyces pombe*). *Chin Sci Bull* **49**, (1) 23
- Magherini F, Gamberi T, Paoli P, Marchetta M, Biagini M, Raugei G, Camici G, Ramponi G, Modesti A*. (2004). *Univ Florence, Dipt Sci Biochim, Florence, Italy. The *in vivo* tyrosine phosphorylation level of yeast immunophilin Fpr3 is influenced by the LMW-PTP Ltp1. *Biochem Biophys Res Commun* **321**, (2) 424
- Matsuo Y, Matsuura Y, Tanaka K, Matsuda H, Kawamukai M*. (2004). *Shimane Univ, Fac Life & Environm Sci, Dept Appl Biosci & Biotechnol, 1060 Nishikawatsu, Matsue 690 8504, Japan. Chr4, a *Schizosaccharomyces pombe* homologue of the *Saccharomyces cerevisiae* Chs4p/Skt5p protein, is related to septum formation and is required for the proper localization of Chs2. *Yeast* **21**, (12) 1005
- Mitra P, Zhang YJ, Rameh LE, Ivshina MP, McCollum D, Nunnari JJ, Hendricks GM, Kerr ML, Field SJ, Cantley LC, Ross AH*. (2004). *Univ Massachusetts Sch Med, Dept Biochem & Mol Pharmacol, 394 Plantation St, Rm 819, Worcester, Ma 01605, USA. A novel phosphatidylinositol(3,4,5)P₃ pathway in fission yeast. *J Cell Biol* **166**, (2) 205
- Miwa T, Takagi Y, Shinozaki M, Yun CW, Schell WA, Perfect JR, Kumagai H, Tamaki H*. (2004). *Kyoto Univ, Grad Sch Biostudies, Div Integrated Life Sci, Sakyo ku, Kyoto 606 8502, Japan. Gpr1, a putative G-protein-coupled receptor, regulates morphogenesis and hypha formation in the pathogenic fungus *Candida albicans*. *Eukaryot Cell* **3**, (4) 919
- Parrish WR, Stefan CJ, Emr SD*. (2004). *Univ Calif San Diego Sch Med, Dept Cellular & Mol Med, La Jolla, Ca 92093, USA. Essential role for the myotubularin-related phosphatase Ymrlp and the synaptotagmin-like phosphatases Sjl2p and Sjl3p in regulation of phosphatidylinositol 3-phosphate in yeast. *Mol Biol Cell* **15**, (8) 3567
- Roggero R, Zufferey R, Minca M, Richier E, Calas M, Vial H, Ben Mamoun C*. (2004). *Univ Connecticut Hlth Ctr, Dept Genet & Dev Biol, Ctr Microbial Pathogenesis, 263 Farmington Ave, Farmington, Ct 06030, USA. Unraveling the mode of action of the antimalarial choline analog G25 in *Plasmodium falciparum* and *Saccharomyces cerevisiae*. *Antimicrob Agents Chemother* **48**, (8) 2816
- Sato T, Ueno Y, Watanabe T, Mikami T, Matsumoto T*. (2004). *Tohoku Pharmaceut Univ, Dept Microbiol, Sendai, Miyagi 981 8558, Japan. Role of Ca²⁺ calmodulin signaling pathway on morphological development of *Candida albicans*. *Biol Pharm Bull* **27**, (8) 1281
- Schneider R, Brugger B, Amann CM, Prestwich GD, Epan RF, Zellnig G, Wieland FT, Epan RM. (2004). Univ Fribourg, Dept Med, Div Biochem, Chemin Musee 5, CH-1700 Fribourg, Switzerland. Identification and biophysical characterization of a very-long-chain-fatty-acid-substituted phosphatidylinositol in yeast subcellular membranes. *Biochem J* **381**, (3) 941
- Shiflett SL, Vaughn MB, Huynh D, Kaplan J, Ward DM*. (2004). *Univ Utah Hlth Sci Ctr, Dept Pathol, Div Cell Biol & Immunol, Salt Lake City, Ut 84132, USA. Bph1p, the *Saccharomyces cerevisiae* homologue of CHS1/Beige, functions in cell wall formation and protein sorting. *Traffic* **5**, (9) 700
- Suzuki K, Noda T, Ohsumi Y*. (2004). *Nat Inst Basic Biol, Dept Cell Biol, Nishigonaka 38, Myodaiji-cho, Okazaki 444 8585, Japan. Interrelationships among Atg proteins during autophagy in *Saccharomyces cerevisiae*. *Yeast* **21**, (12) 1057
- Suzuki M, Asada Y, Watanabe D, Ohya Y*. (2004). *Univ Tokyo, Grad Sch Frontier Sci, Dept Integrat Biosci, Bldg FSB-101, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277 8562, Japan. Cell shape and growth of budding yeast cells in restrictive microenvironments. *Yeast* **21**, (12) 983
- Vink E, Rodriguez-Suarez RJ, Gerard-Vincent M, Ribas JC, De Nobel H, Van den Ende H, Duran A, Klis FM, Bussey H*. (2004). *McGill Univ, Dept Biol, 1205 Ave Dr Penfield, Montreal, Quebec, Canada H3A 1B1. An *in vitro* assay for (1 → 6)-β-D-glucan synthesis in *Saccharomyces cerevisiae*. *Yeast* **21**, (13) 1121

- Walters DE, Kaplan RS. (2004). Rosalind Franklin Univ Med & Sci, Dept Biochem & Mol Biol, 3333 Green Bay Rd, Nth Chicago, IL 60064, USA. Homology-modeled structure of the yeast mitochondrial citrate transport protein. *Biophys J* **87**, (2) 907
- Wang P, Cutler J, King J, Palmer D. (2004). Childrens Hosp, Res Inst Children, Dept Pediat, 200 Henry Clay Ave, New Orleans, La 70118, USA. Mutation of the regulator of G protein signaling Crg1 increases virulence in *Cryptococcus neoformans*. *Eukaryot Cell* **3**, (4) 1028
- Wicky S, Schwarz H, Singer-Kruger B*. (2004). *Univ Stuttgart, Inst Biochem, Pfaffenwaldring 55, DE-70569 Stuttgart, Germany. Molecular interactions of yeast Neo1p, an essential member of the Drs2 family of aminophospholipid translocases, and its role in membrane trafficking within the endomembrane system. *Mol Cell Biol* **24**, (17) 7402
- ## 6 Gene expression
- Abruzzi KC, Lacadie S, Rosbash M*. (2004). *Brandeis Univ, Howard Hughes Med Inst, Dept Biol, MS008, 415 South St, Waltham, Ma 02454, USA. Biochemical analysis of TREX complex recruitment to intronless and intron-containing yeast genes. *EMBO J* **23**, (13) 2620
- Brand A, MacCallum DM, Brown AJP, Gow NAR, Odds FC*. (2004). *Inst Med Sci, Sch Med Sci, Aberdeen Fungal Grp, Aberdeen AB25 2ZD, Scotland. Ectopic expression of *URA3* can influence the virulence phenotypes and proteome of *Candida albicans* but can be overcome by targeted reintegration of *URA3* at the *RPS10* locus. *Eukaryot Cell* **3**, (4) 900
- Bulmer R, Pic-Taylor A, Whitehall SK, Martin KA, Millar JBA, Quinn J, Morgan BA*. (2004). *Univ Newcastle upon Tyne, Fac Med Sci, Inst Cell & Mol Biosci, Newcastle upon Tyne NE2 4HH, England. The forkhead transcription factor Fkh2 regulates the cell division cycle of *Schizosaccharomyces pombe*. *Eukaryot Cell* **3**, (4) 944
- Bungard D, Reed M, Winter E*. (2004). *Thomas Jefferson Univ, Dept Biochem & Mol Pharmacol, 233 Sth 10th St, Rm 228, Philadelphia, Pa 19107, USA. *RSC1* and *RSC2* are required for expression of mid-late sporulation-specific genes in *Saccharomyces cerevisiae*. *Eukaryot Cell* **3**, (4) 910
- Carvin CD, Kladden MP*. (2004). *Texas A&M Univ, Dept Biochem & Biophys, TAMU 2128, College Station, Tx 77843, USA. Effectors of lysine 4 methylation of histone H3 in *Saccharomyces cerevisiae* are negative regulators of *PHO5* and *GAL1-10*. *J Biol Chem* **279**, (32) 33057
- Chen LY, Widom J*. (2004). *Northwestern Univ, Dept Biochem Mol Biol & Cell Biol, 2153 Sheridan Rd, Evanston, IL 60208, USA. Molecular basis of transcriptional silencing in budding yeast. *Biochem Cell Biol* **82**, (4) 413
- Chung WH, Kim KD, Cho YJ, Roe JH*. (2004). *Seoul Natl Univ, Sch Biol Sci, Inst Microbiol, Mol Microbiol Lab, Seoul 151 742, South Korea. Differential expression and role of two dithiol glutaredoxins Grx1 and Grx2 in *Schizosaccharomyces pombe*. *Biochem Biophys Res Commun* **321**, (4) 922
- De Felipe KS, Carter BT, Althoff EA, Cornish VW*. (2004). *Columbia Univ, Dept Chem, New York, NY 10027, USA. Correlation between ligand-receptor affinity and the transcription readout in a yeast three-hybrid system. *Biochemistry* **43**, (32) 10353
- Doherty K, Pramanik A, Pride L, Lukose J, Wood Moore C*. (2004). *City Univ New York Med Sch & Sophie Davis Sch Biomed Educat, Dept Microbiol & Immunol, Sci Bldg & Harris Hall, Convent Ave 138th St, New York, NY 10031, USA. Expression of the expanded *YFL007w* ORF and assignment of the gene name *BLM10*. *Yeast* **21**, (12) 1021
- Evangelisti AM, Wagner A*. (2004). *Univ New Mexico, Dept Biol, 167 Castetter Hall, Albuquerque, NM 87131, USA. Molecular evolution in the yeast transcriptional regulation network. *J Exp Zool* **302B**, (4) 392
- Fuglsang A. (2004). Danish Univ Pharmaceut Sci, Inst Pharmacol, Universitetspk 2, DK-2100 Copenhagen O, Denmark. Bioinformatic analysis of the link between gene composition and expressivity in *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*. *Antonie van Leeuwenhoek* **86**, (2) 135
- Garcia-Martinez J, Aranda A, Perez-Ortin JE*. (2004). *Univ Valencia, Dept Bioquim & Biol Mol, ES-46100 Burjassot, Spain. Genomic run-on evaluates transcription rates for all yeast genes and identifies gene regulatory mechanisms. *Mol Cell* **15**, (2) 303
- Gunji L, Kai T, Takahashi Y, Maki Y, Kurihara W, Utsugi T, Fujimori F, Murakami Y*. (2004). *Sci Univ Tokyo, Fac Ind Sci & Technol, Dept Biol Sci & Technol, 2641 Yamazaki, Noda, Chiba 278 8510, Japan. Global analysis of the regulatory network structure of gene expression in *Saccharomyces cerevisiae*. *DNA Res* **11**, (3) 163
- Ionescu CN, Origanti S, McAlear MA*. (2004). *Wesleyan Univ, Mol Biol & Biochem Dept, Middletown, Ct 06459, USA. The yeast rRNA biosynthesis factor Ebp2p is also required for efficient nuclear division. *Yeast* **21**, (14) 1219
- Izawa S, Takemura R, Inoue Y. (2004). Kyoto Univ, Grad Sch Agr, Mol Microbiol Lab, Kyoto 611 0011, Japan. Gle2p is essential to induce adaptation of the export of bulk poly(A)⁺ mRNA to heat shock in *Saccharomyces cerevisiae*. *J Biol Chem* **279**, (34) 35469
- Karababa M, Coste AT, Rognon B, Bille J, Sanglard D*. (2004). *CHU Vaudois, Inst Microbiol, rue Bugnon 48, CH-1011 Lausanne, Switzerland. Comparison of gene expression profiles of *Candida albicans* azole-resistant clinical isolates and laboratory strains exposed to drugs inducing multidrug transporters. *Antimicrob Agents Chemother* **48**, (8) 3064
- Kim J, Jeon S, Yang YS, Kim J. (2004). Chungnam Natl Univ, Sch Biosci & Biotechnol, Dept Microbiol, Taejon 305 764, South Korea. Posttranscriptional regulation of the karyogamy gene by Kem1p/Xrm1p exoribonuclease and Rok1p RNA helicase of *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun* **321**, (4) 1032
- Kolaczowski M, Kolaczowska A, Gaigg B, Schneider R, Moyer-Rowley WS*. (2004). *Univ Iowa, Dept Physiol & Biophys, 6-530 Bowen Sci Bldg, Iowa City, Ia 52242, USA. Differential regulation of ceramide synthase components *LAC1* and *LAG1* in *Saccharomyces cerevisiae*. *Eukaryot Cell* **3**, (4) 880
- Korber P, Horz W*. (2004). *Univ Munich, Adolf Butenandt Inst, Schillerstr 44, DE-80336 Munich, Germany. *In vitro* assembly of the characteristic chromatin organization at the yeast *PHO5* promoter by a replication-independent extract system. *J Biol Chem* **279**, (33) 35113
- Li X, Zaman S, Langdon Y, Zengel JM, Lindahl L*. (2004). *UMBC, Dept Biol Sci, 1000 Hilltop Circle, Baltimore, Md 21250, USA. Identification of a functional core in the RNA component of RNase MRP of budding yeasts. *Nucleic Acids Res* **32**, (12) 3703
- Luo G, Samaranyake LP*, Cheung BPK, Tang G. (2004). *Univ Hong Kong, Fac Dent, 34 Hosp Rd, Hong Kong, Peoples Rep China. Reverse transcriptase polymerase chain reaction (RT-PCR) detection of *HLP* gene expression in *Candida glabrata* and its possible role in *in vitro* haemolysin production. *APMIS* **112**, (4-5) 283
- Malys N, Carroll K, Miyayama J, Tollervy D, McCarthy JEG*. (2004). *UMIST, Dept Biomol Sci, Post-Transcript Control Grp, Manchester M60 1QD, England. The 'scavenger' m⁷GpppX pyrophosphatase activity of Dcs1 modulates nutrient-induced responses in yeast. *Nucleic Acids Res* **32**, (12) 3590
- Martin AM, Pouchnik DJ, Walker JL, Wyrick JJ*. (2004). *Washington State Univ, Sch Mol Biosci, Fulmer Hall 675, Pullman, Wa 99164, USA. Redundant for histone H3 N-terminal lysine residues in subtelomeric gene repression in *Saccharomyces cerevisiae*. *Genetics* **167**, (3) 1123
- Morlando M, Ballarino M, Greco P, Caffarelli E, Dichtl B, Bozzoni I*. (2004). *Univ Roma la Sapienza, Inst Pasteur Cenci Bolognietti, Dept Genet & Mol Biol, Piazzale Aldo Moro 5, IT-00185 Rome, Italy. Coupling between snoRNP assembly and 3' processing controls box C/D snoRNA biosynthesis in yeast. *EMBO J* **23**, (12) 2392
- Noble CG, Walker PA, Calder LJ, Taylor IA*. (2004). *Natl Inst Med Res, Div Prot Struct, Mill Hill, London NW7 1AA, England. Rna14-Rna15 assembly mediates the RNA-binding capability of *Saccharomyces cerevisiae* cleavage factor IA. *Nucleic Acids Res* **32**, (11) 3364
- Okazaki H, Ono B, Ohsumi Y*, Ohsumi M. (2004). *Teikyo Univ Sci & Technol, Dept Biosci, Yamanashi 409 0193, Japan. *apg15-1*, a UGA mutant allele in the *Saccharomyces cerevisiae* *APG16* gene, and its suppression by a cytoplasmic factor. *Biosci Biotechnol Biochem* **68**, (7) 1541

- Pinson B, Merle M, Franconi JM, Daignan-Fornier B. (2004). Univ Bordeaux 2, CNRS UMR 5095, Inst Biochim & Genet Cellulaires, 1 rue Camille St Saens, FR-33077 Bordeaux, France. Low affinity orthophosphate carriers regulate *PHO* gene expression independently of internal orthophosphate concentration in *Saccharomyces cerevisiae*. *J Biol Chem* **279**, (34) 35273
- Rubin-Bejerano I, Sagee S, Friedman O, Pnueli L, Kassir Y*. (2004). *Technion Israel Inst Technol, Dept Biol, IL-32000 Haifa, Israel. The *in vivo* activity of Ime1, the key transcriptional activator of meiosis-specific genes in *Saccharomyces cerevisiae*, is inhibited by the cyclic AMP protein kinase A signal pathway through the glycogen synthase kinase 3- β homolog Rim11. *Mol Cell Biol* **24**, (16) 6967
- Rustici G, Mata J, Kivinen K, Lio P, Penkett CJ, Burns G, Hayles J, Brazma A, Nurse P, Bahler J*. (2004). *Wellcome Trust Sanger Inst, Cambridge CB10 1SA, England. Periodic gene expression program of the fission yeast cell cycle. *Nat Genet* **36**, (8) 809
- Ryabinkova NA, Sopova YV, Polozkov GV, Savelova MV, Ryabinkova NA. (2004). St Petersburg State Univ, Dept Genet & Breeding, RU-119034 St Petersburg, Russia. Frameshift suppression through inactivation of translation termination in yeast *Saccharomyces cerevisiae*: Significance of the local context. *Russ J Genet* **40**, (7) 717
- Ryan DE, Kim CH, Murray JB, Adams CJ, Stockley PG, Abelson J*. (2004). *UCSF Mission Bay, Box 2200, 600 16th St, San Francisco, Ca 94143, USA. New tertiary constraints between the RNA components of active yeast spliceosomes: A photo-crosslinking study. *RNA* **10**, (8) 1251
- Suntharalingam M, Alcazar-Roman A, Wente SR*. (2004). *Vanderbilt Univ Med Ctr, Dept Cell & Dev Biol, 3120 MRBIII, 465 21st Ave Sth, Nashville, Tn 37232, USA. Nuclear export of the yeast mRNA-binding protein Nab2 is linked to a direct interaction with Gfd1 and to Gle1 function. *J Biol Chem* **279**, (34) 35384
- Tanny JC, Kirkpatrick DS, Gerber SA, Gygi SP, Moazed D*. (2004). *Harvard Univ Sch Med, Dept Cell Biol, 240 Longwood Ave, Boston, Ma 02115, USA. Budding yeast silencing complexes and regulation of Sir2 activity by protein-protein interactions. *Mol Cell Biol* **24**, (16) 6931
- Walker SC, Avis JM*. (2004). *UMIST, Dept Biomol Sci, POB 88, Manchester M60 1QD, England. A conserved element in the yeast RNase MRP PNA subunit can participate in a long-range base-pairing interaction. *J Mol Biol* **341**, (2) 375
- Wu PYJ, Ruhlmann C, Winston F, Schultz P*. (2004). *CNRS/INSERM/ULP, Inst Genet & Biol Mol & Cellulaire, 1 rue Laurent Fries, BP 163, FR-67404 Illkirch Graffenstaden, France. Molecular architecture of the *S. cerevisiae* SAGA complex. *Mol Cell* **15**, (2) 199
- Yin ZK, Stead D, Selway L, Walker J, Riba-Garcia I, McInerney T, Gaskell S, Oliver SG, Cash P, Brown AJP*. (2004). *Univ Aberdeen, Inst Med Sci, Aberdeen Proteome Facil, Aberdeen AB25 2ZD, Scotland. Proteomic response to amino acid starvation in *Candida albicans* and *Saccharomyces cerevisiae*. *Proteomics* **4**, (8) 2425
- Yoon CN, Han SK, Kim HY. (2004). Chungbuk Natl Univ, Basic Res Inst, Cheongju 361 763, South Korea. Linear network model of gene regulation for the yeast cell cycle. *J Korean Phys Soc* **44**, (3) 638
- 7 Genetics & genomics**
- Andalis AA, Storchova Z, Styles C, Galitski T*, Pellman D, Fink GR. (2004). *MIT, Whitehead Inst Biomed Res, 9 Cambridge Ctr, Cambridge, Ma 02142, USA. Defects arising from whole-genome duplications in *Saccharomyces cerevisiae*. *Genetics* **167**, (3) 1109
- Andreishcheva EN, Kunkel JP, Gemmill TR, Trimble RB*. (2004). *New York State Dept Hlth, Wadsworth Ctr Labs & Res, POB 509, Albany, NY 12201, USA. Five genes involved in biosynthesis of the pyruvylated Gal β 1,3-epitope in *Schizosaccharomyces pombe* N-linked glycans. *J Biol Chem* **279**, (34) 35644
- Ben-Aroya S, Mieczkowski PA, Petes TD, Kupiec M*. (2004). *Tel Aviv Univ, Dept Mol Microbiol & Biotechnol, IL-69978 Tel Aviv, Israel. The compact chromatin structure of a Ty repeated sequence suppresses recombination hotspot activity in *Saccharomyces cerevisiae*. *Mol Cell* **15**, (2) 221
- Bhattacharyya S, Lahue RS*. (2004). *Univ Nebraska Med Ctr, Eppley Inst Res Canc & Allied Dis, Box 986805, Omaha, Ne 68198, USA. *Saccharomyces cerevisiae* Srs2 DNA helicase selectively blocks expansions of trinucleotide repeats. *Mol Cell Biol* **24**, (17) 7324
- Boer E, Wartmann T, Luther B, Manteuffel R, Bode R, Gellissen G, Kunze G*. (2004). *Inst Pflanzengen et & Kulturpflanzenforsch, Corrensstr 3, DE-06466 Gatersleben, Germany. Characterization of the *AINV* gene and the encoded invertase from the dimorphic yeast *Arxula adenivorans*. *Antonie van Leeuwenhoek* **86**, (2) 121
- Capiaghi C, Ho TV, Thoma F*. (2004). *ETH Honggerberg, Inst Zellbiol, CH-8093 Zurich, Switzerland. Kinetochores prevent repair of UV damage in *Saccharomyces cerevisiae* centromeres. *Mol Cell Biol* **24**, (16) 6907
- Chappell AS, Lundblad V*. (2004). *Baylor Coll Med, Dept Mol & Human Genet, Houston, Tx 77030, USA. Structural elements required for association of the *Saccharomyces cerevisiae* telomerase RNA with the Est2 reverse transcriptase. *Mol Cell Biol* **24**, (17) 7720
- Chikashige Y, Kurokawa R, Haraguchi T, Hiraoka Y*. (2004). *Natl Inst Informat & Commun Technol, Kansai Adv Res Ctr, Cell Biol Grp, Nishi ku, Kobe, Hyogo 651 2492, Japan. Meiosis induced by inactivation of Pat1 kinase proceeds with aberrant nuclear positioning of centromeres in the fission yeast *Schizosaccharomyces pombe*. *Genes Cells* **9**, (8) 671
- Ciudad T, Andaluz E, Steinberg-Neifach O, Lue NF, Gow NAR, Calderone RA, Larriba G*. (2004). *Univ Extremadura, Fac Ciencias, Dept Microbiol, ES-06071 Badajoz, Spain. Homologous recombination in *Candida albicans*: Role of CaRad52p in DNA repair, integration of linear DNA fragments and telomere length. *Mol Microbiol* **53**, (4) 1177
- Douville J, David J, Fortier PK, Ramotar D*. (2004). *Univ Montreal, Hop Maisonneuve Rosemont, Ctr Rech, 5415 Blvd Assomption, Montreal, Quebec, Canada H1J 2M4. The yeast phosphotyrosyl phosphatase activator protein, yPtp1/Rrd1, interacts with Sit4 phosphatase to mediate resistance to 4-nitroquinoline-1-oxide and UVA. *Curr Genet* **46**, (2) 72
- Eason RG, Pourmand N, Tongprasit W, Herman ZS, Anthony K, Jejelowo O, Davis RW, Stolc V*. (2004). *NASA, Ames Res Ctr, Ctr Nanotechnol, Mail Stop 229-3, Moffett Field, Ca 94035, USA. Characterization of synthetic DNA bar codes in *Saccharomyces cerevisiae* gene-deletion strains. *Proc Natl Acad Sci U S A* **101**, (30) 11046
- Eirich LD, Craft DL, Steinberg L, Asif A, Eschenfeldt WH, Stols L, Donnelly MI, Wilson CR. (2004). Cognis Corp, Res & Technol Biotechnol Grp, Bldg 53, 4900 Este Ave, Cincinnati, Oh 45232, USA. Cloning and characterization of three fatty alcohol oxidase genes from *Candida tropicalis* strain ATCC 20336. *Appl Environ Microbiol* **70**, (8) 4872
- Fraschini R, Bilotta D, Lucchini G, Piatti S*. (2004). *Univ Milan Biococca, Dipt Biotechnol & Biosci, Piazza Sci 2, IT-20126 Milan, Italy. Functional characterization of Dma1 and Dma2, the budding yeast homologues of *Schizosaccharomyces pombe* Dma1 and human Chfr. *Mol Biol Cell* **15**, (8) 3796
- Friberg M, Von Rohr P, Gonnet G. (2004). Inst Computat Sci, ETH-Zentrum, HRS H30, CH-8092 Zurich, Switzerland. Limitations of codon adaptation index and other coding DNA-based features for prediction of protein expression in *Saccharomyces cerevisiae*. *Yeast* **21**, (13) 1083
- Gallardo-Moreno AM, Gonzalez-Martin ML*, Perez-Giraldo C, Bruque JM, Gomez-Garcia AC. (2004). *Univ Extremadura, Dept Phys, ES-06071 Badajoz, Spain. The measurement temperature: An important factor relating physicochemical and adhesive properties of yeast cells to biomaterials. *J Colloid Interface Sci* **271**, (2) 351
- Ghosh SK, Sau S, Lahiri S, Lohia A, Sinha P*. (2004). *Bose Inst, Dept Biochem, IN-700054 Calcutta, India. The Iml3 protein of the budding yeast is required for the prevention of precocious sister chromatid separation in meiosis I and for sister chromatid disjunction in meiosis II. *Curr Genet* **46**, (2) 82
- Hughes AL, Friedman R. (2004). Univ Sth Carolina, Dept Biol Sci, Coker Life Sci Bldg, 700 Sumter St, Columbia, SC 29205, USA. Transposable element distribution in the yeast genome reflects a role in repeated genomic rearrangement events on an evolutionary time scale. *Genetica* **121**, (2) 181

- Ishigami M, Nakagawa Y, Hayakawa M, Imura Y*. (2004). *Yamanashi Univ, Div Biotechnol, Interdisciplinary Grad Sch Med & Engn, 4-3-11 Takeda, Kofu, Yamanashi 400 85, Japan. *FLO11* is essential for flor formation caused by the C-terminal deletion of *NRG1* in *Saccharomyces cerevisiae*. *FEMS Microbiol Lett* **237**, (2) 425
- Itoh T, Toh-e A, Matsui Y*. (2004). *Univ Tokyo, Grad Sch Sci, Dept Biol Sci, Tokyo 113 0033, Japan. Mmr1p is a mitochondrial factor for Myo2p-dependent inheritance of mitochondria in the budding yeast. *EMBO J* **23**, (13) 2520
- Khodjakov A, La Terra S, Chang F. (2004). Wadsworth Ctr, New York State Dept Hlth, POB 509, Albany, NY 12201, USA. Laser microsurgery in fission yeast: Role of the mitotic spindle midzone in anaphase B. *Curr Biol* **14**, (15) 1330
- Kimura Y, Koitabashi S, Kakizuka A, Fujita T. (2004). Tokyo Metropolitan Inst Med Sci, Lab Frontier Sci, 3-18-22 Honkomagome, Bunkyo ku, Tokyo 113 8613, Japan. The role of pre-existing aggregates in Hsp104-dependent polyglutamine aggregate formation and epigenetic change of yeast prions. *Genes Cells* **9**, (8) 685
- Kohler F, Zimmermann A, Hager M, Sippel AE. (2004). Univ Freiburg, Max Planck Inst Immunobiol, Inst Biol Mol Immunol III, Stuebweg 51, DE-79108 Freiburg, Germany. A genetic, non-transcriptional assay for nuclear receptor ligand binding in yeast. *Gene* **337**, 113
- Lamas-Maceiras M, Cerdan ME, Lloret A, Freire-Picos MA*. (2004). *Univ Coruna, Fac Ciencias, Dip Biol Celular & Mol, Campus A Zapateiro s/n, ES-15071 A Coruna, Spain. Characterization of a gene similar to *BIK1* in the yeast *Kluyveromyces lactis*. *Yeast* **21**, (13) 1067
- Maciaszczyk E, Ulaszewski S, Lazowska J*. (2004). *CNRS, Ctr Genet Mol, FR-91198 Gif-sur-Yvette, France. Intragenic suppressors that restore the activity of the maturase encoded by the second intron of the *Saccharomyces cerevisiae* *cyt b* gene. *Curr Genet* **46**, (2) 67
- Maloisel L, Bhargava J, Roeder GS*. (2004). *Yale Univ, Howard Hughes Med Inst, Dept Mol Cellular & Dev Biol, POB 208103, New Haven, Ct 06520, USA. A role for DNA polymerase in gene conversion and crossing over during meiosis in *Saccharomyces cerevisiae*. *Genetics* **167**, (3) 1133
- Masciadri B, Areces LB, Carpinelli P, Foiani M, Draetta GF*. Fiore F. (2004). *European Inst Oncol, Dept Expt Oncol, 435 Via Ripamonti, IT-20141 Milan, Italy. Characterization of the *BUD31* gene of *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun* **320**, (4) 1342
- Matsuo Y, Tanaka K, Nakagawa T, Matsuda H, Kawamukai M*. (2004). *Shimane Univ, Fac Life & Environm Sci, Dept Appl Biosci & Biotechnol, 1060 Nishikawatsu, Matsue, Shimane 690 8504, Japan. Genetic analysis of *chs1+* and *chs2+* encoding chitin synthases from *Schizosaccharomyces pombe*. *Biosci Biotechnol Biochem* **68**, (7) 1489
- Melloy PG, Holloway SL. (2004). Princeton Univ, Dept Mol Biol, Lewis Thomas Lab, Princeton, NJ 08544, USA. Changes in the localization of the *Saccharomyces cerevisiae* anaphase-promoting complex upon microtubule depolymerization and spindle checkpoint activation. *Genetics* **167**, (3) 1079
- Melo RGM, Leitao AC, Padula M*. (2004). *Univ Fed Rio de Janeiro, Fac Farm, Dept Farm, Lab Microbiol Ind & Avaliacao Genotoxica, Sala 15, BR-21941-970 Rio de Janeiro, Brazil. Role of *OGG1* and *NTG2* in the repair of oxidative DNA damage and mutagenesis induced by hydrogen peroxide in *Saccharomyces cerevisiae*: Relationships with transition metals iron and copper. *Yeast* **21**, (12) 991
- Miyake T, Reese J, Loch CM, Auble DT, Li R*. (2004). *Univ Virginia Sch Med, Dept Biochem & Mol Genet, POB 800733, Charlottesville, Va 22908, USA. Genome-wide analysis of ARS (autonomously replicating sequence) binding factor 1 (Abf1p)-mediated transcriptional regulation in *Saccharomyces cerevisiae*. *J Biol Chem* **279**, (33) 34865
- Nishida Y, Katoh H, Andoh M, Ono B*. (2004). *Ritsumeikan Univ, Fac Sci & Engn, Dept Biotechnol, 1-1-1 Noji-higashi, Kusatsu 525 8577, Shiga, Japan. Mating-induced mating-type cassette conversion in *Saccharomyces cerevisiae*. *Yeast* **21**, (13) 1095
- Park YN, Strauss A, Morschhauser J*. (2004). *Univ Wurzburg, Inst Mol Infekt Biol, Rontgenring 11, DE-97070 Wurzburg, Germany. The white-phase-specific gene *WH11* is not required for white-opaque switching in *Candida albicans*. *Mol Genet Genomics* **272**, (1) 88
- Pidoux AL, Allshire RC. (2004). Univ Edinburgh, Wellcome Trust Ctr Cell Biol, Kings Bldg, Mayfield Rd, Edinburgh EH9 3JR, Scotland. Kinetochore and heterochromatin domains of the fission yeast centromere. *Chromosome Res* **12**, (6) 521
- Pujol C, Daniels KJ, Lockhart SR, Srikantha T, Radke JB, Geiger J, Soll DR*. (2004). *Univ Iowa, Dept Sci Biol, 302 BBE, Iowa City, Ia 52242, USA. The closely related species *Candida albicans* and *Candida dubliniensis* can mate. *Eukaryot Cell* **3**, (4) 1015
- Ruiz A, Munoz I, Serrano R, Gonzalez A, Simon E, Arino J*. (2004). *Univ Autonoma Barcelona, Fac Vet, Dept Bioquim & Biol Mol, Ed 5, ES-08193 Barcelona, Spain. Functional characterization of the *Saccharomyces cerevisiae* *VHS3* gene - A regulatory subunit of the Ppz1 protein phosphatase with novel, phosphatase-unrelated functions. *J Biol Chem* **279**, (33) 34421
- Salmon TB, Evert BA, Song BW, Doetsch PW*. (2004). *Emory Univ Sch Med, Dept Biochem, Atlanta, Ga 30322, USA. Biological consequences of oxidative stress-induced DNA damage in *Saccharomyces cerevisiae*. *Nucleic Acids Res* **32**, (12) 3712
- Sanyal K, Baum M, Carbon J*. (2004). *Univ Calif Santa Barbara, Dept Mol Cellular & Dev Biol, Santa Barbara, Ca 93106, USA. Centromeric DNA sequences in the pathogenic yeast *Candida albicans* are all different and unique. *Proc Natl Acad Sci U S A* **101**, (31) 11374
- Talarek N, Louis EJ, Cullin C, Aigle M. (2004). IBGC-CNRS UMR 5095, 1 rue Camille Saint-Saens, FR-33077 Bordeaux, France. Developing methods and strains for genetic studies in the *Saccharomyces bayanus* var. *uvarum* species. *Yeast* **21**, (14) 1195
- Tange Y, Fujita A, Toda T, Niwa O*. (2004). *Kazusa DNA Res Inst, 2-6-7 Kazu Sakamatari, Chiba 292 0818, Japan. Functional dissection of the γ -tubulin complex by suppressor analysis of *gth1* and *alp4* mutations in *Schizosaccharomyces pombe*. *Genetics* **167**, (3) 1095
- Wang XA, Ira G, Tercero JA, Holmes AM, Diffley JFX, Haber JE*. (2004). *Brandeis Univ, Rosenstiel Ctr, Waltham, Ma 02454, USA. Role of DNA replication proteins in double-strand break-induced recombination in *Saccharomyces cerevisiae*. *Mol Cell Biol* **24**, (16) 6891
- Washington MT, Minko IG, Johnson RE, Haracska L, Harris TM, Lloyd RS, Prakash S, Prakash L*. (2004). *Univ Texas, Med Branch, Sealy Ctr Mol Sci, 6-104 Blocker Med Res Bldg, 11th & Mechanic St, Galveston, Tx 77555, USA. Efficient and error-free replication past a minor-groove N^2 -guanine adduct by the sequential action of yeast Rev1 and DNA polymerase ζ . *Mol Cell Biol* **24**, (16) 6900
- Wysocicki R, Kron SJ*. (2004). *Univ Chicago, Ctr Mol Oncol, 924 East 57th St, Rm R320, Chicago, Il 60637, USA. Yeast cell death during DNA damage arrest is independent of caspase or reactive oxygen species. *J Cell Biol* **166**, (3) 311
- Yamada Y, Nakagawa T, Masukata H*. (2004). *Osaka Univ, Grad Sch Sci, Toyonaka, Osaka 560 0043, Japan. A novel intermediate in initiation complex assembly for fission yeast DNA replication. *Mol Biol Cell* **15**, (8) 3740
- Zelter A, Bencina M, Bowman BJ, Read ND*. (2004). *Univ Edinburgh, Inst Cell Biol, Fungal Cell Biol Grp, Edinburgh EH9 3JH, Scotland. A comparative genomic analysis of the calcium signaling machinery in *Neurospora crassa*, *Magnaporthe grisea*, and *Saccharomyces cerevisiae*. *Fungal Genet Biol* **41**, (9) 827

8 Physiology

- Fukuda K, Jin-Shan C, Kawano M, Sudo K, Gunge N*. (2004). *Sojo Univ, Dept Appl Microbial Technol, Ikeda 4-22-1, Kumamoto 860 0082, Japan. Stress responses of linear plasmids from *Debaryomyces hansenii*. *FEMS Microbiol Lett* **237**, (2) 243
- Hazan R, Levine A, Abeliovich H*. (2004). *Hebrew Univ Jerusalem, Dept Biochem & Food Sci, IL-76100 Rehovot, Israel. Benzoic acid, a weak organic acid food preservative, exerts specific effects on intracellular membrane trafficking pathways in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* **70**, (8) 4449
- Mercier-Bonin M, Ouazzani K, Schmitz P*, Lorthois S. (2004). *INSA, UMR INRA 792, UMR CNRS 5504, Lab Biotechnol & Bioprocés, 135 Ave Rangueil, FR-31077 Toulouse 4, France. Study of bioadhesion on a flat plate with a yeast/glass model system. *J Colloid Interface Sci* **271**, (2) 342

- Mercier RW, Rabinowitz NM, Ali R, Gaxiola RA, Berkowitz GA*. (2004). *Univ Connecticut, Dept Plant Sci, Agr Biotechnol Lab, 1390 Storrs Rd U-4163, Storrs, Ct 06269, USA. Yeast hygromycin sensitivity as a functional assay of cyclic nucleotide gated cation channels. *Plant Physiol Biochem* **42**, (6) 529
- Munhoz DC, Netto LES*. (2004). *University Sao Paulo, Inst Biociencias, Department Biol, BR-05508-900 Sao Paulo, Brazil. Cytosolic thioredoxin peroxidase I and II are important defenses of yeast against organic hydroperoxide insult - Catalases and peroxidoxins cooperate in the decomposition of H₂O₂ by yeast. *J Biol Chem* **279**, (34) 35219
- Pelling AE, Sehati S, Gralla EB, Valentine JS, Gimzewski JK*. (2004). *Univ Calif Los Angeles, Dept Chem & Biochem, 607 Charles E Young Dr East, Los Angeles, Ca 90095, USA. Local nanomechanical motion of the cell wall of *Saccharomyces cerevisiae*. *Science* **305**, (5687) 1147
- Ricci M, Martini S, Bonechi C, Trabalzini L, Santucci A, Rossi C*. (2004). *Univ Siena, Dept Chem & Biosyst Sci, Via Aldo Moro 2, IT-53100 Siena, Italy. Inhibition effects of ethanol on the kinetics of glucose metabolism by *S. cerevisiae*: NMR and modelling study. *Chem Phys Lett* **387**, (4-6) 377
- Tani A, Zhang D, Duine JA, Kawai F*. (2004). *Okayama Univ, Bioresources Res Inst, 2-20-1 Chuo, Okayama 710 0046, Japan. Treatment of the yeast *Rhodotorula glutinis* with AlCl₃ leads to adaptive acquisition of heritable aluminum resistance. *Appl Microbiol Biotechnol* **65**, (3) 344
- Tsukamoto I, Yim B, Stavarache CE, Furuta M, Hashiba K, Maeda Y. (2004). Univ Osaka Prefecture, Grad Sch Engn, 1-1 Gakuencho, Sakai, Osaka 599 8531, Japan. Inactivation of *Saccharomyces cerevisiae* by ultrasonic irradiation. *Ultrason Sonochem* **11**, (2) 61
- Zhou WD, Ryan JJ, Zhou HL*. (2004). *Univ Calif San Diego, Ludwig Inst Canc Res, 9500 Gilman Dr, CMM-East, Rm 3050, La Jolla, Ca 92093, USA. Global analyses of sumoylated proteins in *Saccharomyces cerevisiae* - Induction of protein sumoylation by cellular stresses. *J Biol Chem* **279**, (31) 32262
- ## 9 Medical mycology
- Barker KS, Crisp S, Wiederhold N, Lewis RE, Bareither B, Eckstein J, Barbuch R, Bard M, Rogers PD*. (2004). *Lebonheur Childrens Hospital & Med Center, Childrens Fdn Res Ctr, Room 304, West Patient Tower, 50 Nth Dunlap St, Memphis, Tn 38103, USA. Genome-wide expression profiling reveals genes associated with amphotericin B and fluconazole resistance in experimentally induced antifungal resistant isolates of *Candida albicans*. *J Antimicrob Chemother* **54**, (2) 376
- D'Souza CA, Hagen F, Boekhout T, Cox GM, Heitman J*. (2004). *Duke Univ Med Ctr, Dept Mol Genet & Microbiol, Durham, NC 27710, USA. Investigation of the basis of virulence in serotype A strains of *Cryptococcus neoformans* from apparently immunocompetent individuals. *Curr Genet* **46**, (2) 92
- De Lima RF, Brito MMD, Schaffer GMV, De Lima OC, Borba CD*. (2004). *Fiocruz MS, Fundacao Oswaldo Cruz, Insts Oswaldo Cruz, Dept Micol, CP 926, BP-21045-900 Rio de Janeiro, Brazil. Evaluation of the *in vitro* and *in vivo* dimorphism of *Sporothrix schenckii*, *Blastomyces dermatitidis*, and *Paracoccidioides brasiliensis* isolates after preservation in mineral oil. *Can J Microbiol* **50**, (6) 445
- Diaz MR, Fell JW. (2004). Univ Miami, Rosenstiel Sch Marine & Atmosphere Sci, Div Marine Biol & Fisheries, Miami, Fl 33149, USA. High-throughput detection of pathogenic yeasts of the genus *Trichosporon*. *J Clin Microbiol* **42**, (8) 3696
- Echt S, Bauer S, Steinbacher S, Huber R, Bacher A, Fischer M*. (2004). *Max Planck Inst Biochem, Abt Strukturforsch, DE-82512 Martinsried, Germany. Potential anti-infective targets in pathogenic yeasts: Structure and properties of 3,4-dihydroxy-2-butanone 4-phosphate synthase of *Candida albicans*. *J Mol Biol* **341**, (4) 1085
- Gusiano GE, Mangiaterra M, Rojas F, Gomez V. (2004). Univ Nacl Nordeste, Inst Med Reg, Dept Micol, Avda Heras 727, AR-3500 Resistencia, Chaco, Argentina. Yeasts species distribution in neonatal intensive care units in northeast Argentina. *Mycoses* **47**, (7) 300
- Gokulshankar S, Ranganathan S*, Ranjith MS, Ranjithsingh AJA. (2004). *21 Old 6 Shivaji St, IN-600017 Madras, India. Prevalence, serotypes and mating patterns of *Cryptococcus neoformans* in the pellets of different avifauna in Madras, India. *Mycoses* **47**, (7) 310
- McClelland EE, Adler FR, Granger DL, Potts WK. (2004). Albert Einstein Coll Med, Dept Med, 702 Golding, 1300 Morris Pk Ave, Bronx, NY 10461, USA. Major histocompatibility complex controls the trajectory but not host-specific adaptation during virulence evolution of the pathogenic fungus *Cryptococcus neoformans*. *Proc R Soc Lond Ser B Biol Sci* **271**, (1548) 1557
- McCullough M, Jorge J, Lejbkovicz F, Lefler E, Nassar F, Clemons K, Stevens D. (2004). Univ Melbourne, Sch Dent Sci, 711 Elizabeth St, Melbourne, Vic 3000, Australia. Genotypic differences of *Candida albicans* and *C. dubliniensis* isolates related to ethnic/racial differences within the same geographic area. *Mycopathologia* **158**, (1) 39
- Okamoto Y, Aoki S*, Mataga I. (2004). *Nippon Dent Univ, Adv Res Ctr, 1-8 Hamaura cho, Niigata 951 8580, Japan. Enhancement of amphotericin B activity against *Candida albicans* by superoxide radical. *Mycopathologia* **158**, (1) 9
- Pereira LA, Pereira M, Felipe MSS, Zancope-Oliveira RM, Soares CMD*. (2004). *Univ Fed Goias, ICBI/ICB2, Mol Biol Lab, Campus 2, BR-74001-970 Goiania, GO, Brazil. Proteomic identification, nucleotide sequence, heterologous expression and immunological reactivity of the triosephosphate isomerase of *Paracoccidioides brasiliensis*. *Microbes Infect* **6**, (10) 892
- Pfaller MA, Hazen KC, Messer SA, Boyken L, Tendolkar S, Hollis RJ, Diekema DJ. (2004). Univ Iowa Coll Med, Dept Pathol, Med Microbiol Div, C606 GH, Iowa City, Ia 52242, USA. Comparison of results of fluconazole disk diffusion testing for *Candida* species with results from a central reference laboratory in the ARTEMIS global antifungal surveillance program. *J Clin Microbiol* **42**, (8) 3607
- San Miguel LG, Pla J, Cobo J, Navarro F, Sanchez-Sousa A, Alvarez ME, Martos I, Moreno S. (2004). Hosp Ramon y Cajal, Infect Dis Serv, ES-28034 Madrid, Spain. Morphotypic and genotypic characterization of sequential *Candida parapsilosis* isolates from an outbreak in a pediatric intensive care unit. *Diagn Microbiol Infect Dis* **49**, (3) 189
- Shin JH, Kim MN, Shin DH, Jung SI, Kim KJ, Cho D, Kee SJ, Shin MG*, Suh SP, Ryang DW. (2004). *Chonnam Natl Univ, Sch Med, Dept Lab Med, 8 Hakdong dongku, Kwangju 501 757, South Korea. Genetic relatedness among *Candida tropicalis* isolates from sporadic cases of fungemia in two university hospitals in Korea. *Infect Control Hosp Epidemiol* **25**, (8) 634
- Sriburee P, Khayhan S, Khamwan C, Panjaisae S, Tharavichitkul P. (2004). Chiang Mai Univ, Fac Med, Dept Microbiol, TH-50200 Chiang Mai, Thailand. Serotype and PCR-fingerprints of clinical and environmental isolates of *Cryptococcus neoformans* in Chiang Mai, Thailand. *Mycopathologia* **158**, (1) 25
- Sugita T, Tajima M, Takashima M, Amaya M, Saito M, Tsuboi R, Nishikawa A. (2004). Meiji Pharmaceut Univ, Dept Microbiol, TH-5022-1 Noshio, Tokyo 204 8588, Japan. A new yeast, *Malassezia yamatoensis*, isolated from a patient with seborrheic dermatitis, and its distribution in patients and healthy subjects. *Microbiol Immunol* **48**, (8) 579
- Yang Q, Wang L*, Lu DN, Xing SY, Yin D, Zhu XJ. (2004). *NE Normal Univ, Inst Genet & Cytol, CN-130024 Changchun, Peoples Rep China. Phage displaying epitope of *Candida albicans* HSP90 and serodiagnosis. *Chin Med J (Engl)* **117**, (7) 1098
- ## 10 Recombinant DNA technology
- Schaerer-Brodbeck C, Barberis A. (2004). ESBATech AG, Wagistr 21, CH-8952 Schlieren, Switzerland. Coupling homologous recombination with growth selection in yeast: A tool for construction of random DNA sequence libraries. *Biotechniques* **37**, (2) 202
- Terentiev Y, Pico AH, Boer E, Wartmann T, Klabunde J, Breuer U, Babel W, Suckow M, Gellissen G, Kunze G*. (2004). *Inst Pflanzen-genet & Kulturpflanzenforsch, Corrensstr 3, DE-06466 Gaterleben, Germany. A wide-range integrative yeast expression vector system based on *Arxula adenivorans*-derived elements. *J Ind Microbiol Biotechnol* **31**, (5) 223