

# Prioritizing Infection Prevention and Control Activities for SARS-CoV-2 (COVID-19): A Multi-Criteria Decision-Analysis Method

This article was published in the following Dove Press journal:  
*Journal of Healthcare Leadership*

Adnan Sarwar  
Muhammad Imran

Department of Economics and Business  
Management, University of Veterinary  
and Animal Sciences, Lahore, Pakistan

**Purpose:** The outbreak of COVID-19 in China has spread to the whole world within a short period. The spread of COVID-19 has caused innumerable economic and social losses. The coronavirus spread directly from individual to individual. WHO and other governing bodies made guidelines and remedial activities to minimize the spread of COVID-19. This study aims to analyze and prioritize infection prevention and control activities for SARS-CoV-2.

**Methods:** An analytical study was conducted between medical representatives and scientists to check the impact of these precautionary measures (social distancing, shaking hands and hugging peoples, unnecessary touching things, hygiene, and sharing personal items) on COVID-19. The Analytical Hierarchical Process (AHP) method has been utilized to evaluate and prioritize the preventive measures of COVID-19.

**Results:** The results of this investigation shows, unnecessary travel, 3Cs, and touching own body parts should be avoided. Wearing a mask and proper hand washing are important to reduce the spread of coronavirus.

**Conclusion:** The SARS-CoV-2 has affected more than 11 million people around the world. The second spike of COVID-19 is also alarming for people. It is important to follow the instruction provided by the WHO, CDC, and other national and international governing bodies. The AHP method was used to check various measures that are spreading the coronavirus disease. It demonstrated that 3Cs (spaces that are closed, crowded, and involve close contacts) and travel, shaking hand, unnecessary touching own body parts should be avoided.

**Keywords:** pandemic COVID-19, preventive measure, decision making, AHP, SARS-CoV-2

## Introduction

COVID-19 which was first found in Wuhan city of China was soon listed as a public health emergency of international concern by the Emergency Committee of the World Health Organization (WHO). The virus is transmitted directly from infected people (saliva and discharge of nose). Individual may also be infected by touching their own body parts (eyes, nose, and mouth) and touching the surfaces of contaminated.<sup>1</sup> COVID-19 is also spread through liquid particles such as coughing, speaking, shouting, and breathing.<sup>2,3</sup> It did not take a long time to spread to other parts of the world. According to John Hopkins University, to date, it has affected 37 million people globally and caused 1 million deaths. The swift spread of COVID has not only affected the lives of millions of people, but it has also resulted in countless economic and social losses. So far, the United States, India, and Brazil are

Correspondence: Adnan Sarwar  
Department of Economics and Business  
Management, University of Veterinary and  
Animal Sciences, Lahore, Pakistan  
Tel +92 3007699710  
Email adnansarwar@live.com

badly affected by its spread. While in some countries, cases are declining, and some other countries are seeing spikes. The countries like Austria, China, and Singapore which have uplifted lockdowns and eased the rule of social distancing are placing to prevent coronavirus. Some countries are under lockdown again, faced with the threat of a second wave of COVID.<sup>4,5</sup> Experts are warning about the possibilities of the second wave attributed mainly to human behavior. People were frustrated after being at home for a long time and now breaking all the social rules. Analysis of cellphone data shows a declined social distancing. As a result, some experts have warned that the second wave could be more severe. Most of the predictions about COVID-19 by experts are based on studying the behavior of other viruses such as the 1918 flu pandemic and the 2009 H1N1 flu epidemic.<sup>6</sup>

In earlier days, many countries in the world announced lockdowns to curb the spread of virus. However, from an economic standpoint, there's a greater trade-off between saving lives and saving livelihood. Shutting the economy threatens the livelihood of the hundreds of millions of poor, especially in developing countries where government support has been minimal. Policy decisions made during the first wave of COVID-19, have varying impacts in different countries. Due to differences in actions the magnitude of impacts has also varied a lot between countries. Some of the countries have been successful in controlling the spread of disease and some have been worst hit by the virus. A Pew Research Center survey of more than 14,000 adults across 14 advanced economies in Europe, Asia, North America, and Australia, found 73% thought their own country had done a good job of tackling the coronavirus outbreak. Countries like Australia and Denmark got the highest response approval rating from the citizens. Due to delayed action in combating COVID-19 countries like the UK and the US received less emphatic support from their citizens (WEF, 2020).

A renaissance of COVID-19 is a threat to the world according to the World Health Organization. With the country's uplifting lockdown, educational institutes being opened, international travel being resumed, there's a greater threat of a resurgence of COVID-19 due to many unwitting coronavirus carriers in different areas. It's a challenge because now it will make it harder to detect and control transmission, unlike the previous, when we had the opportunity to contact, trace, and quarantine (John Hopkins University). While some countries are facing a second wave and some countries are at risk to

follow the same pattern, national and global health bodies are constantly updating their guidelines to curb the spread of COVID-19. Researchers around the globe are working day and night to find vaccines for COVID-19. Unprecedented cooperation has been seen between public and private health organizations during the pandemic. Many health organizations are working with biopharmaceutical companies to accelerate the process of finding a drug or vaccine for novel coronavirus. Up to now, different vaccines have been produced and being approved by health bodies of countries across the globe.<sup>7</sup> However, universal access to the vaccine is a major challenge for many countries especially resource-poor countries. Additionally, a new variant of the virus is posing threat to scientific achievement against COVID-19. Taking into consideration the above situation, following early mentioned precautionary measures is the wise thing to do, until, we reach universal vaccination.

Various health bodies like WHO, CDC, etc. are calling for curbing further spread of coronavirus by adopting different measures. Global and national health bodies have been advising to adopt different strategies to slow down the spread. One earliest and common advice was to avoid unnecessary travel, 3Cs, and contact with people having COVID-19 symptoms. Authorities have been recommending wearing a face mask, frequently wash and sanitize hands, and use hand gloves, etc.<sup>8</sup> Massive awareness campaigns have been launched by government bodies to educate citizens about safety measures during the pandemic. Information technology companies have also developed applications to provide up-to-date information about coronavirus outbreak.<sup>9</sup>

National Governments have been continuously advised by global (WHO, UNICEF, CDC, etc.) and national health bodies to adopt/enforce different measures to control the spread of COVID-19. These activities include identification, diagnosis, and management of the cases, control and prevention in healthcare settings, implementation of health measures for travelers, and risk communication.<sup>10</sup> The spread of COVID-19 can also be control by public intelligence and social awareness. Various social and physical measures such as social and physical distancing, outdoor use of face masks, restraining from unnecessary travel, hygiene, proper nutrition, health monitoring, etc. have been recommended by global health bodies.<sup>10</sup> However, the adoption of all recommendations at an equal level by people is an unrealistic hope. Therefore, there is a need to priorities those measures which are more helpful in


curtailing the spread as compared to those which have less impact.<sup>11</sup> In this situation, decision making is influenced by several precautionary measures. Therefore, it is important to consider those decision-making methods that help to analyze and prioritize precautionary measures. This is a complex decision-making process to determine priority order.<sup>12</sup> The most appropriate way to achieve this is the Multi-Criteria Decision Making Analysis technique such as Analytical Hierarchical Process (AHP). It is one of the most important methods that has been applied in the presence of several criteria.<sup>13</sup> The AHP method is utilized to

construct the hierarchal framework of a decision problem using different preventive measures recommended by global health bodies. The hierarchical structure of preventive measure is presented in Figure 1.

## Method

### The Analytical Hierarchal Process

The Analytical Hierarchal Process method was introduced by Saaty.<sup>14</sup> It is a powerful method for analyzing complex decision-making problems based on mathematics. It has been extensively studied and refined. It



**Figure 1** The hierarchal structure of preventive measures.  
**Note:** Data from Gao and Yu,<sup>1</sup> US CDC,<sup>2</sup> and WHO.<sup>3</sup>

is an accurate method to develop a rational framework of the problems and quantify its criteria for evaluation.<sup>15</sup> AHP method consists of two phases: 1) Develop the hierarchal structure of decision-making problems, 2) assign the weights to the decision criteria to determine the priorities of alternatives.

An AHP and F-VIKOR were used to develop an integrated management system while considering the needs and expectations of the stakeholders.<sup>16</sup> A fuzzy AHP-TOPSIS was utilized to conduct a comparison of five hydrometallurgical processes to recover the copper.<sup>17</sup> The authors proposed five decision-making methods AHP, FAHP, TOPSIS, FTOPSIS, and PROMETHEE evaluate road segments and the safety level of the roads. The AHP, FAHP, and PROMETHEE methods show better performance as compare to others.<sup>18</sup> This methodology helps the decision-maker to rationalize select and rank complex criteria. The procedure of AHP methodology is as follow:

- Construct the hierarchal framework of a decision problem
- Develop a pairwise comparison matrix of decision problem using Saaty's scale
- Calculate the consistency index (CI)
- To calculate the CI of the pairwise comparison matrix, Lambda max (eigenvalue) is required, and n represents the number of criteria. The CI formula is presented in Equation 1.

$$CI = \frac{\lambda_{\max} - n}{n - 1} \quad (1)$$

Compute consistency ratio (CR). The consistency ratio can be computed by Equation 2.

$$CR = \frac{CI}{RI} \quad (2)$$

**Table 1** Main Criteria and Sub-Criteria

Criteria	Sub-Criteria
Social distance (SD)	Travel SD1, 3Cs SD2, shaking hands and hugging people SD3
Hygiene (H)	Wearing a mask H1, hand wash H2, workplace sanitization H3
Sharing personal items (SPI)	Phone SPI1, combs SPI2, crockery SPI3, sanitary items SPI4, stationery SPI5
Unnecessary touching things (UTT)	Public objects UTT1, own body parts UTT2, Doorknob/stair bannisters UTT3
Healthy Diet (HD)	Mix whole grain foods HD1, avoid processed, baked, and fried foods HD2, drink enough water HD3
Immunity/Fitness (I/F)	Exercise regularly I/F1, get adequate sleep I/F2, minimize stress I/F3, maintain a healthy weight I/F4
Monitor your health (MH)	Shortness of breath MH1, fever MH2, sore throat MH3, congestion/runny nose MH4.

**Note:** Data from Gao and Yu,<sup>1</sup> US CDC,<sup>2</sup> and WHO.<sup>3</sup>

According to Saaty the value of CR must be within the limit of 0.1. If it exceeds revision of the pairwise comparison matrix is needed.

## Results

For this research, the pandemic COVID-19 prevention measure has been selected. As we know WHO and other governing bodies of various countries have issued guidelines for precautionary measures of COVID-19. The Pakistani Govt have also provided various guidelines to curb the spread of the virus. The preventive measures (criteria) and sub-criteria are shown in Table 1. The prevention measure consists of seven main criteria along with twenty-six sub-criteria. A multi-criteria decision-making method has been used to construct the pairwise

**Table 2** Pairwise Comparison Matrix of Preventive Measure, Ranking, and CR

Preventive Measures	SD	H	SPI	UTT	HD	I/F	MH	Priority Weights	Rank	
Social distance (SD)	1	4	3	5	3	5	4	0.36525	1	CI =0.062963 CR= 0.047699 RI= 1.32
Hygiene (H)		1	1	0.50	3	2	2	0.11736	4	
Sharing personal items (SPI)			1	0.50	3	1	3	0.12494	3	
Unnecessary touching things (UTT)				1	3	5	2	0.17976	2	
Healthy diet (HD)					1	0.50	0.33	0.05052	7	
Immunity/fitness (I/F)						1	0.33	0.06663	6	
Monitor your health (MH)							1	0.09553	5	

**Table 3** COVID-19 Preventive Measures (PM), Their Ranking/Priorities, and a Brief Description

PM Codes	Criteria	Weight of Main Criteria	Weight of Sub-Criteria	Global Weights	Rank	Brief Description
SD	<b>Social distance</b>	0.36525				Social distancing also called physical distancing, means keeping a safe distance between you and yourself. WHO and CDC strongly recommend a safe distance to minimize the spread of COVID-19.
SD1	Travel		0.59167	0.21611	1	Unnecessary travel of people is a violation of social distance and may increase the chance of getting and spreading COVID-19.
SD2	3Cs		0.29173	0.10655	2	WHO recommends avoiding 3Cs: spaces that are <u>closed</u> , <u>crowded</u> , and involve <u>close</u> contacts.
SD3	Shaking hands and hugging people		0.11660	0.04259	7	Shaking hands and hugging people is a way of transferring the virus from one person to another. People should avoid hugging to reduce infection.
H	<b>Hygiene</b>	0.11736				Hygiene is extremely important in limiting the transmission of the COVID-19 virus.
H1	Wearing a mask		0.52233	0.06130	4	Wearing a mask is a very important prevention measure to limit the spread of COVID-19.
H2	Hand wash		0.24542	0.02880	11	Proper handwashing prevents infections from one person to others and keeps you healthy.
H3	Workplace sanitization		0.12046	0.01414	19	Cleaning and disinfecting the workplace helps to remove germs.
H4	Cover cough and sneezes		0.11179	0.01312	21	Covering cough and sneeze can help to prevent respiratory diseases like COVID-19.
SPI	<b>Sharing personal items</b>	0.12494				People should avoid sharing their things with others. It is very difficult to identify how a person is infected.
SPI1	Phone		0.12098	0.01512	18	The surface of your phone can be a source of infection. So it is important to stop sharing your phone prevent coronavirus.
SPI2	Combs		0.09851	0.01231	23	People should not share their combs with others to prevent infection.
SPI3	Crockery		0.24302	0.03036	10	Do not share the crockery items or eating utensils.
SPI4	Sanitary items		0.42934	0.05364	5	Sanitary items should not be shared because it could be a source of COVID.
SPI5	Stationary		0.10815	0.01351	20	Limit sharing your stationery items reduce the spread of germs.
UTT	<b>Unnecessary touching things</b>	0.17976				Be mindful to avoid touching unnecessary things.
UTT1	Public objects		0.26752	0.04809	6	People must be careful while touching public objects such as ATM and currency notes.
UTT2	Own body parts		0.53770	0.09666	3	It is extremely important to avoid touching your body parts, such as the mouth, eyes, and nose. Often hands touch surfaces, may pick up the virus, and infect your body.

(Continued)

Table 3 (Continued).

PM Codes	Criteria	Weight of Main Criteria	Weight of Sub-Criteria	Global Weights	Rank	Brief Description
UTT3	Doorknob/stair bannisters		0.19478	0.03501	8	Touching doorknob, lift button, and stair banisters must be avoided.
HD	<b>Healthy Diet</b>	0.05052				It is very important to eat a healthy diet during the COVID-19 pandemic. A healthy diet enables the body to fight, prevent, and recover from infections.
HD1	Mix whole grain foods		0.30376	0.01535	17	Eat mix wholegrain foods such as maize, wheat, beans, and fresh fruits. These foods are rich in fiber and reduce the risk of infections.
HD2	Avoid processed, baked, and fried foods					People must avoid processed foods because it includes artificial ingredients, trans fat, and refined carbohydrates. It may increase the risk of infections
HD3	Drink enough water		0.52421	0.02648	13	Drinking enough water keeps your mouth moist that can prevent coronavirus.
I/F	<b>Immunity/ Fitness</b>	0.06663				The immune system is a network of units that work together to combat invaders, including bacteria, and viruses. Boosting immunity helps to prevent COVID-19.
I/F1	Exercise regularly		0.29364	0.01957	16	The experts recommend do exercising moderately boost immunity to fight against COVID-19.
I/F2	Get adequate sleep		0.32599	0.02172	15	Getting enough sleep support the immunity system. It minimizes risk infections and can improve health for fighting the viruses.
I/F3	Minimize stress		0.19573	0.01304	22	The coronavirus may cause stress for people because it disrupts daily routines. Reducing stress minimizes the risk of infections.
I/F4	Maintain a healthy weight		0.18464	0.01230	24	Eating a balanced diet to maintain a healthy weight during COVID. Overweight may put you at risk for diseases.
MH	<b>Monitor your health</b>	0.09533				Frequently monitoring your health is an important aspect to prevent COVID.
MH1	Shortness of breath		0.11943	0.01141	25	Shortness of breath is also called dyspnea. The coronavirus attacks the lungs and respiratory system. So it is important to perform an exercise to avoid such infections.
MH2	Fever		0.28028	0.02678	12	Fever is also known as elevated temperature. If body temperature is higher than normal. Frequently check the body temperature may help to prevent the virus.
MH3	Sore throat		0.36246	0.03464	9	According to CDC sore throat is an early symptom of COVID-19, because the virus inhaled through the nose and throat. So, people must cover your nose and mouth to prevent the ailment.
MH4	Congestion/ runny nose		0.23764	0.02270	14	Congestion/runny nose is also considered as hallmarks of the novel COVID-19.

comparison matrix of the main criteria as well as sub-criteria. Pairwise comparison matrices have been developed to determine the weights of every criterion. AHP method was utilized to calculate the weights of pandemic COVID-19 preventive measures. The results of this research are presented in Table 2. The major preventive measures: social distance hold the highest weight (0.36525), unnecessary touching things (0.17976), sharing personal items (0.12494), hygiene (0.11736), monitor your health (0.09553), immunity/fitness (0.06663), and healthy diet (0.05052). The priority weights of main criteria along with sub-criteria have been presented in Table 3. The global weights/cumulative weights of each sub-criteria have been computed, to prioritize them. The consistency ratio has also been checked and the results are tolerable. Further, the importance of every criterion has been discussed.

## Discussions

The new virus known as COVID-19 began in China. This disease spread from infected person to person. It also transmits from people's nose droplets or cough. A multi-criteria decision making analysis is used to analyze and prioritize the COVID-19 precautionary measures.

Social distancing is ranked top. It means people should keep 1 meter (3 feet) distance between them. It is also important people should avoid unnecessary travel, hand-shaking, hugging, and crowded areas.

Hygiene plays a vital role to limit the spread of the virus. Wearing a face mask is mandatory. Regular hand washing/sanitizing and cover the mouth while sneezing or cough. Proper cleaning and disinfection of the workplace are essential to prevent this virus. Sharing personal items such as phone, comb, crockery, stationery, and sanitary items should be discouraged. People may catch the virus by touching unnecessary objects such as public objects, their body parts, doorknob/bannister. So, avoid unnecessary touching these objects can defeat COVID-19.

A balanced diet helps the body to combat, avoid, and recover from infections. Drinking enough water and getting adequate sleep also improve health. Doing regular exercise enhances immunity systems to fight against this virus. People should monitor their health properly, anyone who is feeling sick/shortness of breath, sore throat, and runny nose. They must consult the physician to prevent this novel coronavirus.

## Implications

Overall, the significant strength of this research was to analyze and prioritize the various SARS-CoV-2 precautionary measure to reduce transmission of this infection. For this purpose, a multi-criteria decision-making method was applied. Our findings demonstrated that social distancing, unnecessary travel, 3Cs, and unnecessary touching own body part should be avoided. Wearing a face mask, proper hand washing/sanitizing, and maintain a healthy diet is important to prevent the coronavirus.

This scientific research prioritizes the infection prevention and controls precautionary of SARS-CoV-2 informed by the World Health Organization (WHO), CDC, and UNICEF, etc. However, the researchers made a possible effort to consider theoretical and methodological insights in this study.

## Conclusion

The COVID-19 epidemic has affected 215 countries around the world. It is approximately said to have affected more than 11 million people around the globe. The WHO, CDC, and other governing bodies provided guidelines to stop the spread of coronavirus. These guidelines and precautionary measures have been selected in this research. A multi-criteria decision-making method was utilized to prioritize the various preventive measures to determine the impact, importance, and role of spreading the coronavirus disease.

Unnecessary travel and 3Cs (spaces that are closed, crowded, and involve close contacts) have been found, to have the greatest influence and the most significant factor in the spread of coronavirus. Unnecessary touching own body parts, shaking hands, and hugging peoples are the subsequent important issues in the same. All main preventive measures (criteria) along with sub-criteria have been ranked according to their importance. Many other preventive measures as well as other decision-making methods may also be considered in future research to assess the spread of coronavirus.

## Ethical Approval

Not required. The COVID-19 precautionary measure/prevention guidelines are considered for human care and freely available at WHO, CDC, UNICEF websites.

## Funding

There is no funding to report.

## Disclosure

The authors declare no conflicts of interest for this work.

## References

- Gao X, Yu J. Public governance mechanism in the prevention and control of the COVID-19: information, decision-making and execution. *J Chin Governance*. 2020;5(2):178–197. doi:10.1080/23812346.2020.1744922
- How COVID-19 spreads. U.S. Centers for Disease Control and Prevention (CDC); September 18, 2020. Retrieved September 20, 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-covid-spreads.html>.
- Coronavirus disease (COVID-19): How is it transmitted? World Health Organization (WHO); 2020. Retrieved December 6, 2020. Available from: <https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-how-is-it-transmitted>.
- Chatterjee SS, Vora M, Malathesh BC, Bhattacharyaa R. Worried well and Covid-19: re-emergence of an old quandary. *Asian J Psychiatr*. 2020;78:100738.
- Sun P, Lu X, Xu C, Sun W, Pan B. Understanding of COVID-19 based on current evidence. *J Med Virol*. 2020;92(6):548–551. doi:10.1002/jmv.25722
- Maragakis LL. First and second waves of coronavirus. John Hopkins University; 2020. Available from: <https://www.hopkinsmedicine.org/health/conditions-and-diseases/coronavirus/first-and-second-waves-of-coronavirus>. Accessed February 17, 2021.
- Regulatory Affairs Professionals Society. COVID-19 vaccine tracker; 2020. Available from: <https://www.raps.org/news-and-articles/news-articles/2020/3/covid-19-vaccine-tracker>. Accessed February 17, 2021.
- Sohrabi C, Alsafi Z, Neill N, et al. World health organization declares global emergency: a review of 2019 the novel coronavirus. *Int J Surg*. 2020;76:71–760. doi:10.1016/j.ijssu.2020.02.034
- Singh R, Avikal S. COVID-19: a decision-making approach for prioritization of preventive activities. *Int J Healthc Manag*. 2020;13(3):257–262. doi:10.1080/20479700.2020.1782661
- Coronavirus Disease. (COVID-19) situation report – 43 (2020); 2019b. World Health Organization; 2020. Available from: [https://www.who.int/docs/default-source/coronavirus/situation-reports/20200303-sitrep-43-covid-19.pdf?sfvrsn=76e425ed\\_2](https://www.who.int/docs/default-source/coronavirus/situation-reports/20200303-sitrep-43-covid-19.pdf?sfvrsn=76e425ed_2).
- Wu D, Lu J, Liu Y, Zhang Z, Luo L. Positive effects of COVID-19 control measures on influenza prevention. *Int J Infect Dis*. 2020;95:345–346. doi:10.1016/j.ijid.2020.04.009
- Ferrari Neto G, Leal GCL, Galdamez EVC, Souza RCTD. Prioritization of occupational health and safety indicators using the Fuzzy-AHP method. *Production*. 2020;30:1–13.
- Barbosa C, Azevedo R, Rodrigues MA. Occupational safety and health performance indicators in SMEs: a literature review. *Work*. 2019;64(2):217–227. doi:10.3233/WOR-192988
- Emshoff JR, Saaty TL. Applications of the analytic hierarchy process to long range planning processes. *Eur J Oper Res*. 1982;10(2):131–143. doi:10.1016/0377-2217(82)90153-9
- Saaty TL, Vargas LG. *Prediction, Projection, and Forecasting: Applications of the Analytic Hierarchy Process in Economics, Finance, Politics, Games, and Sports*. Kluwer Academic Pub; 1991.
- Ikram M, Zhang Q, Sroufe R. Developing integrated management systems using an AHP-Fuzzy VIKOR approach. *Bus Strat Environ*. 2020;29(1):2265–2283. doi:10.1002/bse.2501
- Yousefzadeh S, Yaghmaeian K, Mahvi AH, Nasser S, Alavi N, Nabizadeh R. Comparative analysis of hydrometallurgical methods for the recovery of Cu from circuit boards: optimization using response surface and selection of the best technique by two-step fuzzy AHP-TOPSIS method. *J Clean Prod*. 2020;249:119401. doi:10.1016/j.jclepro.2019.119401
- Sarraf R, McGuire MP. Integration and comparison of multi-criteria decision making methods in safe route planner. *Expert Syst Appl*. 2020;154:113399. doi:10.1016/j.eswa.2020.113399

Journal of Healthcare Leadership

Dovepress

### Publish your work in this journal

The Journal of Healthcare Leadership is an international, peer-reviewed, open access journal focusing on leadership for the health profession. The journal is committed to the rapid publication of research focusing on but not limited to: Healthcare policy and law; Theoretical and practical aspects healthcare delivery; Interactions between healthcare and society and evidence-based practices; Interdisciplinary decision-making;

Philosophical and ethical issues; Hazard management; Research and opinion for health leadership; Leadership assessment. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-healthcare-leadership-journal>