

Treatment of chronic hepatitis C regimens containing with recombinant interferon in patients with sustained virological response predicts risk of hepatocellular carcinoma

A meta-analysis

Chien-Feng Li, MD^{a,b}, Shih-Ming Tsao, MD, PhD^{a,b}, Hsien-Hua Liao, MD, PhD^{a,c}, Shiu-Chih Chen, MD, PhD^{a,d}, Yuan-Ti Lee, MD, PhD^{a,b,*}

Abstract

Given that evidence supporting chronic hepatitis C (CHC) infection developed chance for hepatocellular carcinoma (HCC) following antiviral agents therapy is controversial. We conducted a meta-analysis to examine the risk.

We evaluated 20 retrospective and prospective cohort studies published up to 31 December 2017 which investigated the association between sustained virological response (SVR) and incidence of HCC patients treated with monotherapy interferon (IFN) or IFN plus ribavirin (RBV) therapy. The primary outcome of the study was the cumulative incidence of HCC. Odds ratio (OR) was used to evaluate the index of effect size for the association between SVR and treatment with IFN alone or IFN/RBV in CHC patients.

SVR patients demonstrated a lower incidence of HCC compared to non-SVR patients. Non-SVR patients had greater odds of HCC incidence compared to SVR patients in the treatment of IFN plus RBV (pooled OR = 7.405, 95% CI = 4.689 to 11.694, $P < .001$). Non-SVR patients had greater odds of HCC incidence compared to SVR patients in the treatment of IFN monotherapy (pooled OR = 4.135, 95% CI = 3.009 to 5.682, $P < .001$). Lack of SVR to IFN therapy was significantly associated with greater risk of HCC incidence (pooled OR = 5.035, 95% CI = 3.915 to 6.474, $P < .001$).

SVR could be as a predictor of HCC in CHC patients treated with IFN or IFN plus RBV, and have important implications during HCC screening, whereby patients who fail to achieve SVR need to be screened more rigorously.

Abbreviations: CHC = chronic hepatitis C, DAA = direct-acting antivirals, HAI = histology activity index, HCC = hepatocellular carcinoma, HCV = hepatitis C virus, I2 = inconsistency index, IFN = Interferon, IR = incomplete response, NR = nonresponse, OR = odds ratio, Peg-interferon = pegylated interferon, PR = partial response, RBV = ribavirin, RCTs = randomized controlled trials, SVR = sustained virologic response, SVR = sustained virological response, TVR = transient virological response.

Keywords: chronic hepatitis C, direct-acting antiviral agents, hepatitis C virus infection, hepatocellular carcinoma, peg-interferon, ribavirin, sustained virological response

1. Introduction

Hepatitis C virus (HCV) infection is a major health concern, with approximately 71 million people infected worldwide and it was estimated that in 2016, approximately 399,000 people died from HCV infection, mostly from cirrhosis and hepatocellular carcinoma (HCC).^[1] Chronic hepatitis C (CHC) infection can

progress to liver fibrosis, cirrhosis, and HCC, which is currently the seventh most common malignancy, and the third cause of cancer-related deaths worldwide.^[2] Necro-inflammation plays an important role in hepatic carcinogenesis, and the occurrence of HCC has been shown to be associated with the degree and severity of liver fibrosis and cirrhosis.^[3,4] The goal of CHC

Editor: Akhilanand Chaurasia.

Funding Source: Chung Shan Medical University Hospital; Award ID: CSH-2020-C-011; Recipient: Yuan-Ti Lee. The funding source played no role in study design and conduct, data collection, analysis or interpretation, writing of the manuscript, or the decision to submit it for publication.

The authors have no conflicts of interest to disclose.

The datasets generated during and/or analyzed during the current study are publicly available.

^a School of Medicine, Chung Shan Medical University, ^b Division of Infectious Diseases, Department of Internal Medicine, ^c Department of Plastic Surgery, ^d Department of Family and Community Medicine, Chung Shan Medical University Hospital, Taichung, Taiwan.

* Correspondence: Yuan-Ti Lee, School of Medicine, Chung Shan Medical University, Taichung, Taiwan, Division of Infectious Diseases, Department of Internal Medicine, Chung Shan Medical University Hospital, Number 110, Section 1, Jianguo N. Road, Taichung City 40201, Taiwan (e-mail: leey521@gmail.com).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Li CF, Tsao SM, Liao HH, Chen SC, Lee YT. Treatment of chronic hepatitis C regimens containing with recombinant interferon in patients with sustained virological response predicts risk of hepatocellular carcinoma: A meta-analysis. *Medicine* 2020;99:40(e22435).

Received: 7 April 2020 / Received in final form: 20 July 2020 / Accepted: 24 August 2020

<http://dx.doi.org/10.1097/MD.00000000000022435>

treatment is virus elimination as well as prevention of HCC. Interferon (IFN) therapy has been shown to reduce inflammation and necrosis scores, prevent progression to HCC, and achieve viral clearance in CHC patients.^[4,5]

A combination of pegylated interferon α (peg-IFN α) and a nucleos(t)ide analog has been used successfully to reduce viral load, and improve inflammation and fibrosis in the liver.^[6,7] As previous studies reported CHC patients are treated with a combination of peg-IFN α along with ribavirin (RBV), which has been shown to arrest disease progression. IFN therapy was reported to be associated with a reduction in the incidence of HCC, especially in patients who achieve a sustained virological response (SVR).^[8–10] SVR is defined as aviremia at 24 weeks after completion of antiviral therapy.^[10,11] CHC patients on IFN therapy who achieved SVR have been shown to have a negligible risk of relapse.^[12]

Patients who achieved SVR or a transient virologic response (TVR) also had a lower incidence rate of HCC within 5 years after the end of treatment compared to patients who did not achieve SVR.^[13] A number of other studies validated these findings, and reported that SVR was significantly associated with a reduced risk of HCC. Particularly, patients with cirrhosis composed of life-threatening complications which may occur HCC even after SVR.^[13–18]

Currently, HCV direct-acting antiviral (DAA) agents are the milestone for the standard treatment of CHC infection.^[19,20] Although the present results have been already confirmed that the present results have been already confirmed that HCV patients who undergoing the DAA treatment can decrease all-cause mortality and liver-related adverse effects, including end-stage liver disease and HCC.^[20] However, therapy cost and access has served as a major barrier to low and middle-income countries in real-life clinical practice.^[21] Although current studies showed that CHC-related cirrhosis patients who received DAA treatment to eliminate HCV and achieving SVR, but need further long-term follow-up study to evaluate the risk of HCC.^[20]

A previous meta-regression analysis showed that a higher SVR rate was a predictor of the efficacy of antiviral therapy in preventing HCC occurrence in CHC patients.^[6] However, the risk of developing HCC can persist in some patients with HCV-related cirrhosis who achieve SVR after completion of IFN therapy.^[10] Additionally, there was no significant difference in the mean interval from the time of completion of IFN therapy until the detection of HCC between SVR and non-SVR HCC patients.^[22]

Based on the available data, it is clear that the link between SVR and the risk of HCC is not fully understood. This meta-analysis aimed to investigate the association between SVR and risk of HCC in CHC patients treated with IFN-based antiviral regimens (IFN alone or IFN plus RBV).

2. Methodology

2.1. Study selection

We performed a comprehensive literature search of the PubMed, Cochrane, and Embase databases using the following combinations of search terms: (interferon OR ribavirin) AND (hepatocellular carcinoma OR cirrhosis) AND hepatitis C, (interferon OR ribavirin) AND (hepatocellular carcinoma OR HCC) AND (hepatitis C). Studies published up to March, 2015 were identified.

The inclusion criteria for this meta-analysis were: all randomized controlled trials (RCTs) or 2-arm prospective studies; retrospective studies, studies evaluating patients with hepatitis C virus infection, studies in which patients were treated with IFN therapy: peg-IFN or peg-IFN and RBV, studies in which virological response was measured, and studies that reported the incidence of HCC in patients who achieved SVR and those who did not. Exclusion criteria were reviews, letters, comments, editorials, case reports, proceedings, and personal communications; studies designed for Hepatitis B infection; studies that did not publish in English; and studies with no quantitative primary outcome.

Data collected from the studies included treatment details, patient demographics, duration of treatment, virologic response, and presence of cirrhosis. Data extraction was performed by 2 independent reviewers, and a third reviewer was consulted for resolution of disagreement. In case the information on study design or data or outcomes was ambiguous, we would contact the original authors by email for clarification. We further updated the references of the relevant studies between April 2015 and July 2019 which complied with the inclusion criteria through hand searching. This study was approved by the institutional review board of the Chung Shan Medical University Hospital, Taichung, Taiwan (CSMUH No: CS18193). Raw patient data and private information were neither required nor used in the present review, and therefore informed consent from study subjects was waived.

2.2. Data analysis

The primary outcome of the study was the cumulative incidence of HCC. Odds ratio (OR) was used to evaluate the index of effect size for the association between SVR and treatment with IFN alone or IFN/RBV in CHC patients. It was interpreted as the ratio of the odds of HCC development in patients without SVR (non-SVR) to the odds of HCC in patients with SVR; an OR greater than 1 indicated that non-SVR patients had a higher risk of HCC than SVR patients on interferon therapy. The DerSimonian and Laird random-effects model was used to determine pooled estimates of OR and 95% confidence interval (95% CI). A 2-sided P value $< .05$ was considered statistically significant.

The Cochran Q and I² statistics were calculated to indicate the presence of heterogeneity across studies. For the Q statistics, a P value $< .10$ was considered statistically significant for heterogeneity. I² statistics were defined as the percentage of the observed between-study variability which occurred due to heterogeneity rather than by chance; heterogeneity was assessed as follows: no heterogeneity (I² = 0%–25%), moderate heterogeneity (I² = 25%–50%), large heterogeneity (I² = 50%–75%), and extreme heterogeneity (I² = 75%–100%).

Subgroup analysis was performed according to the treatment regimens (i.e., IFN plus RBV, IFN alone, or other). “Other” means that the patients were treated with IFN alone or IFN plus RBV, but the subgroup data (IFN alone or IFN plus RBV) were not reported. We also performed additional subgroup analyses according to the country of studies and according to follow-up durations. Sensitivity analysis was carried out for the outcomes using the leave one-out approach. All analyses were performed using Comprehensive Meta-Analysis statistical software, version 2.0 (Biostat, Englewood, NJ).

2.3. Quality assessment

Quality evaluation of the included studies was performed as previously described^[23] using 6 sources of bias related to study

participation, study attrition, measurement of prognostic factors, measurement of and controlling for confounding variables, measurement of outcomes, and analysis approaches. The quality of included studies was independently appraised by 2 reviewers. Disagreements were resolved by a third reviewer.

3. Results

Based on the search terms, and the inclusion and exclusion criteria, this meta-analysis identified a total of 389 studies, of which 336 were excluded for nonrelevance. Nonrelevance criteria included reviews, letters, comments, editorials, case reports; proceedings, personal communications, studies designed

for Hepatitis B infection, studies that did not include IFN therapy, studies not designed to evaluate SVR and incidence of HCC. We removed duplicate studies when searching (n=4). Of the remaining 49 studies, 29 were excluded because they did not include a comparison of outcomes between the SVR and non-SVR groups (n=15), absence of IFN response data at the virologic level (n=5), studies focusing on post HCC IFN therapy (n=3), and studies reporting HCC incidence which could not be pooled with other studies (n=6) (Fig. 1).

Of the 20 studies that were included, 4 studies were prospective studies^[13,15,24,25] and 16 were retrospective studies.^[4,11,16,17,22,26-36] Treatment regimens in 5 of the included studies consisted of combination therapy with peg-IFN and




Figure 1. Flow diagram of study selection. ALT=alanine aminotransferase, HCC=hepatocellular carcinoma, IFN=Interferon, SVR=sustained virological response.

RBV.^[4,11,24,26–28,31,33,34] The duration of treatment was at least 24 weeks in most of the included studies, and the total number of enrolled patients in each study ranged from 130 to 1654. The mean or median age ranged from 46.9 to 63.7 years, the proportion of male patients ranged from 37.6% to 73.4%, and the mean or median follow-up duration ranged from 34.6 months to 10.7 years (Table 1). The Knodell Histological Activity Index score (HAI score, denoted as scoring of necro-inflammatory activity in chronic hepatitis based on histological examination by pathologist), and percentage of patients with cirrhosis are also listed in Table 1. It is important to note that not all the included studies provided the number of incident cases of HCC. We therefore entered data as event rates and total sample size with the Comprehensive Meta-Analysis software, and the number of events was then generated by the software. As a result, there might be a difference in the HCC event number between what we presented and the actual event number.

3.1. Meta-analysis

Forest plots were used to determine the predictive effect of SVR on the incidence of HCC in patients receiving IFN therapy (Fig. 2). In 5 of the included studies where patients were treated with IFN plus RBV, the pooled odds ratio was 7.405 (95% CI=4.689 to 11.694; $P<.001$), without heterogeneity between studies ($Q=0.81$, $P=.937$, $I^2=0\%$). In 10 of the studies where patients were treated with IFN monotherapy, there was no evidence supporting heterogeneity between studies ($Q=7.51$, $P=.585$, $I^2=0\%$). The pooled results showed that non-SVR patients had greater odds of HCC incidence compared to SVR patients (pooled OR=4.135, 95% CI=3.009 to 5.682, $P<.001$). There was extreme heterogeneity across 5 studies where patients were treated with other IFN-related therapeutic regimens ($Q=31.43$, $P<.001$, $I^2=87.3\%$). Lack of SVR to IFN therapy was significantly associated with greater risk of HCC incidence (pooled OR=5.035, 95% CI=3.915 to 6.474, $P<.001$).

Results of sensitivity analyses showed that regardless of subgroups regarding treatment regimen, the direction and magnitude of this association did not change considerably when individual study removed one at a time (Fig. 3).

Subgroup analysis was performed based on the country where the study was conducted (Fig. 4). The results demonstrated that regardless of country, non-SVR patients had greater odds of HCC occurrence compared to SVR patients (Asian countries: pooled OR=5.697, 95% CI=3.870 to 8.385, $P<.001$; Non-Asian countries: pooled OR=3.842, 95% CI=2.182 to 6.763, $P<.001$) (Fig. 4). Furthermore, the pooled results were consistent across subgroups defined according to follow-up duration (<10 years: pooled OR=6.365, 95% CI=4.395 to 9.218, $P<.001$; ≥10 years: pooled OR=4.915, 95% CI=2.933 to 8.234, $P<.001$) (Fig. 5).

Quality assessment was performed using 6 sources of bias which could have impacted the quality of the studies. The data showed that all the included studies were of moderate-to-high quality (Fig. 6).

4. Discussion

In this meta-analysis, we evaluated the data extracted from 20 studies to investigate the predictive value of SVR on risk of HCC in CHC patients treated with IFN therapy. Our analysis showed that CHC patients who achieved SVR after IFN or IFN plus RBV

therapy had a lower risk of HCC compared to patients who failed to achieve SVR, suggesting that SVR may predict HCC risk in this group of patients.

IFN has been shown to reverse CHC-mediated inflammatory and regenerative processes in the liver.^[37] Normalization of aminotransferase levels after IFN therapy was associated with decreased HCC incidence, even in patients who failed to clear HCV RNA from the serum,^[31] and IFN α and IFN β were shown to have a similar efficacy at reducing the incidence of HCC.^[24] These studies were consistent with a meta-analysis of 3 RCTs and 6 prospective cohort studies which reported that single-course IFN therapy suppressed inflammation and fibrosis and significantly reduced the incidence of HCC even in virologic non-responders.^[38] Importantly, it has been reported that although some cirrhotic CHC patients who achieved SVR with IFN therapy developed HCC, none of the patients with histological regression of cirrhosis developed HCC.^[39] Even though, the actual mechanisms of HCV infection on hepatocarcinogenesis are currently not fully understood, similarly hepatitis B virus related to HCC, it is assumed that the hepatic inflammation and injury attributed frequently to host immune response and possibly provides the hepatocarcinogenesis.^[40] These results suggested that the preventive effect of IFN therapy on hepatocarcinogenesis might associate with its anti-inflammatory effects.^[40]

Although it would be interesting to understand if SVR reduces the risk of HCC regardless of the degree of liver fibrosis, we could not do this analysis because we did not have patient-level data for the degree of liver fibrosis. However, this issue has been described in 3 of our included studies that evaluated the incidence of HCC in cirrhotic versus noncirrhotic patients who achieved SVR. Patients who achieved SVR were shown to have a significantly lower risk of progression from a noncirrhotic to a cirrhotic state compared to patients who did not achieve SVR.^[17] Among patients who achieved SVR, the 5-year cumulative incidence of HCC in non-cirrhotic patients was 1.7%, and cirrhosis patients was 18.9%.^[13] This was consistent with another study that showed a significantly lower HCC incidence rate in noncirrhotic patients who achieved SVR compared to cirrhotic patients (0.09% vs 2.67%).^[29]

A previous study suggested that the reduced incidence of HCC and improved survival rates in SVR patients after IFN therapy were strongly correlated with virus eradication.^[29] Recent studies have shown that high SVR rates can predict a low risk of HCC occurrence in CHC patients treated with antiviral therapy.^[11,13,15–17,30,32,35,36] A meta-analysis on CHC patients showed IFN therapy can reduce the 3- and 5-year cumulative incidence of HCC and SVR in these patients was a predictor of superior preventive efficacy.^[6] Another meta-analysis study showed that although the effect of antiviral therapy on reduced risk of HCC was unrelated to virologic response, the effect was more pronounced among patients who achieved SVR compared to those who did not.^[14] Additionally, although IFN monotherapy reduced the risk of HCC among patients aged under 60 years old, suppression of HCC by IFN among patients aged over 60 years old was only observed among those patients who achieved SVR.^[34] Previous studies on the impact of TVR have also been shown to correlate with a lower cumulative incidence rate of HCC when compared with nonresponders.^[13,26,27] Patients who achieved SVR or TVR had significantly lower rates of HCC incidence compared to those who did not show a virologic response to IFN monotherapy,^[11,26–28] or to peg-IFN

Table 1**Study characteristics including treatment details, HAI-Kondell score, and proportion of cirrhosis.**

Study name, year	Study design	Treatment details	Treatment duration	Group of virologic response	No. of patients	Mean age (yr)	Male (%)	HAI score	Cirrhosis (%)	Mean follow-up duration	References
Kashiwagi, 2003	Prospective	Natural IFN α (74.1%), or natural IFN β (25.9%)	20 wk	SVR	66	55.7	66.7	9.9 (4.7)	13.7	5.7 yr	[24]
Shiratori, 2005	Prospective	IFN α -2a (n=157), or natural IFN α (n=114)	39 wk	Non-SVR	194	Median: 57	37.6			Median: 6.8 yr	[25]
Dohmen, 2013	Prospective	peg-IFN + RBV	24–72 wk	SVR	64						
Ogawa, 2013	Prospective	peg-IFN α -2b + RBV	47 (24–48) wk	TVR NVR SVR	285	55.5	48.5			Median: 56 mo	[15]
Imai, 1998	Retrospective	Recombinant IFN α -2a (n=149), or recombinant IFN α -2b (n=94)	6 mo	TVR NVR SVR	116 73 557	Median: 58	49.2		14.8	Median: 3.6 yr	[13]
Kasahara, 1998	Retrospective	Recombinant IFN α -2a, recombinant IFN α -2b, natural IFN β , or natural IFN α	24–52 wk	Relapse NR SR	151	<60 yr: 71% ≥60 yr: 29%	67.0	<10: n=254, ≥10: n=165		Median: 47.6 mo	[11]
Tanaka, 2000	Retrospective	Human lymphoblastoid IFN, recombinant IFN α -2a, or recombinant IFN α -2b	6 mo	TR NR SR	120 148 313	52.9	67.4	9.0 (3.4)		38.9 mo	[27]
Takimoto, 2002	Retrospective	IFN- α (natural or recombinant) or natural IFN- β	8–24 wk	TR NR SVR	304 405 175	51.7	66.4 67.9 69.0	8.9 (3.6) 9.6 (3.6)		36.6 mo 36.9 mo 59.6 mo	[26]
Yu, 2005	Retrospective	Recombinant IFN α -2a (n=40), IFN α -2b (n=102), or lymphoblastoid IFN α -n1 (n=72)	24 wk	TR NR SVR	165 254 201	51.3	58.4			57.3 mo 55.5 mo 54.8 mo	[4]
Yu, 2006	Retrospective	IFN α only (n=297) or IFN α + RBV (n=760)	20–48 wk	IR NR Non-SVR SVR	95 356 194 64	46.5	53.1	4.21 (2.31)		6.81 yr	[28]
Hung, 2006	Retrospective	IFN α -2b + RBV	24 or 48 wk	Non-SVR SVR	136 715	46.9	50.7 60.5	4.10 (2.59)	15.6	5.18 yr	[29]
Ikeda, 2006	Retrospective	Natural or recombinant IFN α (n=1238), natural IFN β (n=386), or both (n=30)	24 wk	Non-SVR SVR	342 73	56.1	57.5	6.8 (2.5)		Median: 37 mo	[30]
Arase, 2007	Retrospective	IFN α , IFN β , or IFN + RBV	Median 165 d	Non-SVR SVR	59 606	Median: 50	45.8 67.1	6.9 (2.6)		Median: 10.7 yr	[31]
Kobayashi, 2007	Retrospective	Natural IFN α , recombinant IFN α -2a, recombinant IFN α -2b, IFN- β , or recombinant IFN α -2b + RBV	24–48 wk	BR NR SVR	266 782 140	63.7	59.3	9.3 (3.4)		7.0 yr	[32]
Bruno, 2007	Retrospective	IFN only	1 yr	Non-SVR SVR	360 373	51.0	46.7 62.7	9.8 (2.7)		7.7 yr Median: 66 mo	[22]
Imai, 2010	Retrospective	Human lymphoblastoid IFN, recombinant IFN α 2a, or recombinant IFN α 2b	6 mo	Non-SVR, nonaged SVR, aged	751 124	54.7	59.9 73.4			96.1 mo	[33]
Cardoso, 2010	Retrospective	peg-IFN + RBV (82%), peg-IFN monotherapy (7%), conventional IFN with or without RBV (11%)	43 wk 38 wk	Non-SVR, nonaged SVR, aged	759 134	48.0	61.7 67.8	<10: n=257, ≥10: n=153			[34]
Velosa, 2011	Retrospective	IFN only or IFN + RBV	43 wk 38 wk	Non-SVR, aged SVR	276 41	63.5	54.4	<10: n=84 ≥10: n=74	53	Median: 3.5 yr	[32]
				Non-SVR SVR	117 103	55.0	70.0				[32]
				Non-SVR SVR	204 39	51.8	66.0 77.0		61	7.1 yr	[35]
				Non-SVR	91		68.0			6.2 yr	

(continued)

Table 1
(continued).

Study name, year	Study design	Treatment details	Treatment duration	Group of virologic response	No. of patients	Mean age (yr)	Male (%)	HAI score	Cirrhosis (%)	Mean follow-up duration	References
Harada, 2014	Retrospective	peg-IFNα2b + RBV	24–72 wk	SVR	454	56.7	67.0			36.2 mo	[16]
				Relapse	191						
				NR	164						
Moon, 2015	Retrospective	Peg-IFNα + RBV	24 or 48 wk	SVR	300	50.9	47.0		9.7	Median: 37.0 mo median: 34.6 mo	[17]
				Non-SVR	163		49.1		32.5		

BR = breakthrough response, HAI = histology activity index, IFN = interferon, IR = incomplete response, NR = nonresponse, NVR = nonvirological response, peg-IFN = pegylated interferon, PR = partial response, RBV = ribavirin, SVR = sustained virological response, TR = transient response, TVR = transient virological response.

plus RBV therapy.^[13,15,16,41] A recent study reported that the risk of developing HCC in patients with HCV-related cirrhosis who achieve SVR can persist for up to 8 years.^[42] In this study, we used the data extracted from the included studies to do a subgroup analysis for risk identification based on <10 years and ≥10 years, and found that the pooled results were consistent across subgroups defined according to follow-up duration.

Addition of DAA therapy to peg-IFN plus RBV therapy was associated with higher SVR rates, and virological cures were observed after 12 weeks of DAA treatment even in cirrhotic patients who were previously nonresponders.^[43,44] The current DAA-based IFN-free regimens were also shown to achieve very high rates of SVR and less adverse effect events.^[19,20,44] The recent studies reported an unexpectedly high rate of tumor recurrence coinciding with HCV clearance after DAA therapy,^[45,46] and this was attributed to the significant difference in the kinetics of viral suppression between DAA-treated patients and patients treated with IFN-based regimens. Rapid viral clearance

after DAA therapy is accompanied by a reduction in inflammation signals, leading to abrogation of immune-mediated inhibition of tumor progression.^[46] The slower viral suppression seen with IFN-based therapies may be associated with IFN-mediated modulation of the immune system and promotion of antitumor immunity.^[40] But, recent meta-analysis studies found no evidence for differential HCC occurrence or recurrence risk following SVR from DAA therapy compared with IFN-based therapy among CHC patients with cirrhosis.^[47] Hence, DAA and IFN-based therapy associated with SVR and the risk of HCC require further investigation.

Although the current standard of care for CHC did not consist of peg-IFN plus RBV,^[20,44] our study provided exclusive information which SVR could help to predict HCC risk in CHC patients with cirrhosis. Our meta-analysis included only 5 studies that used this regimen, while the rest of the studies used IFN monotherapy. Data from our present meta-analysis, which compared the pooled OR of HCC between the SVR and non-SVR




Figure 2. Forest plots to determine the effect of sustained virological response on incidence of hepatocellular carcinoma. 95% CI = 95% confidence interval, IFN = interferon, OR = odds ratio.




Figure 3. Sensitivity-analysis for the effect of sustained virological response on incidence of hepatocellular carcinoma. (A) IFN + ribavirin; (B) IFN monotherapy; (C) other. 95% CI=95% confidence interval, IFN=interferon, OR=odds ratio.

groups from 20 studies, showed that non-SVR patients had greater odds of HCC incidence compared to SVR. This effect is likely due to a combination of decreased inflammation and necrosis, and virus eradication, factors that modulate the progression of liver disease in CHC patients.^[8,31] It will be interesting to investigate differences between IFN monotherapy

regimens and IFN plus RBV regimens on SVR rates and rates of HCC incidence. The role of IFN therapy in CHC patients on hepatocarcinogenesis remains pivotal.

We performed a subgroup analysis based on the country where the study was conducted. The Asia region included Japan,^[4,11,13,15,16,22,24-27,31,32,34] Taiwan,^[28-30] and Korea.^[17]




Figure 4. Subgroup analysis to evaluate the effect of sustained virological response on incidence of hepatocellular carcinoma according to the countries where studies were conducted. 95% CI=95% confidence interval, IFN=interferon, OR=odds ratio.

The non-Asia region included France,^[35] Italy,^[33] and Portugal.^[36] Our analysis showed that regardless of country, patients who failed to achieve SVR had greater odds of HCC occurrence compared to SVR patients. A number of the studies included in our meta-analysis compared survival/mortality rates in the SVR and non-SVR groups. Lack of SVR was shown to increase the risk of liver-related mortality in CHC patients who were treated with IFN therapy.^[25,26,32,33,36] A prospective study investigating the association between mortality and virologic

response to peg-IFN α and RBV reported that patients with SVR or TVR had a significantly lower mortality rate, and lower rates of decompensated liver disease and HCC compared to non-responders.^[41] SVR and TVR were also shown to correlate with decreased mortality rates in chronic HCV patients treated with peg-IFN α plus RBV therapy,^[13,29,35] suggesting that virologic response was a predictor of liver-related mortality. However, the heterogeneity of the data in the included studies made it a challenge to pool the survival data for this analysis. Sensitivity




Figure 5. Subgroup analysis to evaluate the effect of sustained virological response on incidence of hepatocellular carcinoma according to follow-up duration. 95% CI=95% confidence interval, IFN=interferon, OR=odds ratio.




Figure 6. Quality assessment.

analyses for the outcomes using the leave-one-out approach showed that no single study had a significant impact on the pooled results.

Eventhough IFN-free DAA therapy of CHC became mainstream, it seems does not alter the short-term risk for HCC in patients with liver cirrhosis.^[20,47] This meta-analysis notably focused on the incidence of HCC in patients who achieved SVR compared with the patients who did not achieve SVR after IFN therapy. Despite the strengths, there are several limitations associated with our study. First, this meta-analysis was that most of the included studies were retrospective. Only 4 prospective studies were included in the analysis. More well-designed prospective studies are needed to confirm our findings. Secondary, the response rates to IFN-based therapy vary significantly between the different HCV genotypes,^[48,49] we did not have enough information to perform a subanalysis based on HCV genotype, since only one of our included studies provided genotype information.^[30] Also, the results in some subgroups require cautious interpretation due to the heterogeneity observed across the studies included. Another important limitation was the lack of subgroup analysis of important HCC-related factors such as cirrhosis and age due to insufficient data. In future studies, it will be preferable to perform meta-analysis of individual patient data rather than aggregate-level data.

Finally, we did not include the studies published in non-English language.

5. Conclusion

Our data showed a significant correlation between SVR to IFN therapy and reduced incidence of HCC, and suggested that SVR could be as a predictor of HCC in CHC patients treated with IFN or IFN + RBV. These results have important implications for the decision-making process during HCC screening. CHC patients

who fail to achieve SVR after IFN therapy are at high risk for progressing to HCC, and require rigorous follow-up, while patients who achieve SVR would require less frequent screening.

Author contributions

Conceptualization: Yuan-Ti Lee.

Data curation: Chien-Feng Li, Shih-Ming Tsao.

Formal analysis: Hsien-Hua Liao, Shiu-an-Chih Chen, Shih-Ming Tsao.

Investigation: Chien-Feng Li.

Writing – original draft: Chien-Feng Li, Hsien-Hua Liao, Shiu-an-Chih Chen, Shih-Ming Tsao, Yuan-Ti Lee.

Writing – review & editing: Yuan-Ti Lee.

References

- [1] Geneva: World Health Organization (2017) Hepatitis C. Available at: <https://www.who.int/news-room/fact-sheets/detail/hepatitis-c> Accessed July 9, 2019.
- [2] Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- [3] Hayashi J, Furusyo N, Ariyama I, et al. A relationship between the evolution of hepatitis c virus variants, liver damage, and hepatocellular carcinoma in patients with hepatitis C viremia. *J Infect Dis* 2000;181:1523–7.
- [4] Takimoto M, Ohkoshi S, Ichida T, et al. Interferon inhibits progression of liver fibrosis and reduces the risk of hepatocarcinogenesis in patients with chronic hepatitis C: a retrospective multicenter analysis of 652 patients. *Dig Dis Sci* 2002;47:170–6.
- [5] Yoshida H, Arakawa Y, Sata M, et al. Interferon therapy prolonged life expectancy among chronic hepatitis C patients. *Gastroenterology* 2002;123:483–91.
- [6] Shen YC, Hsu C, Cheng CC, et al. A critical evaluation of the preventive effect of antiviral therapy on the development of hepatocellular carcinoma in patients with chronic hepatitis C or B: a novel approach by using meta-regression. *Oncology* 2012;82:275–89.

- [7] Hung HC, Liao HH, Chen SC, et al. Maintenance interferon therapy in chronic hepatitis C patients who failed initial antiviral therapy: a meta-analysis. *Medicine* 2019;98:e15563.
- [8] Singal AG, Volk ML, Jensen D, et al. A sustained viral response is associated with reduced liver-related morbidity and mortality in patients with hepatitis C virus. *Clin Gastroenterol Hepatol* 2010;8:280–8. 288.e1.
- [9] Morgan RL, Baack B, Smith BD, et al. Eradication of hepatitis c virus infection and the development of hepatocellular carcinoma: a meta-analysis of observational studies. *Ann Intern Med* 2013;158:329–37.
- [10] Pinzone MR, Zanghi AM, Rapisarda L, et al. Cirrhotic patients are still at risk of developing hepatocellular carcinoma despite Interferon-induced sustained virological response. *Eur Rev Med Pharmacol Sci* 2014;18:11–5.
- [11] Imai Y, Kawata S, Tamura S, et al. Relation of interferon therapy and hepatocellular carcinoma in patients with chronic hepatitis c. *Ann Intern Med* 1998;129:94–9.
- [12] European Association for the Study of the Liver. EASL Clinical Practice Guidelines: Management of hepatitis C virus infection. *J Hepatol* 2011;55:245–64.
- [13] Ogawa E, Furusyo N, Kajiwara E, et al. Efficacy of pegylated interferon alpha-2b and ribavirin treatment on the risk of hepatocellular carcinoma in patients with chronic hepatitis C: a prospective, multicenter study. *J Hepatol* 2013;58:495–501.
- [14] Kimer N, Dahl EK, Gluud LL, et al. Antiviral therapy for prevention of hepatocellular carcinoma in chronic hepatitis C: systematic review and meta-analysis of randomised controlled trials. *BMJ Open* 2012;2:e001313.
- [15] Dohmen K, Kawano A, Takahashi K, et al. The incidence and risk factors for the development of hepatocellular carcinoma after peginterferon plus ribavirin therapy for chronic hepatitis C. *Hepatogastroenterology* 2013;60:2034–8.
- [16] Harada N, Hiramatsu N, Oze T, et al. Risk factors for hepatocellular carcinoma in hepatitis C patients with normal alanine aminotransferase treated with pegylated interferon and ribavirin. *J Viral Hepat* 2015; 21:357–65.
- [17] Moon C, Jung KS, Kim DY, et al. Lower incidence of hepatocellular carcinoma and cirrhosis in hepatitis C patients with sustained virological response by pegylated interferon and ribavirin. *Dig Dis Sci* 2015;60: 573–81.
- [18] Honda T, Ishigami M, Masuda H, et al. Effect of peginterferon alfa-2b and ribavirin on hepatocellular carcinoma prevention in older patients with chronic hepatitis C. *J Gastroenterol Hepatol* 2015;30:321–8.
- [19] Geddawy A, Ibrahim YF, Elbahie NM, et al. Direct acting anti-hepatitis C virus drugs: clinical pharmacology and future direction. *J Transl Int Med* 2017;5:8–17.
- [20] AASLD-IDSA HCV Guidance Panel. Hepatitis C Guidance 2018 Update: AASLD-IDSA Recommendations for Testing, Managing, and Treating Hepatitis C Virus Infection. *Clin Infect Dis* 2018;67:1477–92.
- [21] Yilmaz H, Yilmaz EM, Leblebicioglu H. Barriers to access to hepatitis C treatment. *J Infect Dev Ctries* 2016;10:308–16.
- [22] Kobayashi S, Takeda T, Enomoto M, et al. Development of hepatocellular carcinoma in patients with chronic hepatitis C who had a sustained virological response to interferon therapy: a multicenter, retrospective cohort study of 1124 patients. *Liver Int* 2007;27:186–91.
- [23] Hayden JA, Côté P, Bombardier C. Evaluation of the quality of prognosis studies in systematic reviews. *Ann Intern Med* 2006;144: 427–37.
- [24] Kashiwagi K, Kubo N, Nakashima H, et al. A prospective comparison of the effect of interferon-alpha and interferon-beta treatment in patients with chronic hepatitis C on the incidence of hepatocellular carcinoma development. *J Infect Chemother* 2003;9:333–40.
- [25] Shiratori Y, Ito Y, Yokosuka O, et al. Antiviral therapy for cirrhotic hepatitis C: association with reduced hepatocellular carcinoma development and improved survival. *Ann Intern Med* 2005;142:105–14.
- [26] Tanaka H, Tsukuma H, Kasahara A, et al. Effect of interferon therapy on the incidence of hepatocellular carcinoma and mortality of patients with chronic hepatitis C: a retrospective cohort study of 738 patients. *Int J Cancer* 2000;87:741–9.
- [27] Kasahara A, Hayashi N, Mochizuki K, et al. Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. *Hepatology* 1998; 27:1394–402.
- [28] Yu ML, Dai CY, Chen SC, et al. High versus standard doses interferon-alpha in the treatment of naïve chronic hepatitis C patients in Taiwan: a 10-year cohort study. *BMC Infect Dis* 2005;5:27.
- [29] Yu ML, Lin SM, Chuang WL, et al. A sustained virological response to interferon or interferon/ribavirin reduces hepatocellular carcinoma and improves survival in chronic hepatitis C: a nationwide, multicentre study in Taiwan. *Antivir Ther* 2006;11:985–94.
- [30] Hung CH, Lee CM, Lu SN, et al. Long-term effect of interferon alpha-2b plus ribavirin therapy on incidence of hepatocellular carcinoma in patients with hepatitis C virus-related cirrhosis. *J Viral Hepat* 2006;13:409–14.
- [31] Ikeda K, Arase Y, Saitoh S, et al. Anticarcinogenic impact of interferon on patients with chronic hepatitis C: a large-scale long-term study in a single center. *Intervirology* 2006;49:82–90.
- [32] Arase Y, Ikeda K, Suzuki F, et al. Long-term outcome after interferon therapy in elderly patients with chronic hepatitis C. *Intervirology* 2007;50:16–23.
- [33] Bruno S, Stroffolini T, Colombo M, et al. Sustained virological response to interferon- α is associated with improved outcome in HCV-related cirrhosis: a retrospective study. *Hepatology* 2007;45:579–87.
- [34] Imai Y, Tamura S, Tanaka H, et al. Reduced risk of hepatocellular carcinoma after interferon therapy in aged patients with chronic hepatitis C is limited to sustained virological responders. *J Viral Hepat* 2010; 17:185–91.
- [35] Cardoso AC, Moucari R, Figueiredo-Mendes C, et al. Impact of peginterferon and ribavirin therapy on hepatocellular carcinoma: incidence and survival in hepatitis C patients with advanced fibrosis. *J Hepatol* 2010;52:652–7.
- [36] Velosa J, Serejo F, Marinho R, et al. Eradication of hepatitis C virus reduces the risk of hepatocellular carcinoma in patients with compensated cirrhosis. *Dig Dis Sci* 2011;56:1853–61.
- [37] Hoofnagle JH, Mullen KD, Jones DB, et al. Treatment of chronic non-A, non-B hepatitis with recombinant human alpha interferon. *N Engl J Med* 1986;315:1575–8.
- [38] Miyake Y, Iwasaki Y, Yamamoto K. Meta-analysis: reduced incidence of hepatocellular carcinoma in patients not responding to interferon therapy of chronic hepatitis C. *Int J Cancer* 2010;127:989–96.
- [39] Mallet V, Gilgenkrantz H, Serpaggi J, et al. Brief communication: the relationship of regression of cirrhosis to outcome in chronic hepatitis C. *Ann Intern Med* 2008;149:399–403.
- [40] Oh JK, Weiderpass E. Infection and cancer: global distribution and burden of diseases. *Ann Glob Health* 2014;80:384–92.
- [41] Morgan TR, Ghany MG, Kim HY, et al. Outcome of sustained virological responders with histologically advanced chronic hepatitis C. *Hepatology* 2010;52:833–44.
- [42] Aleman S, Rahbin N, Weiland O, et al. A risk for hepatocellular carcinoma persists long-term after sustained virologic response in patients with hepatitis C-associated liver cirrhosis. *Clin Infect Dis* 2013;57:230–6.
- [43] Hsu YC, Wu CY, Lin JT. Hepatitis C virus infection, antiviral therapy, and risk of hepatocellular carcinoma. *Semin Oncol* 2015;42:329–38.
- [44] Pawlotsky JM, Negro F, Aghemo A, et al. EASL recommendations on treatment of hepatitis C 2018. *J Hepatol* 2018;69:461–511.
- [45] Minami T, Tateishi R, Nakagomi R, et al. The impact of direct-acting antivirals on early tumor recurrence after radiofrequency ablation in hepatitis C-related hepatocellular carcinoma. *J Hepatol* 2016;65:1272–3.
- [46] Reig M, Mariño Z, Perelló C, et al. Unexpected high rate of early tumor recurrence in patients with HCV-related HCC undergoing interferon-free therapy. *J Hepatol* 2016;65:719–26.
- [47] Waziry R, Hajarizadeh B, Grebely J, et al. Hepatocellular carcinoma risk following direct-acting antiviral HCV therapy: a systematic review, meta-analyses, and meta-regression. *J Hepatol* 2017;67:1204–12.
- [48] Fried MW, Shiffman ML, Reddy KR, et al. Peginterferon Alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–82.
- [49] McHutchison JG, Lawitz EJ, Shiffman ML, et al. Peginterferon Alfa-2b or Alfa-2a with ribavirin for treatment of hepatitis C infection. *N Engl J Med* 2009;361:580–93.