

## CORRECTION

# Correction: The Presence of the Y-Chromosome, Not the Absence of the Second X-Chromosome, Alters the mRNA Levels Stored in the Fully Grown XY Mouse Oocyte

Baozeng Xu, Yayoi Obata, Feng Cao, Teruko Taketo

The probe ID numbers given in Table S1 and Table S2 are incorrect. The correct probe ID numbers are available from: <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE45668>.

## Reference

1. Xu B, Obata Y, Cao F, Taketo T (2012) The Presence of the Y-Chromosome, Not the Absence of the Second X-Chromosome, Alters the mRNA Levels Stored in the Fully Grown XY Mouse Oocyte. PLoS ONE 7(7): e40481. doi: [10.1371/journal.pone.0040481](https://doi.org/10.1371/journal.pone.0040481) PMID: [22792347](https://pubmed.ncbi.nlm.nih.gov/22792347/)



---

## OPEN ACCESS

**Citation:** Xu B, Obata Y, Cao F, Taketo T (2015) Correction: The Presence of the Y-Chromosome, Not the Absence of the Second X-Chromosome, Alters the mRNA Levels Stored in the Fully Grown XY Mouse Oocyte. PLoS ONE 10(11): e0143788. doi:10.1371/journal.pone.0143788

**Published:** November 23, 2015

**Copyright:** © 2015 Xu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.