



ARTICLE

The Ratio of Dietary n-3 Polyunsaturated Fatty Acids Influences the Fat Composition and Lipogenic Enzyme Activity in Adipose Tissue of Growing Pigs

Chang Hyun Song¹, Seung Min Oh², SuHyup Lee³, YoHan Choi⁴, Jeong Dae Kim¹, Aera Jang^{5,*}, and JinSoo Kim^{1,*}

¹College of Animal Life Sciences, Kangwon National University, Chuncheon 24341, Korea

²Gyeongbuk Livestock Research Institute, Yeongju 63052, Korea

³Department of Swine Science, Korea National College of Agriculture and Fisheries, Jeonju 54874, Korea

⁴Swine Division, National Institute of Animal Science, Rural Development Administration, Cheonan 31000, Korea

⁵Department of Applied Animal Science, BK21 Plus Program, Kangwon National University, Chuncheon 24341, Korea

OPEN ACCESS

Received October 9, 2019

Revised December 23, 2019

Accepted January 16, 2020

*Corresponding author : Aera Jang
Department of Applied Animal Science,
BK21 Plus Program, Kangwon National
University, Chuncheon 24341, Korea
Tel: +82-33-250-8643
Fax: +82-33-251-7719
E-mail: ajang@kangwon.ac.kr

JinSoo Kim
College of Animal Life Sciences,
Kangwon National University,
Chuncheon 24341, Korea
Tel: +82-33-250-8614
Fax: +82-33-259-5572
E-mail: kjs896@kangwon.ac.kr

*ORCID
Chang Hyun Song
<https://orcid.org/0000-0003-0431-3373>
Seung Min Oh
<https://orcid.org/0000-0001-8848-8028>
SuHyup Lee
<https://orcid.org/0000-0001-8996-3740>
YoHan Choi
<https://orcid.org/0000-0003-4710-4731>
Jeong Dae Kim
<https://orcid.org/0000-0001-8761-319X>
Aera Jang
<https://orcid.org/0000-0003-1789-8956>
JinSoo Kim
<https://orcid.org/0000-0002-9518-7917>

Abstract Currently, there is a growing interest among consumers in selecting healthier meat with a greater proportion of essential fatty acids (FA). This experiment was conducted to evaluate the role of different ratios of dietary n-6:n-3 on growth performance, FA profile of *longissimus dorsi* (LD), relative gene expression of cytokines, meat quality, and blood parameters in finishing pigs. A total of 108 finishing pigs was randomly allotted to three treatments including a control (basal diet) and low ratios (4:1 and 2:1) of n-6:n-3. The 4:1 and 2:1 diets decreased the overall stearic acid in LD. There were reductions in the content of stearic acid, palmitoleic acid, total saturated acid, and n-6:n-3 ratio of LD in pigs fed 4:1 and 2:1 diet compared with the control diet. The 4:1 and 2:1 diets increased the concentration of α -Linolenic acid and polyunsaturated FA in the LD of pigs. Acetyl-CoA carboxylase enzyme gene was down-regulated in pigs fed 2:1 diet compared with finishing pigs fed the control or 4:1 diets. The relative expression of hormone-sensitive lipase was increased in pigs fed 2:1 and 4:1 ratio diets. Lower total cholesterol of plasma was observed in finishing pigs fed 2:1 and 4:1 diets. The cooking loss ratio of meat was lower in pigs fed the 2:1 and 4:1 diets compared with the control diet. Pigs fed the 4:1 and 2:1 diets had greater final body weight. In conclusion, the 2:1 and 4:1 diets have the potential to increase the meat quality and growth performance of pigs.

Keywords linseed, unsaturated fatty acids, finishing pigs, meat quality

Introduction

Polyunsaturated fatty acids (PUFA) are well recognized essential fatty acids (FA) because they are the major components of cell membranes, which play an important role to control cardiovascular diseases in humans (Sargent, 1997; Sanders, 2000). The

correct balance of essential FA in the human diet can increase the consumption of these beneficial n-3 FA in order to enhance health status. Oily fishes are the main resource of PUFA (Howe et al., 2002; Sargent, 1997) but not among the popular foodstuffs in most of the countries. Pork is the most popular meat worldwide in terms of consumption rate, but mainly contains only a small amount of n-3 FA (Olsson and Pickova, 2005). A strategy to enhance the n-3 FA concentration of the diets would be an acceptable way to enrich the foodstuffs with high popularity such as pork. There has been a considerable increase in the n-6 consumption over the past decades due to a higher intake of vegetable oils from sunflower seeds, soybeans, and corn, which contain considerable amounts of n-6 PUFAs (Corino et al., 2002; Sanders, 2000). Furthermore, animal diets are based on grains with a high content of n-6 PUFAs, which led to the production of high n-6 content meat. In addition, animals are not able to produce n-3 PUFA by converting n-6 PUFA due to the lack of n-3 FA desaturase genes. It is logical to increase the ratio of essential n-3 FA in pork by providing high n-3 in pigs diet.

Linseed, as a good α -linolenic acid source, has gained attention as an alternative to fish oil in swine diets. The n-6:n-3 ratio in the food is reduced by supplementing linseed into pig feed, which possibly can reduce the n-6:n-3 ratio in the meat as well. However, the high content of unsaturated FA may decrease the meat quality by increasing in carcass fat softness and also changing the flavor and odor by decreasing the oxidative stability. It is well known that FA oxidation reaction in meat initiates by unsaturated FA (Wood et al., 2008). The oxidation reaction can adversely affect color, flavor, of meat (Corino et al., 2002; Howe et al., 2002). Thus, it seems essential to produce a diet containing an appropriate ratio of linseed over a period in order to improve the required requirement of n-3 in tissue without adverse effects on meat shelf-life, flavor and texture. The aim of this experiment was to evaluate the influences of different n-6:n-3 on the FA composition of adipose tissue, lipogenic enzyme activity, meat quality, and carcass characteristics.

Materials and Methods

The protocol for this study was approved by the ethics of the Institutional Animal Care and Use Committee of Kangwon National University. This study was conducted at the swine research station at the Kangwon National University, Chuncheon, Korea.

Animals and experimental design

A total of 108 finisher pigs (Landrace \times Yorkshire \times Duroc) was assigned to 3 dietary groups including control (basal diet; 18:1) and low ratios (4:1 and 2:1) of n-6:n-3 with average body weight (BW) of 84.2 ± 0.74 kg. The treatments were divided into 6 replicate pens of 6 pigs for each treatment. The pigs in all the treatments were fed isoenergetic diets with different ratios of n-6:n-3, prepared using 3.00%, 1.50%, and 0% of expended linseed to replace of animal fat to set the dietary n-6:n-3 ratios of the 3 diets about 18:1, 4:1, and 2:1 respectively. The experimental diets (meal form) were formulated to meet or exceed the current nutrient requirements for finisher pigs (NRC, 2012). Table 1 and Table 2 show the ingredient, chemical composition, and FA composition of the diets. The finisher pigs were housed (2.80 m \times 5.00 m) in partially slatted and concrete floor pens. Each pen was equipped with an automatic nipple waterer.

Experimental procedures, measurements, and analyses

All animals were weighed before being placed into pens, and daily feed intake (FI) was measured to calculate ADG, FI, and gain to feed ratio (GF). Relative gene expression levels of enzymes including hormone-sensitive lipase (HSL), acetyl

Table 1. Ingredient and chemical composition of experimental diets

Item	Control	Expanded linseed diet	
		4:1	2:1
Ingredients (%)			
Corn	58.79	58.54	58.29
Wheat	10.00	10.00	10.00
SBM (45%)	21.02	20.20	19.38
Rape seed meal	2.00	2.00	2.00
Animal fat	3.07	2.62	2.18
Linseed	-	1.50	3.00
Molasses	3.00	3.00	3.00
L-Lysine HCl (78%)	0.20	0.22	0.23
DL-Methionine (100%)	0.05	0.05	0.05
L-Threonine	0.05	0.05	0.06
Limestone	0.30	0.30	0.30
Tricalcium phosphate	0.92	0.92	0.91
Salt	0.35	0.35	0.35
Vitamin premix ¹⁾	0.10	0.11	0.11
Mineral premix ²⁾	0.10	0.10	0.10
Phytase	0.05	0.05	0.05
Chemical composition (%)			
ME (kcal/kg)	3,400	3,400	3,400
Crude protein	16.37	16.28	16.20
Lysine	0.97	0.97	0.97
Methionine+cysteine	0.60	0.61	0.61
Calcium	0.55	0.55	0.55
Available phosphorus	0.31	0.31	0.31

¹⁾ Supplied per kilogram of diet: 16,000 IU vitamin A, 3,000 IU vitamin D₃, 40 IU vitamin E, 5.0 mg vitamin K₃, 5.0 mg vitamin B₁, 20 mg vitamin B₂, 4 mg vitamin B₆, 0.08 mg vitamin B₁₂, 40 mg pantothenic acid, 75 mg niacin, 0.15 mg biotin, 0.65 mg folic acid.

²⁾ Supplied per kilogram of diet: 45 mg Fe, 0.25 mg Co, 50 mg Cu, 15 mg Mn, 25 mg Zn, 0.35 mg I, 0.13 mg Se.
SBM, soybean meal; ME, metabolizable energy.

CoA carboxylase (ACC), fatty acid synthase (FAS), and lipoprotein lipase (LPL) were investigated in FA tissue of two pigs per pen in the range of average BW on d 35 of study. About 50 g adipose tissue were separately collected from the *longissimus dorsi* (LD; 10–11th rib) of pigs and stored at -80°C to evaluate FA composition and gene expression of enzymes (HSL, ACC, FAS, and LPL).

Fatty acid composition

All FA samples from adipose tissues were converted to methyl esters as described previously (Lepage and Roy, 1986). The prepared methyl esters were analyzed to separate the FA by gas chromatography (Shimadzu, GC-17A, Kyoto, Japan). The initial temperature of oven was set at 175°C for a period of 30 min, constantly increased to 235°C . The temperature (260°C)

Table 2. Fatty acid composition of diets

Fatty acid composition (g/100 g)	Control	Expanded linseed diet	
		4:1	2:1
C8:0	0.42	0.32	0.20
C10:0	0.27	0.17	0.09
C12:0	0.29	0.24	0.28
C14:0	0.91	0.78	0.71
C16:0	19.24	18.64	17.70
C16:1n-9	1.58	1.68	1.68
C18:0	9.16	7.64	6.95
C18:1n-9	28.68	28.75	29.21
C18:2n-6	32.84	31.33	27.41
C18:3n-3	1.92	7.35	12.41
n-6 PUFA:n-3 PUFA	17.69	4.39	2.31

PUFA, poly unsaturated fatty acids.

of injector and detector was kept constant. The identification of FA methyl ester samples was operated by methyl ester standards.

RNA extraction of adipose tissue

Total ribonucleic acid (RNA) extraction was performed using TRIZOL reagent on adipose tissue samples (Invitrogen, Carlsbad, CA, USA) as suggested in the manufacturer's guideline. The ratio of absorption (260/280) for all samples was in a range of 1.6 to 1.8 using a Nanodrop 1000 (NanoDrop Technologies, Wilmington, DE, USA). Aliquot of polyadenylated RNA samples were electrophoresed to assess integrity. Real-time reverse transcription was operated by using ImProm-II™ kit (Promega, Madison, WI, USA). The reverse transcription was conducted on 1 µg of total RNA into cDNA in 20 µL reaction volume including 1 µL of random oligonucleotide primer. Reverse transcription was performed by a thermal program of 75°C for 5 min and 4°C for 5 min, followed by 25°C for 5 min, 42°C for 60 min, and 70°C for 15 min.

Real-time RT-PCR

The cDNA samples were mixed with 10.0 µL SYBR® Green quantitative real-time polymerase chain reaction (Toyobo, Osaka, Japan). The mixtures were used in the presence of 0.50 µL of forward and reverse primers (10 pmol/µL) for porcine LPL, FAS, ACC, and HPL, then were used for qPCR under a standard condition. As a control, the similar reverse reaction mixes were selected to PCR using the pairs of porcine 18S rRNA primers. The sequences of primers (Kim et al., 2014) are given in Table 3. A Real-time PCR (Bio-Rad, Hercules, CA, USA) was used for mixtures incubation to perform in 20 µL of total reaction volume, according to the manufacturer's guideline. An initial denaturation step PCR conditions were conducted at 95°C for 1 min, then 40 cycles of denaturation/primer annealing/elongation (at 95°C for 15 s, at 58°C for 30 s, and at 95°C for 15 s, respectively) with a final extension at 72°C for 10 min. Melting curve analysis was constructed on all reactions as a straightway for real-time PCR product specification and identification. Agarose gel electrophoresis was applied to confirm specificity. cDNA was synthesized, and control PCR assay was conducted in the absence of reverse transcriptase to excluded

Table 3. Specific primers used for real time quantitative PCR

Gene name	Sequence	Ta	Product size (BP)	Gene bank accession no
18S rRNA	Forward: 5-GCGGCTTTGGTGA CTCTA-3 Reverse: 5-CTGCCTCCTTGGATGTG-3	60	194	NR 002170.3
Acetyl CoA carboxylase	Forward: 5-ATG TTT CGG CAGTCC CTG AT-3 Reverse: 5-TGT GGA CCA GCTGAC CTT GA-3	60	133	EF618729
Fatty acid synthase	Forward: 5-AGC CTA ACT CCTCGC TGC AAT-3 Reverse: 5-TCC TTG GAA CCGTCT GTG TTC-3	58	196	AY183428
Lipoprotein lipase	Forward: 5-AAC TTG TGG CTGCCC TAT-3 Reverse: 5-GAC CCT CTG GTGAAT GTG-3	55	367	X62984
Hormone sensitive lipase	Forward: 5-GCT CCC ATC GTCAAG AAT C-3 Reverse: 5-TAA AGC GAA TGCGGT CC-3	55	262	AJ000482

RNA, ribonucleic acid.

contamination in the total RNA preparation. In all samples, no further bands were detected. $\Delta\Delta$ Ct method was used to analyze the results (fold changes). An 18S rRNA was considered as a ‘housekeeping gene’ for various gene expressions of the samples.

Hematological traits

On d 35, blood samples (10-mL from jugular vein puncture) was taken by a vacutainer tube coated with sodium heparin (Becton Dickinson, Franklin, NJ, USA) from the selected pigs (2 pigs per pen around the average BW). The plasma was separated by centrifugation (3,000×g for 15 min at 4°C) and stored immediately at –20°C until required for further analysis including total cholesterol, high-density lipoprotein (HDL), low-density lipoprotein (LDL), triglyceride, white blood cell, red blood cell, lymphocytes, and cortisol). Blood profile analyzed using blood analyzers (HEMAVET, Drew Scientific, Oxford, CT, USA).

Carcass and meat quality

The 12 finisher pigs from each treatment (around the average BW; 2 pigs per replicate) were weighed, then euthanized and prepared for carcass evaluation. Carcass samples were split down from the dorsal midline, weighed, and chilled overnight. After chilling at 4°C for 24 h, LD between the 10th and 11th ribs area and backfat thickness (10th rib; three-quarters of the distance along the LD toward the belly) were examined. After slaughtering, initial pH (pH 45 min) in the muscle (muscularis LD) was evaluated at the last rib by a pH meter (IstekNeoMet 77P, Istek Inc, Seoul, Korea). The ultimate pH was obtained at 24 h and 48 h after slaughter. Minolta CR-310 (Yasuda Seiko Instrument Co., Tokyo, Japan) was applied to evaluate the color score among visual scores, redness (a*), lightness (L*), and yellowness (b*) at 24 h of refrigerated storage (Mohammadi Gheisar et al., 2016). The drip loss measurement was carried out on the sample slices (approximately 100 g weight and 2.54 cm thickness). Slice samples were weighed and then kept in sealed polyethylene bags at 4±0.8°C. The samples were used for drip loss decantation, and after this, the samples were weighed again to measure final drip loss as a percentage (Shim et al., 2018). Cooking loss was evaluated by water cooking (in vacuum pack bags) of the meat samples (200 g and 2.54 cm thickness) at 81°C with an inner temperature of 73°C. Before weighing, the meats were chilled and kept at

25°C. The calculation of cooking loss was performed according to the weight of the initial sample as explained by Hosseindoust et al. (2016). To evaluate the shear force, porks were cooked at 80°C. After this, the porks were kept at 25°C. Five rectangular blocks cut (1 cm×1 cm) with parallel direction to the muscle fibers were divided from cooked samples. A texture analyzer was applied to measure the required force (TA-XT2i, Stable Microsystems Ltd., UK).

Statistical analysis

The experimental values were analyzed by GLM procedure of SAS Statistics (SAS Institute Inc., Cary, NC, US). The difference of means was tested by Tukey's multiple range test. A significant difference was expressed either $p < 0.01$ or $p < 0.05$, however p -values 0.05 to 0.1 were sometimes given to indicate if the values are tended to differ.

Results

Fatty acid composition

Among saturated FA (Table 4), no difference was detected for the content of lauric acid, myristic acid, and palmitic acid ($p < 0.05$). However, with the increase of n-6:n-3 ratio in the pigs diet, the content of stearic acid was increased ($p < 0.05$) in adipose tissue. Among monounsaturated FA, a decreased ($p < 0.01$) concentration of palmitoleic acid was observed in LD muscle of pigs fed linseed supplemented diets (ratios of 4:1 and 2:1). The concentration of oleic acid in adipose tissue was not influenced by the treatments. There was no significant difference between the treatments in the linoleic acid concentration of LD muscle. However, pigs fed dietary n-6:n-3 ratio of 4:1 and 2:1 had greater ($p < 0.05$) α -linolenic acid concentration of

Table 4. Effects of dietary n-6:n-3 ratio on fatty acid concentrations (g/100 g) of adipose tissue in finishing pigs

Characteristic	Control	n-6:n-3 ratio		SEM	p-value
		4:1	2:1		
Saturated fatty acids (SFA)					
Lauric acid (C12:0)	0.221	0.207	0.221	0.01	0.272
Myristic acid (C14:0)	2.185	2.095	2.12	0.03	0.087
Palmitic acid (C16:0)	22.94	22.32	21.26	0.46	0.059
Stearic acid (C18:0)	13.62 ^a	12.02 ^b	11.90 ^b	0.27	0.001
Monounsaturated fatty acids (MUFA)					
Palmitoleic acid (C16:1n-9)	2.51 ^a	2.32 ^b	2.26 ^b	0.04	0.002
Oleic acid (C18:1c-9)	40.17	39.59	38.82	0.70	0.371
Polyunsaturated fatty acids (PUFA)					
Linoleic acid (C18:2n-6)	13.55	14.24	14.75	0.42	0.159
α -Linolenic acid (C18:3n-3)	0.91 ^c	3.51 ^b	4.38 ^a	0.12	<0.001
Σ SFA	38.97 ^a	36.63 ^b	35.50 ^b	0.54	0.001
Σ MUFA	42.68	41.41	41.08	0.71	0.278
Σ PUFA	14.54 ^c	17.75 ^b	19.13 ^a	0.39	<0.001
n-6/n-3	15.81 ^a	4.06 ^b	3.41 ^b	1.09	<0.001

^{a-c} Means within a column with different letters are significantly different ($p < 0.05$).

LD muscle than the control pigs. The concentration of short-chain fatty acids (SFA) was decreased ($p<0.01$) in pigs fed linseed supplemented diets (ratios of 4:1 and 2:1). However, the total MUFA concentration was unaffected. Additionally, pigs fed dietary n-6:n-3 ratio of 4:1 and 2:1 had greater ($p<0.05$) concentration of PUFA and lower n-6:n-3 ratio than that of the control pigs.

Related gene expression of lipid metabolism enzymes

The expression of ACC, LPL, FAS, and HSL in adipose tissue is presented in Fig. 1. The relative expression level of the ACC gene was decreased ($p<0.05$) in pigs fed 4:1 or 2:1 diets compared with the control. However, no difference was shown for the relative expression of the FAS and LPL genes ($p<0.05$). The relative gene expression level of HSL was greater ($p<0.05$) in pigs fed 4:1 or 2:1 diets compared with the control.

Blood profile

In pigs fed 2:1 and 4:1 diets, total cholesterol levels increased (Table 5). However, the concentration of HDL, LDL, and triglyceride in blood were not affected by the treatments.

Meat quality

Carcass characteristics (dressing percentage, backfat thickness, and Loin eye area) were not influenced by different n-6:n-3




Fig. 1. Relative expression of (a) acetyl CoA carboxylase (ACC), (b) fatty acid synthase (FAS), (c) lipoprotein lipase (LPL), and (d) Hormone sensitive lipase (HSL) genes in adipose tissue of pigs with dietary n-6:n-3 ratio: animal fat (18:1), supplemented with 1.5% Expanded linseed (4:1), supplemented, supplemented with 3% Expanded linseed (2:1). ^{a,b} Means within a column with different letters are significantly different ($p<0.05$).

Table 5. Effect of dietary n-6:n-3 ratio on blood profiles in finishing pigs

	Control	n-6:n-3 ratio		SEM	p-value
		4:1	2:1		
Total cholesterol	94.50 ^a	87.50 ^b	85.33 ^b	1.41	0.006
HDL cholesterol	40.83	42.17	41.17	2.33	0.916
LDL cholesterol	52.67	51.67	50.83	2.16	0.837
Triglyceride	48.83	44.17	42.33	1.97	0.087

^{a,b} Means within a column with different letters are significantly different ($p < 0.05$). HDL, high-density lipoprotein; LDL, low-density lipoprotein.

ratios (Table 6). *Longissimus muscle* pH (pH at 45 min, 24 h, and 48 h post-slaughter) and color parameters (Lightness, redness, and yellowness) showed no difference between treatments. Meat drip loss and shear force were not influenced by the diets, however, meat cooking loss was decreased in linseed supplemented diets (2:1 and 4:1 ratios).

Growth performance

The ADG, FI, and GF of finishing pigs fed different dietary n-6:n-3 ratio is shown in Table 7. Supplementing the dietary n-6:n-3 ratios of 4:1 and 2:1 showed greater ($p < 0.05$) final BW and ADG than pigs fed the control diet. No significant variation was detected in FI and GF among the treatments.

Table 6. Effect of dietary n-6:n-3 ratio on carcass and meat quality in finishing pigs

	Control	n-6:n-3 ratio		SEM	p-value
		4:1	2:1		
Carcass characteristics					
Dressing percentage	78.09	78.29	78.47	0.48	0.862
Backfat thickness (mm)	23.37	23.45	23.12	0.37	0.806
Loin eye area (cm ²)	50.65	50.80	51.32	0.53	0.662
<i>Longissimus muscle</i> pH					
pH _{45 min} (at 45 min post-slaughter)	6.13	6.17	6.14	0.12	0.961
pH _{24 h} (at 24 h post-slaughter)	5.55	5.42	5.37	0.28	0.892
pH _{48 h} (at 48 h post-slaughter)	5.43	5.37	5.39	0.06	0.807
Color parameters					
Lightness (L*)	51.91	49.39	49.65	0.70	0.082
Redness (a*)	7.26	7.25	7.44	0.49	0.951
Yellowness (b*)	6.06	6.98	6.34	0.31	0.137
Water holding capacity					
Drip loss (%)	6.42	6.17	5.65	0.36	0.325
Cooking loss (%)	19.80 ^a	18.19 ^b	17.80 ^b	0.43	0.012
Shear force (kg/cm ²)	33.40	32.39	34.59	1.72	0.824

^{a,b} Means within a column with different letters are significantly different ($p < 0.05$).

Table 7. Effects of dietary n-6:n-3 ratio on average daily gain (ADG), average daily feed intake (FI) and gain to feed ratio (GF) in finishing pigs

	Control	n-6:n-3 ratio		SEM	p-value
		4:1	2:1		
Initial body weight (kg)	82.10	82.12	82.21	0.37	0.979
Final body weight (kg)	106.43 ^b	108.61 ^a	109.08 ^a	0.42	0.001
ADG (g)	695 ^b	757 ^a	768 ^a	11.77	0.001
FI (g)	2,482	2,562	2,714	71.84	0.099
GF	3.58	3.39	3.54	0.01	0.521

^{a,b} Means within a column with different letters are significantly different ($p < 0.05$).

Discussion

Several researchers have previously identified that the source of fat in the diet influences the composition of FA in tissues (Corino et al., 2002; Enser et al., 2000). Among the oil sources, it is believed that the supplementation of linseed oil changes the FA composition of adipose tissues in pigs (Guillevic et al., 2009; Kim et al., 2014; Wood et al., 2008). Therefore, accurate information about the ratio of FA in tissues and physical characteristics of meat are required. In this study, the 4:1 and 2:1 treatments showed a higher α -linolenic acid concentration nearly 4 fold in adipose tissue, whereas palmitoleic acid (C16:1c-9) concentration of SFA was reduced accordingly in comparison to pigs fed the control diet. In agreement, Leikus et al. (2018) reported an improvement in the content of essential FA in adipose tissues of finishing pigs fed linseed. Additionally, Duran-Montgé et al. (2009) indicated that the addition of linseed into diet showed a greater ratio of linolenic acid in adipose tissue of pigs. A higher ratio of PUFA and α -linolenic acid in adipose tissues of pigs fed 30 g/kg linseed may be attributed to the higher dietary content of PUFA. Because of the much higher α -linolenic acid concentration of adipose tissue in linseed supplemented diets, it can be reasonably postulated that this nutritional improvement may significantly affect human health. Therefore, it is possible to manipulate the composition of FA in adipose tissue even in a short period of 35 days. In our study, even the 4:1 ratio significantly decreased n6:n3 ratio. Our results show that linseed even at a low level (15 g/kg) can be added to the diet to alter the FA content of adipose tissue in meat to produce healthier food for consumers.

The rate of FA biosynthesis in adipose tissue can be alerted by regulating the expression of lipolytic enzymes (LPL and HSL) or lipogenic enzymes (FAS and ACC) (Clarke 1993; Zhao et al., 2010). Acetyl CoA carboxylase is a determinant factor in FA oxidation or synthesis through the *de-novo* pathway (Doran et al., 2006); whereas, FAS is a crucial factor to maximize the *de-novo* FA production in tissues (Bee 2001; Clarke 1993). This study showed that the change in dietary n6:n3 ratio did not affect the expression of FAS genes; however, low n6:n3 ratio diets down-regulated the ACC expression in adipose tissue. Duran-Montgé et al. (2009) reported that supplementing 100 g/kg linseed to pig diet showed a higher expression of ACC in adipose tissue compared to high dietary n6:n3 ratio, but no effect was observed on FAS gene expression in adipose tissue. The down-regulated ACC gene expression indicates that there is a reduced capacity of *de-novo* FA synthesis in pigs fed diets with a high n6:n3 ratio. In addition, LPL is a key enzyme for LDL and triglycerides hydrolysis in chylomicrons to utilize free FA in tissue; but HSL is associated with FA catabolism for oxidation and exportation of intracellular triacylglycerol (Kim et al., 2014). The current results obtained in this study indicated that the dietary supplementation of linseed resulted in an up-regulated expression of HSL in pigs, but the source of fat in the diet showed no change was the gene expression of LPL in adipose tissue. Therefore, it can be suggested that the reduction of dietary n6:n3 ratio increases the rate of free FA release in

adipose tissue and regulates the lipolysis and mobilization of stored FA in adipose tissues.

The result of the current study showed lower blood total cholesterol in pigs fed linseed (4:1 and 2:1 diets). The positive association between PUFA and blood total cholesterol in the present study is compatible with those of human studies that used high dietary PUFA to decrease blood cholesterol (Horrobin et al., 1983), although the results of adding linseed into the diet among animal-based studies have been inconsistent (Kouba et al., 2003). Therefore, the lower blood cholesterol may theoretically imply the better health status of pigs.

The present study showed no clear relationship between dietary linseed and carcass characteristics such as dressing percentage, backfat thickness, and loin eye area, but linseed supplementation to the diet significantly decreased cooking losses systematically across both level treatment groups. These effects of dietary linseed inclusion are in contrast to the results reported for gilts and barrows by Juárez et al. (2011) who showed no difference in pork cooking loss between treatments with different dietary linseed supplementation. The significant cooking loss differences for linseed-included diets were unexpected and no studies were showed similar results with similar dietary treatments for cooking loss parameter. The difference shown in this study was small, however, it would be expected to have a positive effect on meat quality. Overall, as there was no difference between treatments for the other meat quality characteristics such as meat color, pH, drip loss, and shear force, it can be concluded that dietary linseed at 3% had no adverse impact on the meat quality characteristics measured.

This study confirms the possibility of increasing n-3 PUFA content in adipose tissues with supplementation of linseed to the diets without any adverse influences on meat quality characteristics or gain to GF of finisher pigs. In addition, dietary supplementation of linseed showed even a greater final BW in pigs. These findings are in agreement with the reports of a previous study (Juárez et al., 2011), which showed an improved ADG and FI in pigs fed linseed. Conversely, several studies reported no influences of dietary linseed on ADG, FI, or GF of pigs (Guillevic et al., 2009; Kim et al., 2014; Upadhaya et al., 2015). These variations may be related to the differences in the composition of diet, feeding periods, or levels of linseed. Several studies have shown that the low levels of linseed (less than 5%) or short feeding duration (less than 60 d) may not improve growth performance (Guillevic et al., 2009; Kouba et al., 2003). This is in contrast to the result of the present study that showed a significantly greater ADG when 3% linseed (2:1) was added to the diet in a period of 35 days. A slightly greater FI in pigs fed linseed may imply that the palatability can also be considered as another effective factor in improving final growth performance.

Conclusions

Decreasing dietary linoleic acid to linolenic acid ratios (4:1 and 2:1) improves the growth performance, linolenic acid content of adipose tissue, and cooking loss. Therefore, linseed, as a dietary linolenic source, can be a suitable alternative to animal fat. Moreover, regarding the insignificant growth performance difference between 4:1 and 2:1 ratios, the 4:1 ratio can be recommended to reduce the cost of diet.

Conflicts of Interest

The authors declare no potential conflict of interest.

Acknowledgements

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry

(IPET) through Agri-Bio Industry Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (315022-3).

Author Contributions

Conceptualization: Song CH, Kim JD, Kim JS. Data curation: Song CH, Oh SM, Kim JS. Formal analysis: Lee HS, Jang A. Methodology: Song CH. Software: Lee HS, Choi YH, Kim JS. Validation: Oh SM. Investigation: Song CH, Kim JS. Writing - original draft: Kim JS. Writing - review & editing: Song CH, Oh SM, Lee SH, Choi YH, Kim JD, Jang A, Kim JS.

Ethics Approval

The experiment was approved by the Institutional Animal Care and Use Committee (IACUC: KW-170519-1), Kangwon National University (KNU), Korea.

References

- Bee G. 2001. Dietary conjugated linoleic acids affect tissue lipid composition but not *de novo* lipogenesis in finishing pigs. *Anim Res* 50:383-399.
- Clarke SD. 1993. Regulation of fatty acid synthase gene expression: An approach for reducing fat accumulation. *J Anim Sci* 71:1957-1965.
- Corino C, Magni S, Pagliarini E, Rossi R, Pastorelli G, Chiesa LM. 2002. Effects of dietary fats on meat quality and sensory characteristics of heavy pig loins. *Meat Sci* 60:1-8.
- Doran O, Moule SK, Teye GA, Whittington FM, Hallett KG, Wood JD. 2006. A reduced protein diet induces stearoyl-CoA desaturase protein expression in pig muscle but not in subcutaneous adipose tissue: Relationship with intramuscular lipid formation. *Br J Nutr* 95:609-617.
- Duran-Montgé P, Theil PK, Lauridsen C, Esteve-Garcia E. 2009. Dietary fat source affects metabolism of fatty acids in pigs as evaluated by altered expression of lipogenic genes in liver and adipose tissues. *Animal* 3:535-542.
- Enser M, Richardson RI, Wood JD, Gill BP, Sheard PR. 2000. Feeding linseed to increase the n-3 PUFA of pork: Fatty acid composition of muscle, adipose tissue, liver and sausages. *Meat Sci* 55:201-212.
- Guillevic M, Kouba M, Mourot J. 2009. Effect of a linseed diet or a sunflower diet on performances, fatty acid composition, lipogenic enzyme activities and stearoyl-CoA-desaturase activity in the pig. *Livest Sci* 124:288-294.
- Horrobin DF, Manku MS. 1983. How do polyunsaturated fatty acids lower plasma cholesterol levels? *Lipids* 18:558-562.
- Hosseindoust A, Park JW, Kim IH. 2016. Effects of *Bacillus subtilis*, Kefir and β -glucan supplementation on growth performance, blood characteristics, meat quality and intestine microbiota in broilers. *Korean J Poult Sci* 43:159-167.
- Howe PR, Downing JA, Grenyer BF, Grigonis-Deane EM, Bryden WL. 2002. Tuna fishmeal as a source of DHA for n-3 PUFA enrichment of pork, chicken, and eggs. *Lipids* 37:1067-1076.
- Juárez M, Dugan MER, Aldai N, Aalhus JL, Patience JF, Zijlstra RT, Beaulieu AD. 2011. Increasing omega-3 levels through dietary co-extruded flaxseed supplementation negatively affects pork palatability. *Food Chem* 126:1716-1723.
- Kim JS, Ingale SL, Lee SH, Choi YH, Kim EH, Lee DC, Kim YH, Chae BJ. 2014. Impact of dietary fat sources and feeding level on adipose tissue fatty acids composition and lipid metabolism related gene expression in finisher pigs. *Anim Feed*

Sci Technol 196:60-67.

- Kouba M, Enser M, Whittington FM, Nute GR, Wood JD. 2003. Effect of a high-linolenic acid diet on lipogenic enzyme activities, fatty acid composition, and meat quality in the growing pig. *J Anim Sci* 81:1967-1979.
- Leikus R, Juskiene V, Juska R, Juodka R, Stankeviciene D, Nainiene R, Siukscius A. 2018. Effect of linseed oil sediment in the diet of pigs on the growth performance and fatty acid profile of meat. *Rev Bras Zootec* 47:e20170104.
- Lepage G, Roy CC. 1986. Direct transesterification of all classes of lipids in a one-step reaction. *J lipid Res* 27:114-120.
- Mohammadi Gheisar M, Hosseindoust A, Kim IH. 2016. Effects of thermo-resistant non-starch polysaccharide degrading multi-enzyme on growth performance, meat quality, relative weights of body organs and blood profile in broiler chickens. *J Anim Physiol Anim Nutr* 100:499-505.
- National Research Council [NRC]. 2012. Nutrient requirements of swine. 11th ed. National Academy Press, Washington, DC, USA. p 420.
- Olsson V, Pickova J. 2005. The influence of production systems on meat quality, with emphasis on pork. *AMBIO: J Hum Environ* 34:338-343.
- Sanders TA. 2000. Polyunsaturated fatty acids in the food chain in Europe. *Am J Clin Nutr* 71:176S-178S.
- Sargent JR. 1997. Fish oils and human diet. *Br J Nutr* 78:S5-S13.
- Shim Y, Kim J, Hosseindoust A, Choi Y, Kim M, Oh S, Ham H, Kumar A, Kim K, Jang A, Chae B. 2018. Investigating meat quality of broiler chickens fed on heat processed diets containing corn distillers dried grains with solubles. *Korean J Food Sci Anim Resour* 38:629-635.
- Upadhaya SD, Kim JC, Mullan BP, Pluske JR, Kim IH. 2015. Vitamin E and omega-3 fatty acids independently attenuate plasma concentrations of proinflammatory cytokines and prostaglandin E2 in *Escherichia coli* lipopolysaccharide-challenged growing-finishing pigs. *J Anim Sci* 93:2926-2934.
- Wood JD, Enser M, Fisher AV, Nute GR, Sheard PR, Richardson RI, Hughes SI, Whittington FM. 2008. Fat deposition, fatty acid composition and meat quality: A review. *Meat Sci* 78:343-358.
- Zhao S, Wang J, Song X, Zhang X, Ge C, Gao S. 2010. Impact of dietary protein on lipid metabolism-related gene expression in porcine adipose tissue. *Nutr Metab* 7:6.