


 Cite this: *RSC Adv.*, 2021, 11, 5451

 Received 11th November 2020  
 Accepted 23rd January 2021

DOI: 10.1039/d0ra09588e

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

## Cobalt-catalyzed annulation of styrenes with $\alpha$ -bromoacetic acids†

 Tung T. Nguyen,<sup>ID</sup>\*<sup>ab</sup> Bao H. T. Ngo,<sup>ab</sup> Huy X. Le,<sup>ab</sup> Linh N. P. Vu,<sup>ab</sup> Tuong A. To,<sup>ID</sup><sup>ab</sup>  
 Anh N. Q. Phan<sup>ab</sup> and Nam T. S. Phan<sup>ID</sup>\*<sup>ab</sup>


We report a method for addition of  $\alpha$ -bromophenylacetic acids to vinyl C=C bonds in styrenes to afford  $\gamma$ -lactones. Reactions employed a simple cobalt catalyst  $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  in the presence of dipivaloylmethane (dpm) ligand. Many functionalities including halogen, ester, and nitro groups were compatible with reaction conditions. If  $\alpha$ -bromoesters were used, vinylacetates were the major products.

$\gamma$ -Lactones are ubiquitously found in many natural products and bioactive molecules.<sup>1</sup> Among the available synthetic methods to obtain these five-membered rings, those using transition metals for functionalization of unsaturated carbon-carbon bonds, such as alkenes and alkynes, with oxygen-containing nucleophiles have been substantially developed.<sup>2</sup> Notable reports have used scarce noble metals and/or photo-redox conditions.<sup>3</sup> Some examples employing abundant first-row metals to obtain  $\gamma$ -lactones from  $\alpha$ -bromoacetic acids or isosteric esters have been reported.<sup>4</sup> A seminal method for copper(0)-promoted hydrocarboxylation of alkenes with  $\alpha$ -iodocarboxylates was introduced by Metzger and Mahler.<sup>4a</sup> Following that, copper(i) salts have been developed for catalytic annulation of  $\alpha$ -bromoacetamides with styrenes.<sup>4b,4c</sup> Methods for using low valent complexes of other non-noble metals such as those of Ni(0) or Fe(0) are known as well.<sup>4d,4e</sup> It should be noted that the reported transformations were presumably triggered by a homolytic cleavage of carbon-halogen bonds in  $\alpha$ -halocarboxylates.<sup>4d</sup> We envisioned that such an activation could be obtained if a suitable cobalt complex is used.

Using cobalt adducts in the organic transformations that included alkyl halides was known.<sup>5</sup> Most of the known examples involved the formation of cobalt(i) species, thus usually suffering from the presence of strong reductants and/or tailored ligands. Meanwhile, a direct elaboration of cobalt(ii) complexes for incorporation of alkyl electrophiles is much rarer. It was hypothesized that a Co(ii)/Co(iii) catalytic cycle could be exploited since single-electron-transfer oxidation of Co(ii) was feasible.<sup>6</sup> Herein we report our attempts to annulate styrenes with  $\alpha$ -bromophenylacetic acids for synthesis of  $\gamma$ -lactones

using a simple cobalt(ii) catalyst (Scheme 1). The method has allowed for the use of  $\alpha$ -halocarboxylic acids that are not limited to those containing quaternary centers.<sup>4d,4e</sup>

We firstly investigated the coupling of diphenylethylene **1a** and  $\alpha$ -bromophenylacetic acid **2a** to afford the  $\gamma$ -lactone **3aa**. Brief results and control experiments are shown in Table 1.<sup>7</sup> Optimization was carried out with respect to cobalt salt, ligand, base, and solvent. Using  $\text{Co}(\text{acac})_2$  as catalyst for the annulation afforded **3aa** in 69% yield (entry 1). If a cobalt(iii) precursor was employed, a trace amount of the lactone was detected (entry 2). Since  $\text{Co}(\text{OAc})_2$  was inferior to  $\text{Co}(\text{acac})_2$  (entry 3), our hypothesis was that 1,3-diketone may play an important role in the coupling.<sup>8</sup> Running the reaction with  $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  catalyst and acetylacetone ligand (**L1**) furnished **3aa** in a comparable yield with respect to that of  $\text{Co}(\text{acac})_2$  (entry 4). A better yield of the product was observed if dipivaloylmethane (dpm) ligand (**L2**) was used (entry 5). Only 27% yield of **3aa** was obtained in case of an electron-poor 1,3-diketone (**L3**, entry 6). An unsymmetrical ligand bearing a phenyl moiety (**L4**) was also attempted, furnishing the lactone **3aa** in moderate yield (entry 7). Sodium carbonate was an inferior base to potassium carbonate (entry 8). To our surprise, adding a stoichiometric amount of DMF helped to increase the yield of **3aa** (entry 9). 1,4-Dioxane could be used as the solvent for this coupling (entry 10). Notably, a small amount of by-product derived from **2a** and  $\text{CH}_3\text{CN}$  could be observed. Thus, pivalonitrile solvent was




**Scheme 1** Metal-catalyzed synthesis of  $\gamma$ -lactones via activation of  $\alpha$ -bromoacetic acids.

<sup>a</sup>Faculty of Chemical Engineering, Ho Chi Minh City University of Technology (HCMUT), 268 Ly Thuong Kiet, District 10, Ho Chi Minh City, Vietnam. E-mail: tungtn@hcmut.edu.vn; ptsnam@hcmut.edu.vn


<sup>b</sup>Vietnam National University Ho Chi Minh City, Linh Trung Ward, Thu Duc District, Ho Chi Minh City, Vietnam

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra09588e



Table 1 Studies of reaction conditions<sup>a</sup>


Entry	[Co]	Base	Solvent	Yield of 3aa (%)
1	Co(acac) <sub>2</sub>	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	69
2	Co(acac) <sub>3</sub>	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	Trace
3	Co(OAc) <sub>2</sub>	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	57
4 <sup>b</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	70
5 <sup>c</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	78
6 <sup>d</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	27
7 <sup>e</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	52
8 <sup>c</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	Na <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	67
9 <sup>c,f</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	84
10 <sup>c,f</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	1,4-Dioxane	75
11 <sup>c,f</sup>	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	K <sub>2</sub> CO <sub>3</sub>	(CH <sub>3</sub> ) <sub>3</sub> CCN	91




<sup>a</sup> **1a** (0.1 mmol), **2a** (0.2 mmol), [Co] catalyst (0.02 mmol), ligand (0.05 mmol), base (0.1 mmol), solvent (1 mL), at 140 °C for 24 h. Yields of **3aa** are GC yields using diphenyl ether internal standard. <sup>b</sup> **L1** was used. <sup>c</sup> **L2** was used. <sup>d</sup> **L3** was used. <sup>e</sup> **L4** was used. <sup>f</sup> DMF (0.1 mmol) was added. Abbreviation: acac = acetylacetonate.

attempted to avoid such addition (entry 11), affording 91% yield of the lactone **3aa**.

Scope of the reaction with respect to reagents containing C=C bonds next studied. The results are shown in Scheme 2. Notably, some products were obtained in excellent diastereoselectivities in comparison with two examples reported by Nishihara and Iwasaki.<sup>4e</sup> A triphenyl-substituted  $\gamma$ -lactone **3aa** was isolated in 85% yield. The simplest styrene coupled with  $\alpha$ -bromophenylacetic acid to furnish **3ba** in 81% yield. Functionalities including bromo (**3ca**), chloro (**3da**), ester (**3ea**), and nitro (**3fa**, **3ga**) groups were compatible with reaction conditions. Some vinyl arenes containing polyaromatics were also attempted. 2-Vinyl naphthalene was reacted to give the lactone **3ha** in 88% yield. Cyclization of 9-vinylanthracene yielded a trace amount of the desired product, somewhat implying the importance of steric effect. Heterocycles, such as thiazole (**3ia**), were competent substrates. Other  $\alpha$ -bromoarylacetic acids such as those containing chloro substituents could successfully annulate with C=C bonds. Thus, the lactones (**3hb**, **3ac**) were obtained in moderate to good yields. At this moment, very electron-poor styrenes (*i.e.* 3,5-bis(trifluoromethyl)styrene) or internal alkenes (*i.e.* indene) were not tolerant of reaction conditions.


To show the generality of our method, annulation of styrenes with  $\alpha$ -bromoisobutyric acid **4** was attempted. The conditions




Scheme 2 Annulation of styrenes and  $\alpha$ -bromo arylacetic acids. Conditions: **1a–1j** (0.5 mmol), **2a–2c** (1.0 mmol), Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O (0.1 mmol), **L2** (0.25 mmol), K<sub>2</sub>CO<sub>3</sub> (0.5 mmol), DMF (0.5 mmol), (CH<sub>3</sub>)<sub>3</sub>CCN (5 mL), at 140 °C for 24 h. Yields are isolated yields. Diastereomeric ratios (d.r.) were determined using <sup>1</sup>H NMR spectrum.

were relatively milder than those of  $\alpha$ -bromophenylacetic acids, presumably due to the presence of the quaternary carbon that help cyclization step. Scope of styrenes is presented in Scheme 3. An excellent yield of the lactone (**5a**) was obtained in case of diphenylethylene substrate. Reactions of molecules containing bromo (**5c**), ester (**5e**), and nitro (**5f**) substituents occurred without any difficulties. Similar to the iron-catalyzed method,<sup>4e</sup> 2-vinyl naphthalene was suitable for the annulation (**5h**). Vinyl heteroarenes were competent substrates (**5i**). Electron-rich arenes were also attempted, furnishing the lactones in good yields (**5k**, **5l**).

Using  $\alpha$ -bromoacetates to couple with styrenes could yield lactones.<sup>4d,4e</sup> However, only vinylacetates **7a–7c** were obtained in our conditions if similar substrates were employed (Scheme 4). The results were indicative of a slow hydrolysis during reaction course. It should be noted that the presence of quaternary carbons in esters was crucial to afford the addition products.




**Scheme 3** Annulation of styrenes and  $\alpha$ -bromoisobutyric acid. Conditions: styrenes (0.5 mmol), **4** (1.0 mmol),  $\text{Co(NO}_3\text{)}_2\cdot 6\text{H}_2\text{O}$  (0.1 mmol), **L2** (0.25 mmol),  $\text{K}_2\text{CO}_3$  (0.5 mmol),  $(\text{CH}_3\text{)}_3\text{CCN}$  (5 mL), at 100 °C for 24 h. Yields are isolated yields.



**Scheme 4** Scope of  $\alpha$ -bromoacetates. Conditions: **1a** (0.5 mmol),  $\alpha$ -bromoacetates **6a-6c** (1.0 mmol),  $\text{Co(NO}_3\text{)}_2\cdot 6\text{H}_2\text{O}$  (0.1 mmol), **L2** (0.25 mmol),  $\text{K}_2\text{CO}_3$  (0.5 mmol), DMF (0.5 mmol),  $(\text{CH}_3\text{)}_3\text{CCN}$  (5 mL), at 140 °C for 24 h. Yields are isolated yields.

Based on previous studies,<sup>4d,4e</sup> a plausible mechanism was proposed as that shown in Scheme 5. Homolytic cleavage of  $\text{C}(\text{sp}^3)\text{-Br}$  bond in **2a** would afford the benzyl radical **8**. Radical addition to **1a** generated **9** followed by an oxidation that could afford a benzylic cation **10**. The resulting species underwent

cyclization, in the presence of base, to eventually furnish the lactone **3aa**. To support the hypothesis, some control experiments were carried out. Firstly, addition of radical quenchers such as TEMPO or BHT stopped the annulation, somewhat confirming the involvement of radical species during the



Scheme 5 Mechanistic considerations.

reaction course. Secondly, running the reaction in absence of **1a** gave an anhydride **11** (detected by GC-MS). Such the dead-end intermediate could be obtained *via* a homo-combination of the radical **8** followed by a dehydration.

## Conclusions

In conclusion, we have developed a general method to couple styrene C=C bonds with  $\alpha$ -bromoarylacetic acids to afford  $\gamma$ -lactones. The transformation employed a simple cobalt precursor  $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  facilitated by dipivaloylmethane ligand. Reaction conditions were well compatible with a diversified array of functional groups. Preliminary control experiments supported a radical-involved mechanism that could be triggered by the cobalt-catalyzed activation of carbon–bromide bond.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We would like to thank the Vietnam National Foundation for Science and Technology Development (NAFOSTED) for supporting this research (under grant number 104.01-2019.44).

## Notes and references

- (a) E. J. Kang and E. Lee, *Chem. Rev.*, 2005, **105**, 4348; (b) G. Lin, S. S. K. Chan, H. S. Chung and S. L. Li, *Stud. Nat. Prod. Chem.*, 2005, **32**, 611.
- N. T. Patil, R. D. Kavthe and V. S. Shinde, *Tetrahedron*, 2012, **68**, 8079.
- (a) N. G. Kundu and M. Pal, *J. Chem. Soc., Chem. Commun.*, 1993, 86; (b) H.-Y. Liao and C.-H. Cheng, *J. Org. Chem.*, 1995, **60**, 3711; (c) R. C. Larock, M. J. Doty and X. Han, *J. Org. Chem.*, 1999, **64**, 8770; (d) X.-J. Wei, D.-T. Yang, L. Wang, T. Song, L.-Z. Wu and Q. Liu, *Org. Lett.*, 2013, **15**, 6054; (e) I. Triandafillidi, M. G. Kokotou and C. G. Kokotos, *Org. Lett.*, 2018, **20**, 36.
- (a) J. O. Metzger and R. Mahler, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 902; (b) T. Nishikata, K. Itonaga, N. Yamaguchi and M. Sumimoto, *Org. Lett.*, 2017, **19**, 2686; (c) Y. Lv, W. Pu, S. Mao, X. Ren, Y. Wu and H. Cui, *RSC Adv.*, 2017, **7**, 41723; (d) D. Liu, S. Tang, H. Yi, C. Liu, X. Qi, Y. Lan and A. Lei, *Chem. Eur. J.*, 2014, **20**, 15605; (e) M. Iwasaki, N. Miki, Y. Ikemoto, Y. Ura and Y. Nishihara, *Org. Lett.*, 2018, **20**, 3848; (f) B. Wei, K.-W. Li, Y.-C. Wu, S.-Q. Tong and R.-J. Song, *Synthesis*, 2020, **52**, 3855.
- (a) A. Guérinot and J. Cossy, *Acc. Chem. Res.*, 2020, **53**, 1351; (b) K. Gao and N. Yoshikai, *Acc. Chem. Res.*, 2014, **47**, 1208.

- 6 For recent examples, see:(a) L. Grigorjeva and O. Daugulis, *Angew. Chem., Int. Ed.*, 2014, **53**, 10209; (b) S. Maity, R. Kancherla, U. Dhawa, E. Hoque, S. Pimparkar and D. Maiti, *ACS Catal.*, 2016, **6**, 5493.
- 7 Please see the ESI† for full optimization studies.
- 8 For recent examples of the cobalt complex ligated by dipivaloylmethane, see:(a) T. T. Nguyen, L. Grigorjeva and O. Daugulis, *ACS Catal.*, 2016, **6**, 551; (b) L. Lukasevics, A. Cizikovs and L. Grigorjeva, *Org. Lett.*, 2020, **22**, 2720.