Erratum for Mirhendi et al., the first case of onychomycosis in a koala (*Phascolarctos cinereus*) due to atypical isolates of *Microsporum gypseum*, a diagnostic challenge

Mirhendi H^{1, 2*}, Nishiyama Y³, Rezaei-Matehkolaei A⁴, Satoh K⁵, Makimura K⁵

ERRATUM

Volume 2, no. 2, 45-50, 2016, DOI: 10.18869/acadpub.cmm.2.2.2 [1]. The original version of this article contained incorrect dose of terbinafine. This erratum adds the correct dose as shown below. Terbinafine dosage should change from 250 mg/kg to 250 mg/day.

▶ How to cite this paper

Mirhendi H, Nishiyama Y, Rezaei-Matehkolaei A, Satoh K, Makimura K. The first case of onychomycosis in a koala (*Phascolarctos cinereus*) due to atypical isolates of *Microsporum gypseum*: a diagnostic challenge. Curr Med Mycol. 2018; 4(1): 35. DOI: 10.18502/cmm.4.1.34

References

1. Mirhendi H, Nishiyama Y, Rezaei-Matehkolaei A, Satoh K, Makimura K. The first case of onychomycosis in a koala (*Phascolarctos cinereus*) due to atypical isolates of *Microsporum gypseum*: a diagnostic challenge. Curr Med Mycol. 2016; 2(2): 45-50. DOI: 10.18869/acadpub.cmm.2.2.2

¹ Department of Medical Parasitology & Mycology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

² Department of Medical Parasitology & Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³ Teikyo University Institute of Medical Mycology, Tokyo, Japan

⁴ Department of Medical Mycology, School of Medicine, Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁵ General Medical Education and Research Center, Teikyo University, Tokyo, Japan