

Prostaglandin E1 plus methylcobalamin combination therapy versus prostaglandin E1 monotherapy for patients with diabetic peripheral neuropathy

A meta-analysis of randomized controlled trials

De-Qi Jiang, PhD^a, Shi-Hua Zhao, MSc^a, Ming-Xing Li, MSc^b, Li-Lin Jiang, BS^a, Yong Wang, PhD^c, Yan Wang, MSc^{d,*}

Abstract

Background: Prostaglandin E1 (P) or methylcobalamin (M) treatment has been suggested as a therapeutic approach for diabetic peripheral neuropathy (DPN) in many clinical trial reports. However, the combined effects of 2 drugs still remain dubious.

Objective: The aim of this report was to evaluate the efficacy of M plus P (M+P) for the treatment of DPN compared with that of P monotherapy, in order to provide a reference resource for rational drug use.

Methods: Randomized controlled trials (RCTs) of M+P for DPN published up to September 2017 were searched. Risk ratio (RR), mean difference (MD), and 95% confidence interval (CI) were calculated and heterogeneity was assessed with the I^2 test. Subgroup and sensitivity analyses were also performed. The outcomes measured were as follows: the clinical efficacy, median motor nerve conduction velocities (MNCV), median sensory nerve conduction velocity (SNCV), peroneal MNCV, peroneal SNCV, and adverse effects.

Results: Sixteen RCTs with 1136 participants were included. Clinical efficacy of M+P combination therapy was significantly better than P monotherapy (fifteen trials; RR 1.25, 95% CI 1.18–1.32, $P < .00001$, $I^2 = 27\%$). Compared with P monotherapy, the pooled effects of M+P combination therapy on nerve conduction velocity were (MD 6.29, 95% CI 4.63–7.94, $P < .00001$, $I^2 = 90\%$) for median MNCV, (MD 5.68, 95% CI 3.53–7.83, $P < .00001$, $I^2 = 94\%$) for median SNCV, (MD 5.36, 95% CI 3.86–6.87, $P < .00001$, $I^2 = 92\%$) for peroneal MNCV, (MD 4.62, 95% CI 3.48–5.75, $P < .00001$, $I^2 = 86\%$) for peroneal SNCV. There were no serious adverse events associated with drug intervention.

Conclusions: M+P combination therapy was superior to P monotherapy for improvement of neuropathic symptoms and NCVs in DPN patients. Moreover, no serious adverse events occur in combination therapy.

Abbreviations: CI = confidence interval, DM = diabetes mellitus, DPN = diabetic peripheral neuropathy, FE = fixed-effect, M = methylcobalamin, MNCV = motor nerve conduction velocity, P = prostaglandin E1, RCTs = randomized controlled trials, RR = risk ratio, SNCV = sensory nerve conduction velocity.

Keywords: diabetic peripheral neuropathy, efficacy, meta-analysis, methylcobalamin, nerve conduction velocity, prostaglandin E1

Editor: Zelena Dora.

This study was supported by grants from the Doctoral Scientific Research Foundation of Yulin Normal University of China (No. G2016006).

The authors have no conflicts of interest to disclose.

^a College of Biology and Pharmacy, Guangxi Key Laboratory of Agricultural Resources Chemistry and Biotechnology, Yulin Normal University, Yulin,

^b Department of Pharmacy, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou, ^c Department of Pharmacy, Zhujiang Hospital of Southern Medical University, Guangzhou, ^d Department of Pharmacy, Guangdong Province Hospital of Integrated Traditional Chinese and Western Medicine, Foshan, China.

* Correspondence: Yan Wang, No.16 5th Road, GuiCheng street South, Nanhai, Foshan, Guangdong Province, 528200, China (e-mails: wy6507364@163.com, yongwh2005@hotmail.com).

Copyright © 2018 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Medicine (2018) 97:44(e13020)

Received: 31 May 2018 / Accepted: 8 October 2018

<http://dx.doi.org/10.1097/MD.0000000000013020>

1. Introduction

As one of the most common complications of diabetes mellitus (DM), diabetic peripheral neuropathy (DPN) carries complicated pathogenesis which mainly lies on microcirculatory disturbance caused by impaired endothelial function,^[1] while the endothelial dysfunction can be aggravated by elevated advanced glycation end products resulted from long-term hyperglycemia.^[2] Therefore, peripheral neuropathy occurred at 5 to 10 years after the onset of type 2 diabetes, the atherosclerosis of major vessels, especially in the lower extremity arteries, also appeared in diabetic patients.^[1] Which is to say, a vicious circle of “ischemia-inflammation” is set up, then, the onset and progression of DPN are accelerated.^[3,4] Current therapeutic options for the treatment of DPN include good glycaemic control, nerve nurturing, oxidative-stress suppressing, microcirculation improving and others.^[5] As there is no effective single therapy currently existing for DPN, combination therapy with multiple drugs is generally performed. Clinically, prostaglandin E1 (P) is mainly used to relax vessels, decrease hematic viscosity, inhibit platelet aggregation and improve microcirculation.^[6] Methylcobalamin (M), a kind of endogenous coenzyme, promotes axon regeneration and

myelinogenesis by increasing nucleic acid, protein and phosphatidylcholine synthesis.^[7,8] M also can speed up nerve conduction velocities (NCVs) through accelerating delayed nerve impulse conduction.^[7,9] By far, clinical performances of the 2 medications in the treatment of DPN have been demonstrated by various studies.^[7,10,11]

Deng H et al suggested that treatment with M plus P (M + P) for patients with DPN was safe and could gain better outcomes in neuropathic symptoms and NCVs compared with M alone by meta-analysis.^[12] Furthermore, compared with P monotherapy, the efficacy and safety of M+P combination therapy have also been explored by numerous studies in mainland China.^[13–15] In order to understand the effects of M+P for DPN comprehensively, the present meta-analysis identified the efficacy and safety of M+P in DPN more precisely by retrieving data published in the randomized controlled trials (RCTs).

2. Methods

2.1. Search strategy

We retrieved the electronic databases of PubMed, Embase, Web of Science, Cochrane Library, Chinese BioMedical Database, Chinese National Knowledge Infrastructure Database and Wanfang Database (last search date September 2017) without language restrictions. The key terms used in this search were (diabetic peripheral neuropathy or diabetic neuropathy or diabetic neuropathies or DPN) and (M or mecobalamin or vitamin B12) and (P or alprostadil).

2.2. Study selection criteria

All the following inclusion criteria must be met for this study at the same time:

- (1) study design was RCT.
- (2) Patients had DM and distal symmetrical sensorimotor polyneuropathy of the limbs, the diagnostic criteria included standardized DM criteria of World Health Organization,^[16] clinical assessments and nerve conduction.^[12]
- (3) Patients were treated with combination therapy (M+P) versus monotherapy (P).
- (4) Data on symptoms and (or) NCVs could be extracted,
- (5) treatment duration was ≥ 14 days, and a full-text publication was available.

The exclusion criteria included:

- (1) sensorimotor polyneuropathy caused by other factors.
- (2) Trials with some deficiencies in data or study design.
- (3) Patients with DPN received oral administration of M and (or) P.

2.3. Data extraction

All potentially relevant data including patient baseline characteristics, trial durations, daily doses of M and P along with outcomes were extracted independently by the investigators from the collected studies. The primary outcomes of interest were clinical efficacy, median motor nerve conduction velocity (MNCV), median sensory nerve conduction velocity (SNCV), peroneal MNCV, and peroneal SNCV. Clinical efficacy was divided into 3 categories including markedly effective (disappearance of subjective symptoms, recovered tendon reflex, and NCV increased by at

least 5 m/s), effective (alleviated subjective symptoms, improved tendon reflex, and NCV increased by at least 3 m/s) and ineffective (no improvement in symptoms, tendon reflex and NCV).^[17] Moreover, secondary outcomes included adverse events.

2.4. Quality assessment

The established Jadad scale was used to assess the methodological quality of included studies by the authors.^[18] Four to 7 points implied high-quality trials, and 0 to 3 points implied poor or low-quality trials.^[16,19] The disputes during quality assessment were solved by consensus.

2.5. Ethical approval

All the data in present meta-analysis were extracted from the previous published studies, no ethical approval or patient consent was required.

2.6. Statistical analysis

Dichotomous data (efficacy) were expressed as risk ratio (RR) and 95% confidence intervals (95% CIs), and the weighted mean difference (MD) and 95% CIs were estimated for continuous data (NCVs). The statistical heterogeneity between trials was assessed by the Q-statistic and I^2 -test.^[20] A significant Q-statistic ($P \leq .10$) indicated heterogeneity across studies. The random-effect (RE) model was used to pool the data when heterogeneity was confirmed ($P \leq .10$ or $I^2 \geq 50\%$ suggested significant heterogeneity among studies),^[21] otherwise, the fixed-effect (FE) model was employed. Funnel plot was used to detect the possibility of publication bias. Sensitivity analysis was performed by excluding 1 trial at a time, starting from the trial with lower-quality, to further study the effect of a single trial on pooled data. Subgroup analyses were also conducted based on the treatment duration (≥ 28 days or < 28 days). All tests were 2-sided and a value of $P < .05$ was regarded as statistically significant. Statistical analysis was performed using Revman Manager 5.3 software (Cochrane Collaboration, Oxford, UK).

3. Results

3.1. Description of the studies

The process of the study selection and literature search was displayed in Figure 1. The 205 potentially relevant articles were identified from the initial searches, but only 16 trials^[12–14,22–34] satisfying the inclusion and exclusion criteria were selected for this meta-analysis. The key characteristics of the 16 RCTs and Jadad scores were presented in Table 1. The 576 DPN patients were included in the M+P combination therapy group and 560 DPN patients were included in the P monotherapy group. The daily doses of M were 0.1 mg or 0.5 mg or 1.0 mg, daily doses of P were 10 μg or 20 μg or 100 μg , respectively. The routes of drug administration included intravenous bolus injection or intravenous drip infusion; furthermore, intramuscular injection was also used for M administration. The treatment durations varied from 14 to 28 days in most studies except 1 trial^[22] for 30 days and the other trial^[25] for 56 days. Only 2 studies^[14,25] with 4 points were of high quality and the remaining 14 trials with 3 or lower points were all of low quality. Five studies^[24,25,27–29] reported the DM duration. Seven trials^[12,14,22,25,27,30,33] did not differentiate the type of diabetes.




Figure 1. Map of the literature search and selection process.

Table 1

Characteristics of the studies included in the meta-analysis.

Study	Number M + P/P	Age M + P/P	Gender male/female	Type of diabetes (n)	DM duration (year) M + P/P	Treatment duration/days	Treatment drugs sig/day			Outcomes ①②③④⑤	Quality
							M + P		P		
							M	P			
An XQ et al, 2016	46/46	NR	NR	NR	NR	14	0.5mg im	20 µg ivgtt	20 µg ivgtt	①②③④⑤	2
Chen RL et al, 2014	45/45	69.8/71.5	61/29	2	NR	21	0.1 mg ivgtt	10 µg iv	10 µg iv	①②③④⑤	2
Dai HB, 2011	40/40	NR	NR	NR	NR	30	1.0mg ivgtt	10 µg iv	10 µg iv	①②③④⑤	2
Feng LH et al, 1999	38/40	57/55	32/46	2	NR	14	0.5mg im	100 µg ivgtt	100 µg ivgtt	①	2
Fu QL et al, 2012	35/35	65/64	41/29	2	18.5/16.5	28	0.1 mg ivgtt	10 µg ivgtt	10 µg ivgtt	①④⑤	3
Li HJ, 2014	41/41	54.9/54.2	43/39	NR	9/8.7	56	0.5mg iv	10 µg ivgtt	10 µg ivgtt	①②③④⑤	4
Li HY et al, 2013	34/34	NR	NR	2	NR	14	0.5mg im	10 µg ivgtt	10 µg ivgtt	①	2
Liu XL, 2015	31/31	45/46	35/27	NR	8/9	14	0.5mg im	10 µg ivgtt	10 µg ivgtt	①②③④⑤	2
Liu Y et al, 2012	30/30	42.2/43.1	NR	1 or 2	7/7.2	28	0.5mg iv	10 µg ivgtt	10 µg ivgtt	①②③④⑤	2
Niu XH et al, 2009	29/28	70.2/68.2	33/24	2	11.1/10.2	21	1.0mg ivgtt	10 µg iv	10 µg iv	①④⑤	3
Pan LY et al, 2015	42/42	68.7/69.5	45/39	NR	NR	21	0.1 mg ivgtt	10 µg ivgtt	10 µg ivgtt	②③④⑤	4
Peng WD, 2013	48/48	49.7/50.9	57/39	NR	NR	21	0.5mg ivgtt	100 µg ivgtt	100 µg ivgtt	①	3
Wang ZH et al, 2009	38/31	NR	NR	2	NR	28	0.5mg iv	10 µg iv	10 µg iv	①	2
Yi LJ et al, 2005	21/23	50.6/47.8	19/25	1 or 2	NR	28	0.5mg iv	10 µg ivgtt	10 µg ivgtt	①②③④⑤	2
Yu JT et al, 2014	26/26	53.5/50.4	21/31	NR	NR	14	0.5mg im	10 µg iv	10 µg iv	①	3
Zhu XP et al, 2001	32/20	NR	NR	2	NR	28	0.5mg im	100 µg ivgtt	100 µg ivgtt	①②③④⑤	2

DM=diabetes mellitus, DPN=diabetic peripheral neuropathy, im=intramuscular injection, iv=intravenous bolus injection, ivgtt=intravenously guttae, ①=efficacy, ②=median MNCV, ③=median SNCV, ④=peroneal MNCV, ⑤=peroneal SNCV, M=methylcobalamin, NR=not report, P=prostaglandin E1, Quality was assessed by the established Jadad scale and 4-7 points implied high-quality trials.

3.2. Efficacy

Fifteen trials^[12,13,22–34] involving 1052 patients measured the efficacy (534 patients received M+P combination therapy and 518 patients received P monotherapy). As shown in Figure 2, the FE model was used because insignificant heterogeneity between studies for the 2 groups was observed ($P=.16$, $I^2=27\%$). Compared with P monotherapy, M+P combination therapy for DPN significantly enhanced the efficacy (RR 1.25, 95% CI 1.18–1.32, $P<.00001$). The subgroup with ≥ 28 days of study duration showed moderate heterogeneity in efficacy outcome ($I^2=51\%$, $P=.05$). Figure 3 showed the funnel shape was not perfectly symmetrical, indicating a potential publication bias.

3.3. Median MNCV

Nine trials^[12–14,22,25,27,28,32,34] involving 646 patients measured the median MNCV. Heterogeneity was significant for the analysis ($P<.00001$, $I^2=90\%$), the RE model was used. Compared with P monotherapy, median MNCV showed significant improvement in the M+P combination therapy group (MD 6.29, 95% CI 4.63–7.94, $P<.00001$) (Fig. 4A). On sensitivity analyses, after excluding the study reported by Li HJ,^[25] the I^2 value ranged from 90% to 71% and the overall effect ranged from 7.42 to 10.38. The subgroup with < 28 days of study duration showed moderate heterogeneity in median MNCV outcome ($I^2=57\%$, $P=.07$).

3.4. Median SNCV

Nine trials^[12–14,22,25,27,28,32,34] involving 646 patients measured the median SNCV. As shown in Figure 4B, the RE model was used because significant heterogeneity between studies for the 2 groups was observed ($P<.00001$, $I^2=94\%$). Compared with P monotherapy, M+P combination therapy increased median SNCV significantly (MD 5.68, 95% CI 3.53–7.83, $P<.00001$). On sensitivity analyses, we found the I^2 value ranged from 89% to 95%, which indicated the result, was robust.

3.5. Peroneal MNCV

Eleven trials^[12–14,22,24,25,27–29,32,34] involving 773 patients measured the peroneal MNCV. As shown in Figure 5A, the RE model was used because significant heterogeneity between studies for the two groups was observed ($P<.00001$, $I^2=92\%$). Compared with P monotherapy, M+P combination therapy accelerated peroneal MNCV significantly (MD 5.36, 95% CI 3.86–6.87, $P<.00001$). The sensitivity analyses showed that the I^2 value ranged from 88% to 93% and the overall effect ranged from 6.07 to 7.60, which indicated the result was robust.

3.6. Peroneal SNCV

Eleven trials^[12–14,22,24,25,27–29,32,34] involving 773 patients measured the peroneal SNCV. As shown in Figure 5B, the RE model was used because significant heterogeneity between studies for the 2 groups was observed ($P<.00001$, $I^2=86\%$). Compared with P monotherapy, M+P combination therapy improved peroneal SNCV significantly (MD 4.62, 95% CI 3.48–5.75, $P<.00001$). On sensitivity analyses, we found the I^2 value ranged from 80% to 88%, which indicated the result, was robust.

3.7. Safety

Eight studies^[12,14,24–26,28,29,31] reported the adverse events, there were no serious treatment-related side effects during the treatment period in both M+P combination therapy group and P monotherapy group. Only some mild adverse effects including facial blushing (4 cases),^[12,14,25] local skin redness (3 cases),^[29] pain at the injection site (9 cases),^[24,28] gastrointestinal discomfort (1 case),^[25] dizziness (4 cases),^[14,25,29] abdominal distention (3 cases),^[12] limb burning (3 cases),^[31] anepithymia (2 cases),^[12] headache (2 cases)^[25,28] and transient orthostatic hypotension (1 case)^[26] in M+P combination therapy group, and facial blushing (2 cases),^[14,29] pain at the injection site (12 cases),^[24,25,28,29] abdominal distention (1 case),^[12] limb burning (1 case),^[31] anepithymia (2 cases),^[12] headache (1 case),^[28]




Figure 2. Comparison of M+P combination therapy and P monotherapy in the efficacy for diabetic peripheral neuropathy, the former showed a better effect. P= prostaglandin E1, M=methylcobalamin.




Figure 3. Funnel plot was not perfectly symmetrical, indicating the presence of between-study heterogeneity and a potential publication bias.

transient orthostatic hypotension (1 case),^[26] and dizziness (2 cases)^[14,25] in P monotherapy group were reported. Because most studies did not report these side effects in detail, we were unable to analyze the rates of adverse events.

4. Discussion

With the raised prevalence of diabetes, the occurrence of DPN increases significantly and has become a leading cause of




Figure 4. M+P combination therapy improved the median MNCV (A) and median SNCV (B) significantly for treatment of diabetic peripheral neuropathy compared with P monotherapy. P=prostaglandin E1, M=methylcobalamin, MNCV= motor nerve conduction velocity, SNCV=sensory nerve conduction velocity.




Figure 5. Compared with P monotherapy, M+P combination therapy increased the peroneal MNCV (A) and peroneal SNCV (B) significantly for treatment of diabetic peripheral neuropathy. P=prostaglandin E1, M=methylcobalamin, MNCV= motor nerve conduction velocity, SNCV=sensory nerve conduction velocity.

diabetes-related disability. Sensory neuropathy is a principal form of DPN, whose pathological changes include demyelination of nerve fibers, axonal degeneration, cell hyperplasia, and then fading-away of myelinated fibers.^[3] Currently, DPN is believed to be closely related to various factors, including genetic predisposition, glucose toxicity, abnormal aldose reductase activity, oxidative stress.^[5,35] Besides that, factors such as diabetic microangiopathy-induced hypoxic-ischemic neuronal death and altered hemodynamics, as well as disorder of intrinsic clotting, also play a vital role in the occurrence of DPN.^[1,36] Therefore, regulating blood glucose homeostasis, improving the microcirculation of peripheral nerve endings, and suppressing oxidative stress are cures of DPN, moreover, significant treatment options for diabetic complications.

P, an efficient biological activator, induces vasodilation of blood vessels through activating intracellular adenylate cyclase, enhances erythrocyte deformability, improves microcirculation disturbance, protects against ischemia-hypoxia injury in peripheral nerve tissue, and finally ameliorates the symptoms of peripheral nervous system involvement in the form of sensory impairment and diminished tendon jerks suggesting the presence of neuropathy.^[37] In addition, P can reactivate Na(+)-K(+)-ATPase at the surfaces of nerve cells, improve neuronal metabolism and inhibit oxidation of the plasma membrane of a cell,^[10,37] all of which contribute to the improvements of DPN. It has been demonstrated that P significantly improve the clinical symptoms of DPN and increase the conduction velocity of sensory and motor fibres in human median, peroneal nerves.^[10,11,38]

M, a vitamin B12 analog, is involved in methyl transfer reactions in vivo by methylation. M is distributed to organelles in axons of nerve cells easily after being absorbed into the body, promotes nucleic acid and protein synthesis, and axon regeneration.^[39,40] It also can stimulate phosphatidylcholine synthesis to increase myelinogenesis, and then speed up the motor and sensory NCVs.^[41,42] Additionally, M accelerates NCVs directly by improving blocked nerve impulse conduction and decreased neurotransmitter levels.^[43] Many studies suggested that M monotherapy or polytherapy with other drugs is an effective and safe therapy for patients with DPN.^[25,44-46]

Our findings showed that, after P monotherapy and M+P combination therapy, DPN patients all had improvement in clinical symptoms and NCVs, while patients who received the later therapy showed significant higher-level improvement. Moreover, the results also indicated that synergistic potential existed in the course of combination therapy without severe adverse events. We executed subgroup and sensitivity analyses in order to minimize the influence of a particular study or an inferior study design. Results of subgroup analyses according to the study duration suggested that the efficacy and 4 NCVs benefits were seen in 2 subgroups (Table 2).

Our analysis also has several limitations that must be taken into consideration when interpreting the results. First, the sample size of 3 trials was small.^[29,32,33] Second, a reporting bias existed in our meta-analysis, due to only the data from published trials were included and the unpublished statistically nonsignificant results were excluded, but it would be very difficult to gain access

Table 2
Subgroup analyses for efficacy and NCVs according to the treatment duration by meta-analysis.

Treatment duration	No. of patients and studies					RR (95% CI)		MD (95% CI)					I ² (%)					P value				
	①	②	③	④	⑤	①	②	③	④	⑤	①	②	③	④	⑤	①	②	③	④	⑤		
<28 days	595, 8	328, 4	328, 4	385, 5	385, 5	1.28 (1.18, 1.39)	6.48 (5.26, 7.70)	6.42 (3.87, 8.97)	6.14 (3.96, 8.31)	5.63 (3.12, 8.14)	0	57	91	87	87	<0.001	<0.001	<0.001	<0.001	<0.001		
≥28 days	457, 7	318, 5	318, 5	388, 6	388, 6	1.17 (1.05, 1.31)	6.11 (3.04, 9.17)	5.04 (1.33, 8.76)	4.80 (2.88, 6.72)	4.05 (2.67, 5.44)	51	95	96	93	88	0.004	<0.001	0.008	<0.001	<0.001		

CI=confidence interval, ①=efficacy, ②=median MNCV, ③=median SNCV, ④=peroneal MNCV, ⑤=peroneal SNCV, MD=mean difference, MNCV=motor nerve conduction velocity, NCVs=nerve conduction velocities, SNCV=sensory nerve conduction velocity.

to data from the unpublished studies. Third, because this study was a study-level meta-analysis, individual patient data were not included in the analysis, thus, we could not adjust for patient-level confounders. In addition, the small-study effect, insufficient number of trials, and significant statistical heterogeneity may result in the asymmetry of funnel plot, which indicated the likelihood of publication bias.

In summary, this meta-analysis suggests that DPN patients with M+P combination therapy have significant higher-level improvement in clinical symptoms and NCVs compared with P monotherapy. Moreover, the results also indicate that no serious adverse events occur during M+P combination therapy. But, due to poor methodological quality of the studies included, strong and definitive recommendations cannot be made for patients with DPN and further large-scale, well-designed RCTs are urgently needed.

Author contributions

Investigation: De-Qi Jiang, Shi-Hua Zhao, Ming-Xing Li.

Methodology: Shi-Hua Zhao, Li-Lin Jiang, Yong Wang.

Writing – original draft: De-Qi Jiang.

Writing – review & editing: Yan Wang.

References

[1] Flynn MD, Tooke JE. Diabetic neuropathy and the microcirculation. *Diabet Med* 1995;12:298–301.

[2] Kilo S, Berghoff M, Hilz M, et al. Neural and endothelial control of the microcirculation in diabetic peripheral neuropathy. *Neurology* 2000;54:1246–52.

[3] Dyck PJ, Giannini C. Pathologic alterations in the diabetic neuropathies of humans: a review. *J Neuropathol Exp Neurol* 1996;55:1181–93.

[4] Wang Y, Schmeichel AM, Iida H, et al. Enhanced inflammatory response via activation of NF-kappaB in acute experimental diabetic neuropathy subjected to ischemia-reperfusion injury. *J Neurol Sci* 2006;247:47–52.

[5] Tesfaye S, Stevens LK, Stephenson JM, et al. Prevalence of diabetic peripheral neuropathy and its relation to glycaemic control and potential risk factors: the EURODIAB IDDM complications Study. *Diabetologia* 1996;39:1377–84.

[6] Wilkens JH, Wilkens H, Elger B, et al. Cardiac and microcirculatory effects of different doses of prostaglandin E1 in man. *Eur J Clin Pharmacol* 1987;33:133–7.

[7] Kuwabara S, Nakazawa R, Azuma N, et al. Intravenous methylcobalamin treatment for uremic and diabetic neuropathy in chronic hemodialysis patients. *Intern Med* 1999;38:472–5.

[8] Jiang DQ, Li MX, Wang Y, et al. Effects of prostaglandin E1 plus methylcobalamin alone and in combination with lipoic acid on nerve conduction velocity in patients with diabetic peripheral neuropathy: A meta-analysis. *Neurosci Lett* 2015;594:23–9.

[9] Sun Y, Lai MS, Lu CJ. Effectiveness of vitamin B12 on diabetic neuropathy: systematic review of clinical controlled trials. *Acta Neurol Taiwan* 2005;14:48–54.

[10] Akahori H, Takamura T, Hayakawa T, et al. Prostaglandin E1 in lipid microspheres ameliorates diabetic peripheral neuropathy: clinical usefulness of Semmes-Weinstein monofilaments for evaluating diabetic sensory abnormality. *Diabetes Res Clin Pract* 2004;64:153–9.

[11] Wu JD, Tao S, Jin X, et al. PGE1 improves diabetic peripheral neuropathy in patients with type 2 diabetes. *Prostaglandins Other Lipid Mediat* 2016;126:24–8.

[12] Deng H, Yin J, Zhang J, et al. Meta-analysis of methylcobalamin alone and in combination with prostaglandin E1 in the treatment of diabetic peripheral neuropathy. *Endocrine* 2014;46:445–54.

[13] An XQ, Sun JF, Jiang JR. Efficacy and safety of mecobalamine combined with alprostadil in treatment of diabetic peripheral neuropathy. *J Clin Med Pract* 2016;20:50–2.

[14] Chen RL, Sun ZL. Analysis on mecobalamin combined with alprostadil in the treatment of elderly type 2 diabetic peripheral neuropathy. *China Mod Doct* 2014;52:132–4.

[15] Pan LY, Liu R. Clinical effects observation of prostaglandin E1 and mecobalamin in the treatment of patients with diabetic peripheral neuropathy. *J Commun Med* 2015;13:41–2.

[16] Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539–53.

[17] Xu Q, Pan J, Yu J, et al. Meta-analysis of methylcobalamin alone and in combination with lipoic acid in patients with diabetic peripheral neuropathy. *Diabetes Res Clin Pract* 2013;101:99–105.

[18] Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary. *Control Clin Trials* 1996;17:1–2.

[19] Moher D, Pham B, Jones A, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses. *Lancet* 1998;352:609–13.

[20] Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954;10:101–29.

[21] Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.

[22] Dai HB. Clinical observation of mecobalamin combined with alprostadil in the treatment of diabetic peripheral neuropathy. *Guide China Med* 2011;9:216–7.

[23] Feng LH, Feng YS, Li CX, et al. Combined therapeutic effects of mecobalamin and prostaglandin E1 on treating patients with peripheral neuropathy. *Henan J Pract Nerv Dis* 1999;2:25–6.

[24] Fu QL, Rao GF. Effects and safety of methylcobalamin combined with alprostadil in the treatment of type 2 diabetes mellitus patients with peripheral neuropathy. *Chin J Clin Pharmacol* 2012;28:490–2.

[25] Li HJ. Clinical study of prostaglandin E1 plus mecobalamin combination therapy for patients with diabetic neuropathy. *Lab Med Clin* 2014;11:485–6.

[26] Li YH, Sun HB, Wang Q. Clinical efficacy of alprostadil combined with methylcobalamin in the treatment of diabetic patients with peripheral neuropathy. *Guide China Med* 2013;11:185–6.

[27] Liu XL. Clinical observation of prostaglandin E1 combined with mecobalamin in the treatment of patients with diabetic neuropathy. *Diabet New World* 2015;35:41–3.

[28] Liu Y, Zhou L, Xie XM. Clinical study of prostaglandin E1 combined with mecobalamin in the treatment of diabetic neuropathy. *Baotou Med* 2012;36:65–6.

[29] Niu XH, Si QQ, Song Q. Combined therapeutic effects of mecobalamin and prostaglandin E1 for patients with diabetic peripheral neuropathy. *J Clin Intern Med* 2009;26:707–8.

[30] Peng WD. Clinical analysis of alprostadil combined with methylcobalamin treatment for 48 cases of diabetic neuropathy. *China Prac Med* 2013;8:156–7.

[31] Wang ZH, Chen HY, Guo RQ. Combined therapeutic effect of alprostadil and mecobalamin on 128 patients with diabetic peripheral neuropathy. *Chin J Pract Med* 2009;36:2–3.

- [32] Yi LJ, Yue YH. Clinical efficacy of comprehensive treatment for patients with diabetic peripheral neuropathy. *J Clin Exp Med* 2005;4:25–7.
- [33] Yu JT, Xu CC. Efficacy analysis of alprostadil combined with mecobalamin in the treatment of diabetic peripheral neuropathy. *Chin Foreign Med Res* 2014;12:27–8.
- [34] Zhu XP, Zhu ZG. Clinical observation of combined therapeutic effect of prostaglandin E1 and mecobalamin on diabetic peripheral neuropathy. *Bull Hunan Med Univ* 2001;26:343–4.
- [35] Boulton AJ, Gries FA, Jervell JA. Guidelines for the diagnosis and outpatient management of diabetic peripheral neuropathy. *Diabet Med* 1998;15:508–14.
- [36] Brooks B, Delaney-Robinson C, Molyneaux L, et al. Endothelial and neural regulation of skin microvascular blood flow in patients with diabetic peripheral neuropathy: effect of treatment with the isoform-specific protein kinase C beta inhibitor, ruboxistaurin. *J Diabetes Complications* 2008;22:88–95.
- [37] Toyota T, Hirata Y, Ikeda Y, et al. Lipo-PGE1, a new lipid-encapsulated preparation of prostaglandin E1: placebo-and prostaglandin E1-controlled multicenter trials in patients with diabetic neuropathy and leg ulcers. *Prostaglandins* 1993;46:453–68.
- [38] Hong L, Zhang J, Shen J. Clinical efficacy of different doses of lipo-prostaglandin E1 in the treatment of painful diabetic peripheral neuropathy. *J Diabetes Complications* 2015;29:1283–6.
- [39] Okada K, Tanaka H, Tempurin K, et al. Methylcobalamin increases Erk1/2 and Akt activities through the methylation cycle and promotes nerve regeneration in a rat sciatic nerve injury model. *Exp Neurol* 2010;222:191–203.
- [40] Watanabe T, Kaji R, Oka N, et al. Ultra-high dose methylcobalamin promotes nerve regeneration in experimental acrylamide neuropathy. *J Neurol Sci* 1994;122:140–3.
- [41] Jian-bo L, Cheng-ya W, Jia-wei C, et al. The preventive efficacy of methylcobalamin on rat peripheral neuropathy influenced by diabetes via neural IGF-1 levels. *Nutr Neurosci* 2010;13:79–86.
- [42] Kikuchi M, Kashii S, Honda Y, et al. Protective effects of methylcobalamin, a vitamin B12 analog, against glutamate-induced neurotoxicity in retinal cell culture. *Invest Ophthalmol Vis Sci* 1997;38:848–54.
- [43] Yagihashi S, Tokui A, Kashiwamura H, et al. In vivo effect of methylcobalamin on the peripheral nerve structure in streptozotocin diabetic rats. *Horm Metab Res* 1982;14:10–3.
- [44] Yaqub BA, Siddique A, Sulimani R. Effects of methylcobalamin on diabetic neuropathy. *Clin Neurol Neurosurg* 1992;94:105–11.
- [45] Li S, Chen X, Li Q, et al. Effects of acetyl-L-carnitine and methylcobalamin for diabetic peripheral neuropathy: A multicenter, randomized, double-blind, controlled trial. *J Diabetes Investig* 2016;7:777–85.
- [46] Han Y, Wang M, Shen J, et al. Differential efficacy of methylcobalamin and alpha-lipoic acid treatment on symptoms of diabetic peripheral neuropathy. *Minerva Endocrinol* 2018;43:11–8.