

SCIENTIFIC REPORTS



OPEN

Deficiency of *Mkrn2* causes abnormal spermiogenesis and spermiation, and impairs male fertility

Received: 24 June 2016
Accepted: 14 November 2016
Published: 23 December 2016

Xu Qian^{1,2,*}, Lin Wang^{3,*}, Bo Zheng^{1,*}, Zhu-Mei Shi^{1,2,4}, Xin Ge^{1,2}, Cheng-Fei Jiang^{1,2}, Ying-Chen Qian^{1,2}, Dong-Mei Li^{1,2}, Wei Li^{1,2}, Xue Liu^{1,2}, Yu Yin^{1,2}, Ji-Tai Zheng^{1,2}, Hua Shen⁵, Min Wang^{1,2}, Xue-Jiang Guo¹, Jun He⁶, Marie Lin⁷, Ling-Zhi Liu⁶, Jia-Hao Sha¹ & Bing-Hua Jiang^{1,2,6}

Although recent studies have shed insights on some of the potential causes of male infertility, new underlying molecular mechanisms still remain to be elucidated. *Makorin-2* (*Mkrn2*) is an evolutionarily conserved gene whose biological functions are not fully known. We developed an *Mkrn2* knockout mouse model to study the role of this gene, and found that deletion of *Mkrn2* in mice led to male infertility. *Mkrn2* knockout mice produced abnormal sperms characterized by low number, poor motility, and aberrant morphology. Disruption of *Mkrn2* also caused failure of sperm release (spermiation failure) and misarrangement of ectoplasmic specialization (ES) in testes, thus impairing spermiogenesis and spermiation. To understand the molecular mechanism, we found that expression of *Odf2*, a vital protein in spermatogenesis, was significantly decreased. In addition, we found that expression levels of *Odf2* were decreased in *Mkrn2* knockout mice. We also found that *MKRN2* was prominently expressed in the sperm of normal men, but was significantly reduced in infertile men. This result indicates that our finding is clinically relevant. The results of our study provided insights into a new mechanism of male infertility caused by the *MKRN2* downregulation.

Infertility is a severe health problem leading to tremendous social consequences. Approximately 10–15% of all couples experience fertility-related problems during their childbearing years¹ and around 50% of such cases are caused by male infertility². Sperm carries paternal genetic information delivering heredity to offspring. Spermatogenesis can be divided into three stages: mitosis of spermatogonia, meiosis of spermatocytes, and spermiogenesis^{3–5}. Although many genes, including more than 20 male germ cell-specific genes, have been identified to be essential for spermatogenesis⁶, the underlying mechanisms of spermatogenesis remain largely unknown.

The makorin (*MKRN*) gene family encodes putative ribonucleoproteins with a distinctive array of zinc finger domains⁷. Makorin-1 (*MKRN1*) is the most extensively studied member of this family of proteins^{7–9}. Makorin-2 (*MKRN2*), formerly designated as *HSPC070*, is a new member of the Makorin family, which was first identified in human CD34+ hematopoietic stem/progenitor cells¹⁰. *MKRN2* is a zinc finger protein containing the typical C₃HC₄ protein-protein interaction motif (termed the RING domain) found in most E3 ubiquitin ligases^{11,12}. It is ubiquitously expressed in various tissues and cell lines^{10,13}, but *MKRN2* function is not known yet in mammals. We previously demonstrated that forced expression of *Mkrn2* in *Xenopus* produced impaired tadpoles with dorso-posterior deficiencies and small-head/short-tail phenotype, whereas knockdown of *Mkrn2* by morpholino

¹State Key Laboratory of Reproductive Medicine, Nanjing Medical University, Nanjing, Jiangsu 210029, China.

²Department of Pathology, and Cancer Center, Nanjing Medical University, Nanjing, Jiangsu 210029, China.

³Institute of Medical and Pharmaceutical Sciences, Zhengzhou University, Henan 450052, China. ⁴Department of Neurosurgery, The First Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu 210029, China.

⁵Department of Oncology, The First Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu 210029, China. ⁶Center for Molecular Carcinogenesis, Department of Pathology, Anatomy and Cell Biology, Thomas Jefferson University, Philadelphia, PA 19107, USA. ⁷Biomedical Engineer Research Center, Kunming Medical University, Kunming, Yunnan 65000, China. *These authors contributed equally to this work. Correspondence and requests for

materials should be addressed to J.H.S. (email: shajh@njmu.edu.cn) or B.H.J. (email: binghjiang@yahoo.com)




Figure 1. Expression profile of Mkrn2 in mice tissues. (a) Mkrn2 mRNA levels in different tissues were analyzed by RT-qPCR and normalized to those in liver tissue. Data in the figure are represented as means \pm SE. from four mice. (b) The Mkrn2 protein levels in mouse organs were detected by immunoblotting. β -Tubulin was used as a loading control. The blotting density was normalized to that of the liver. (c) The location of Mkrn2 in the testis of 4-week mice was determined using immunohistochemistry. Scale bar = 20 μ m.

antisense oligonucleotides induced a double-axis phenotype during neurogenesis¹⁴; however, further studies are required to fully determine the biological function of Mkrn2 in mammals.

To investigate the function of the *MKRN2* gene, we generated *Mkrn2*-knockout mice. Based on the results of our previous study, we expected that deletion of *Mkrn2* in mice may result in deficiency in the development of neurogenesis¹⁴. Surprisingly, *Mkrn2*-knockout mice were viable with detectable abnormal neurogenesis, but the male mice were sterile. As mice and human share similar process of spermiogenesis and spermiation¹⁵, these infertile *Mkrn2*-knockout mice could serve as an *in vivo* model to investigate the biological function of MKRN2 in human spermiogenesis and spermiation.

Results

Expression pattern of Mkrn2 in mouse testes. Mkrn2 mRNA and protein expression levels were evaluated using qRT-PCR and immunoblotting analysis, respectively, in different tissue samples of mice. The *Mkrn2* mRNA levels were low in the brain, thymus, heart, lung, liver, spleen, kidney, ovary, uterus, and seminal vesicle; but were very high in the testis with 30-fold higher than those in the brain and heart tissues (Fig. 1a). Similarly, Mkrn2 protein was ubiquitously expressed at low levels in different tissues, but preferentially expressed in the testis (Fig. 1b), indicating that Mkrn2 may play a vital role in testes. Furthermore, immunohistochemical analysis revealed that Mkrn2 protein was mainly localized in the Sertoli cells and spermatids (Fig. 1c). These data implied that Mkrn2 may play an important role in spermiogenesis and spermiation.

MKRN2 expression in infertile human sperms. MKRN2 protein levels were lower in human sperm samples from patients of oligoasthenoatozoospermia (OAT) than in the normal samples (Fig. 2a, Supplementary Figure S1). Similarly, the mRNA levels of *MKRN2* in the OAT patient sperms were significantly lower by more than two-fold than those in the normal group (Fig. 2b). To validate this finding in an independent cohort, we obtained MKRN2 expression data from sperm samples (including 8 OAT patients and 13 normal subjects) of American populations which was submitted to the database of European Bioinformatics Institute (EBI) by Platts and colleagues (E-GEOD-6872)¹⁶. By analyzing the raw data, we found that the *MKRN2* mRNA levels were significantly decreased by 5-fold in the OAT group than in the corresponding normal group (Fig. 2c). These two independent cohorts demonstrated that low MKRN2 expression levels in sperm are strongly correlated with human male infertility.




Figure 2. Expression of MKRN2 in human sperm. (a) MKRN2 protein expression levels from human sperm samples were subjected to immunoblotting analysis. (b) The MKRN2 mRNA levels from human sperm samples were measured by RT-qPCR. (c) MKRN2 mRNA expression levels of sperm samples from the US cohort were compared to available raw data extracted from the European Bioinformatics Institute (EBI) database. OAT, oligoasthenozoospermia. * And ** significant difference at $P < 0.05$ and $P < 0.01$, respectively (two-tailed Student's t test).

Generation of *Mkrn2* knockout mice. Model animals are very useful tools to study the functions and mechanisms of target genes¹⁷. To better investigate the role of *Mkrn2* gene *in vivo*, we generated *Mkrn2* knockout mice using a Cre-LoxP system (Fig. 3a). The *Mkrn2* gene in mice contains eight exons, in which the functional domains are mainly encoded by Exons 2–6. Thus, we disrupted *Mkrn2* gene by deleting these exons and introducing frameshift mutations in Exons 7 and 8, thus obtaining maximum knockout effect. The mutant allele could be transmitted from both *Mkrn2* heterozygous male and female mice, and homozygous mice could be obtained by interbreeding these heterozygous mice. *Mkrn2* heterozygous and homozygous offspring were identified using a PCR-based genotyping assay (Fig. 3b). Immunoblotting analysis using testis extracts verified the absence of Mkrn2 protein in *Mkrn2* knockout mice (Fig. 3c). Amino acid sequence alignment showed that the MKRN2 protein in mice and human shared 88% identity (Supplementary Figure S2), suggesting that *Mkrn2* knockout mice may be useful to explore the biological function of MKRN2 in human.

Characteristics of *Mkrn2* knockout mice. Both male and female *Mkrn2* knockout mice were viable, and had no gross abnormalities except that their weights were lighter than wild-type counterparts (Supplementary Figure S3). *Mkrn2* knockout male mice could mate with female mice, evidenced by the existing of vaginal plugs after mating. During the four-month duration of the mating test, the average litter size of wild-type male and female mice and *Mkrn2* heterozygous male and female mice were 8.6 and 8.9 pups, respectively. However, no pups were born between *Mkrn2* knockout male mice and female mice (including wild-type, heterozygous, and knockout), indicating that *Mkrn2* knockout male mice were completely sterile (Table 1). Note that female *Mkrn2* knockout mice were fertile, but displayed reduced fecundity with an average litter size of 5.9 pups. We then determined why *Mkrn2* deletion leads to male infertility. First, the lower sperm-producing ability of male *Mkrn2* knockout mice was confirmed by examining their cauda epididymis. Although a few of *Mkrn2* knockout mice produced no sperm, most of them produced sperms, but with abnormal heads and tails (Fig. 4a). The abnormality rates in the sperm heads and tails of *Mkrn2* knockout sperm were 13.2% and 38.9%, respectively, which were remarkably higher than the corresponding rates of 2.3% and 4.1% in wild-type sperms (Fig. 4b). Computer-assisted sperm analysis (CASA) showed that sperm amount was significantly reduced (Fig. 4c), and that approximately 70% of sperms lost motile ability (Fig. 4d) in the *Mkrn2* knockout mice. Although the remaining 30% of the sperms in the *Mkrn2* knockout group were motile, their mean, linear, and curvilinear velocities were significantly lower than those sperms from the wild-type mice (Fig. 4e). The observation of high rate of abnormality and low motile ability of *Mkrn2* knockout sperms is very similar to those in the sperm samples obtained from OAT patients (Supplementary video 1 and 2). Thus, male *Mkrn2* knockout mice produced




Figure 3. Generation of *Mkrn2* knockout mice. (a) Strategy for generating *Mkrn2* knockout mice. *Mkrn2*-floxed mice whose Exons (numbered black boxes) 2–6 of the *Mkrn2* gene were flanked by loxP sites (arrow) were crossed with *EIIa-Cre* transgenic mice to generate *Mkrn2* knockout mice. E, exon; f, Floxed; +, wild-type; –, mutant. (b) *Mkrn2* gene disruption was confirmed by PCR genotyping. Tail genomic DNA was amplified with specific primers for the wild-type (694 bp) and mutant (570 bp) *Mkrn2* alleles. (c) Immunoblotting analysis showing the absence of *Mkrn2* in the protein extracts of *Mkrn2* knockout testes. β -Tubulin expression was used as an internal control to indicate equal amount of loading proteins. +/+, wild-type; +/-, heterozygous; -/-, homozygous (knockout).

Male (n)	Female (n)	Litter number	Pups	Litter Size (mean \pm SD)
+/+ (3)	+/+ (6)	11	95	8.6 \pm 1.8
+/- (11)	+/- (20)	40	354	8.9 \pm 1.5
+/+, +/- (7)	-/- (7)	9	53	5.9 \pm 1.3
-/- (17)	+/+ (7)	0		0
	+/- (5)	0		0
	-/- (11)	0		0

Table 1. Reproductive Defects in *Mkrn2* Knockout Mice. The mean litter size was calculated by dividing the number of live pups born by the number of litters. The results are from the mating of mice less than 6 months of age.

abnormal sperms with a low count, reduced motility, and deformed morphology, which are similar to the most common type of male infertility in human^{18,19}.

Failure of spermiation in the testes of *Mkrn2* knockout mice. The internal organs of the mice were weighed and normalized to their body weight. The relative organ weights were similar between *Mkrn2* knockout and wild-type mice in most of organs including brain, thymus, heart, lung, liver, spleen, kidney, and stomach (Supplementary Table S1); while the relative weights and volumes of testis were significantly lower in *Mkrn2* knockout mice compared to those of the wild-type counterparts (Fig. 5a,b). In the testes of *Mkrn2* knockout mice, all 12 stages of spermatogenesis were present (Supplementary Figure S4). In Stage VIII seminiferous tubules of *Mkrn2* knockout mice, all cell types can be observed, but the number of round spermatids was significantly decreased (Supplementary Figure S5a,b). And elongated spermatids (EI) were lining in the center of the tubules prior to spermiation, similar to that observed in the testis of wild-type mice (Supplementary Figure S5a). However, a distinguishable defect was found in *Mkrn2* knockout testes at Stage X (Fig. 5c). High-power magnification fields showed that *Mkrn2* knockout seminiferous tubules retained elongated spermatids (termed as spermiation failure) among germ cells, which should be released at this stage like those in the Stage X seminiferous tubules of the wild-type mice. These data indicated that *Mkrn2* is important in the regulation of spermiation.

Deletion of *Mkrn2* damages arrangement of actin filaments in the ectoplasmic specialization. The ectoplasmic specialization (ES) is a specialized adhesion junction found in Sertoli cells at sites of attachment to elongated spermatids or neighboring Sertoli cells in the testes²⁰. Spermiation failure can be indicative of disruption of ES²¹. We examined the ultrastructure of the testes using transmission electron microscopy (TEM). In




Figure 4. *Mkrn2* knockout mouse sperm structure and characteristics. (a) Sperm obtained from the cauda epididymis were stained by hematoxylin and eosin (H&E). Sperm from *Mkrn2* knockout mice were deformed: the sperm tails bent forward in the midpiece and the heads were round and irregular (arrow). Scale bar = 50 μm . (b) The proportions of abnormal sperm were calculated from five randomly selected microscopic fields in a double-blinded manner. Data are represented as the means \pm SE. of five random microscopic fields. The sperm count (c), motility (d), and velocity (e) were measured by computer-assisted sperm analysis (CASA) from six different mice of corresponding phenotypes. Data are represented as the means \pm SE. of six mice. Error bars represent SE. * $P < 0.05$, ** $P < 0.01$ (two-tailed Student's *t* test).

normal testes, the ES synchronously stretches along with the acrosome, and is characterized by the presence of actin filament bundles sandwiched between endoplasmic reticulum (ER), and the two apposite plasma membranes of elongating spermid and Sertoli cells (Fig. 6a, left panel). In the *Mkrn2* knockout testes, however, the ES arrangement was disrupted, as evidenced by desynchronization with the acrosome (Fig. 6a, left and middle panels). The ES broke across off the boundary of the acrosome, and extended further, preventing proper elongation of round spermatids by Sertoli cells in *Mkrn2* knockout testes. We next examined the expression levels of the ES marker Espin. The Espin protein and mRNA levels in *Mkrn2* knockout testes were decreased to one-third of the levels detected in the wild-type group (Fig. 6b,c). Immunohistochemical analysis revealed that Espin was expressed in the region between the ES and the heads of elongating/elongated spermatids in the wild-type group, but its expression was barely detected in *Mkrn2* knockout testis (Fig. 6d). These data indicate that *Mkrn2* regulates the accurate arrangement of the ES and Espin expression.

Disruption of *Mkrn2* results in sperm abnormalities. To further define the nature of the abnormalities in *Mkrn2* knockout sperms, we compared the ultrastructure of sperm from the *Mkrn2* knockout and wild-type groups using TEM. Results revealed that spermatozoa were normal with well-shaped heads and tails in the wild-type mice (Fig. 7a,i). However, sperms from the *Mkrn2* knockout mice showed deformed heads, disorganized axonemal structure, and disorganized flagellar structure (Fig. 7a, ii–iv). The shapes of the sperm heads were aberrant, resulting in abnormal or missing acrosomes (Fig. 7a, ii, asterisk). Normal sperm tails contain a highly ordered structure termed the “9 + 2” axoneme²². The “9 + 2” axoneme arrangement in the tail flagellae of *Mkrn2*-deficient sperm exhibited two types of abnormalities (Fig. 7a, iii–iv, arrows): First, complete loss of the axoneme doublets in one side of the fibrous sheath; and second, disordered assembly of the “9+2” arrangement. These data demonstrated that *Mkrn2* is required for the normal sperm development.




Figure 5. Analysis of *Mkrn2* knockout mouse testes. (a) Macroscopic appearance of testes. Scale bar = 2 mm. (b) Testis weights. The net testis weight was normalized to body weight. Data are represented as the means \pm SE. From six testes for each phenotype. Error bars represent SE. ** $P < 0.01$ (two-tailed Student's *t* test). (c) Hematoxylin & eosin (H&E) staining sections of Stage X testicular tubules. Arrows indicate the spermiation failure spermatids in *Mkrn2* knockout testes (right); these sperm should have been released at Stage X, as observed in the wild-type testes (left). Scale bar: Upper panel, 100 μ m; Bottom panel, 20 μ m.

Next, we investigated the molecular mechanism of *Mkrn2* deficiency leading to male infertility. Interestingly, we found that the outer dense fiber (Odf), which is an important component of flagellae (Fig. 7a, iv, arrowheads), was either absent or improperly arranged in the *Mkrn2* knockout epididymal sperms. This finding is consistent with the reduced expression of Odf2 protein, which is important for the regulation of sperm morphology (Fig. 7b). In addition, Odf2 mRNA and protein levels were significantly lower in *Mkrn2* knockout sperms and testes than those of the wild-type (Fig. 7b,c). This result suggests that *Mkrn2* is required for both Odf2 protein and mRNA expression.

Discussion

Mkrn2 is a new member of the highly evolutionarily conserved makorin family, and it may have originally been created by gene duplication of *MKRN1* approximately 450 million years ago¹³. *MKRN1* plays multiple functions such as modulating gene transcription, controlling protein stabilization, and regulating tumorigenesis^{23–25}. However, the biological function of *Mkrn2* in mammals still remains unknown. We previously found that *Xenopus mkrn2* inhibited neurogenesis during early embryonic development¹⁴. To further explore the *in vivo* biological function of *Mkrn2*, we generated a strain of *Mkrn2* knockout mice by using EIIa-Cre method. EIIa-Cre is widely expressed and highly efficient in gene knockout because it is expressed as early as the zygote stage^{26,27}. Since the biological function of *Mkrn2* is largely unknown, we firstly employed EIIa-Cre method to efficiently delete *Mkrn2* in the whole body of mice, and to investigate the phenotype caused by *Mkrn2* knockout. According to our previous study on a *Xenopus* model, we anticipated that disruption of *Mkrn2* would result in prenatal death or deficiency in neurogenesis in mice. Surprisingly, *Mkrn2* knockout mice were viable with no obvious phenotype associated with neurodeficiency. After fecundity tests, male *Mkrn2* knockout mice were found to be sterile, and no pups were born during the 4-month period of mating with female mice. This finding revealed the potential role of *Mkrn2* in fertility of male mice. Further analysis indicated that *Mkrn2* knockout mice either did not have sperm, or produced sperm with a low count, poor motility, and/or abnormal shapes. This condition is termed oligoasthenoteratozoospermia (OAT) in human, which is the most common cause of infertility in human. Mice are commonly used to study human diseases because of the genetic similarities between human and mice^{15,28,29}. Nevertheless, the results from the animal model studies need to be confirmed in human. In this study, we showed that human *MKRN2* protein and mRNA levels in the infertile OAT sperm were lower than those in the fertile sperm from a human cohort. This finding was validated by the results of the second cohort of sperm samples obtained from the EBI, in which the *MKRN2* mRNA levels were downregulated in OAT sperm compared to their normospermic counterparts in the same cohort. These two independent cohorts provide direct evidence indicating that *MKRN2* is significantly correlated with infertility in men. Together, the results obtained from both *Mkrn2* knockout mice and human cohorts suggest that *MKRN2* is important for regulating fertility and that *Mkrn2* knockout mice may serve as a useful animal model for further studies.

To understand why disruption of *Mkrn2* impairs fertility, we examined the distribution of *Mkrn2* in mouse organs. While *Mkrn2* was expressed at low levels in all the organs examined, it was preferentially overexpressed




Figure 6. Analysis of *Mkrn2* knockout mouse ectoplasmic specialization (ES) structure. (a) TEM analysis of ES structure. Arrows indicate the boundary of the acrosome, and arrowheads indicate the improper arrangement of ES. Nu, nucleus; Ac, acrosome; ER, endoplasmic reticulum. Data are represented as means \pm SE. From six different testis tissues for each phenotype. Error bars represent SE. ** $P < 0.01$ (two-tailed Student's t test). Scale bar = 1 μ m. (b) Espin expression levels in the testes were analyzed by immunoblotting. β -Tubulin was used as the internal control. (c) The Espin mRNA levels in the testes were analyzed by RT-qPCR and normalized to that of β -Tubulin. Data are represented as means \pm SE. From six different testis tissues for each phenotype. Error bars represent SE. ** $P < 0.01$ (two-tailed Student's t test). (d) Immunohistochemical staining of Espin in the testes shows co-localization of Espin at the apical ES and in the heads of elongated spermatids in wild-type testes (left), but no obvious staining can be observed in the tubules of the *Mkrn2* knockout testes. Scale bar: Upper panel, 50 μ m; Bottom panel, 20 μ m.

in the testis with up to 30-fold higher mRNA levels than those in other organ tissues. Furthermore, *Mkrn2* was expressed in Sertoli cells and spermatids. Based on these results, we speculated that *Mkrn2* may play an important role in testis development and in the process of spermatogenesis. As expected, the sizes and weights of the testes in *Mkrn2* knockout mice were lower than those in normal mouse testes. TEM analysis revealed that impaired ES structure in *Mkrn2* knockout mice^{20,30,31}. ES is mainly characterized by the presence of a unique junction plaque containing parallel actin bundles sandwiched between the plasma membrane of Sertoli cells and affiliated endoplasmic reticulum^{32,33}. ES is considered important for orienting the positions and shapes of the spermatid head and for regulating spermiation^{34,35}, thus playing a vital role in spermiogenesis and spermiation. Histological examination revealed abnormal spermiogenesis and spermiation in the Stage X seminiferous tubules of the knockout mice, observed by retention of elongated spermatids and sloughing in the different compartments of germ cells. This is a very interesting phenomenon termed spermiation failure, which can be indicative of changes in the components of the ES^{21,36}. Ultrastructure analyses revealed that the ES in the knockout mouse testes was desynchronized with the acrosome and stretched beyond the boundary of the acrosome. Immunoblot analysis showed that Espin was rarely detected in *Mkrn2* knockout mouse testis, indicating that the ES was severely impaired in these mice. As a specialized structure formed by Sertoli cells, ES plays a vital role in spermiogenesis and spermiation. Our results indicated that deletion of *Mkrn2* in the testes, especially in Sertoli cells, impaired ES functions, thus leading to sperm abnormalities.

Except for some cases where the *Mkrn2* knockout male mice produced no sperm, most knockout mice were capable of producing lower numbers of sperm. The low sperm count maybe due to the reduced number of round




Figure 7. Analysis of *Mkrn2* knockout mouse sperm structures and *Odf2* expression. (a) TEM analysis of epididymal sperm. i, Note the normal head and axoneme with typical “9 + 2” microtubule structure (nine pairs of peripheral and two central microtubules) in wild-type mice. ii–iv, Abnormalities in *Mkrn2* knockout mouse sperm were evaluated by the appearance of deformed heads (ii, asterisk), deformed “9 + 2” structures with missing and/or misarranged microtubules (iii–iv, arrows) and lack of Odf (iv, arrowhead). Scale bar = 1 μ m. (b) The *Odf2* protein levels in wild-type and *Mkrn2* knockout mouse sperm and testes, respectively. β -Tubulin was used as the loading control. (c) The *Odf2* mRNA expression levels in the testes were determined by RT-qPCR and normalized to that of β -Tubulin. Data are represented as the means \pm SE. From six different testes tissue samples for each phenotype. Error bars represent SE. * $P < 0.05$ (two-tailed Student’s *t* test).

spermatids in Stage VIII and/or the spermiation failure in stage X seminiferous tubules. Furthermore, deletion of the *Mkrn2* gene led to severe impairment of sperm motility, which strongly indicated abnormality in sperm tails. The same phenomenon was observed in the sperm of human patients with OAT. Sperm from the *Mkrn2* knockout group were morphologically abnormal, with deformed heads and tails bent toward the head at the principle piece. Ultrastructure analyses of cauda epididymal sperm revealed a drastically deformed nucleus and improperly arranged sperm flagellae in *Mkrn2* knockout mice. The “9 + 2” structure was partially lost or disordered. These results implied that deletion of *Mkrn2* in the testes, especially in germ cells, would impair some essential components during sperm flagellar assembly. After further exploration, we found that Odf was not assembled correctly. *Odf2*, an important member of Odf family, has been reported to be associated with flagellar assembly^{37,38}. The *Odf2* expression levels were greatly reduced both in the sperm and testes of *Mkrn2* knockout mice, indicating that *Mkrn2* may control the expression of *Odf2* directly or indirectly, consequently modulating sperm flagellar assembly. It was reported that *Odf2* KO results in preimplantation lethality, suggesting that *Odf2* may be an important downstream protein of MKRN2 for spermiogenesis and spermiation³⁹.

This study uses an *Mkrn2* knockout mouse model and two independent cohorts of human sperm samples to demonstrate the important role of *Mkrn2* in male fertility. Our results demonstrate that *Mkrn2* is essential for spermiogenesis and spermiation. We found that deletion of *Mkrn2* in somatic Sertoli cells results in disruption of ES, leading to abnormalities in sperm heads and spermiation failure. This deletion also suppresses *Odf2* expression in germ cells, disrupting the flagellar assembly of sperm tails. Furthermore, the *Mkrn2* expression levels were significantly downregulated in the infertile male subjects in our study. *Mkrn2* knockout mice may serve as a model for investigating the involvement of *Mkrn2* in spermiogenesis and spermiation in the future.

Material and Methods

Sperm sample collection and analysis. Human sperm samples have been collected for several years by the Center of Clinical Reproductive Medicine, the First Affiliated Hospital of Nanjing Medical University. Informed consents were obtained from all subjects participating in the study. Ejaculates were evaluated according to World Health Organization criteria⁴⁰. These sperm samples and other related samples have been kept in the tissue bank of the First Affiliated Hospital of Nanjing Medical University. These samples have been prepared for cases in the First Affiliated Hospital of Nanjing Medical University biorepository, and clinical annotation is available through a database using caBIG. Cases will be classified and selected based on diagnosis using the CoPath Anatomic Pathology system as well as caBIG. No information regulated by The Health Insurance Portability and Accountability Act was included in the study, which qualifies for the status of NIH Exemption # 4. The sperm samples based on the database containing 33 infertile men diagnosed as having oligoasthenoteratozoospermia (OAT) and 33 fertile men were used for this study, and the results are showed in Supplementary Figure S1.

Ethics statement. All animal experiments were approved by the Committee of Laboratory Animal Experimentation of Nanjing Medical University. All methods were performed in accordance with the relevant guidelines and regulations. The mice used in this study were housed in a controlled specific pathogen-free (SPF) environment and cared according to the approved protocol.

Generation of *Mkrn2* knockout mice. To generate *Mkrn2* floxed mice, a targeting vector containing *Mkrn2* exons 2–6 flanked by *Loxp* sites was linearized by digestion with *NotI* was introduced into embryonic stem cells derived from mice with a 129P2/OlaHsd background by electroporation. Southern blotting analysis was used to confirm the resulting cells harboring floxed alleles of *Mkrn2*. The targeted embryonic stem cell clones were microinjected into C57BL/6J blastocysts of female mice, which gave birth to chimeric mice. The offspring were subsequently crossed with C57BL/6J mice to generate F1 *Mkrn2*-floxed offspring.

To generate *Mkrn2* knockout mice, *Mkrn2*-floxed mice that had been backcrossed with C57BL/6J for more than five generations were crossed with transgenic *EIIa-cre* mice with a C57BL/6J background (Jackson Laboratory) to obtain mosaic mice with the *Mkrn2*^{+/fl}-*EIIa-cre* genotype. The resulting mice were crossed with C57BL/6J to obtain *Mkrn2* heterozygotes. *Mkrn2* knockout mice were generated by sibmating of heterozygotes. Genomic DNA isolated from the tails was genotyped with oligonucleotides listed in Supplementary Table S2, yielding 694-bp (wild-type) and 570-bp (mutant) products.

Morphological and immunohistochemical analysis. The testes and epididymis tissues were dissected, fixed in Bouin's fixatives overnight, and embedded in paraffin. Sections were stained by hematoxylin and eosin (H&E) for histological examination or immunohistochemically using antibodies against MKRN2 (Novus Biologicals) and Espin (BD Biosciences). The epididymis sections were dissected, incubated in PBS, and epididymal sperm morphology was analyzed by observing the sections on a slide with methanol fixation and H&E staining.

Sperm analysis. Sperm motility and count were measured using a CASA system as described previously⁴¹. Mature sperm were isolated from the cauda epididymis dissected from sexually mature mice. The tissue was incised at four sites and placed in 0.5 mL of PBS for 15 min in a 37 °C incubator. Debris was removed, and 20 μL aliquots of sperm were placed in an 80-μm-deep chamber for the assessment of sperm motility, and the sperm count was obtained by CASA using an IVOS Sperm Analyzer (Hamilton Thorne Biosciences).

Ultrastructure analysis. For electron microscopy, testicular and cauda epididymal sperm from adult mice were fixed in 5% (v/v) and 2.5% (v/v) glutaraldehyde at 4 °C, respectively. The specimens were embedded according to standard procedures, sectioned, and examined with a JEM-1010 transmission electron microscope (JEOL).

RNA analysis. Total RNAs were prepared using TRIzol reagent (Invitrogen) according to the manufacturer's instructions. First-strand cDNAs were synthesized and subjected to real-time RT-PCR analysis (Vazyme). The relative amounts of target genes were calculated by the comparative threshold cycle (Ct) method. β-Tubulin mRNA levels were used as the internal control. The oligonucleotide primer sequences used here are listed in Supplemental Table 2.

Immunoblotting. Tissues were homogenized in ice-cold RIPA buffer supplemented with protease inhibitor cocktail (Sigma). The cell suspensions were centrifuged for 15 minutes at 15,000 rpm at 4 °C. The sperm samples were collected and homogenized in lysis buffer containing 7 M urea (Thermal Scientific), 2 M thiourea (Sigma-Aldrich), 4% (w/v) CHAPS (Sigma-Aldrich), 65 mM dithiothreitol (SunShine Biotechnology), and 1% (v/v) Protease Inhibitor Cocktail for sperm protein analysis, as described in a previous study⁴². Briefly, homogenates were subjected to four repetitions of ultrasound every 15 minutes, and then centrifuged for 1 h at 15,000 rpm at 4 °C. The extracted proteins were subjected to immunoblot analysis for detecting MKRN2 (Novus Biologicals), Espin or Odf2 (BD Biosciences). β-Tubulin (Cell Signaling Technology) was used as the loading control.

Statistical analysis. Data were presented as the means ± SE. of at least three independent experiments unless otherwise indicated. The Student's unpaired *t* test was used for comparison of the wild-type and knockout mice. Values were considered significantly different at *P* < 0.05.

References

- De Kretser, D. M. & Baker, H. W. Infertility in men: recent advances and continuing controversies. *The Journal of clinical endocrinology and metabolism* **84**, 3443–3450, doi: 10.1210/jcem.84.10.6101 (1999).
- Hirsh, A. Male subfertility. *Bmj* **327**, 669–672, doi: 10.1136/bmj.327.7416.669 (2003).
- Eddy, E. M. Male germ cell gene expression. *Recent progress in hormone research* **57**, 103–128 (2002).
- Zhang, Z. *et al.* MEIG1 is essential for spermiogenesis in mice. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 17055–17060, doi: 10.1073/pnas.0906414106 (2009).
- Alves, M. G. *et al.* Molecular mechanisms beyond glucose transport in diabetes-related male infertility. *Biochimica et biophysica acta* **1832**, 626–635, doi: 10.1016/j.bbdis.2013.01.011 (2013).
- Jamsai, D. & O'Bryan, M. K. Mouse models in male fertility research. *Asian journal of andrology* **13**, 139–151, doi: 10.1038/aja.2010.101 (2011).
- Omwancha, J. *et al.* Makorin RING finger protein 1 (MKRN1) has negative and positive effects on RNA polymerase II-dependent transcription. *Endocrine* **29**, 363–373, doi: 10.1385/ENDO:29:2:363 (2006).
- Kim, J. H. *et al.* Ubiquitin ligase MKRN1 modulates telomere length homeostasis through a proteolysis of hTERT. *Genes & development* **19**, 776–781, doi: 10.1101/gad.1289405 (2005).
- Ko, A. *et al.* Acceleration of gastric tumorigenesis through MKRN1-mediated posttranslational regulation of p14ARF. *Journal of the National Cancer Institute* **104**, 1660–1672, doi: 10.1093/jnci/djs424 (2012).

10. Zhang, Q. H. *et al.* Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34+ hematopoietic stem/progenitor cells. *Genome research* **10**, 1546–1560 (2000).
11. Deshaies, R. J. & Joazeiro, C. A. RING domain E3 ubiquitin ligases. *Annual review of biochemistry* **78**, 399–434, doi: 10.1146/annurev.biochem.78.101807.093809 (2009).
12. Riley, B. E. *et al.* Structure and function of Parkin E3 ubiquitin ligase reveals aspects of RING and HECT ligases. *Nature communications* **4**, 1982, doi: 10.1038/ncomms2982 (2013).
13. Gray, T. A. *et al.* Phylogenetic conservation of the makorin-2 gene, encoding a multiple zinc-finger protein, antisense to the RAF1 proto-oncogene. *Genomics* **77**, 119–126, doi: 10.1006/geno.2001.6627 (2001).
14. Yang, P. H. *et al.* Makorin-2 is a neurogenesis inhibitor downstream of phosphatidylinositol 3-kinase/Akt (PI3K/Akt) signal. *The Journal of biological chemistry* **283**, 8486–8495, doi: 10.1074/jbc.M704768200 (2008).
15. Church, D. M. *et al.* Lineage-specific biology revealed by a finished genome assembly of the mouse. *PLoS biology* **7**, e1000112, doi: 10.1371/journal.pbio.1000112 (2009).
16. Platts, A. E. *et al.* Success and failure in human spermatogenesis as revealed by teratozoospermic RNAs. *Human molecular genetics* **16**, 763–773, doi: 10.1093/hmg/ddm012 (2007).
17. Baddack, U. *et al.* A chronic model of arthritis supported by a strain-specific periarticular lymph node in BALB/c mice. *Nature communications* **4**, 1644, doi: 10.1038/ncomms2625 (2013).
18. Wei, T. C., Huang, W. J., Lin, A. T. & Chen, K. K. The role of hormones on semen parameters in patients with idiopathic or varicocele-related oligoasthenoteratozoospermia (OAT) syndrome. *Journal of the Chinese Medical Association: JCMA* **76**, 624–628, doi: 10.1016/j.jcma.2013.07.005 (2013).
19. Jungwirth, A. *et al.* European Association of Urology guidelines on Male Infertility: the 2012 update. *European urology* **62**, 324–332, doi: 10.1016/j.eururo.2012.04.048 (2012).
20. Parreira, G. G., Melo, R. C. & Russell, L. D. Relationship of sertoli-sertoli tight junctions to ectoplasmic specialization in conventional and en face views. *Biology of reproduction* **67**, 1232–1241 (2002).
21. Borg, C. L., Wolski, K. M., Gibbs, G. M. & O'Bryan, M. K. Phenotyping male infertility in the mouse: how to get the most out of a 'non-performer'. *Human reproduction update* **16**, 205–224, doi: 10.1093/humupd/dmp032 (2010).
22. Porter, M. E. & Sale, W. S. The 9 + 2 axoneme anchors multiple inner arm dyneins and a network of kinases and phosphatases that control motility. *The Journal of cell biology* **151**, F37–42 (2000).
23. Lee, E. W. *et al.* Differential regulation of p53 and p21 by MKRN1 E3 ligase controls cell cycle arrest and apoptosis. *The EMBO journal* **28**, 2100–2113, doi: 10.1038/emboj.2009.164 (2009).
24. Miroci, H. *et al.* Makorin ring zinc finger protein 1 (MKRN1), a novel poly(A)-binding protein-interacting protein, stimulates translation in nerve cells. *The Journal of biological chemistry* **287**, 1322–1334, doi: 10.1074/jbc.M111.315291 (2012).
25. Shimada, H. *et al.* Identification of Makorin 1 as a novel SEREX antigen of esophageal squamous cell carcinoma. *BMC cancer* **9**, 232, doi: 10.1186/1471-2407-9-232 (2009).
26. Holzenberger, M. *et al.* Cre-mediated germline mosaicism: a method allowing rapid generation of several alleles of a target gene. *Nucleic acids research* **28**, E92 (2000).
27. Lakso, M. *et al.* Efficient *in vivo* manipulation of mouse genomic sequences at the zygote stage. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 5860–5865 (1996).
28. Cox, R. D. & Brown, S. D. Rodent models of genetic disease. *Current opinion in genetics & development* **13**, 278–283 (2003).
29. Kamberov, Y. G. *et al.* Modeling recent human evolution in mice by expression of a selected EDAR variant. *Cell* **152**, 691–702, doi: 10.1016/j.cell.2013.01.016 (2013).
30. O'Donnell, L., Nicholls, P. K., O'Bryan, M. K., McLachlan, R. I. & Stanton, P. G. Spermiation: The process of sperm release. *Spermatogenesis* **1**, 14–35, doi: 10.4161/spmg.1.1.14525 (2011).
31. Li, M. W., Mruk, D. D., Lee, W. M. & Cheng, C. Y. Connexin 43 is critical to maintain the homeostasis of the blood-testis barrier via its effects on tight junction reassembly. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 17998–18003, doi: 10.1073/pnas.1007047107 (2010).
32. Cheng, C. Y. & Mruk, D. D. A local autocrine axis in the testes that regulates spermatogenesis. *Nature reviews. Endocrinology* **6**, 380–395, doi: 10.1038/nrendo.2010.71 (2010).
33. Bartles, J. R., Wierda, A. & Zheng, L. Identification and characterization of espin, an actin-binding protein localized to the F-actin-rich junctional plaques of Sertoli cell ectoplasmic specializations. *Journal of cell science* **109** (Pt 6), 1229–1239 (1996).
34. Yan, H. H., Mruk, D. D., Lee, W. M. & Cheng, C. Y. Ectoplasmic specialization: a friend or a foe of spermatogenesis? *BioEssays: news and reviews in molecular, cellular and developmental biology* **29**, 36–48, doi: 10.1002/bies.20513 (2007).
35. Mruk, D. D. & Cheng, C. Y. Cell-cell interactions at the ectoplasmic specialization in the testis. *Trends in endocrinology and metabolism: TEM* **15**, 439–447, doi: 10.1016/j.tem.2004.09.009 (2004).
36. Lee, N. P. & Cheng, C. Y. Ectoplasmic specialization, a testis-specific cell-cell actin-based adherens junction type: is this a potential target for male contraceptive development? *Human reproduction update* **10**, 349–369, doi: 10.1093/humupd/dmh026 (2004).
37. Tarnasky, H. *et al.* Gene trap mutation of murine outer dense fiber protein-2 gene can result in sperm tail abnormalities in mice with high percentage chimaerism. *BMC developmental biology* **10**, 67, doi: 10.1186/1471-213X-10-67 (2010).
38. Mariappa, D. *et al.* Inhibition of tyrosine phosphorylation of sperm flagellar proteins, outer dense fiber protein-2 and tektin-2, is associated with impaired motility during capacitation of hamster spermatozoa. *Molecular reproduction and development* **77**, 182–193, doi: 10.1002/mrd.21131 (2010).
39. Salmon, N. A., Reijo Pera, R. A. & Xu, E. Y. A gene trap knockout of the abundant sperm tail protein, outer dense fiber 2, results in preimplantation lethality. *Genesis* **44**, 515–522, doi: 10.1002/dvg.20241 (2006).
40. Menkveld, R. Clinical significance of the low normal sperm morphology value as proposed in the fifth edition of the WHO Laboratory Manual for the Examination and Processing of Human Semen. *Asian journal of andrology* **12**, 47–58, doi: 10.1038/aja.2009.14 (2010).
41. Yildiz, Y. *et al.* Mutation of beta-glucosidase 2 causes glycolipid storage disease and impaired male fertility. *The Journal of clinical investigation* **116**, 2985–2994, doi: 10.1172/JCI29224 (2006).
42. Guo, X. *et al.* Proteomic analysis of proteins involved in spermiogenesis in mouse. *Journal of proteome research* **9**, 1246–1256, doi: 10.1021/pr900735k (2010).

Acknowledgements

This work was supported in part by grants from the National Key Basic Research Program of China (2011CB944304, 2011CB504003), and by the Chinese Natural Science Funds (81270736, 81222006, 81302182, 81472944, 81320108019), and by the National Institutes of Health grant R01ES020868 and R01CA193511. We thank Model Animal Research Center of Nanjing University for generation of the *Mkfn2* floxed mice.

Author Contributions

X.Q., L.W. and B.Z. designed and performed experiments, analyzed data and wrote the manuscript; L.W., Z.M.S., X.G. and C.F.J. maintained mice and performed IHC and qRT-PCR assay; Y.C.Q., D.M.L., X.L., W.L., Y.Y. and

J.T.Z. acquired data, reporter assay and immunoblotting analysis; H.S., M.W, M.L. and X.J.G. analyzed TEM data and did statistical analysis; J.H. and L.Z.L. prepared the manuscript. J.H.S. and B.H.J. designed and supervised the project.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Qian, X. *et al.* Deficiency of *Mkrrn2* causes abnormal spermiogenesis and spermiation, and impairs male fertility. *Sci. Rep.* **6**, 39318; doi: 10.1038/srep39318 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016