

# Review of Impact of COVID-19 on Maternal, Neonatal Outcomes, and Placental Changes

Review began 08/12/2022  
Review ended 08/22/2022  
Published 08/31/2022

© Copyright 2022

Tanna et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Resham Tanna<sup>1</sup>, Henry J. Nava Dugarte<sup>2</sup>, Sowjanya Kurakula<sup>3,4,5,6</sup>, Vandana Muralidharan<sup>7</sup>, Arghadip Das<sup>8</sup>, Sri Padma Ravali Kanigalpula<sup>9</sup>, Ileana Elita Mendez<sup>10</sup>, Munaza Afaq<sup>11</sup>, Radhika Bassi<sup>12</sup>, Kinjal Shah<sup>13</sup>, Zainab Saddiq<sup>14</sup>

1. Obstetrics and Gynaecology, Spartan Health Sciences University, Vieux Fort, LCA 2. Obstetrics and Gynaecology, Rejuvenating Fertility Centre, New York, USA 3. Obstetrics and Gynecology, Mamta Institute of Medical Sciences, Khammam, IND 4. Obstetrics and Gynecology, Sekgoma Memorial Hospital, Serowe, BWA 5. Obstetrics and Gynecology, Nyangabgwe Referral Hospital, Francistown, BWA 6. Obstetrics and Gynecology, Gandhi Medical College, Musheerabad, IND 7. Obstetrics and Gynaecology, Lifeline Medical Associates, New Jersey, USA 8. Internal Medicine, Nilratan Sircar Medical College and Hospital, Kolkata, IND 9. Obstetrics and Gynaecology, Ascension Health via Christi Hospital, Manhattan, USA 10. Medical Sciences Department, Universidad Autonoma de Centro America (UACA), San José, CRI 11. Obstetrics and Gynaecology, Government Medical College Srinagar, Srinagar, IND 12. Obstetrics and Gynaecology, Ross University School of Medicine, Bridgetown, BRB 13. Health Administration, Edward J. Bloustein School of Planning and Public Policy, New Jersey, USA 14. Obstetrics and Gynecology, Annotto Bay Hospital, St.Mary, JAM

Corresponding author: Resham Tanna, reshamtannam@gmail.com

## Abstract

Coronavirus disease (COVID-19), caused by SARS-CoV-2, is a disease that has caused a global impact. COVID-19 is transmitted through airborne droplets, respiratory secretions, and direct contact. The pandemic has affected individuals of different ages, and studying the impact of COVID-19 on maternal and newborn outcomes is critical. In this review, we highlight the impact of COVID-19 infection in pregnancy and its repercussion in the maternal-fetal binomial. Physiological changes that occur during pregnancy have significant effects on the immune system, cardiopulmonary system, and coagulation, and these changes can result in an altered response to COVID-19 infection. The symptoms, risk factors, and maternal health consequences of COVID-19 were discussed. In addition, the impact of newborns born to mothers with COVID-19 was reviewed. Finally, placental changes and vertical transmission of COVID-19 during pregnancy were also discussed in this review.

**Categories:** Obstetrics/Gynecology, Infectious Disease, Epidemiology/Public Health

**Keywords:** placental changes, neonatal outcomes, maternal outcomes, pregnancy, covid-19

## Introduction And Background

The first outbreak of coronavirus disease (COVID-19) caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) was reported in December 2019 in Wuhan, China. It was declared a global pandemic in March 2020 with a total of 505,817,953 confirmed cases and 6,213,876 deaths reported by the World Health Organization (WHO) as of April 2022 [1]. COVID-19 illness is transmitted through airborne droplets, respiratory secretions, and direct contact [2]. While the symptoms related to COVID-19 disease are primarily respiratory, the disease has been reported to have multisystemic effects. COVID-19 illness symptoms can be asymptomatic, mild, moderate, severe, or critical [3,4,5]. Fever, cough, dyspnea, and myalgia were the most common mild symptoms [6-10]. The most common imaging findings were ground-glass opacities, and bilateral infiltrates, and the most common laboratory findings were leucopenia, thrombocytopenia, and elevated CRP [6,10]. Less common symptoms were nausea, vomiting, and diarrhea [11].

The Centers for Disease Control and Prevention (CDC) defines critical illness in a person with COVID-19 as a person who may require hospitalization, admission to an intensive care unit, and mechanical ventilation [3]. In addition, pregnancy is also listed as an underlying medical condition that increases the risk of contracting COVID-19 illness [3]. Multiple studies have found that pregnant women with severe or critical COVID-19 disease require oxygen, mechanical ventilation, and ICU admissions [4,12-15]. A variety of studies have shown that maternal mortality rates range from 0.3%, 0.74%, 1.37%, 1.60%, one in 80, and seven in nine women with critical COVID-19 illness in pregnancy [5,13,16-19]. Conflicting results were reported in a study reporting that there is no significant correlation between COVID-19 pregnancy and mortality [20].

The effects of the COVID-19 pandemic are not limited to its associated morbidity and mortality, it has also led to social and economic disruption around the world and has through ripple effects impacted the healthcare system in many ways, affecting the quality of patient care. The maternal health care system which includes prenatal care, obstetrics, and postpartum care, has also undergone major changes in its structure [21]. During social distancing measures in some parts of the UK, women were provided with blood pressure

### How to cite this article

Tanna R, Dugarte H, Kurakula S, et al. (August 31, 2022) Review of Impact of COVID-19 on Maternal, Neonatal Outcomes, and Placental Changes. Cureus 14(8): e28631. DOI 10.7759/cureus.28631

monitors and urine dipsticks to do prenatal screening themselves [22]. Kotlar et al., reported that fewer prenatal visits, a strained healthcare infrastructure, and potentially harmful policies were implemented with limited evidence during the pandemic [23]. On the other hand, telemedicine has evolved and has been extremely useful for prenatal consultations, although underutilized in low-income countries [22].

Pregnancy is generally considered to be a high-risk condition associated with infectious diseases because the immunological changes of pregnancy can increase susceptibility to pathogens and associated complications [11]. Pregnant women are listed as a high-risk population for COVID-19 disease [24]. Physiological changes that occur during pregnancy have significant effects on the immune system, cardiopulmonary system, and coagulation, and these changes can result in an altered response to COVID-19 infection [25].

The impact of the disease on maternal and fetal outcomes has been questioned since the pandemic began. In this review, we explore the latest research on the maternal and neonatal consequences of COVID-19. We also highlight the peculiar placental changes that occur as a result of this infection.

## Review

### Impact of COVID-19 on maternal outcomes

According to the WHO, COVID-19 infection is more severe in pregnant women than in non-pregnant women [26]. Some ethnic groups, such as Hispanics, Blacks, and Asians, had higher rates of moderate and severe COVID-19 illness [3,27]. Additionally, the CDC found that pregnant women of the same ethnic group are at increased risk for COVID-19 illness due to health inequalities they face [28]. Son et al. however, found no differences in pregnancy-related outcomes between pre- and during-pandemic deliveries and those testing positive or negative for COVID-19 disease in geographically diverse US cohorts [29].

Several studies have compared pregnant women who tested positive for COVID-19 disease with those who tested negative [12,13,20,27,29,30]. Pregnant women with COVID-19 disease were more likely to develop serious outcomes if they had comorbidities such as chronic obstructive pulmonary disease (COPD), asthma, type 1 and type 2 diabetes, and obesity [4,10,12,17,27,31,32]. A variety of studies have identified maternal age > 40, hypertension, and autoimmune disorders as additional risk factors for contracting SARS-CoV-2 infection in pregnancy [3-5,12,27,31,33]. Wei et al. found that severe COVID-19 disease is strongly associated with preeclampsia [30]. There were high rates of preeclampsia and chronic hypertension, preterm birth, and cesarean births in pregnant patients with a critical COVID-19 illness [4,8,10,13,14,27,34]. An increased likelihood of placental abruption due to SARS-CoV-2 infection was noted by Marta and colleagues [35]. In one study, fetal growth restriction and premature prelabor rupture of membranes (PPROM) were observed in 11.7% and 20.7%, respectively [36]. It is however reported that the clinical course and severity of COVID-19 pneumonia and severe COVID-19 illness during pregnancy are similar to those of non-pregnant women [33,34,37,38]. In addition, various organ system involvement in COVID-19 disease in obstetric populations was discussed by Syeda et al., which is represented in Table 1 [11]. It is however important to note that some of these findings can also be due to direct obstetric complications [11].

Impact of COVID-19 on various maternal organ systems				
Cardiac	Pulmonary	Renal	Hepatic	Hematological
Cardiomyopathy	Silent hypoxia	Acute Kidney Injury	Elevated transaminases	Neutrophilia
	Risk of pulmonary edema due to combined acute kidney injury and pulmonary injury		Findings mimicking HELLP syndrome and severe preeclampsia	Lymphopenia
	Mixed and complete consolidations on chest imaging			

**TABLE 1: Impact of COVID-19 in pregnancy on various maternal organ systems. Original figure drawn by authors.**

Abbreviations: HELLP = Hemolysis, Elevated Liver enzymes, Low Platelets

A notable increase in mental illness in the form of self-harm, depression, and anxiety have also been observed in pregnant women with COVID-19 illness [16,39,40]. Domestic violence increased among pregnant women during the pandemic, they had lower incomes than men, and working mothers had to deal with increased childcare needs [22,23]. One study had reported lower vitamin D levels in COVID-19 affected pregnancies compared to healthy pregnancies [41].

## Impact of COVID-19 on neonatal outcomes

The impact of maternal SARS-CoV-2 infection on the fetus and subsequently the neonate remains poorly understood [42]. The increasing number of reports of neonates born to women with COVID-19 showing signs of early-onset infection is suggestive of a scenario of transplacental transmission of SARS-CoV-2 [43-45]. However, vertical transmission of the virus remains controversial as clinical and laboratory evidence considers it a rare event [9,44,46-51]. In particular, a cause-and-effect relationship between viral infection and the adverse neonatal outcome has not been formally established, since other factors causing postpartum complications cannot be ruled out [43,49]. Therefore, it remains uncertain whether early neonatal SARS-CoV-2 infection occurs in utero, during birth, or shortly after birth [52].

Pregnant women with COVID-19 have high rates of spontaneous preterm birth, with studies reporting rates of 21% - 40% spontaneous preterm birth in their cohorts, which is two to four times the average rate of 10% [36,53-55]. Approximately 3% of pregnancies in women with COVID-19 end in stillbirth, with the usual rate being < 1% in pregnant patients without COVID-19 illness [56,57]. Studies have shown that placental damage from SARS-CoV-2 infection is sufficient to cause severe morbidity and even mortality in newborns [51,58,59]. This remains controversial as reports show that stillborn newborns and their placenta are negative for SARS-CoV-2 infection despite extensive placental damage and funisitis [51,58,59]. In addition, the absence of symptoms in the infected mother does not rule out the death of the newborn [51,58,59]. A case study by Poisson et al. reported a SARS-CoV-2-positive asymptomatic pregnant woman at 32 weeks gestation, went on to have a still birth after 35 weeks gestation. The stillborn baby showed no abnormalities, but 75% of the placenta showed extensive fetal vascular insufficiency and parenchymal infarction [60]. In another study, two newborns of symptomatic COVID-19 women tested positive for nasopharyngeal swabs shortly after birth and seven days after birth, but neither developed infection-related symptoms [61]. The placenta in both cases was positive for SARS-CoV-2 by in situ hybridization and Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR) analysis [61]. Facchetti et al. reported a case of a SARS-CoV-2-positive pregnant woman with pneumonia and severe thrombocytopenia requiring induction of labor. The newborn was positive for the presence of the virus and developed pneumonia and severe shortness of breath 24 hours after birth [62]. In summary, their observations provide evidence of the development of intrauterine vertical transmission of SARS-CoV-2 infection and have infrequent clinical effects on newborns [62].

In a multinational cohort study, the risk of major neonatal complications, including a stay in the neonatal intensive care unit (NICU) of at least seven days, as well as the summary index of major neonatal morbidity and its individual components, were also significantly higher in the group of women diagnosed with COVID-19 disease [13]. The increased neonatal risk persisted after adjustment for earlier preterm births and preterm births in the index pregnancy; therefore, a direct impact of COVID-19 on the newborn is likely. They found that 12.1% of newborns born to test-positive women also tested positive, a higher number than in a recent systematic review. Reassuringly, since SARS-CoV-2 has not been isolated from human milk, breastfeeding was not associated with an increase in the rate of test-positive newborns [63]. Many studies have not detected the SARS-CoV-2 virus in breast milk and have also confirmed that SARS-CoV-2 antibodies are transmitted through breast milk [47,64,65]. The WHO states that breastfeeding is the optimal form of nutrition for babies and recommends that women infected with COVID-19 breastfeed their babies if possible [66].

## Placental changes due to COVID-19 infection

Pregnancy is a time of immune regulation that allows the fetus to develop in the womb while protecting the mother from infection. The state of placental immune protection for the fetus in a normal pregnancy could potentially have an adverse effect on pregnancy affected by COVID-19 [67].

During pregnancy, many fetal and neonatal pathologies are associated with cross-placental transmission of viruses such as rubella, varicella, cytomegalovirus, dengue, and zika, among others. Evidence of fetal and maternal vascular malperfusion, as well as evidence of inflammation in the placenta following maternal SARS-CoV-1 infection, have been reported [68].

Histopathological examination of placental tissue can provide important information about the condition of the mother and fetus [51]. Laresgoiti-Servitje et al. 2021, reported that the placentas of women infected with SARS-CoV-2 had a higher rate of fibrinoid deposition, a clinical feature of maternal vascular malperfusion (MVM), when compared with controls [69]. Additionally, it was noted by Shanes and Col., that relative to controls, COVID-19 placentas showed an increased prevalence of decidual arteriopathy and other features of MVM, a pattern of placental injury that reflects abnormalities in oxygenation within the intervillous space associated with adverse perinatal outcomes [51].

The angiotensin-converting enzyme 2 (ACE2) and the transmembrane protease serine 2 (TMPRSS2) are SARS-CoV-2 specific entry mediators that are highly expressed in the human placenta in early pregnancy, but their expression decreases significantly as the pregnancy progresses [70]. SARS-CoV-2 infection present in the maternal circulation has the potential to invade the maternal blood-soaked syncytiotrophoblast and infect the placenta via ACE2 binding. The higher levels of ACE2 mRNA in the placenta at earlier stages of pregnancy raise the possibility of higher susceptibility to SARS-CoV-2 infection in the placenta in the first

trimester. It has been demonstrated that there is no evidence of vertical transmission when the infection manifests itself in the third trimester of pregnancy [9].

Pathological studies have shown that syncytiotrophoblasts are commonly infected with SARS-CoV-2, but fetuses are not always infected [71]. These findings suggest the presence of a placental barrier [71]. Facchetti et al., reported for the first time that SARS-CoV-2 S and N proteins were highly expressed in the placenta of a pregnant COVID-19 woman whose newborn tested positive for viral RNA and developed COVID-19 respiratory symptoms soon after birth [62]. The report revealed SARS-CoV-2 virus products and/or particles in villous syncytiotrophoblast, endothelial cells, fibroblasts, and maternal macrophages that contribute to inflammatory infiltration, as well as Hoffbauer cells. Surprisingly, particles morphologically consistent with the coronavirus localized into the fetal circulating mononuclear cells [62].

It is important to mention that other studies were found with no evidence of placental infection or vertical transmission of SARS-CoV-2 [72,73]. Such as that presented by Levitan, where after evaluation of placentas from 65 polymerase chain reaction women, proven SARS-CoV-2 infection and histological evaluation indicated that there is no characteristic histopathology in most placentas [72]. The same results were obtained from Smithgall in the study of 51 placentas of SARS-CoV-2 positive third-trimester mothers, where found non-specific histomorphologic changes that point to indicated maternal/fetal vascular malperfusion. All newborns tested negative for SARS-CoV-2 infection and all mothers recovered clinically [73]. This could be related to the fact that the transplacental transfer of anti-SARS-CoV-2 antibodies was inefficient and factors such as lack of viremia, reduced coexpression, and colocalization of placental angiotensin-converting enzyme 2 and transmembrane serine protease 2 as protection could serve mechanisms against vertical transmission [74].

Interpretation of placental changes and vertical transmission requires caution as there may be other undefined mechanisms linking obstetric outcomes of SARS-CoV-2 under normal and pathologic conditions.

## Conclusions

COVID-19 disease is an evolving disease with varying reports of a significant impact of severe COVID-19 illness on maternal outcomes. Various studies have shown conflicting results on the neonatal consequences of COVID-19 infection. The concept of placental changes and vertical transmission also remains inconclusive. Therefore, it is important to continuously investigate the impact of COVID-19 disease on maternal and neonatal outcomes, and its effect on vertical transmission and placental changes.

## Additional Information

### Disclosures

**Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

## References

1. WHO Coronavirus (COVID-19) dashboard. (2020). Accessed: 22nd April 2022: <https://covid19.who.int>.
2. Li Q, Guan X, Wu P, et al.: Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. 2020, 382:1199-207. [10.1056/NEJMoa2001316](https://doi.org/10.1056/NEJMoa2001316)
3. Brandt JS, Hill J, Reddy A, et al.: Epidemiology of coronavirus disease 2019 in pregnancy: risk factors and associations with adverse maternal and neonatal outcomes. *Am J Obstet Gynecol*. 2021, 224:389.e1-9. [10.1016/j.ajog.2020.09.043](https://doi.org/10.1016/j.ajog.2020.09.043)
4. Khoury R, Bernstein PS, Debolt C, et al.: Characteristics and outcomes of 241 births to women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection at five New York City medical centers. *Obstet Gynecol*. 2020, 136:273-82. [10.1097/AOG.0000000000004025](https://doi.org/10.1097/AOG.0000000000004025)
5. Metz TD, Clifton RG, Hughes BL, et al.: Disease severity and perinatal outcomes of pregnant patients with coronavirus disease 2019 (COVID-19). *Obstet Gynecol*. 2021, 137:571-80. [10.1097/AOG.0000000000004339](https://doi.org/10.1097/AOG.0000000000004339)
6. Andrikopoulou M, Madden N, Wen T, et al.: Symptoms and critical illness among obstetric patients with coronavirus disease 2019 (COVID-19) infection. *Obstet Gynecol*. 2020, 136:291-9. [10.1097/AOG.0000000000003996](https://doi.org/10.1097/AOG.0000000000003996)
7. Afshar Y, Gaw SL, Flaherman VJ, et al.: Clinical presentation of coronavirus disease 2019 (COVID-19) in pregnant and recently pregnant people. *Obstet Gynecol*. 2020, 136:1117-25. [10.1097/AOG.0000000000004178](https://doi.org/10.1097/AOG.0000000000004178)
8. Juan J, Gil MM, Rong Z, Zhang Y, Yang H, Poon LC: Effect of coronavirus disease 2019 (COVID-19) on maternal, perinatal and neonatal outcome: systematic review. *Ultrasound Obstet Gynecol*. 2020, 56:15-27. [10.1002/uog.22088](https://doi.org/10.1002/uog.22088)
9. Yan J, Guo J, Fan C, et al.: Coronavirus disease 2019 in pregnant women: a report based on 116 cases. *Am J Obstet Gynecol*. 2020, 223:111.e1-111.e14. [10.1016/j.ajog.2020.04.014](https://doi.org/10.1016/j.ajog.2020.04.014)
10. Jafari M, Pormohammad A, Sheikh Neshin SA, et al.: Clinical characteristics and outcomes of pregnant

- women with COVID-19 and comparison with control patients: A systematic review and meta-analysis. *Rev Med Virol.* 2021, 31:1-16. [10.1002/rmv.2208](https://doi.org/10.1002/rmv.2208)
11. Syeda S, Baptiste C, Breslin N, Gyamfi-Bannerman C, Miller R: The clinical course of COVID in pregnancy. *Semin Perinatol.* 2020, 44:151284. [10.1016/j.semperi.2020.151284](https://doi.org/10.1016/j.semperi.2020.151284)
  12. Allotey J, Stallings E, Bonet M, et al.: Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ.* 2020, 370:m3520. [10.1136/bmj.m3520](https://doi.org/10.1136/bmj.m3520)
  13. Villar J, Ariff S, Gunier RB, et al.: Maternal and neonatal morbidity and mortality among pregnant women with and without COVID-19 infection: the INTERCOVID multinational cohort study. *JAMA Pediatr.* 2021, 175:817-26. [10.1001/jamapediatrics.2021.1050](https://doi.org/10.1001/jamapediatrics.2021.1050)
  14. Huntley BJ, Huntley ES, Di Mascio D, Chen T, Berghella V, Chauhan SP: Rates of maternal and perinatal mortality and vertical transmission in pregnancies complicated by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection: a systematic review. *Obstet Gynecol.* 2020, 136:303-12. [10.1097/AOG.0000000000004010](https://doi.org/10.1097/AOG.0000000000004010)
  15. Savasi VM, Parisi F, Patanè L, et al.: Clinical findings and disease severity in hospitalized pregnant women with coronavirus disease 2019 (COVID-19). *Obstet Gynecol.* 2020, 136:252-8. [10.1097/AOG.0000000000003979](https://doi.org/10.1097/AOG.0000000000003979)
  16. Chmielewska B, Barratt I, Townsend R, et al.: Effects of the COVID-19 pandemic on maternal and perinatal outcomes: a systematic review and meta-analysis. *Lancet Glob Health.* 2021, 9:e759-72. [10.1016/S2214-109X\(21\)00079-6](https://doi.org/10.1016/S2214-109X(21)00079-6)
  17. Lokken EM, Huebner EM, Taylor GG, et al.: Disease severity, pregnancy outcomes, and maternal deaths among pregnant patients with severe acute respiratory syndrome coronavirus 2 infection in Washington State. *Am J Obstet Gynecol.* 2021, 225:77.e1-77.e14. [10.1016/j.ajog.2020.12.1221](https://doi.org/10.1016/j.ajog.2020.12.1221)
  18. Hantoushzadeh S, Shamshirsaz AA, Aleyasin A, et al.: Maternal death due to COVID-19. *Am J Obstet Gynecol.* 2020, 223:109.e1-109.e16. [10.1016/j.ajog.2020.04.030](https://doi.org/10.1016/j.ajog.2020.04.030)
  19. Rodrigues C, Baía I, Domingues R, Barros H: Pregnancy and breastfeeding during COVID-19 pandemic: a systematic review of published pregnancy cases. *Front Public Health.* 2020, 8:558144. [10.3389/fpubh.2020.558144](https://doi.org/10.3389/fpubh.2020.558144)
  20. Qeadan F, Mensah NA, Tingey B, Stanford JB: The risk of clinical complications and death among pregnant women with COVID-19 in the Cerner COVID-19 cohort: a retrospective analysis. *BMC Pregnancy Childbirth.* 2021, 21:305. [10.1186/s12884-021-03772-y](https://doi.org/10.1186/s12884-021-03772-y)
  21. Altman MR, Gavin AR, Eagen-Torkko MK, Kantrowitz-Gordon I, Khosa RM, Mohammed SA: Where the system failed: the COVID-19 pandemic's impact on pregnancy and birth care. *Glob Qual Nurs Res.* 2021, 8:23333936211006397. [10.1177/23333936211006397](https://doi.org/10.1177/23333936211006397)
  22. Martins O I, Alexander S R, Khan M M, et al.: Challenges to women's health during pregnancy in COVID era-Review Article. *Biomed J Sci & Tech Res.* 38:2021-0061346.
  23. Kotlar B, Gerson E, Petrillo S, Langer A, Tiemeier H: The impact of the COVID-19 pandemic on maternal and perinatal health: a scoping review. *Reprod Health.* 2021, 18:10. [10.1186/s12978-021-01070-6](https://doi.org/10.1186/s12978-021-01070-6)
  24. People with certain medical conditions. (2020). Accessed: 22nd April 2022: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>.
  25. Wastnedge EA, Reynolds RM, van Boeckel SR, Stock SJ, Denison FC, Maybin JA, Critchley HO: Pregnancy and COVID-19. *Physiol Rev.* 2021, 101:303-18. [10.1152/physrev.00024.2020](https://doi.org/10.1152/physrev.00024.2020)
  26. Coronavirus disease (COVID-19): Pregnancy, childbirth and the postnatal period. (2020). Accessed: 20th April, 2022: <https://www.who.int/news-room/questions-and-answers/item/coronavirus-disease-covid-19-pregnancy-and-childbirth#:~:tex...>
  27. Khan DS, Hamid LR, Ali A, Salam RA, Zuberi N, Lassi ZS, Das JK: Differences in pregnancy and perinatal outcomes among symptomatic versus asymptomatic COVID-19-infected pregnant women: a systematic review and meta-analysis. *BMC Pregnancy Childbirth.* 2021, 21:801. [10.1186/s12884-021-04250-1](https://doi.org/10.1186/s12884-021-04250-1)
  28. Pregnant and recently pregnant people. (2019). Accessed: 20th April 2022: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/pregnant-people.html>.
  29. Son M, Gallagher K, Lo JY, et al.: Coronavirus disease 2019 (COVID-19) pandemic and pregnancy outcomes in a U.S. population. *Obstet Gynecol.* 2021, 138:542-51. [10.1097/AOG.0000000000004547](https://doi.org/10.1097/AOG.0000000000004547)
  30. Wei SQ, Bilodeau-Bertrand M, Liu S, Auger N: The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis. *CMAJ.* 2021, 193:E540-8. [10.1503/cmaj.202604](https://doi.org/10.1503/cmaj.202604)
  31. Boushra MN, Koyfman A, Long B: COVID-19 in pregnancy and the puerperium: a review for emergency physicians. *Am J Emerg Med.* 2021, 40:193-8. [10.1016/j.ajem.2020.10.055](https://doi.org/10.1016/j.ajem.2020.10.055)
  32. Grechukhina O, Greenberg V, Lundsberg LS, et al.: Coronavirus disease 2019 pregnancy outcomes in a racially and ethnically diverse population. *Am J Obstet Gynecol MFM.* 2020, 2:100246. [10.1016/j.ajogmf.2020.100246](https://doi.org/10.1016/j.ajogmf.2020.100246)
  33. Pettrosso E, Giles M, Cole S, Rees M: COVID-19 and pregnancy: a review of clinical characteristics, obstetric outcomes and vertical transmission. *Aust N Z J Obstet Gynaecol.* 2020, 60:640-59. [10.1111/ajo.13204](https://doi.org/10.1111/ajo.13204)
  34. Narang K, Enninga EA, Gunaratne MD, et al.: SARS-CoV-2 infection and COVID-19 during pregnancy: a multidisciplinary review. *Mayo Clin Proc.* 2020, 95:1750-65. [10.1016/j.mayocp.2020.05.011](https://doi.org/10.1016/j.mayocp.2020.05.011)
  35. Rodríguez-Díaz M, Alonso-Molero J, Cabero-Perez MJ, Llorca J, Dierssen-Sotos T, Gómez-Acebo I, The Moacc-Group: Pregnancy and birth outcomes during the early months of the COVID-19 pandemic: The MOACC-19 cohort. *Int J Environ Res Public Health.* 2021, 18: [10.3390/ijerph182010931](https://doi.org/10.3390/ijerph182010931)
  36. Di Mascio D, Khalil A, Saccone G, et al.: Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis. *Am J Obstet Gynecol MFM.* 2020, 2:100107. [10.1016/j.ajogmf.2020.100107](https://doi.org/10.1016/j.ajogmf.2020.100107)
  37. Wang CL, Liu YY, Wu CH, Wang CY, Wang CH, Long CY: Impact of COVID-19 on pregnancy. *Int J Med Sci.* 2021, 18:763-7. [10.7150/ijms.49923](https://doi.org/10.7150/ijms.49923)
  38. Maleki Dana P, Kolahdooz F, Sadoughi F, Moazzami B, Chaichian S, Asemi Z: COVID-19 and pregnancy: a review of current knowledge. *Infez Med.* 2020, 28:46-51.
  39. Ahmad M, Vismara L: The psychological impact of COVID-19 pandemic on women's mental health during

- pregnancy: a rapid evidence review. *Int J Environ Res Public Health*. 2021, 18: [10.3390/ijerph18137112](https://doi.org/10.3390/ijerph18137112)
40. Wu Y, Zhang C, Liu H, et al.: Perinatal depressive and anxiety symptoms of pregnant women during the coronavirus disease 2019 outbreak in China. *Am J Obstet Gynecol*. 2020, 223:240.e1-9. [10.1016/j.ajog.2020.05.009](https://doi.org/10.1016/j.ajog.2020.05.009)
  41. Sinaci S, Ocal DF, Yucel Yetiskin DF, et al.: Impact of vitamin D on the course of COVID-19 during pregnancy: a case control study. *J Steroid Biochem Mol Biol*. 2021, 215:105964. [10.1016/j.jsmb.2021.105964](https://doi.org/10.1016/j.jsmb.2021.105964)
  42. Arthurs AL, Jankovic-Karasoulos T, Roberts CT: COVID-19 in pregnancy: what we know from the first year of the pandemic. *Biochim Biophys Acta Mol Basis Dis*. 2021, 1867:166248. [10.1016/j.bbadis.2021.166248](https://doi.org/10.1016/j.bbadis.2021.166248)
  43. Schwartz DA, Dhaliwal A: Infections in pregnancy with covid-19 and other respiratory RNA virus diseases are rarely, if ever, transmitted to the fetus: experiences with coronaviruses, HPIV, hMPV RSV, and influenza. *Arch Pathol Lab Med*. 2020, [10.5858/arpa.2020-0211-SA](https://doi.org/10.5858/arpa.2020-0211-SA)
  44. Zaigham M, Andersson O: Maternal and perinatal outcomes with COVID-19: a systematic review of 108 pregnancies. *Acta Obstet Gynecol Scand*. 2020, 99:823-9. [10.1111/aogs.13867](https://doi.org/10.1111/aogs.13867)
  45. Alzamora MC, Paredes T, Caceres D, Webb CM, Valdez LM, La Rosa M: Severe COVID-19 during pregnancy and possible vertical transmission. *Am J Perinatol*. 2020, 37:861-5. [10.1055/s-0040-1710050](https://doi.org/10.1055/s-0040-1710050)
  46. Yang Z, Wang M, Zhu Z, Liu Y: Coronavirus disease 2019 (COVID-19) and pregnancy: a systematic review. *J Matern Fetal Neonatal Med*. 2022, 35:1619-22. [10.1080/14767058.2020.1759541](https://doi.org/10.1080/14767058.2020.1759541)
  47. Zhu H, Wang L, Fang C, et al.: Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl Pediatr*. 2020, 9:51-60. [10.21037/tp.2020.02.06](https://doi.org/10.21037/tp.2020.02.06)
  48. Liu W, Wang J, Li W, Zhou Z, Liu S, Rong Z: Clinical characteristics of 19 neonates born to mothers with COVID-19. *Front Med*. 2020, 14:193-8. [10.1007/s11684-020-0772-y](https://doi.org/10.1007/s11684-020-0772-y)
  49. Schwartz DA: An analysis of 38 pregnant women with COVID-19, their newborn infants, and maternal-fetal transmission of SARS-CoV-2: maternal coronavirus infections and pregnancy outcomes. *Arch Pathol Lab Med*. 2020, 144:799-805. [10.5858/arpa.2020-0901-SA](https://doi.org/10.5858/arpa.2020-0901-SA)
  50. Baergen RN, Heller DS: Placental pathology in Covid-19 positive mothers: preliminary findings. *Pediatr Dev Pathol*. 2020, 23:177-80. [10.1177/1093526620925569](https://doi.org/10.1177/1093526620925569)
  51. Shanes ED, Mithal LB, Otero S, Azad HA, Miller ES, Goldstein JA: Placental pathology in COVID-19. *Am J Clin Pathol*. 2020, 154:23-32. [10.1093/ajcp/aqaa089](https://doi.org/10.1093/ajcp/aqaa089)
  52. Zeng L, Xia S, Yuan W, Yan K, Xiao F, Shao J, Zhou W: Neonatal early-onset infection with SARS-CoV-2 in 33 neonates born to mothers with COVID-19 in Wuhan, China. *JAMA Pediatr*. 2020, 174:722-5. [10.1001/jamapediatrics.2020.0878](https://doi.org/10.1001/jamapediatrics.2020.0878)
  53. Chen L, Li Q, Zheng D, et al.: Clinical characteristics of pregnant women with Covid-19 in Wuhan, China. *N Engl J Med*. 2020, 382:e100. [10.1056/NEJMc2009226](https://doi.org/10.1056/NEJMc2009226)
  54. Mullins E, Evans D, Viner RM, O'Brien P, Morris E: Coronavirus in pregnancy and delivery: rapid review. *Ultrasound Obstet Gynecol*. 2020, 55:586-92. [10.1002/uog.22014](https://doi.org/10.1002/uog.22014)
  55. Chawanpaiboon S, Vogel JP, Moller AB, et al.: Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *Lancet Glob Health*. 2019, 7:e37-46. [10.1016/S2214-109X\(18\)30451-0](https://doi.org/10.1016/S2214-109X(18)30451-0)
  56. Delahoy MJ, Whitaker M, O'Halloran A, et al.: Characteristics and maternal and birth outcomes of hospitalized pregnant women with laboratory-confirmed COVID-19 - COVID-NET, 13 states, March 1- August 22, 2020. *MMWR Morb Mortal Wkly Rep*. 2020, 69:1347-54. [10.15585/mmwr.mm6938e1](https://doi.org/10.15585/mmwr.mm6938e1)
  57. Hoyert DL, Gregory EC: Cause of fetal death: data from the fetal death report, 2014. *Natl Vital Stat Rep*. 2016, 65:1-25.
  58. Baud D, Greub G, Favre G, Gengler C, Jatton K, Dubruc E, Pomar L: Second-trimester miscarriage in a pregnant woman with SARS-CoV-2 infection. *JAMA*. 2020, 323:2198-200. [10.1001/jama.2020.7233](https://doi.org/10.1001/jama.2020.7233)
  59. Richtmann R, Torloni MR, Oyamada Otani AR, et al.: Fetal deaths in pregnancies with SARS-CoV-2 infection in Brazil: a case series. *Case Rep Womens Health*. 2020, 27:e00245. [10.1016/j.crwh.2020.e00245](https://doi.org/10.1016/j.crwh.2020.e00245)
  60. Poisson TM, Pierone G Jr: Placental pathology and fetal demise at 35 weeks of gestation in a woman with SARS-CoV-2 infection: a case report. *Case Rep Womens Health*. 2021, 30:e00289. [10.1016/j.crwh.2021.e00289](https://doi.org/10.1016/j.crwh.2021.e00289)
  61. Patanè L, Morotti D, Giunta MR, et al.: Vertical transmission of coronavirus disease 2019: severe acute respiratory syndrome coronavirus 2 RNA on the fetal side of the placenta in pregnancies with coronavirus disease 2019-positive mothers and neonates at birth. *Am J Obstet Gynecol*. 2020, 2:100145. [10.1016/j.ajogmf.2020.100145](https://doi.org/10.1016/j.ajogmf.2020.100145)
  62. Facchetti F, Bugatti M, Drera E, et al.: SARS-CoV2 vertical transmission with adverse effects on the newborn revealed through integrated immunohistochemical, electron microscopy and molecular analyses of placenta. *EBioMedicine*. 2020, 59:102951. [10.1016/j.ebiom.2020.102951](https://doi.org/10.1016/j.ebiom.2020.102951)
  63. Chambers C, Krogstad P, Bertrand K, Contreras D, Tobin NH, Bode L, Aldrovandi G: Evaluation for SARS-CoV-2 in breast milk from 18 infected women. *JAMA*. 2020, 324:1347-8. [10.1001/jama.2020.15580](https://doi.org/10.1001/jama.2020.15580)
  64. Wang X, Zhou Z, Zhang J, Zhu F, Tang Y, Shen X: A case of 2019 novel coronavirus in a pregnant woman with preterm delivery. *Clin Infect Dis*. 2020, 71:844-6. [10.1093/cid/ciaa200](https://doi.org/10.1093/cid/ciaa200)
  65. Robertson CA, Lowther SA, Birch T, et al.: SARS and pregnancy: a case report. *Emerg Infect Dis*. 2004, 10:345-8. [10.3201/eid1002.030736](https://doi.org/10.3201/eid1002.030736)
  66. Breastfeeding and COVID-19. (2022). Accessed: August 22: <https://www.who.int/news-room/commentaries/detail/breastfeeding-and-covid-19>.
  67. Liu H, Wang LL, Zhao SJ, Kwak-Kim J, Mor G, Liao AH: Why are pregnant women susceptible to COVID-19? An immunological viewpoint. *J Reprod Immunol*. 2020, 159:103122. [10.1016/j.jri.2020.103122](https://doi.org/10.1016/j.jri.2020.103122)
  68. Sharps MC, Hayes DJL, Lee S, et al.: A structured review of placental morphology and histopathological lesions associated with SARS-CoV-2 infection. *Placenta*. 2020, 101:13-29. [10.1016/j.placenta.2020.08.01](https://doi.org/10.1016/j.placenta.2020.08.01)
  69. Laresgoiti-Servitje E, Cardona-Pérez JA, Hernández-Cruz RG, et al.: COVID-19 infection in pregnancy: PCR cycle thresholds, placental pathology, and perinatal outcomes. *Viruses*. 2021, 13: [10.3390/v13091884](https://doi.org/10.3390/v13091884)
  70. Bloise E, Zhang J, Nakpu J, et al.: Expression of severe acute respiratory syndrome coronavirus 2 cell entry genes, angiotensin-converting enzyme 2 and transmembrane protease serine 2, in the placenta across gestation and at the maternal-fetal interface in pregnancies complicated by preterm birth or preeclampsia.

- Am J Obstet Gynecol. 2021, 224:298.e1-8. [10.1016/j.ajog.2020.08.055](https://doi.org/10.1016/j.ajog.2020.08.055)
71. Komine-Aizawa S, Takada K, Hayakawa S: Placental barrier against COVID-19. *Placenta*. 2020, 99:45-9. [10.1016/j.placenta.2020.07.022](https://doi.org/10.1016/j.placenta.2020.07.022)
  72. Levitan D, London V, McLaren RA, et al.: Histologic and immunohistochemical evaluation of 65 placentas from women With Polymerase Chain Reaction-Proven Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection. *Arch Pathol Lab Med*. 2021, 145:648-56. [10.5858/arpa.2020-0793-SA](https://doi.org/10.5858/arpa.2020-0793-SA)
  73. Smithgall MC, Liu-Jarin X, Hamele-Bena D, Cimic A, Mourad M, Debelenko L, Chen X: Third-trimester placentas of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-positive women: histomorphology, including viral immunohistochemistry and in-situ hybridization. *Histopathology*. 2020, 77:994-9. [10.1111/his.14215](https://doi.org/10.1111/his.14215)
  74. Edlow AG, Li JZ, Collier AY, et al.: Assessment of maternal and neonatal SARS-CoV-2 viral load, transplacental antibody transfer, and placental pathology in pregnancies during the COVID-19 pandemic. *JAMA Netw Open*. 2020, 3:e2030455. [10.1001/jamanetworkopen.2020.30455](https://doi.org/10.1001/jamanetworkopen.2020.30455)