



OPEN ACCESS

Use of the Rehabilitation Treatment Specification System (RTSS) in the management of nitrous oxide (N₂O)-induced spinal cord injury

Charlotte Buttery,^{1,2} Jonathan Birns,³ Jamie Gibson,^{1,4} Gareth David Jones ^{1,5}

¹Physiotherapy Department, Guy's and St Thomas' NHS Foundation Trust, London, UK

²Physiotherapy Department, Maidstone and Tunbridge Wells NHS Trust, Maidstone, UK

³Department of Ageing & Health, Guy's and St Thomas' NHS Foundation Trust, London, UK

⁴Workforce Transformation, Health Education England (HEE), Leeds, UK

⁵Centre for Human and Applied Physiological Sciences (CHAPS), King's College London, London, UK

Correspondence to

Dr Gareth David Jones; gareth.jones@gstt.nhs.uk

Accepted 5 January 2023

SUMMARY

Nitrous oxide (N₂O) is an inhaled anaesthetic gas and a popular intoxicant. Excessive recreational use can cause spinal cord myelopathy. Previous studies have discussed the medical management. However, none have specified the sensorimotor rehabilitation management. This case report documents the investigations, physical rehabilitation and functional outcomes in two cases of N₂O-associated myelopathy. Both presented with lower limb strength and sensorimotor integration impairments resulting in ataxic ambulation. Dorsal column signal abnormality was observed on T2-weighted MRI in one case. Myelopathy was diagnosed based on clinical presentation and both were treated with vitamin B₁₂. Rehabilitation was conceived and specified using the Rehabilitation Treatment Specification System (RTSS). Both cases achieved independent indoor gait on hospital discharge, and full function at 9 months in one case. Appropriate and timely medical management and reasoned rehabilitation provided excellent functional outcomes for N₂O-related myelopathy. By using the RTSS, reasoned rehabilitation efficacy can be tested in the future.

BACKGROUND

The analgesic and intoxicating effects of the colourless inorganic gas nitrous oxide (N₂O) have been known for over 200 years.^{1,2} As an inhaled analgesic, N₂O acts as an N-methyl-D-aspartate antagonist which decreases excitatory neurotransmission throughout the central nervous system via non-competitive glutamate inhibition.³ When prescribed as a mixture with 50% oxygen, it can be safely administered by trained professionals (eg, nurses, midwives, dentists, paramedics) during painful or distressing procedures.⁴ As an inhaled intoxicant without oxygen, N₂O (colloquially sweet-air, nos, noz, nox, hippie-crack⁵) elicits short-lived feelings of euphoria with pleasurable psychedelic and empathogenic effects.⁶

Recreational N₂O intoxication is popular. The UK lifetime-user prevalence rate from a non-probabilistic sample is 38.6%; typical users appear to be male clubbers in their 20s.⁷ The catering industry's development of small 8 g steel recharging canisters for whipped cream dispensers containing pressurised N₂O (bulbs, whippits, nangs) has made them ubiquitous to buy cheaply (50p–£2 per canister⁸) usually in packs of 100. Use from large cylinders (2–10 kg) has become more prevalent recently.⁹ N₂O remains legal to buy in the UK

although this is now under review¹⁰ not least due to recent media concern about the increase in chronic abuse,¹¹ and the alarming increase in use reported in the Netherlands, which has led it to plan for formal controls in 2023.¹² Consuming N₂O directly from the canister is dangerous—the pressure (~200 kPa) can cause barotrauma,¹³ and the temperature of the gas near the outlet (–55°C¹⁴) can cause frostbite injuries.¹⁵ Instead, the gas is typically transferred into balloons via either the whipped cream dispenser or a secondary device (creamers or crackers) easily available that can dispense the canister gas into balloons inserted over the device spout. The preferred recreational use is to inhale or rebreathe the N₂O via the balloon at parties and festivals.¹⁶

Although recreational N₂O use is relatively safe compared with other drugs, acute harm includes falls, accidental injury, confusion⁷ or vomit aspiration.¹⁷ Acute death is rare. A 2016 systematic review of case reports found evidence of 29 case deaths mostly due to acute asphyxiation due to hypoxia with users consuming N₂O with a mask or plastic bag.¹⁸ In the UK, 17 deaths were reported between 2006 and 2012¹⁹ mainly due to accidental asphyxia when N₂O displaces oxygen if used in enclosed spaces.²⁰ Most users inhale low doses per session (1–2 canisters), but overuse exists (~100 canisters), which can lead to persistent neurological sequelae. Chronic clinical toxicity is due to N₂O's interaction with vitamin B₁₂ converting it from an active to an inactive form by irreversible oxidation.²¹ By impairing methylation reactions and DNA synthesis, oxidation results in an accumulation of homocysteine and impaired maintenance of the myelin sheath.²² Thus, B₁₂ deficiency from excessive N₂O use can cause demyelination of the central and peripheral nervous system and a diagnosis of subacute combined degeneration (SACD). If demyelination presents in the spinal cord, the dorsal columns are classically affected, with sometimes the lateral but rarely the anterior columns. The associated diagnostic procedures, neural impairments (progressive vibratory and proprioceptive sensory abnormalities, ascending paraesthesia, ataxic gait, hyporeflexia or hyper-reflexia, and rarely muscle weakness and loss of sphincter control^{23–25}), and medical treatment options have been described in medical case reports and observational studies of SACD of the spinal cord previously.^{22,25–36}

In contrast, *rehabilitative* treatment options for N₂O abuse SACD are not as pervasive in the



© BMJ Publishing Group Limited 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Buttery C, Birns J, Gibson J, et al. *BMJ Case Rep* 2023;**16**:e252529. doi:10.1136/bcr-2022-252529

literature. It is as if patients are expected to recover naturally with cursory reassurance and good luck. One published case study advocates physicians to be cognisant that tailored rehabilitation for a patient’s functional and neuropsychiatric impairments can limit the extent of neurological damage,³⁷ and another clinical review advises medical clinicians to refer to physiotherapy if a patient’s mobility is affected.³⁸ Yet, rehabilitation treatment is not well specified even in these welcome cases.

This lack of rehabilitation specificity is not unusual in research into rehabilitation treatment efficacy where objective definitions of treatments of interest, or even ‘usual care’,³⁹ are required. For surgical or pharmacological treatments, it is usual for active treatment ingredients to be obvious and be specified as chemical structures or anatomical results. In contrast, rehabilitation specification is challenging for several reasons. A binary definition of rehabilitation is tenuous because there are non-binary categories of rehabilitation,⁴⁰ rehabilitation ingredients are not obvious, rehabilitation treatments attempt to change multiple interacting patient functions (eg, muscle strength, movement fluidity, participation in occupation) and specific treatment ingredient definitions do not exist, let alone their downstream effect on patient function.⁴¹ To date, the active ingredients of treatment at the macro level of acute inpatient rehabilitation are rarely specified, which equates with the ‘black box’ approach where little insight into the active treatment ingredients is given which is a typical criticism in the rehabilitation science and clinical communities.⁴² The recently developed coherent framework (the Rehabilitation Treatment Specification System (RTSS)) addresses the black box problem by developing methods that specify, and ultimately measure, the rehabilitative content (ingredients), that via theoretical or observed processes (mechanisms of action), cause a change in desired systems (targets).⁴³ Because of relative rarity of both the clinical condition and the specification of its rehabilitation using the RTSS, the following report of two cases with spinal cord injury post-N₂O overuse is intended to provide an exemplar of holistic medical and rehabilitation management. It was prepared following the Case Report guidelines.⁴⁴

CASE PRESENTATION

The patients were both male. Case 1 was in his early teens (mass 71.7 kg, height 183.0 cm, body mass index (BMI) 21.4 kg/m²) and case 2 was in his 20s (54.1 kg, 173.5 cm, BMI 18.0 kg/m²) on presentation to our large, urban hospital emergency department. Case 1 presented with a 2-day history of gradually worsening lower leg weakness, sensory disturbance in a stocking distribution, reduced balance and mobility. Case 2 presented with a 3-month history of altered leg sensation/weakness commencing with bilateral distal sensory disturbance progressing to pins and needles in a stocking distribution with bilateral leg weakness and reduced mobility.

Neither disclosed any medical history, nor recalled using other recreational drugs except for case 1 who chose to smoke cannabis occasionally. Both confirmed consuming five to six units of alcohol socially per week. History of N₂O consumption was similar for both cases and varied from 50 to 100 canisters per month (usually with large quantities in a single sitting). However, both cases describe an acute escalation in consumption to 100–200 canisters in the week prior to hospital admission. Case 1 was in full-time education, and case 2 worked part time as a football coach. Both cases lived in rented flats; case 1 with his mother, and case 2 with his parents and siblings.

Both patients presented with normal cognition, orientation and fluent speech. The cranial nerves and upper limb examination were normal. There were no cerebellar signs. The sensorimotor lower limb impairments (table 1) included distal lower limb weakness measured using the Medical Research Council ordinal rating scale⁴⁵ and translated into diminished ambulatory function with both cases requiring aids to transfer or walk. Neither case presented with any urinary or bowel dysfunction.

INVESTIGATIONS

Methylmalonic acid (MMA) serum levels were elevated above the normal threshold level of 243 nmol/L,⁴⁶ more so in case 1 (table 1) indicative of vitamin B₁₂ deficiency. Serum folate levels in both cases were unremarkable (3.8 µg/L and 4.9 µg/L for cases 1 and 2, respectively; both within the normal range of 3.1–20.5 µg/L). Other investigations are summarised in table 1.

DIFFERENTIAL DIAGNOSIS

Thorough history taking identified a gradual onset of symptoms over several days, ruling out a vascular cause of neurological dysfunction such as spinal or cerebral stroke. Further neurological assessment and clinical investigations (table 1), along with a history of excessive N₂O intake, led to a diagnosis of SACD of the spinal cord causing sensorimotor myelopathy.

TREATMENT

Case 1

Intramuscular hydroxocobalamin injections (1 mg) were provided daily for 1 week and then on alternate days for a further week. The patient was advised to cease all recreational drug use including N₂O inhalation. Physiotherapy treatment (total with therapist 445 min) was provided in 11 sessions over a 12-day inpatient stay (mean 40.5 min/session). Clinical assessment revealed bilateral lower limb sensorimotor weakness, and a total score of 7 out of 56 on the Berg Balance Scale (BBS—a 14-item ordinal measure that assesses static balance, functional mobility and falls risk⁴⁷) indicated difficulties in functional mobility and balance particularly a reliance on visual sensory cues, and poor

Table 1 Clinical features and investigation findings at initial assessment

Case	Sensation			Motor		Chemistry	Diagnostics		
	LT	Proprioception	Vibration	Strength*	DTR	MMA (nmol/L)	MRI	Neurophysiology	Ambulatory function
1	Impaired/absent LT L1–S2	Symmetrically absent at 1st metatarsal and ankle, intact at knee	Absent below T4	5/5 hip and knee 2/5 ankle	Absent ankle Reduced knee	16930	Dorsal column signal abnormality: T3–T10 levels confirmed subacute combined degeneration	Sensorimotor neuropathy with mild conduction slowing and axonal loss	Mobile 10m with frame and assistance ×1
2	Impaired/absent LT L1–S2	Symmetrically absent at 1st metatarsal, ankle and knee	Absent below T10	4/5 hip and knee 2/5 ankle	Absent ankle Reduced knee	2629	No intrinsic or compressive cord, cauda equina or neural lesion demonstrated	Not completed	Step round transfer with assistance ×2

All clinical tests shown relate to the lower limbs.
 *Strength testing as per Medical Research Council ordinal rating.⁴⁵
 †Ambulatory function—assistance ×1 relates to one person (×2 relates to two people) being at touch distance necessary to prevent loss of balance.
 DTR, deep tendon reflex; LT, light touch; MMA, methylmalonic acid.

planning of whole body transitional movement and reactions to centre-of-mass excursions. Seeing as the mean (\pm SD) BBS for a sample of ambulatory spinal cord injured-paraplegic participants (American Spinal Injury Association (ASIA) Impairment Scale D (AIS-D))⁴⁸ is 44.8 (\pm 13.0),⁴⁹ case 1's functional mobility and balance deficits at initial assessment were profound, a finding supported by their slow observed average (10m) self-selected gait velocity (0.29 m/s) indicative of a home-only ambulator.⁵⁰ Overall activities of daily living function was moderate (modified Barthel Index⁵¹ 12 out of 20). The aims of physical rehabilitation treatment were safe community ambulation with or without walking aids. Treatment specification was undertaken based on the RTSS⁴³ (table 2) and includes interventions designed to affect volitional rehabilitation behaviour using a behaviour change model that considers three essential conditions: capability, opportunity and motivation (COM-B system).⁵²

Case 2

Intramuscular hydroxocobalamin injections (1 mg) for 1 week were provided which was reduced to alternating days for a further week. He was also prescribed oral paracetamol 1 g four times a day, codeine phosphate 30 mg four times a day and gabapentin 300 mg three times a day for symptom control. Similarly to case 1, the patient was also advised to cease recreational N₂O inhalation. Physiotherapy treatment (370 min) was provided in 12 sessions over a 14-day inpatient stay (30.9 min/session). Assessment revealed similar impairments to case 1: bilateral lower limb sensorimotor weakness (table 1). A combined time of 0/120 s on the modified Clinical Test of Sensory Interaction on Balance (CTSIB-M—a performance measure quantifying postural standing control under four sensory conditions each tolerated for up to 30 s for a maximum score of 120 s^{53 54}) indicated their sensorimotor balance control was profoundly impaired. In fact, case 2's impairments were so profound compared with case 1; they were unable to ambulate at all and required maximal assistance to transfer safely. The aims of treatment were nonetheless the same as in case 1 and remained safe community ambulation with or without walking aids. Treatment specification was undertaken based on the RTSS (table 2).

While folic acid treatment was not indicated to correct abnormal serum folate levels in either of our cases, it should be noted that correcting any diagnosed folate insufficiency should be undertaken only after initial phase B₁₂ replacement to prevent worsening of the neurological spinal cord degeneration.⁵⁵

OUTCOME AND FOLLOW-UP

Cases 1 and 2 improved with treatment and their MMA concentration levels were within the normal range prior to hospital discharge. There were however residual sensorimotor deficits. In case 1, plantarflexor strength was graded at 5/5 on reassessment prior to hospital discharge, but dorsiflexion strength had only recovered to 3/5 bilaterally. Proprioception was no longer impaired at the ankle, but distal impairments remained bilaterally. In addition, there were residual sensory impairments to light touch on the left in an L1–S1 and on the right an L5 dermatomal distribution. He was independently mobile with foot-up splints (due to dorsiflexor fatigue over longer distances) and two elbow crutches up to 45 m but could not ambulate outdoors. Average gait speed increased from 0.29 to 0.41 m/s, an increase of 0.12 m/s, which is a clinically meaningful difference⁵⁶ but still less than the comfortable average gait velocity threshold of 0.49 m/s that delineates community from household-only ambulation.⁵⁰ His sensorimotor recovery translated to a meaningful

improvement in his BBS (47 of 56), and a Barthel score of 17 out of 20 on discharge. The patient was duly referred to local community physiotherapy services available at their discharge residence for treatment as specified.

On discharge, case 2 had residual impaired sensation to light touch bilaterally in an L4–S1 dermatomal distribution. Proprioception at the first metatarsals remained absent bilaterally. He was independently mobile indoors with two elbow crutches up to 60 m and was able to control the knee avoiding recurvatum during stance phases in \geq 90% of gait cycles. Indoor independent gait velocity at discharge was 0.38 m/s. However, he could not ambulate outdoors independently similarly to case 1. His recovery translated to an improved CTSIB-M (71/120 s) revealing residual reliance on visual sensory information, and a Barthel score of 19 out of 20. He too was duly referred for outpatient therapy for treatment as specified; no community therapy options were available at his discharge residence.

Case 1 chose to not participate in further therapy or medical follow-up. In contrast, case 2 attended further physiotherapy and was reviewed in an outpatient medical clinic 3 weeks post-hospital discharge. He was ambulating independently but self-reported mildly impaired balance. While proprioception was intact on clinical examination, he presented with symmetrical, lower limb impaired light touch sensation in a stocking distribution, and could only tolerate the Romberg position (eyes closed)⁵⁷ for 22 s without excessive sway or losing balance.

At a subsequent outpatient medical clinic review at 9 months post-hospital discharge, case 2 reported a full recovery including being able to sprint on a treadmill in his physiotherapy sessions. Repeat neurophysiology tests showed improved conduction velocities and normal sensory studies; however, compound muscle action potentials in the lower limbs remained below the normal range.

DISCUSSION

We observed profound sensorimotor impairment caused by excessive N₂O abuse which led to SACD of the spinal cord in two cases. The cases presented with typical features that collectively confirmed the diagnosis. First, a clinical myelopathy anatomically localised to the spinal cord (confirmed by MRI studies in case 1); second, evidence of vitamin B₁₂ deficiency; and lastly, an absence of identifiable central or peripheral nervous system pathology in keeping with the clinical findings.²³ The lack of confirmatory imaging studies in case 2 is not unusual—a published series of 54 patients with dorsal column injury reported low sensitivity with imaging confirming the diagnosis only in 8 patients (14.8%).⁵⁸ Previous literature has reported excessive N₂O use and vitamin B₁₂ deficiency cases whose main presenting symptoms were weakness, paraesthesia and ataxic gait,^{22 28 32} falls,^{31 33} absent or impaired deep tendon reflexes and diminished vibration sense,^{26 29} hyporeflexia,³⁴ limb proprioceptive abnormalities^{27 30} and wide-based steppage gait,³⁵ not dissimilar to the present cases.

While resolution of symptoms following vitamin B₁₂ therapy is sometimes incomplete in older patients (mean (\pm SD) age 55.9 (\pm 15.5) years), it tends to be complete in younger patients (39.8 (\pm 18.8) years).²³ So it was not surprising that the two young cases presented in this paper progressed in their recovery and made a near full sensorimotor recovery at 9 months, at least in the second case. Because their sensorimotor function returned to near normal, we can presume that the two cases did not re-abuse N₂O while we followed them up. If they had re-abused N₂O inhalation during the follow-up period, then B₁₂ would have

Table 2 Treatment specification based on the RTSS for cases 1 and 2

Description of clinical interaction	Case	Target			Ingredients		
		What/in what way	Group	Volition type	MOA	Ingredient	Dosing parameters
The PT facilitates stretching, and voluntary performance of dorsiflexion and plantarflexion exercises for strengthening and maintenance of musculoskeletal integrity. The PT teaches patient how to practise for outside of therapy sessions.	1 & 2	Ankle dorsiflexion and plantarflexion strength/increase	O	DV	Muscle fibre hypertrophy	▶ Verbal description and demonstration of open-kinematic chain dorsiflexion	▶ N/A
						▶ Active right, left and bilateral open-chain dorsiflexion with PT	▶ 1 × max 10 reps each (or until perceives fatigue), in long sitting or (progression) in sitting
						▶ Verbal description and demonstration of bilateral closed-chain dorsiflexion	▶ N/A
						▶ Standing, back to wall ≥10 cm away from heels, feet at shoulder width—controlled leaning back to wall and return to stand with PT	▶ 3 × 15 reps (or until perceives fatigue), increase distance if no fatigue
						▶ Verbal description and demonstration of bilateral calf raises	▶ N/A
						▶ Active bilateral calf raises with therapist, arm support for balance	▶ 2 × 10 reps (or until perceives fatigue) in standing
						▶ Unsupervised seated open-chain active right, left and bilateral dorsiflexion/plantarflexion	▶ 2 × max 10 reps (daily in long sitting or (progression) in sitting)
						▶ Opportunity to practise technique under supervision of therapist	▶ Until correct technique achieved
						▶ Positive reinforcement feedback from therapist	▶ On correct performance of exercise
						▶ Provision of exercise sheet with images and dosing	▶ N/A

Continued

Table 2 Continued

Description of clinical interaction	Case	Target			MOA	Ingredients	
		What/in what way	Group	Volition type		Ingredient	Dosing parameters
The PT provides the patient with a resting ankle splint to wear in bed and teaches self-directed plantarflexor stretches with towel	1 & 2	Passive ankle range of movement/maintain 90°	0	DV	Maintained sarcomere length	▶ Apply ankle-foot resting splint	▶ 2× daily
						▶ When supine in bed	▶ 1 hour (progress to max 4 hours in increments of 1 hour)
						▶ Passive stretch to calves with towel	▶ 2× daily per foot for 2 min
						▶ Long sitting, manipulating towel behind ball of one foot and toes, holding both ends of towel and applying calf stretch with knee straight	▶ Progression when 3/5 MRC strength—standing (hands on support), heels lowered over a step, knees straight
Self-application of ankle-foot resting splint/achieve independence	S			DV	Learning by doing	▶ Ankle-foot resting splint	▶ N/A
						▶ Provide opportunity to practise donning/doffing	▶ Until independent
Self-application of passive ankle stretching technique/achieve independence						▶ Verbal reinforcing feedback	▶ On each trial
						▶ Access to ~1 m length rolled-up towel	▶ N/A
						▶ Demonstration of long sitting, placing towel behind foot and toes, and applying calf stretch with knee straight	▶ N/A
						▶ Provide opportunity to practise	▶ Until independent
						▶ Physical facilitation and verbal corrective feedback	▶ As required
Independent ankle-foot resting splint programme/perform as directed	R			V	Cognitive and affective information processing	▶ Verbal education about risk of donning/doffing ankle resting splint (check for skin integrity), and not donning (losing range at ankle)	▶ N/A
						▶ Written instruction on dosing and technique	▶ N/A
						▶ Logging record for nursing care staff	▶ Night-shift handover: duration/frequency, skin integrity
Ankle stretching programme/perform as directed	R			V	Cognitive and affective information processing	▶ Verbal explanation of benefits of stretching programme	▶ N/A
						▶ Verbal and written information on technique and dosing	▶ N/A

Continued

Table 2 Continued

Description of clinical interaction	Case	Target		MOA	Ingredients								
		What/in what way	Group		Volition type	Ingredient	Dosing parameters						
The PT rehabilitates the patient's dynamic balance deficits using a bimanual reaching task and changing the context within sessions progressively.	1	Dynamic balance in standing/improve	S	DV	Learning by doing	▶ Football to hold bimanually	▶ Size 5 football						
						▶ Parallel bars & chair to sit on for safety	▶ N/A						
						▶ Therapist demonstration of technique	▶ N/A						
						▶ Patient passes football to therapist up/down, forward/back, left/right directions within reach boundary in reaction to therapist instruction in standing with eyes closed	▶ Maintain ~80% success rate without patient touching parallel bars, opening eyes, or therapist touch to prevent loss of balance						
						▶ Variation of exercises:	▶ Progression of variations:						
						▶ Change foot/ground interface	▶ Progress to foam balance mat if 80% success rate achieved						
						▶ Change ball	▶ Increase mass of ball if 80% success rate achieved						
						▶ Increase/vary passing tempo	▶ Increase tempo if 80% success rate achieved						
						▶ Increase/vary passing distance	▶ Increase passing distance if 80% success rate achieved						
						▶ Change body orientation	▶ Receive ball, turn 180° with eyes open before passing ball						
						▶ Change feet configuration	▶ Progress to tandem stand, single leg if 80% success rate						
						▶ Change height level when receiving ball	▶ Transfer to step-up for high ball, or squat for low ball						
						The PT rehabilitates the patient's standing balance and proximal joint-lower limb strength deficits by facilitating an exercise regime with the patient changing the context within sessions progressively.	2	Knee extension strength/increase	O	DV	Muscle hypertrophy	▶ Leg squats	▶ 3x8 reps (or until fatigue) bilateral, progress—decrease squat height or single-leg squat
												▶ Therapist demonstration of technique	▶ N/A
												▶ Standing with hands on bar, plinth situated behind for safety	▶ N/A
▶ Bridges in supine	▶ 3x10 reps (or until fatigue), progress to single-leg bridge												
▶ Therapist demonstration of technique	▶ N/A												
▶ Lying on plinth bent knee drop-outs	▶ 3x10 reps (or until perceives fatigue)												
▶ Therapist demonstration of technique	▶ N/A												
Balance in stance function/improve	O	DV	Learning by doing	▶ Single leg stands	▶ 3x10 reps (or until needs support for balance)								
				▶ Bilateral standing on foam cushion	▶ 3x10 reps (or until needs support for balance)								
				▶ Stand-by support from PT	▶ N/A								
Hip extension strength/increase	O	DV	Muscle hypertrophy	▶ High plinth in front, chair behind for safety	▶ N/A								

Continued

Table 2 Continued

Description of clinical interaction	Case	Target			Ingredients		
		What/In what way	Group	Volition type	MOA	Ingredient	Dosing parameters
The PT teaches the patient to self-monitor exertional fatigue.	1&2	Ability to monitor fatigue/improve perform as directed	S	DV	Learning by doing	► Opportunity to practise self-rating exertional fatigue using the NRS-FRS ²	► Until patient reports confidence in using scale
						► Verbal information on importance of managing fatigue	► N/A
						► Education on effect of fatigue on function and quality of life	► N/A
The PT provides the patient with bilateral Boxia splints to compensate for foot drop and to increase gait velocity.	1&2	Replacement of antigravity dorsiflexion function of ankle dorsiflexors	R	V	Habituation to passive support of ankle	► Provision of bilateral Boxia foot-up splints	► Use in ambulation until $\geq 3/5$ MRC dorsiflexion strength
						► Provide opportunity to practise donning and doffing splints	► Until independent
						► Physical facilitation	► As required
The PT rehabilitates gait and transfer tasks and provides feedback to provide gait re-education.	1&2	Independence with ambulation on level surface/increase	S	DV	Learning by doing	► Verbal corrective feedback	► As required
						► Provide opportunities to practise ambulatory function within safe exertional fatigue level and balance function independently	► Maximise walking distance or tempo or contexts up to ≥ 2 exertional fatigue levels above resting state
						► Use walking aid(s)	► Progression—frame to elbow crutches to stick to no aid
						► Provide opportunity to practise whole-body transitions with PT	► With therapist, daily, ≥ 15 min
						► Provide verbal evaluative feedback	► Intermittent, delayed knowledge of performance at rest intervals
						► Change direction, obstacles	► Turns, cuts, step-overs, introduce uneven ground
						► Include gait initiation and cessation	► Instruct
						► Include reaching activities	► As purpose to ambulation
						► Include head movements	► Include, vary tempo, specify eye contact to targets, introduce walking and reading/talking/carrying objects
						► Provide opportunity to practise bed, chair and toilet transfers	► Until independent
						The PT rehabilitates transfer methods for functional independence.	1&2
► Verbal performance feedback	► On each attempt						
► Provide opportunity to self-correct knee recurvatum in gait	► N/A						
The PT coaches gait re-education due to poor proprioceptive control and episodes of knee recurvatum.	2	Knee recurvatum during gait stance phase/reduce	S	DV	Learning by doing	► Verbal coaching	► If knee recurvatum observed $> 1 \times$ every 3 gait cycles
						► Provide opportunity to practise stepping with stable knee	► 3×8 reps each leg, then repeat continuous gait practise
						► In parallel bars for bilateral upper limb support	► N/A
						► PT provides manual support at femoral segment and hip	► 3×8 reps each leg, then repeat continuous gait practise

DV, direct target for volitional; FRS, Face Rating Scale; MOA, mechanism of action; MRC, Medical Research Council; N/A, not applicable; NRS, Numerical Rating Scale; O, organ functions; PT, physiotherapist; R, representation; RTSS, Rehabilitation Treatment Specification System; S, skills and habits; V, volitional.

been deactivated very rapidly with a commensurate deterioration in neurology and we would have expected to have observed a relapse in balance and ambulation dysfunction.⁵⁹ Re-abuse could be a sign of addiction to N₂O although N₂O inhalation addiction is contentious within the literature.^{18 60 61} While clinical discourse to persuade cessation of use appears to have been successful in these two cases, clinicians should be aware that patients presenting with acute recreational N₂O toxicity should be provided with necessary psychosocial support if difficulties in self-selected cessation of abuse are suspected.³⁸

The positive outcomes overall in the two cases presented here are almost certainly due to the medical diagnostic procedures and subsequent clinical decision to commence vitamin B₁₂ treatment in a timely way with regular monitoring. What is less clear is whether the equally timely and novel specification of rehabilitation interventions using the RTSS led to a more reasoned and tailored physical therapy approach. To our knowledge, this is the first specified rehabilitation report using the RTSS in cases of this type. By acting as a coherent framework for specifying rehabilitation interventions based on the clinician's treatment theory, the RTSS fractionates multidimensional and progressive rehabilitation interventions into entities (treatment components) that are amenable to clinical adoption (including among clinical colleagues) and empirical research.⁴¹ Thus, the specification described herein acts first as a reasoned approach for fellow rehabilitation clinicians to use with these patients downstream of the acute hospital; second, as an example for other clinicians treating similar cases; and lastly, as a record of the specified treatment to inform researchers. It is worth noting though that the treatment specifications provided are examples peculiar to the two cases presented. They are not meant as a prescriptive regime for all patients presenting with sensorimotor system impairments due to recreational N₂O toxicity and functional B₁₂ deficiency. Nonetheless, we contend that in successfully specifying the treatment in this way, the patient's treatment was reasoned and tailored and thus verifies the RTSS's utility.

The impairments (bilateral lower limb sensorimotor) and aims of treatment (restitution of normal ambulatory function) were specified identically between cases. Treatments were also identical based on organ function targets (muscle strengthening and stretching), and skills and habits targets (function balance and ambulation practise tasks). Representation targets were identical too. They were designed to influence cognitive processes when managing the musculoskeletal integrity and injurious falls risks inherently associated with distal lower limb paralysis, and ensure the consequences of failing to voluntarily manage those risks were understood.

However, there were treatment specification differences between cases because case 2 initially presented with impairments so profound that no ambulatory function was possible. Case 2 presented with proximal lower limb joint symmetrical weakness and more widespread proprioceptive loss. Therefore, therapy time was prioritised towards specifying a more widespread and explicit strengthening programme and more rudimentary static balance tasks compared with case 1.

What's more, when case 2 progressed to ambulating, knee recurvatum (KR) was observed during some, but not all, stance phases. KR is a relatively common impairment observed in other neurological pathologies (eg, 30%–50% of ambulatory hemiplegic stroke survivors^{62 63}). In normal continuous gait, coordination between pretibial muscle contraction (which restrains ankle plantarflexion and allows the tibia to move forward over the foot during weight acceptance) and triceps surae/quadriceps contraction (crucial to restraining tibial forward motion during

mid-stance while the femoral segment progresses forward distal to it with extension at the knee) controls knee flexion and forward tibial movement over the stationary foot in stance phases.⁶⁴

While the main cause for gait KR is premature overactivity or spasticity of the plantarflexors which prevents the knee from flexing during loading response in early stance^{63 65} or vasti hypertonia,^{66 67} we reasoned that KR in case 2 was due to dorsiflexion/plantarflexion muscle weakness, poor proprioception and reliance on forward placed walking aids. This is because no muscle hyper-reflexia was observed on assessment at the ankle or knee. The variability of KR observed in case 2 was probably due to instances when knee hyperextension was simply deployed to stabilise the knee in stance by adopting a quadriceps-avoidance strategy sometimes seen in anterior cruciate ligament deficient gait⁶⁸ in combination with plantarflexion weakness during mid-stance.⁶⁹ Our theory therefore was that treatment ingredients targeting muscle weakness and sensory integration (including proprioception) in static balance would ameliorate the KR in case 2. Passive treatment ingredients designed to target KR more directly to support the approach would include issuing of rigid ankle/foot orthotics which were avoided here. Instead, gait re-education was deployed via instructions based on implicit motor learning to avoid KR with delayed knowledge-of-performance feedback.⁷⁰

Specified treatments were not then all identical between cases and support the argument that treatment was tailored for the two cases. This is one of the advantages of the RTSS. Its coherent framework is intended to administer the complexity of rehabilitation practice in the interactions of clinical impairments, dysfunctions and participation in society in order to specify individuals' rehabilitation treatment.⁴³ Yet, the two cases' rehabilitation treatment aims were similar and were focused on ambulation function. Acute adult inpatient physical rehabilitation often includes stressing motor systems toward safe, independent ambulation because ambulatory function is a key milestone for discharge planning. This is important within the goal of reducing acute hospital length of stay (LOS) which is itself dependent on coherent, equitable and resourced rehabilitation services downstream of an acute setting. But the problem with yoking rehabilitation with minimising LOS is that the rehabilitation aim is often achievement of minimal *adaptive* ambulatory function necessary for a safe discharge. The aim instead should be the *restitution*⁷¹ of ambulatory function, especially since stroke⁷² and spinal cord-injured⁷³ neurorehabilitation patients express it as their primary goal. Restitution of ambulatory performance therefore requires rehabilitative progression towards pre-morbid performance, not merely an assessment of current ambulatory performance.

The RTSS includes accurate specification of rehabilitation progression and was included throughout the two cases described. For ambulatory function, a separate treatment ingredient was included so the two cases were proficient in self-monitoring their exertional fatigue—a common symptom burden for patients with acute spinal cord injury.⁷⁴ That allowed the ambulatory treatment's progression to be specified in both cases by attributing the therapeutic dose of ambulatory practise (distance or time walked) with the patient's increase in self-rated exertional fatigue (using a dual Numerical and Face Rating Scale⁷⁵) during ambulatory treatment ingredients.

Similar separate treatment ingredients were included in our specification designed for the cases to attain skills and habits in managing the risk to their musculoskeletal integrity by applying ankle resting splints and foot-up devices when ambulating. We contend that without the RTSS's coherent framework, there

would be a risk that therapists might not have clinically reasoned tailored treatments as thoroughly, nor been as accurate in specifying it.

In conclusion, the combination of timely medical treatment and specific and clinically reasoned rehabilitation in a structured format using the RTSS provided excellent functional outcomes in both the acute and subacute phases post-injury in two individuals who sustained N₂O-induced spinal cord injury. Stronger conclusions can be drawn when future studies determine the impact of rehabilitation on clinical outcomes at the impairment, activity and participation levels in this patient cohort by virtue of using the RTSS.

Patient's perspective

Case 2 and his next of kin were able to provide their perspective of their assessment and treatment.

Before the hospital admission we had no idea about the possible side effects of the inhaled canisters. Lots of the young people where we live took them at parties and we believed that they were risk free, we had never heard of others experiencing medical complications from taking them.

The profound loss of ambulation was terrifying particularly because there were no initial explanation why these symptoms had occurred.

When we initially went to hospital the doctors weren't sure what was the cause of the weak legs and the inability to walk. After the scans the doctors explained that there was a problem with the nerves in the back and they weren't totally sure if my son's walking would fully recover. They thought that the gas canisters were the cause because of some of the blood test results. We were all shocked that this had happened and it was scary to hear that someone so young might have long term effects that they would have to live with. It was hard for us not to know what the future may hold for him, we felt very worried.

However, the medical assessment and treatment and the inpatient physical therapy rehabilitation delivered enabled meaningful recovery – a recovery that was complete after receiving longer term community physiotherapy. We feel lucky that my son has made a full recovery, it was slow to begin with and he had to work hard in physio sessions in the hospital. Unfortunately it took quite a long time for him to get seen in the physio clinic after leaving hospital, but he's now managing to do everything he could do before the injury. We are thankful for all of the care that my son received.

Learning points

- ▶ Rates of nitrous oxide recreational use are rising, meaning that incidents of abuse and associated subacute combined degeneration pathology may also rise leading to profound sensorimotor dysfunction.
- ▶ Timely diagnosis and medical treatment to reverse vitamin B₁₂ deficiency in combination with skilled physical rehabilitation led to favourable outcomes.
- ▶ Specified treatment from rehabilitation clinicians in the acute phase after neurological pathology in these two cases shows that timely, tailored and theoretically reasoned treatments can be put into practice if the goal is to reconstitute normal movement function.

methodology, validation, investigation, resources, writing (review and editing). JG—conceptualisation, methodology, validation, resources, writing (original draft), writing (review and editing), visualisation, investigation, writing (review and editing). GDJ—conceptualisation, validation, resources, data curation, writing (original draft), writing (review and editing), visualisation, project administration.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Gareth David Jones <http://orcid.org/0000-0001-5516-9418>

REFERENCES

- 1 Davy SH. *Researches, chemical and philosophical; chiefly concerning nitrous oxide or dephlogisticated nitrous air, and its respiration*. Bristol: Biggs and Cottle, 1800.
- 2 Priestley J. *Experiments and observations on different kinds of air*. London: J Johnson, 1790.
- 3 Jevtović-Todorović V, Todorović SM, Mennerick S, et al. Nitrous oxide (laughing gas) is an NMDA antagonist, neuroprotectant and neurotoxin. *Nat Med* 1998;4:460–3.
- 4 Annequin D. Nitrous oxide (N₂O) angel or devil? *Paediatr Anaesth* 2020;30:388–9.
- 5 Public Health England. Talk to frank - nitrous oxide london: public health england. 2021. Available: <https://www.talktofrank.com/drug/nitrous-oxide> [Accessed 10 Apr 2021].
- 6 Beckman NJ, Zacny JP, Walker DJ. Within-subject comparison of the subjective and psychomotor effects of a gaseous anesthetic and two volatile anesthetics in healthy volunteers. *Drug Alcohol Depend* 2006;81:89–95.
- 7 Kaar SJ, Ferris J, Waldron J, et al. Up: the rise of nitrous oxide abuse. An international survey of contemporary nitrous oxide use. *J Psychopharmacol* 2016;30:395–401.
- 8 Winstock AR, Ferris JA. Nitrous oxide causes peripheral neuropathy in a dose dependent manner among recreational users. *J Psychopharmacol* 2020;34:229–36.
- 9 van Riel AJHP, Hunault CC, van den Hengel-Koot IS, et al. Alarming increase in poisonings from recreational nitrous oxide use after a change in EU-legislation, inquiries to the Dutch poisons information center. *Int J Drug Policy* 2022;100:S0955-3959(21)00437-0:103519..
- 10 HM Government. Nitrous oxide: home secretary's letter to the ACMD: advisory council on the misuse of drugs. London:; 2021. Available: <https://www.gov.uk/government/publications/nitrous-oxide-home-secretarys-letter-to-the-acmd> [Accessed 12 Oct 2021].
- 11 Davis N. Doctors warn of rise in nerve damage linked to nitrous oxide [newspaper article]. the guardian, london, UK: guardian news & media ltd. 2022. Available: <https://www.theguardian.com/society/2022/aug/30/doctors-warn-of-rise-in-nerve-damage-linked-to-nitrous-oxide> [Accessed 3 Sep 2022].
- 12 Sumnall H. Recreational use of nitrous oxide. *BMJ* 2022;378:2297.
- 13 Sellers WFS. Misuse of anaesthetic gases. *Anaesthesia* 2016;71:1140–3.
- 14 Rowbottom SJ. Nitrous oxide abuse. *Anaesth Intensive Care* 1988;16:241–2.
- 15 Hwang JC, Himel HN, Edlich RF. Frostbite of the face after recreational misuse of nitrous oxide. *Burns* 1996;22:152–3.
- 16 Chair of the Advisory Council on the Misuse of Drugs (ACMD). Letter to teresa may (the home secretary) on nitrous oxide abuse [letter]. london, UK: UK government. 2015. Available: <https://www.gov.uk/government/publications/acmd> [Accessed 3 Mar 2021].
- 17 Randhawa G, Bodenham A. The increasing recreational use of nitrous oxide: history revisited. *Br J Anaesth* 2016;116:321–4.
- 18 Garakani A, Jaffe RJ, Savla D, et al. Neurologic, psychiatric, and other medical manifestations of nitrous oxide abuse: A systematic review of the case literature. *Am J Addict* 2016;25:358–69.
- 19 Wolfson S. *Is the growth in nitrous oxide misuse a laughing matter?* London: The Guardian, 2014. Available: <https://www.theguardian.com/society/2014/aug/13/brick-lane-is-the-uks-laughing-gas-megastore-but-for-how-long>
- 20 Ghodse H, Ahmed K, Corkery J, et al. Trends in UK deaths associated with abuse of volatile substances, 1971-2008. report 23. st. georges, university of london: volatile substance abuse (VSA) mortality project, international centre for drug policy. 2010.

Contributors CB—conceptualisation, methodology, validation, investigation, resources, data curation, writing (review and editing). JB—conceptualisation,

- 21 Banks RGS, Henderson RJ, Pratt JM. Reactions of gases in solution. Part III. Some reactions of nitrous oxide with transition-metal complexes. *J Chem Soc, A* 1968;1968:2886.
- 22 Flippo TS, Holder WD. Neurologic degeneration associated with nitrous oxide anesthesia in patients with vitamin B12 deficiency. *Arch Surg* 1993;128:1391–5.
- 23 Vasconcelos OM, Poehm EH, McCarter RJ, et al. Potential outcome factors in subacute combined degeneration: review of observational studies. *J Gen Intern Med* 2006;21:1063–8.
- 24 Xiang Y, Li L, Ma X, et al. Recreational nitrous oxide abuse: prevalence, neurotoxicity, and treatment. *Neurotox Res* 2021;39:975–85.
- 25 Oussalah A, Julien M, Levy J, et al. Global burden related to nitrous oxide exposure in medical and recreational settings: a systematic review and individual patient data meta-analysis. *J Clin Med* 2019;8:551.
- 26 Buizert A, Sharma R, Koppen H. When the laughing stops: subacute combined spinal cord degeneration caused by laughing gas use. *J Addict Med* 2017;11:235–6.
- 27 Butzkueven H, King JO. Nitrous oxide myelopathy in an abuser of whipped cream bulbs. *J Clin Neurosci* 2000;7:73–5.
- 28 Choi C, Kim T, Park KD, et al. Subacute combined degeneration caused by nitrous oxide intoxication: a report of two cases. *Ann Rehabil Med* 2019;43:530–4.
- 29 Hsu C-K, Chen Y-Q, Lung V-Z, et al. Myelopathy and polyneuropathy caused by nitrous oxide toxicity: a case report. *Am J Emerg Med* 2012;30:1016.
- 30 Jiang J, Shang X. Clinical-radiological dissociation in a patient with nitrous oxide-induced subacute combined degeneration: a case report. *BMC Neurol* 2020;20:99.
- 31 Morris N, Lynch K, Greenberg SA. Severe motor neuropathy or neuronopathy due to nitrous oxide toxicity after correction of vitamin B12 deficiency. *Muscle Nerve* 2015;51:614–6.
- 32 Renard D, Dutray A, Remy A, et al. Subacute combined degeneration of the spinal cord caused by nitrous oxide anaesthesia. *Neurol Sci* 2009;30:75–6.
- 33 Samia AM, Nenow J, Price D. Subacute combined degeneration secondary to nitrous oxide abuse: quantification of use with patient follow-up. *Cureus* 2020;12:10.
- 34 Yuan JL, Wang SK, Jiang T, et al. Nitrous oxide induced subacute combined degeneration with longitudinally extensive myelopathy with inverted V-sign on spinal MRI: a case report and literature review. *BMC Neurol* 2017;17.
- 35 Zhao B, Zhao L, Li Z, et al. Subacute combined degeneration induced by nitrous oxide inhalation: two case reports. *Medicine (Baltimore)* 2020;99:18.
- 36 Temple C, Horowitz BZ. Nitrous oxide abuse induced subacute combined degeneration despite patient initiated B12 supplementation. *Clin Toxicol (Phila)* 2022;60:872–5.
- 37 Chin J, Forzani B, Chowdhury N, et al. Rehabilitation essential in the recovery of multifactorial subacute combined degeneration. *Ann Phys Rehabil Med* 2015;58:S1877-0657(15)00004-4:190–2..
- 38 Wong J, Viyasar T, Layton B, et al. The dangers of recreational inhalation of nitrous oxide. *Br J Hosp Med (Lond)* 2021;82:1–8.
- 39 Negrini S, Arienti C, Pollet J, et al. Clinical replicability of rehabilitation interventions in randomized controlled trials reported in main journals is inadequate. *J Clin Epidemiol* 2019;114:S0895-4356(19)30315-4:108–17..
- 40 Wade DT. Defining rehabilitation: an exploration of why it is attempted, and why it will always fail. *Clin Rehabil* 2021;35:1650–6.
- 41 Van Stan JH, Whyte J, Duffy JR, et al. Rehabilitation treatment specification system: methodology to identify and describe unique targets and ingredients. *Arch Phys Med Rehabil* 2021;102:S0003-9993(20)31020-0:521–31..
- 42 Whyte J, Hart T. It's more than a black box; it's a Russian doll: defining rehabilitation treatments. *Am J Phys Med Rehabil* 2003;32:639–52.
- 43 Hart T, Dijkers MP, Whyte J, et al. A theory-driven system for the specification of rehabilitation treatments. *Arch Phys Med Rehabil* 2019;100:S0003-9993(18)31310-8:172–80..
- 44 Riley DS, Barber MS, Kienle GS, et al. Care guidelines for case reports: explanation and elaboration document. *J Clin Epidemiol* 2017;89:S0895-4356(17)30037-9:218–35..
- 45 Medical research Council (MRC). Aids to the examination of the peripheral nervous system (memorandum no.45; superseding war memorandum no.7) london: her majesty's stationery office. 1976. Available: <https://mrc.ukri.org/research/facilities-and-resources-for-researchers/mrc-scales/mrc-muscle-scale> [Accessed 28 Mar 2021].
- 46 Nardin RA, Amick ANH, Raynor EM. Vitamin B (12) and methylmalonic acid levels in patients presenting with polyneuropathy. *Muscle Nerve* 2007;36:532–5.
- 47 Berg K, Wood-Dauphinee S, Williams JI. The balance scale: reliability assessment with elderly residents and patients with an acute stroke. *Scand J Rehabil Med* 1995;27:27–36.
- 48 Kirsblum SC, Burns SP, Biering-Sorensen F, et al. International standards for neurological classification of spinal cord injury (revised 2011). *J Spinal Cord Med* 2011;34:535–46.
- 49 Lemay JF, Nadeau S. Standing balance assessment in Asia D paraplegic and tetraplegic participants: concurrent validity of the Berg balance scale. *Spinal Cord* 2010;48:245–50.
- 50 Fulk GD, He Y, Boyne P, et al. Predicting home and community walking activity poststroke. *Stroke* 2017;48:406–11.
- 51 Collin C, Wade DT, Davies S, et al. The barthel ADL index: a reliability study. *Int Disabil Stud* 1988;10:61–3.
- 52 Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
- 53 Shumway-Cook A, Horak FB. Assessing the influence of sensory interaction of balance. suggestion from the field. *Phys Ther* 1986;66:1548–50.
- 54 Cohen H, Blatchly CA, Gombash LL. A study of the clinical test of sensory interaction and balance. *Phys Ther* 1993;73:346–51;
- 55 Hunt A, Harrington D, Robinson S. Vitamin B12 deficiency. *BMJ* 2014;349:g5226.
- 56 Perera S, Mody SH, Woodman RC, et al. Meaningful change and responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc* 2006;54:743–9.
- 57 Lanska DJ, Goetz CG. Romberg's sign: development, adoption, and adaptation in the 19th century. *Neurology* 2000;55:1201–6.
- 58 Jain KK, Malhotra HS, Garg RK, et al. Prevalence of Mr imaging abnormalities in vitamin B12 deficiency patients presenting with clinical features of subacute combined degeneration of the spinal cord. *J Neurol Sci* 2014;342:S0022-510X(14)00310-4:162–6..
- 59 Zheng R, Wang Q, Li M, et al. Reversible neuropsychiatric disturbances caused by nitrous oxide toxicity: clinical, imaging and electrophysiological profiles of 21 patients with 6-12 months follow-up. *Neuropsychiatr Dis Treat* 2020;16:2817–25.
- 60 van Amsterdam J, Nabben T, van den Brink W. Recreational nitrous oxide use: prevalence and risks. *Regul Toxicol Pharmacol* 2015;73:S0273-2300(15)30101-X:790–6..
- 61 Brunt TM, van den Brink W, van Amsterdam J. Mechanisms involved in the neurotoxicity and abuse liability of nitrous oxide: a narrative review. *Int J Mol Sci* 2022;23:14747:23..
- 62 Hogue RE, McCandless S. Genu recurvatum: auditory biofeedback treatment for adult patients with stroke or head injuries. *Arch Phys Med Rehabil* 1983;64:368–70.
- 63 Knutsson E, Richards C. Different types of disturbed motor control in gait of hemiparetic patients. *Brain* 1979;102:405–30.
- 64 Bulea TC, Kobetic R, Audu ML, et al. Stance controlled knee flexion improves stimulation driven walking after spinal cord injury. *J Neuroeng Rehabil* 2013;10:68.
- 65 Higginson JS, Zajac FE, Neptune RR, et al. Muscle contributions to support during gait in an individual with post-stroke hemiparesis. *J Biomech* 2006;39:1769–77.
- 66 Perry J. *Gait analysis: normal and pathological function*. Thorofare, NJ: SLACK Inc, 1992: 223–44.
- 67 Gross R, Delporte L, Arsenault L, et al. Does the rectus femoris nerve block improve knee recurvatum in adult stroke patients? A kinematic and electromyographic study. *Gait Posture* 2014;39:S0966-6362(13)00642-5:761–6..
- 68 Kawahara K, Sekimoto T, Watanabe S, et al. Effect of genu recurvatum on the anterior cruciate ligament-deficient knee during gait. *Knee Surg Sports Traumatol Arthrosc* 2012;20:1479–87.
- 69 Cooper A, Alghamdi GA, Alghamdi MA, et al. The relationship of lower limb muscle strength and knee joint hyperextension during the stance phase of gait in hemiparetic stroke patients. *Physiother Res Int* 2012;17:150–6.
- 70 Kleynen M, Braun SM, Bleijlevens MH, et al. Using a Delphi technique to seek consensus regarding definitions, descriptions and classification of terms related to implicit and explicit forms of motor learning. *PLoS One* 2014;9:e100227.
- 71 Levin MF, Kleim JA, Wolf SL. What do motor " recovery " and " compensation " mean in patients following stroke? *Neurorehabil Neural Repair* 2009;23:313–9.
- 72 Lord SE, McPherson K, McNaughton HK, et al. Community ambulation after stroke: how important and obtainable is it and what measures appear predictive? *Arch Phys Med Rehabil* 2004;85:234–9.
- 73 Hug A, Spingler T, Hensel C, et al. Goal attainment in mobility after acute rehabilitation of mobility-restricting paralysis syndromes with regard to the ambulatory therapeutic level of participation neuromoves: a German national multicenter observational cohort study. *BMC Neurol* 2021;21:149.
- 74 Anton HA, Miller WC, Townson AF, et al. The course of fatigue after acute spinal cord injury. *Spinal Cord* 2017;55:94–7.
- 75 Chuang L-L, Lin K-C, Hsu A-L, et al. Reliability and validity of a vertical numerical rating scale supplemented with a faces rating scale in measuring fatigue after stroke. *Health Qual Life Outcomes* 2015;13:91.

Copyright 2023 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow