

Review

New Insight into Plant Signaling: Extracellular ATP and Uncommon Nucleotides

Małgorzata Pietrowska-Borek ^{1,*}, Jędrzej Dobrogojski ¹, Ewa Sobieszczuk-Nowicka ² and Sławomir Borek ²

¹ Department of Biochemistry and Biotechnology, Faculty of Agronomy and Bioengineering, Poznań University of Life Sciences, Dojazd 11, 60-632 Poznań, Poland; dobrogojski@gmail.com

² Department of Plant Physiology, Faculty of Biology, Adam Mickiewicz University, Poznań, Uniwersytetu Poznańskiego 6, 61-614 Poznań, Poland; evaanna@amu.edu.pl (E.S.-N.); borek@amu.edu.pl (S.B.)

* Correspondence: malgorzata.pietrowska-borek@up.poznan.pl

Received: 6 January 2020; Accepted: 30 January 2020; Published: 2 February 2020



Abstract: New players in plant signaling are described in detail in this review: extracellular ATP (eATP) and uncommon nucleotides such as dinucleoside polyphosphates (Np_nN's), adenosine 5'-phosphoramidate (NH₂-pA), and extracellular NAD⁺ and NADP⁺ (eNAD(P)⁺). Recent molecular, physiological, and biochemical evidence implicating concurrently the signaling role of eATP, Np_nN's, and NH₂-pA in plant biology and the mechanistic events in which they are involved are discussed. Numerous studies have shown that they are often universal signaling messengers, which trigger a signaling cascade in similar reactions and processes among different kingdoms. We also present here, not described elsewhere, a working model of the Np_nN' and NH₂-pA signaling network in a plant cell where these nucleotides trigger induction of the phenylpropanoid and the isochlorogenic acid pathways yielding metabolites protecting the plant against various types of stresses. Through these signals, the plant responds to environmental stimuli by intensifying the production of various compounds, such as anthocyanins, lignin, stilbenes, and salicylic acid. Still, more research needs to be performed to identify signaling networks that involve uncommon nucleotides, followed by omic experiments to define network elements and processes that are controlled by these signals.

Keywords: adenosine 5'-phosphoramidate; adenosine 5'-tetraphosphate; diadenosine 5',5'''-tetraphosphate; dinucleoside polyphosphates; eATP; eNAD(P)⁺

1. Introduction

Plant signaling is a set of phenomena that enables the transduction of external and internal signals into physiological responses such as modification of enzyme activity, cytoskeleton structure, and gene expression. It is known that in plants there exist mechanisms involved in the signal transduction pathways. Plants have evolved signaling networks providing reactions to environmental stimuli through signaling proteins such as plasma membrane receptors and ion transporters and by cascades of kinases and other enzymes up to effectors. For many years plant hormones were considered to be dominant molecules in plant signaling. Nowadays this term embraces many other compounds including second messengers, such as cytosolic Ca²⁺ [1], reactive oxygen (ROS) and nitrogen species (RNS) [2] or cyclic nucleotides such as adenosine 3',5'-cyclic monophosphate (cAMP) and guanosine 3',5'-cyclic monophosphate (cGMP) [3]. Nowadays there is more and more information about synthesis, degradation, and function of cAMP and cGMP in plant [3–5], and they are currently accepted as key signaling molecules in many processes in plants including growth and differentiation, photosynthesis, and biotic and abiotic defense [6]. However, recently it was shown that nucleotides, such as ATP and (di)nucleoside polyphosphates, also can play signaling roles in plant cells. In this review, we focus on

the extracellular ATP (eATP) and uncommon nucleotides, such as mono- (p_nN_s) and dinucleoside polyphosphates ($Np_nN's$), and their new function as signaling molecules.

2. eATP as a Signaling Molecule

Adenosine 5'-triphosphate (ATP), as well as the other nucleoside triphosphates, are established as agents providing energy in various reactions inside cells, both in animal and plant organisms [7]. As ATP is omnipresent in all living cells, it is often called the essential energy currency molecule. In the 1970s it was first hypothesized that ATP might be released into the extracellular environment and act as a signaling compound for animal cells [8]. However, it is worth mentioning that the first effect of eATP on a cell was noted much earlier, i.e., in 1929, during research on heart muscle contraction [9]. In animals, three possible ways of ATP release from the cells into the extracellular matrix have been proposed; they involve multiple channels, transporters, and exocytosis. The interest in the eATP signaling function accelerated after the first purinoreceptor was cloned and characterized in rat brain tissue [10]. Currently, eATP, as well as some of the other nucleotides, are considered as signaling molecules mediating numerous animal cellular processes. For decades the role of nucleotides as signaling molecule functioning similarly in plants as it was demonstrated in animals was viewed with skepticism. A real breakthrough in this topic came with the discovery of the existence of a plant transmembrane receptor protein with serine/threonine kinase activity having a high affinity for extracellular nucleotides [11]. However, the evidence for the mechanism of ATP release from the cytosol into the extracellular matrix in plants appeared earlier [12–15]. In plants, there are several possible ways of ATP outlet. The ATP release triggered by environmental stimuli appears via the wounded cell membrane [16], exocytosis [12], the p-glycoprotein (PGP1) belonging to the ATP-binding cassette ABC transporters [17], and plasma membrane-localized nucleotide transporters (PM-ANT1) [18]. The receptor-eATP interaction begins a cascade reaction that leads to further downstream physiological changes protecting the plant against both biotic and abiotic stresses but also guarantees proper plant growth and development [19].

In order to maintain proper cell growth and function, eATP concentration must be controlled. During regeneration after the stress factor has disappeared, eATP degradation is conducted by hydrolytic enzymes called apyrases [20]. The human apyrases are the best characterized and described apyrases among different kingdoms. Their cellular localization includes plasma membrane, Golgi apparatus, and endoplasmic reticulum. The human apyrases present in plasma membrane show ecto-apyrases activity mediating regulation of the eATP in the extracellular environment. The *Arabidopsis thaliana* apyrase family consists of seven enzymes among which two closely related ones, APY1 and APY2, are the most extensively characterized. These enzymes mediate the luminal glycosylation and can be a component of regulation of the eATP level. It was demonstrated that these two enzymes are an integral component of the Golgi apparatus membrane where they indirectly control the eATP level by modulating the luminal concentration of ATP in secretory vesicles [21–23] (Figure 1). Both APY1 and APY2 are also essential enzymes for proper plant growth and development. These processes are auxin-dependent. Among various factors, auxin transport also depends on the expression of the genes encoding APY1 and APY2. Suppression of the APY1 and APY2 expression causes dwarfism, impaired polar auxin transport and eATP over-accumulation in *Arabidopsis thaliana* [24,25]. However, it was suggested that some fractions of the APY1 and APY2 population with ecto-apyrases activity might be localized also in the plasma membrane [19]. Although there are many data regarding subcellular localization of *Arabidopsis* apyrases (APY1 and APY2), the apyrases from soybean (GS52), pea (PsAPY1), and potato (StAPY3) occur outside the cell (ecto-apyrases) [26–28]. A study conducted by Wu and co-workers indicated that the externally applied APY1 and APY2 inhibitors cause the increase of eATP and physiological changes typical for the plant reaction to stress [24]. Furthermore, there are some reports showing particular plant species secreting individual apyrase members out of the endomembrane system. Taking all these circumstances into account, the existence of a plasma membrane-localized *Arabidopsis thaliana* apyrase (Figure 1) cannot be excluded [29].




Figure 1. Model of changes occurring in the plant cell triggered by the extracellular ATP (eATP). In this model, three possible ways of ATP release into the extracellular matrix are demonstrated. It considers the wounded cell membrane, exocytosis, and two transporters: the p-glycoprotein (PGP1) belonging to the ATP-binding cassette ABC transporters, and the plasma membrane-localized nucleotide transporters (PM-ANT1). Two apyrases, APY1, and APY2, localized in the Golgi apparatus membrane of *Arabidopsis thaliana* regulate the concentration of eATP. Additionally, the hypothesized apyrase (APY) located at the extracellular surface of the plasma membrane can decrease eATP concentration directly in the extracellular matrix. The released eATP acts as a signaling molecule triggering elevation of the cytosolic Ca^{2+} level by activation of the P2K1 receptor, which in turn activates the Ca^{2+} channel. The hypothetical non-P2K1 receptor, whose binding with eATP leads to activation of the $G\alpha$ subunit of the heterotrimeric G-protein, activates the cell membrane Ca^{2+} channel. High cytosolic Ca^{2+} concentration causes an increase in production of nitric oxide (NO), reactive oxygen species (ROS), and mitogen-activated protein kinases (MAPKs), which finally leads to various physiological responses. The ROS boosted production is due to the activation of the RBOHD subunit of the plasma membrane-localized NADPH oxidase. The contribution of the transcription factors in the regulation of gene expression is of high importance.

2.1. Plant eATP Receptors

The eATP animal receptors were discovered in 1976 [30]. Initially, they were called ‘purinergic receptors’. The name was changed to ‘nucleotide receptors’, as both purine and pyrimidine nucleotides trigger their activation [31]. These receptors belong to two groups, P1 and P2, but only P2 receptors are activated by ATP [32]. P2 receptors are divided into two classes: ligand-gated ion channels (P2X) and G protein-coupled (P2Y) receptors [33]. It is important to emphasize that P2X and P2Y receptors do not exist in plant organisms [34]. For many years it was hard to prove the existence of a similar receptor in plants although there were papers indicating changes in plants’ development, growth and response to stresses under exogenously applied ATP [11]. Research on *dorn1* (DOes not Respond to Nucleotides) mutants of *Arabidopsis thaliana* revealed the first plant receptor with a high affinity to bind eATP. Mutants showed a lower cytosolic Ca^{2+} level, lack of mitogen-activated protein kinases (MAPKs) activation and as a consequence decline of the defense-related gene expression level. Initially, the newly discovered receptor was named DORN1, but later the name was changed to P2K1 because the research showed that the *DORN1* gene encodes the L-type lectin receptor-like kinase I.9 (LecRK-I.9). This receptor has three domains: an extracellular ATP-binding lectin domain, a single transmembrane domain, and an intracellular kinase domain (Figure 1). In the case of *dorn1* mutants defective in the kinase domain, eATP is not able to connect with the receptor and as a result is not able to trigger downstream responses. Decreased expression of the *LecRK-I.9* gene caused weaker eATP responses, while intensified *LecRK-I.9* gene expression enhanced eATP responses [35]. DORN1 is a plant purine receptor that belongs to the lectin-receptor kinase family and is denoted as P2K1 to distinguish it from the animal P2 receptors: P2X and P2Y [11,36,37]. Interestingly, *P2K1* gene expression level is relatively high during the major stages of plant development [37], suggesting the essential role of the eATP in processes such as seedling growth, stomata movement, pollen tube development, root hair growth, gravitropism, and biotic and abiotic stress responses [38].

Recently, the existence of another, unidentified, non-P2K1 receptor with affinity to eATP was suggested [39]. It was reported that *Arabidopsis thaliana dorn1* null mutants demonstrated an increased level of cytosolic Ca^{2+} under the exogenously applied ATP. It was a result of the heterotrimeric G-protein G_{α} subunit activation followed by the opening of a plasma-membrane Ca^{2+} channel. This finding suggests new, P2K1-independent responses to eATP, involved among others in the root-bending mechanism [39]. Moreover DORN1 could underpin several calcium-related responses but it may not be the only receptor for eATP in *Arabidopsis thaliana* [40].

2.2. Plant eATP Signal Transduction Pathway

In plants, eATP takes part in cell signaling as a messenger, which triggers a signaling cascade, when binding to the P2K1 receptor (Figure 1). Cytosolic nitric oxide (NO), Ca^{2+} , and ROS form a secondary messenger trio, which occurs in various signaling pathways leading to the transient phosphorylation of MAPK, especially MPK3 and MPK6, and expression of defense-related genes [41]. The multibranch reaction starts with the elevation of the cytosolic Ca^{2+} level triggered by eATP, leading to the accumulation of MAPK and NO, as well as to the phosphorylation and activation of the RBOHD (respiratory burst oxidase homolog protein D) subunit of the plasma membrane-localized NADPH oxidase [42]. This enzyme catalyzes the synthesis of extracellular ROS like superoxide ($O_2^{\cdot-}$), which is then converted into hydrogen peroxide (H_2O_2) in the extracellular milieu [16,33,43]. Extracellular H_2O_2 crosses the double-layered plasma membrane via various channels among which aquaporins are distinguishable [44]. Subsequently, ROS trigger changes in the expression of nuclear genes, responsible for defense responses [16], root hair growth [12] or regulation of Na^+ , H^+ , and K^+ levels. Consequently, a disturbance in the ion homeostasis may lead to the mitochondria-independent type of programmed cell death by activation of the caspase-like proteases [45].

2.3. eATP Involvement in Plant Responses to Biotic and Abiotic Stresses

Aside from different eATP-induced factors and signaling molecules contributing to the plant resistance against pathogens and various abiotic stresses, classical defense hormones such as jasmonate, ethylene, and salicylic acid are of high importance [38]. Based on the pathogens' lifestyles two groups of these organisms are distinguishable: biotrophs and necrotrophs. Interestingly, the efficiency of the used phytohormone against these two groups of pathogenesis differs. Salicylic acid induces plant defense against biotrophic pathogens, whereas jasmonate and ethylene are indispensable when necrotrophic pathogens and herbivorous insects attack the plant. Furthermore, salicylic acid regulates pathogen-induced systemic acquired resistance (SAR), whereas jasmonate and ethylene play a crucial role in rhizobacteria-mediated induced systemic resistance (ISR) [46]. Transcriptomic research of *Arabidopsis thaliana* mutants defective in the jasmonate, ethylene, and salicylic acid signaling pathway, which were treated with ATP, revealed crosstalk in the signaling of the typical plant hormones and eATP. These results showed that from among all of the defense-related genes, up to 50% were induced by eATP in cooperation with the typical plant defense hormones. This finding suggests a complex network of the plant defense mechanism that needs to be explored more deeply at different levels [38]. The contribution of transcription factors to ATP-responsive transcription is also considered. It was demonstrated that a calmodulin-binding transcription activator (CAMTA3) and MYC transcription factors are required for proper defense-related gene transcription, whose expression is induced by eATP [38].

In order to examine the importance of eATP in the plant reaction to pathogens two approaches might be considered: either alteration of the P2K1 receptor or manipulation at the eATP level. Overexpression of the P2K1 gene resulted in increased plant resistance to the bacterial pathogen *Pseudomonas syringae* and the oomycete pathogen *Phytophthora brassicae* [47,48]. Plants treated with ATP were found to be protected against various organisms such as the fungal pathogen *Botrytis cinerea* [7] and the bacterial pathogen *Pseudomonas syringae* [48]. Although the complete mechanisms of eATP signal transduction through the P2K1 receptor remain unclear, a plenitude of evidence shows the involvement of eATP in plant resistance to biotic stresses.

Studies show that eATP level, as well as P2K1 activity, are also of high importance in the plant-fungus symbiosis. Interactions of the filamentous root endophyte *Serendipita indica* with various experimental plant hosts, including *Arabidopsis thaliana* and *Hordeum vulgare*, have been examined. *Serendipita indica* colonization was found to be beneficial for plants, reflected in plant growth enhancement, assimilation of nitrate and phosphate improvement and better tolerance to both abiotic and biotic stresses [49]. Although *Serendipita indica* penetrates the host's root cells, massive plant cell death does not occur. It is because of an enzymatically active ecto-5'-nucleotidase (E5'NT) enzyme secreting by *Serendipita indica* which is capable of hydrolyzing nucleotides in the apoplast. By the hydrolyzation of ATP, ADP, and AMP, *Serendipita indica* E5'NT modifies the eATP level, leading to switching off the plant defense reactions which promote proper fungal accommodation [50]. Moreover, it was observed that the rhizobial nodule factor stimulates the release of ATP outside the cell by the root hair tips of *Medicago truncatula* [12]. Treatment of plants with eATP caused also the change their susceptibility to pathogen infection [16]. It is hypothesized that ecto-apyrases may decrease the elevated level of eATP upon symbiont infection and thereby prevents the activation of plant defense pathways that could limit symbiont invasion [51].

Several studies indicate the role of eATP in the response to different types of abiotic stresses. In addition to mechanical stimuli caused by wounding or touch, ATP is also released in response to treatment with molecules such as abscisic acid and L-glutamate [15,52]. A similar reaction was observed during both osmotic and salt stress [12,15,53] as well as under cadmium treatment [54]. Consequently, eATP accumulation triggers plant physiological changes leading to enhancement of resistance. It involves rapid closure of leaf stomata [55], probable seedling viability enhancement [56] and modification of root growth direction when encountering an obstacle [34]. In addition, hypertonic salt stress interferes with the photosynthesis machinery by decreasing the levels of maximal efficiency

of photosystem II and depleting the photochemical quenching [54]. Moreover, abiotic stress caused by cadmium triggers a rapid increase in lipid peroxidation but also higher antioxidant and lipoxygenase activities in *Arabidopsis thaliana* cells [54]. These reactions are associated with boosted synthesis of jasmonic acid, which is one of the most important molecules involved in the different stress responses [57]. Nowadays, one of the major anthropogenic pollutants with the highest level of threat to human health is cadmium [58]. It is also considered as one of the most phytotoxic elements among heavy metals, causing a reduction in crop biomass by disrupting photosynthesis and respiration [59]. In addition, cadmium can induce other physiological changes including induction of oxidative stress by boosting the production of ROS [58], modifications in gene expression [60,61], as well as changes in enzyme activity [62], hence activating plant defense.

3. Extracellular Pyridine Nucleotides

The pyridine nucleotides nicotinamide adenine dinucleotide (NAD⁺) and NAD⁺ phosphate (NADP⁺) are commonly occurring electron carriers that are involved in metabolic reactions as well as intracellular signaling [63,64]. It is known that in plants NAD(P)⁺ can be released outside the cell [65]. Extracellular NAD(P)⁺ (eNAD(P)⁺) induces the expression of pathogen-related genes and the resistance to *Pseudomonas syringae* in *Arabidopsis thaliana* through pathways involving calcium- and salicylic acid-mediated defense signaling. It was also indicated that eNAD(P)⁺ induces transcriptional and metabolic changes in *Arabidopsis thaliana* similar to those caused by pathogen infection [66]. Moreover, the expression of the human NAD(P)⁺-hydrolyzing ecto-enzyme CD38 in *Arabidopsis thaliana* partially compromises systemic acquired resistance (SAR) and this suggests that eNAD(P)⁺ can be a SAR signal molecule [67]. Recent studies on the function of eNAD(P)⁺ focus on understanding their role in plants and on identifying their receptor(s). Analysis of transcriptome changes in *Arabidopsis thaliana* evoked by eNAD⁺ identified a lectin receptor kinase (LecRK) LecRK-I.8 as a potential eNAD⁺ receptor. LecRK-I.8 is located in the plasma membrane, has kinase activity, and specifically binds only NAD⁺, but not NADP⁺, ATP, ADP or AMP. Moreover, the expression of LecRK-I.8 is induced by the eNAD⁺ [68]. Another LecRK that can be a potential receptor for eNAD(P)⁺ is LecRK-VI.2, which was identified in *Arabidopsis thaliana*. LecRK-VI.2 is constitutively associated with Brassinosteroid Insensitive1-Associated Kinase1 (BAK1) and it was shown that complex LecRK-VI.2/BAK1 is involved in SAR [69].

4. Uncommon Nucleotides as Signaling Molecules

4.1. Mononucleoside Polyphosphates

4.1.1. Structure and Occurrence of Mononucleoside Polyphosphates

Despite mono- and dinucleoside polyphosphates having been discovered in the middle of the twentieth century, our knowledge about their biological function is still poor, especially in plants. Mononucleoside polyphosphates (p_nNs) contain a nucleoside and oligophosphate chain. Examples of these nucleotides are adenosine 5'-tetraphosphate (p₄A, ppppA, Figure 2) and adenosine 5'-pentaphosphate (p₅A, pppppA). Both of them were discovered in commercial preparations of ATP obtained from bovine cells [70,71], horse muscle [72] and yeast [73]. Additionally, other purine and pyrimidine p₄Ns were found as contamination of various nucleoside triphosphates (NTP) preparations: p₄G [74,75], p₄U [76], and p₄C [77]. The existing of p₄A and p₅A was confirmed in biological materials such as rat liver [78,79], rabbit and horse muscle [80], bovine adrenal medulla [80–83], rabbit thrombocytes [84], and *Saccharomyces cerevisiae* [85]. However, the concentration of p₄A in the above-mentioned animal samples was about 2 μM, but in chromaffin granules from the adrenal medulla it was about 800 μM, and it was even up to 4 orders of magnitude and 300-fold lower than ATP concentration, respectively [83]. Until now there is no information about the content of p_nNs in the plant tissues.




Figure 2. Structure of adenosine 5'-tetraphosphate (p₄A).

4.1.2. Synthesis and Degradation of Mononucleoside Polyphosphates

The level of p_nNs in a cell depends on its biosynthesis and degradation, but also p_nNs may be a product of the degradation of some dinucleoside polyphosphates (N_{p_n}N's). Enzymes that can synthesize p_nNs in vitro can be grouped into two categories: aminoacyl-tRNA synthetases (AARS) and non-aminoacyl-tRNA synthetases (non-AARS) [86]. Among enzymes synthesizing p_nNs only one belongs to the AARS and it is lysyl-tRNA synthetase (LysRS) from *Escherichia coli*, which can synthesize p₄A [87,88]. The majority of the enzymes belong to the non-AARS. Table 1 presents non-plant enzymes synthesizing p_nNs.

Table 1. Non-plant enzymes synthesizing mononucleoside polyphosphates (p_nNs).

Enzyme	Organism	Reaction (E, Enzyme)	References
Lysyl-tRNA synthetase (EC 6.1.1.6)	<i>Escherichia coli</i>	1st step: E + lysine + pppA ↔ ↔ E:lysyl~pA + PP _i 2nd step: E:lysyl~pA + ppp → → ppppA + lysine + E	[87–98]
Luciferase EC 1.13.12.7	<i>Photinus pyralis</i>	1st step: E + luciferin + pppA → → E-luciferin~pA + PP _i 2nd step: E-luciferin~pA + (p)ppp → → (p)ppppA + luciferin + E	[99]
UTP:glucose-1-phosphate uridylyltransferase (EC 2.7.7.9)	<i>Saccharomyces cerevisiae</i>	1st step: glucose-1-P + pppU → → Upp-glucose + PP _i 2nd step: Upp-glucose + (p)ppp → → (p)ppppU + glucose-1-P	[100]
Phosphoglycerate kinase (EC 2.7.2.3)	<i>Saccharomyces cerevisiae</i>	1,3-ppGly + pppA ↔ ↔ 3-pGly + ppppA	[80,101]
Adenylate kinase (EC 2.7.4.3)	Rabbit and pig muscles	ppA + pppA ↔ pA + ppppA	[102]
Succinyl-CoA synthetase (EC 6.2.1.5)	<i>Escherichia coli</i>	E-P + pppA ↔ E + ppppA	[103]
Acetyl-CoA synthetase (EC 6.2.1.1)	<i>Saccharomyces cerevisiae</i>	acetyl~pA + ppp ↔ acetate + ppppA	[104,105]
Acyl-CoA synthetase (EC 6.2.1.3)	<i>Pseudomonas fragi</i>	acyl~pA + ppp ↔ fatty acid + ppppA	[106,107]
DNA ligase (EC 6.5.1.1)	T4 phage, <i>Pyrococcus furiosus</i>	E-pA + ppp ↔ E + ppppA	[105,108–110]
RNA ligase (EC 6.5.1.3)	T4 phage	E-pA + (p)ppp ↔ E + (p)ppppA	[111]
UDP-MurNAc-L-alanine:D- glutamate ligase (EC 6.3.2.9)	<i>Escherichia coli</i>	E:acyl~P + pppA ↔ ppppA + acyl + E	[112]

Among enzymes that can degrade p_nNs some of them exhibit low substrate specificity, for example, alkaline (EC 3.1.3.1) and acid phosphatases (EC 3.1.3.2), that release phosphate residues up to adenosine [113]. Apyrase (EC 3.6.1.5) also can cut off phosphate residues, but only to

AMP [113]. Phosphodiesterase I (EC 3.1.4.1) degrades p₄A to triphosphate and AMP [100]. Additionally, adenosine-phosphate deaminase from *Aspergillus oryzae* and *Helix pomatia* is involved in p₄A metabolism, converting it into inosine 5'-tetraphosphate (p₄I) [114].

As mentioned above, p_nNs can accumulate in the cell as a result of the degradation of Np_nN's. The way in which p₄A can accumulate in a cell is the degradation of diadenosine 5',5'''-pentaphosphate (Ap₅A) and diadenosine 5',5'''-hexaphosphate (Ap₆A). The degrading enzymes include phosphodiesterase I (EC 3.1.4.1) occurring in prokaryotes and eukaryotes [115,116], symmetrical dinucleoside tetraphosphatase (EC 3.6.1.41) from bacteria [117], dinucleoside tetraphosphate phosphorylase (EC 2.7.7.53) from *Saccharomyces cerevisiae* and *Euglena gracilis* [118,119], and dinucleoside triphosphatase (EC 3.6.1.29) among others from *Lupinus luteus* [120].

Despite the lack of knowledge about the occurrence and the concentration of mononucleoside polyphosphates in higher plants, there are described a few enzymes which can synthesize p_nNs. All of them are listed in Table 2. The first described plant enzyme that synthesizes p_nNs is 4-coumarate:CoA ligase (4CL2) from *Arabidopsis thaliana* that catalyzes the reaction of the synthesis of both p₄A and p₅A [121]. This enzyme is a branch point in the phenylpropanoid pathway that leads to the biosynthesis of flavonoids, lignin, and stilbenes. It is known that the phenylpropanoid pathway is involved in plant responses to numerous environmental stimuli, especially under biotic and abiotic stresses [122]. Another enzyme synthesizing p₄A is jasmonate:amino acid synthetase from *Arabidopsis thaliana* (JAR1) [123]. JAR1 is involved in the function of jasmonic acid (JA) as a plant hormone and it catalyzes the synthesis of several JA-amido conjugates, the most important of which appears to be jasmonic acid-isoleucine. Both of the above-mentioned plant enzymes belong to the acyl~adenylate-forming firefly luciferase superfamily [124] which catalyzes a two-step reaction. During the first step, acyl and ATP form an acyl~adenylate intermediate with the simultaneous release of pyrophosphate (PP_i). In the second step, 4CL in the absence of CoA catalyzes the formation of uncommon mononucleoside polyphosphates such as p₄A and p₅A [121].

Table 2. Plant enzymes synthesizing mononucleoside polyphosphates (p_nNs).

Enzyme	Organism	Reaction (E, Enzyme)	References
4-Coumarate:CoA ligase (4CL2) (EC 6.2.1.12)	<i>Arabidopsis thaliana</i>	1st step: E + coumarate + ATP → → E:coumaroyl~pA + PP _i 2nd step: E:coumaroyl~pA + (p)ppp → → (p)ppppA + coumarate + E	[121]
Jasmonate:amino acid synthetase (JAR1) (EC 6.3.2.52)	<i>Arabidopsis thaliana</i>	1st step: E + jasmonate + ATP → → E:jasmonyl~pA + PP _i 2nd step: E:jasmonyl~pA + ppp → → ppppA + jasmonate + E	[123]

In plants there also exists an enzyme degrading p₄A. It is nucleoside tetraphosphate hydrolase (EC 3.6.1.14) occurring in *Lupinus luteus* seeds. This enzyme hydrolyzes both p₄A and p₄G with the same rate while p₅A is degraded up to 200-fold more slowly than p₄A and p₄G [120].

4.2. Adenosine 5'-Phosphoramidate

Structure, Occurrence, and Metabolism of Adenosine 5'-Phosphoramidate

Among naturally occurring uncommon mononucleotides is adenosine 5'-phosphoramidate (NH₂-pA, Figure 3). This compound is believed to occur in all organisms; however, so far NH₂-pA has only been detected among cellular nucleotides purified from the green alga *Chlorella pyrenoidosa* [125]. Similarly to p_nNs, the level of NH₂-pA in a cell is enzymatically controlled. It is known that NH₂-pA can be synthesized by adenylyl sulfate:ammonia adenylyltransferase (EC 2.7.7.51) in the algae *Chlorella pyrenoidosa*, *Euglena gracilis*, amoeba *Dictyostelium discoideum*, bacteria *Escherichia coli*, and in higher plants *Hordeum vulgare*, *Spinacia oleracea* [126], and *Lupinus luteus* [127]. This transferase catalyzes the following reaction: SO₄-pA + NH₄⁺ → NH₂-pA + SO₄²⁻ + 2H⁺.




Figure 3. Structure of adenosine 5'-phosphoramidate (NH₂-pA).

The supposition that NH₂-pA is a ubiquitous compound and that its concentration is enzymatically controlled may be supported by the existence of various enzymes that catalyze the cleavage of NH₂-pA to ammonia and AMP (pA) by hydrolysis [128–133], or to ammonia and ADP (ppA) by phosphorolysis [130]. Both the synthesis [127] and the degradation [131] of NH₂-pA can be controlled by HIT (histidine triad) proteins. One of the proteins belonging to the HIT family proteins is Fhit (fragile histidine triad). Based on the mechanism of the action of Fhit protein, which is able to hydrolyze the P-N bond in adenosine phosphoimidazolidine [134], it is hypothesized that other uncommon nucleotides can be substrates for human and *Arabidopsis thaliana* Fhit. Other proteins belonging to HIT family proteins are Hint protein (having the activity of NH₂-pA hydrolase) and GalT proteins (having the specific activity of nucleoside monophosphate transferase) [135]. The protein Fhit from human and *Arabidopsis thaliana* exhibited the activity attributed to adenylyl sulfate sulfohydrolase and nucleoside phosphoamidases releasing AMP from adenosine 5'-phosphosulfate (SO₄-pA) and NH₂-pA, respectively. Fhit protein also catalyzes the hydrolysis of the P-F bond in the synthetic nucleotide adenosine 5'-monophosphofluoride (F-pA) releasing AMP [136]. Recently it has been shown that NH₂-pA can also be synthesized by Fhit proteins from *Lupinus luteus* seeds and *Arabidopsis thaliana*. It catalyzes the ammonolysis of SO₄-pA, leading to the formation of NH₂-pA [127].

4.3. Dinucleoside Polyphosphates

4.3.1. Structure and Occurrence of Dinucleoside Polyphosphates

Dinucleoside polyphosphates (Np_nN's) consist of an oligophosphate chain that links the two 5'-esterified nucleosides. Among Np_nN's that are most frequently tested are adenine nucleotides: diadenosine triphosphate (Ap₃A) and diadenosine tetraphosphate (Ap₄A, Figure 4). The first reports about the existence of Np_nN's come from the 1950s. It was discovered that during the chemical synthesis of adenosine-uridine monophosphate (ApU) there were synthesized byproducts such as diadenosine diphosphate (Ap₂A) and diuridine diphosphate (Up₂U) [137]. Nevertheless, the first Np_nN discovered in the biological material was diguanosine tetraphosphate (Gp₄G). The presence of Gp₄G and Gp₃G was observed in biological preparations in encysted gastrulae of the brine shrimp and *Artemia salina* in millimolar concentrations [138,139]. The most common Np_nN is Ap₄A, which for the first time was detected in rat liver, and its concentration was estimated at 30 nM [97]. Subsequently Ap₄A and other Np_nN's have been observed in submicromolar concentrations in many investigated cells and tissues [140,141]. Np_nNs were identified in bacteria [142,143], yeast [144,145], and animal and human cells [140,146–150]. So far, Np_nN's have not been identified in plants.




Figure 4. Structure of diadenosine 5', 5'''-tetrphosphate (Ap_4A).

4.3.2. Synthesis and Degradation of Dinucleoside Polyphosphates

The level of Np_nN' s in cells may be a result of the synthesis and degradation of these compounds. Table 3 lists non-plant enzymes synthesizing various Np_nN' s. Enzymes synthesizing Np_nN' s are some ligases [89,92,106,121,151,152], luciferase from *Photinus pyralis* [153], and some transferases [100,119,154]. Among aminoacyl-tRNA synthetases, the most effective in Ap_nN synthesis are lysyl- (EC 6.1.1.6), phenylalanyl- (EC 6.1.1.20), alanyl- (EC 6.1.1.7), and prolyl- (EC 6.1.1.15) t-RNA synthetases [86]. Moreover, the RNA-dependent RNA polymerase elongation complex from hepatitis C virus (HCV) can use in vitro nucleoside triphosphates (NTPs) to excise the terminal nucleotide in nascent RNA and mismatched ATP, UTP, or CTP could mediate excision of 3'-terminal CMP to generate the dinucleoside tetraphosphate products Ap_4C , Up_4C , and Cp_4C , respectively [155].

Table 3. Non-plant enzymes synthesizing dinucleoside polyphosphates (Np_nN' s).

Enzyme	Organism	Reaction (E, Enzyme)	References
Luciferase (EC 1.13.12.7)	<i>Photinus pyralis</i>	1st step: $E + \text{luciferin} + \text{pppA} \rightarrow$ $\rightarrow E\text{-luciferin}\sim\text{pA} + \text{PP}_i$ 2nd step: $E\text{-luciferin}\sim\text{pA} + \text{pppN} \rightarrow$ $\rightarrow \text{AppppN} + \text{luciferin} + E$	[99]
GTP:GTP guanylyltransferase (EC 2.7.7.45)	<i>Artemia salina</i> , <i>Saccharomyces cerevisiae</i>	1st step: $E + \text{pppN} \leftrightarrow E\sim\text{pN} + \text{PP}_i$ 2nd step: $E\sim\text{pN} + \text{pppN} \rightarrow \text{NppppN}' + E$	[154]
UTP:glucose-1-phosphate uridylyltransferase (EC 2.7.7.9)	<i>Saccharomyces cerevisiae</i>	1st step: $\text{glucose-1-P} + \text{pppU} \rightarrow$ $\rightarrow \text{Upp-glucose} + \text{PP}_i$ 2nd step: $\text{Upp-glucose} + \text{pppN} \rightarrow$ $\rightarrow \text{UppppN} + \text{glucose-1-P}$	[100]
Acyl-CoA synthetase (EC 6.2.1.3)	<i>Pseudomonas fragi</i>	1st step: $\text{Fatty acid} + \text{pppA} \leftrightarrow$ $\leftrightarrow \text{acyl}\sim\text{pA} + \text{PP}_i$ 2nd step: $\text{acyl}\sim\text{pA} + \text{pppA} \rightarrow$ $\rightarrow \text{AppppA} + \text{fatty acid}$	[106,107]
RNA ligase (EC 6.5.1.3)	T4 phage	$E\sim\text{pA} + \text{pppN} \leftrightarrow E + \text{AppppN}$	[111]
DNA ligase (EC 6.5.1.1)	T4 phage, <i>Pyrococcus</i> <i>furiosus</i>	$E\sim\text{pA} + \text{pppN} \leftrightarrow E + \text{AppppN}$	[105,108–110]
Lysyl-tRNA synthetase (EC 6.1.1.6)	<i>Escherichia coli</i>	1st step: $E + \text{lysine} + \text{pppA} \leftrightarrow$ $\leftrightarrow \text{lysyl}\sim\text{pA} + \text{PP}_i$ 2nd step: $E:\text{lysyl}\sim\text{pA} + \text{pppN} \rightarrow$ $\rightarrow \text{AppppN} + \text{lysine} + E$	[86]

So far, only three enzymes synthesizing Np_nN' s have been described in plants (Table 4). Among aminoacyl-tRNA synthetases only phenylalanyl- (EC 6.1.1.20) and seryl-tRNA (EC 6.1.1.11) synthetases from *Lupinus luteus* [152] synthesize Ap_nN . The mechanism of the synthesis of Ap_nN s catalyzed by plant aminoacyl-tRNA synthetases is based on the formation of aminoacyl \sim pA and the transfer of

adenylate to pppN [152]. The third enzyme which is involved in the synthesis of Ap_nN in plant cells is 4:coumarate-CoA ligase (EC 6.2.1.12) [121].

Table 4. Plant enzymes synthesizing dinucleoside polyphosphates (Np_nN's).

Enzyme	Organism	Reaction (E, Enzyme)	References
Phenylalanyl-tRNA synthetase (EC 6.1.1.20)	<i>Lupinus luteus</i>	1st step: E + phenylalanine + pppA ↔ ↔ phenylalanyl~pA + PPi 2nd step: E:phenylalanyl~pA + pppN → → AppppN + phenylalanine + E	[152]
Seryl-tRNA synthetase (EC 6.1.1.11)	<i>Lupinus luteus</i>	1st step: E + serine + pppA ↔ ↔ seryl~pA + PPi 2nd step: E:seryl~pA + pppA → → AppppA + serine + E	[152]
4-Coumarate:CoA ligase (4CL2) (EC 6.2.1.12)	<i>Arabidopsis thaliana</i>	1st step: E + coumarate + pppA → → E:coumaroyl~pA + PPi 2nd step: E:coumaroyl~pA + pppN → → NppppN + coumarate + E	[121]

Enzymes degrading Np_nN's may be divided into substrate-specific and substrate non-specific. Substrate-specific enzymes degrading Np_nN's include, among others, *asymmetrical* Np₄N hydrolase (EC 3.6.1.17), *symmetrical* Np₄N hydrolase (EC 3.6.1.41), and dinucleoside triphosphate (Np₃N) hydrolase (EC 3.6.1.29) [156]. In plants, the activity of *asymmetrical* Np₄N hydrolase was detected in *Lupinus luteus* seeds [157], *Helianthus annuus* and marrow (*Cucurbita pepo*) seeds [158], tomato cells [159], *Lupinus angustifolius* seeds [160] and in *Hordeum vulgare* [161]. Among Np₄N's, only Ap₄A can be degraded by the *asymmetrical* Np₄N hydrolase giving ATP. The substrate non-specific enzymes in plants are phosphodiesterase I (EC 3.1.4.1) from *Lupinus luteus* [157] and nucleotide pyrophosphatase (EC 3.6.1.9) from potato tuber [162].

4.4. Function of Uncommon Nucleotides

The cellular level of Np_nN's is modified under various physiological and pathological conditions and the compounds have been suggested as intracellular messengers in diverse cellular processes [163,164]. The increased concentration of Ap₄A was correlated with high cellular proliferation rates [165] or some phases of the cell cycle [165,166]. A dramatic increase of levels of Np₄N and various Np_nN's has been observed in cells subjected to stresses, such as elevated temperature, ethanol or cadmium [143–145,167]. A very interesting function of uncommon nucleotides is their potential function as alarmones [168]. However, so far no clear metabolic or molecular targets or receptors of the postulated alarm signaled by the Np_nN's have been experimentally demonstrated. Alarmones are intracellular signaling molecules that are produced under adverse environmental factors. They regulate gene expression at the transcriptional level. Nucleotides (p)ppGpp and ppGpp have such a role in bacteria [169]. The results of recent studies on the role of uncommon nucleotides focus on microorganisms and animals, including man. It has been demonstrated that Np_nN's may accumulate in cells under stress conditions. In cells of *Saccharomyces cerevisiae* and *Escherichia coli* under elevated temperature and cadmium ions an increased level of Np_nN's was observed. The intracellular concentrations of these molecules vary from 10⁻⁹ M in a basal metabolic state to 10⁻⁴ M when the organisms are subjected to stresses, such as heat shock or exposure to cadmium [143,167]. In cyanobacteria the same effects are a result of thermal shock and heavy metals [145]. In *Salmonella*, the increased synthesis of Np_nN's was caused by ethanol [142]. Studies performed on orange fruit showed that under high temperature enzymes involved in Np_nN' metabolism were activated and an extremely large increase in bis(5'-adenosyl) triphosphatase protein content was observed. These results suggest intensive synthesis of Ap₃A under high temperature [170]. Another group of enzymes involved in Np_nN' metabolism is Nudix (nucleoside diphosphate linked to x) hydrolases (NUDTs). An increase of transcript and protein NUDT7 levels was found in *Arabidopsis thaliana* plants under biotic stress evoked by *Pseudomonas syringae* avrRpt2 [171]. In *Arabidopsis thaliana*, there was also observed a quick

increase in NUDT7 content under the effect of ozone and pathogen infection [172]. Recent studies on AtNUDT19 indicate the involvement of this hydrolase in the photo-oxidative stress response by regulating photosynthesis, the antioxidant system and the synthesis of salicylic acid—a signaling molecule involved in the response to biotic and abiotic stresses [173]. Additionally, it is suggested that barley NUDTs respond to abiotic stress [174]. It is also known that NUDTs bind RNA and participate in the regulation of gene expression in animals and plants [175–177].

Studies on the function of Ap₄A in *Escherichia coli* have shown that it may be a damage metabolite and a few proteins binding Ap₄A in these bacteria were identified [178]. Recent studies on the function of Ap₄A have shown that aminoglycoside antibiotics induced synthesis of Ap₄A in *Escherichia coli* and it caused bacterial cell killing by these antibiotics [179]. Lysyl-tRNA synthetase plays a key role in MITF (microphthalmia-associated transcription factor) transcriptional activity via Ap₄A as an important signaling molecule in mast cells [180,181]. Studies on chronic myelogenous leukemia cells showed that NUDT2 disruption elevates Ap₄A and down-regulates immune response and cancer promotion genes [182]. Ap₃A and Ap₄A induce activation of a signaling pathway that results in increase of proliferation of vascular smooth muscle cells by stimulation of ERK1/2(MAP kinase) [183]. It is known that Np_nN's can act as neurotransmitters and influence the vascular system through purinergic receptors. There is also evidence that Ap₄A inhibits the initiation of DNA replication. It is also proposed that Ap₄A acts as an important inducible ligand in the DNA damage response to prevent the replication of damaged DNA [184]. In contrast, the increase in the ratio of Ap₃A/Ap₄A in human HL60 cells can induce apoptosis [164].

So far there are only a few papers describing the role of uncommon nucleotides in plant cells. It was shown that the phenylpropanoid pathway may be affected by exogenous uncommon nucleotides [185–188]. The phenylpropanoid pathway is a source of secondary metabolites which are very important in plant responses to biotic and abiotic stresses [122]. The first data showed that Ap₃A and Ap₄A regulate the expression of genes and the activity of enzymes involved in the phenylpropanoid pathway in seven-day old *Arabidopsis thaliana* seedlings [185]. Ap₃A and Ap₄A caused an increase in the activity of phenylalanine ammonia-lyase (PAL) and 4-coumarate-CoA ligase (4CL) just 10 min after application. Analysis of *PAL* gene expression showed that there was strong induction (about 70-fold) of *PAL2* gene expression by Ap₃A and Ap₄A. Additionally, PAL activity was significantly enhanced by Ap₃A (about 9-fold). Moreover, it has been shown that the activity of 4CL and the *4CL* gene expression level were higher in seedlings treated with Ap₃A and Ap₄A [185]. These results may indicate a dual role of 4CL in the plant response to stress factors. Firstly, this enzyme synthesizes Np_nN's, i.e., compounds that can play a function as alarmones [121], and secondly, it synthesizes secondary metabolites minimizing the adverse effects of stresses in plant cells [189]. Also, Ap₃A caused changes in the gene expression level and the activity of chalcone synthase (CHS) in *Arabidopsis thaliana* seedlings. This enzyme catalyzes the synthesis of chalcone, which is the precursor of secondary metabolites—flavonoids [185]. These data strongly indicate the signaling role of Np_nN's in plants. The inductive effect of Ap₃A on gene expression of the phenylpropanoid pathway proteins has also been confirmed in a cell suspension of *Vitis vinifera* cv. Monastrell. A synergistic effect evoked by Ap₃A and cyclodextrins in *trans*-resveratrol biosynthesis has been demonstrated [186]. It may suggest the involvement of Np_nN's in the plant response to stress. Interestingly, cyclodextrins act as elicitors in their chemical similarity to the alkyl-derivatized pectic oligosaccharides, which are released from the cell wall during fungal infection [190]. It is known that one of the defense strategies of higher plants against biotic and abiotic stresses is activation of the phenylpropanoid pathway leading to enhanced production of various phenylpropanoid compounds, such as flavonoids [191,192], lignin [193,194], anthocyanins [194–196], and salicylic acid [197] (Figure 5). Recently the differences in the regulation of gene expression of the phenylpropanoid pathway enzymes and phenylpropanoid accumulation by purine or pyrimidine Np_nN's in *Vitis vinifera* suspension cell culture were described. The pyrimidine dinucleotides, such as Cp₃C, Cp₄C, and Ap₄C, markedly (6- to 8-fold) induced the expression of the gene coding the cinnamoyl-CoA:NADP oxidoreductase (CCR) that controls lignin biosynthesis.

The most effective in stilbene accumulation was Up₄U, but other pyrimidine dinucleotides (Cp₃C, Cp₄C, and Ap₄C) strongly inhibited the biosynthesis of these phenylpropanoids [188].




Figure 5. Hypothetical working model of Np_nN' and NH₂-pA signaling network in a plant cell. Dinucleoside polyphosphate (Np_nN) and adenosine 5'-phosphoramidate (NH₂-pA) trigger induction of the phenylpropanoid and the isochlorogenic acid pathways yielding metabolites protecting plant against various types of stresses. Plant cells respond to environmental stimuli by intensification of the production of various compounds, such as anthocyanins, lignin, stilbenes, and salicylic acid. Question marks indicate the hypothetical components of the signaling network.

As described above, in plants, the presence of enzymes degrading and synthesizing another uncommon nucleotide, NH₂-pA, has been described [127,132]. It was evidenced that NH₂-pA regulates the expression of genes coding enzymes of the phenylpropanoid pathway, such as PAL,

cinnamate-4-hydroxylase (C4H), 4CL, CHS, CCR, and isochorismate synthase (ICS) in *Arabidopsis thaliana* seedlings. CCR is an enzyme involved in the biosynthesis of lignin, whereas ICS is involved in the biosynthesis of a signaling molecule—salicylic acid. Among the analyzed genes the strongest induction in gene expression caused by NH₂-pA was observed for CCR2 (approx. 4-fold). This was also accompanied by an increased level of lignin in the seedlings. Another important effect caused by NH₂-pA was an increase in the ICS gene expression and a significant increase (2-fold) in the level of free salicylic acid. In view of the fact that salicylic acid is one of the signaling molecules involved in the response of plants to biotic stress, the induction of the synthesis of salicylic acid by NH₂-pA may suggest the interaction of both these compounds in plant responses to stresses [187].

Another uncommon nucleotide occurring in plants is nicotinic acid adenine dinucleotide phosphate (NAADP). The synthesis of NAADP is not fully understood. However, it is supposed that it can be synthesized by adenosine-5'-diphosphateribosyl-cyclase (EC 3.2.2.5) [198]. This nucleotide is involved in calcium signaling in many organisms including plants [199–201]. NAADP-mediated calcium release has been shown in the microsomal vesicles of red beets (*Beta vulgaris*) and cauliflower (*Brassica oleracea*). Analysis of sucrose gradient-separated cauliflower microsomes revealed that the NAADP-sensitive Ca²⁺ pool was derived from the endoplasmic reticulum [199].

4.5. Uncommon Nucleotide Signal Transduction Pathways in Plants

As yet there has been no information about signal transduction pathways or receptors activated by uncommon nucleotides. Based on the data existing in the literature about the involvement of these molecules in different biochemical and physiological processes in bacteria, fungi, animal, and plant organisms, we propose putative signaling pathways induced by uncommon nucleotides in plant cells (Figure 5). It is hypothesized that uncommon nucleotides may be signal molecules synthesized by pathogens outside the cell and interact with plant cells through unknown plasma membrane receptor(s). These nucleotides can also be transported inside the plant cell by unknown plasma membrane transporter(s). In addition, their level in the plant cell can be regulated by synthesizing and degrading enzymes. Both the extracellular and intracellular uncommon nucleotides may affect the biosynthesis of other plant cell signal molecules, e.g., secondary messengers and hormones that can engage MAPK cascades, which are involved in plant growth and development, cellular responses to hormones, regulation of the cell cycle, and responses to biotic and abiotic stresses, such as pathogen infection, wounding, low temperature, drought, high salinity, metals, and ROS. Moreover, it is known that MAPK cascades can regulate gene transcription by activation or repression of transcription factors. In addition, MAPK transcript levels in *Arabidopsis thaliana* seedlings were shown to increase in a time-dependent manner following exposure to Cu and Cd [202]. Many MAPK cascades respond to hormones such as abscisic acid, jasmonic acid, salicylic acid, ethylene, auxins, and brassinosteroids. Usually, signaling molecules participate in distinct signaling pathways, resulting in the formation of a cross-talking network that co-ordinates responses to different stresses [202–204]. We postulate that MAPK cascades can be involved in the signal transduction pathway triggered by uncommon nucleotides. So far, it is only known that extracellular uncommon nucleotides can induce gene expression of the phenylpropanoid pathway, the accumulation of phenylpropanoids, and the synthesis of salicylic acid. However, components of the nucleotide transduction pathway are still not known (Figure 5).

5. Conclusions

Plants are not able to escape from their abiotic (cold, heat, etc.) or their biotic (herbivores, insects) environment [205]. Furthermore, their food uptake and gas exchange take place through external surfaces (leaves: light and CO₂, roots: ions and water). Plants must, therefore, possess systems to exchange information throughout the entire plant to ensure the coordination of plant development and defense. Evidence suggests that in plants information exchange relies on at least two different systems: one involving molecules that are transported within the plant and another that uses electrical

and/or hydraulic signals to carry the information throughout the entire organism [205]. The systems for transmitting this information are complex and involve multiple components, which are far from being understood. For many years plant hormones were considered to be dominant molecules in plant signaling. Nowadays, this term is applied to many more compounds. It is suggested that dynamic changes in the level of second messengers, such as Ca^{2+} , ROS, and NO, serve as signatures for both intracellular signaling and cell-to-cell communications. These second messenger signatures work in concert with physical signal signatures (such as electrical and hydraulic signals) to create a “lock and key” mechanism that triggers an appropriate response to various stresses [206]. Part of the system, new players in plant signaling, are chemical signals such as extracellular ATP and uncommon nucleotides described and discussed in this work. These molecules are essential for proper coordination of processes in particular organs as well as responses to internal and external signals playing analogical or similar functions in plants and in animals. The data obtained during the last few years suggest that in plants both purine and pyrimidine dinucleotides should be considered as members of a family of naturally occurring stress signaling molecules. So far, our knowledge of these signals in plants is still insufficient to clearly understand their signaling role. The mechanisms of signal perception and transduction are unknown. Further studies are required, which should aim at describing the signaling networks involving uncommon nucleotides, and also perform omic experiments to identify the network components and processes that are regulated by these signals.

Author Contributions: M.P.-B. and E.S.-N. conceived the topic of the article; M.P.-B. and J.D. wrote the majority of the manuscript and designed all figures; S.B. drew all the figures and was involved in the writing of the manuscript; E.S.-N. was involved in the discussion and writing of the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: The publication was co-financed within the framework of the Ministry of Science and Higher Education program as “Regional Initiative Excellence” in years 2019–2022, Project No. 005/RID/2018/19.

Acknowledgments: Compliments to Andrzej Guranowski for his invaluable contribution to knowledge about the metabolism of uncommon nucleotides and special thanks to Professor from M.P.-B. for supervising Ph.D. thesis, many years of fruitful collaboration, inspiration in science and common scientific path. We thank Richard Ashcroft (Bioscience editor, www.anglopolonia.com/home.html) for the professional language editing of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

1,3-ppGly	1,3-Diphosphoglycerate
3-pGly	3-Phosphoglycerate
4CL	4-coumarate:CoA ligase
AARS	Aminoacyl t-RNA synthetase
ABC transporter	ATP-binding cassette ABC transporter
AMP, pA	Adenosine 5'-monophosphate
Ap ₃ A, ApppA	Diadenosine 5',5'''-P ¹ ,P ³ -triphosphate
Ap ₄ A, AppppA	Diadenosine 5',5'''-P ¹ ,P ⁴ -tetraphosphate
Ap ₄ C	P ¹ -(5'-Adenosyl)-P ⁴ -(5'-cytidyl)-tetraphosphate
ApU	Adenosine-uridine monophosphate
APY1 and APY2	Apyrase 1 and 2
C4H	Cinnamate-4-hydroxylase
cAMP	Adenosine 3',5'-cyclic monophosphate
CAMTA3	Calmodulin-binding transcription activator
CCR	Cinnamoyl-CoA:NADP oxidoreductase
cGMP	Guanosine 3',5'-cyclic monophosphate
CHS	Chalcone synthase
Cp ₃ C	Dicytosine 5',5'''-P ¹ ,P ³ -triphosphate
Cp ₄ C	Dicytosine 5',5'''-P ¹ ,P ⁴ -tetraphosphate
DORN1	Does not Respond to Nucleotide

E	Enzyme
eATP	Extracellular ATP
eNAD ⁺	Extracellular NAD ⁺
eNADP ⁺	Extracellular NADP ⁺
Fhit	Fragile histidine triad
F-pA	Adenosine 5'-O-phosphorofluoridate
Gp ₃ G	Diguanosine 5',5'''-P ¹ ,P ³ -triphosphate
Gp ₄ G	Diguanosine 5',5'''-P ¹ ,P ⁴ -tetraphosphate
HIT	Histidine triad
ICS	Isochorismate synthase
ISR	Induced systemic resistance
JAR1	Jasmonate:amino acid synthetase
LecRK-1.9	L-type lectin receptor-like kinase 1.9
LysRS	Lysyl-tRNA synthetase
MAPK	Mitogen-activated protein kinases
MITF	Microphthalmia-associated transcription factor
MYC	Transcription factors
NAADP	Nicotinic acid adenine dinucleotide phosphate
NH ₂ -pA	Adenosine 5'-phosphoramidate
NO	Nitric oxide
Non-AARS	Non-aminoacyl t-RNA synthetase
Np _n N'	Dinucleoside polyphosphate
NTP, pppN	Nucleoside 5'-triphosphate
NUDT	Nudix hydrolase
Nudix	Nucleoside diphosphate linked to x (x: any moiety)
P2K1	P2 receptor kinase 1
P2X	Ligand-gated ion channel
P2Y	G protein-coupled receptor
P ₃ , ppp	Triphosphate
P ₄ , pppp	Tetraphosphate
p ₄ A, ppppA	Adenosine 5'-tetraphosphate
p ₄ C	Cytosine 5'-tetraphosphate
p ₄ G	Guanosine 5'-tetraphosphate
p ₄ U	Uridine 5'-tetraphosphate
p ₅ A, pppppA	Adenosine 5'-pentaphosphate
PAL	Phenylalanine ammonia-lyase
PGP1	p-Glycoprotein 1
PM-ANT1	Plasma membrane-localized nucleotide transporters
p _n N	Nucleoside 5'-polyphosphate
ppGpp	Guanosine 3'-diphosphate 5'-diphosphate
PP ₁	Pyrophosphate
(p)ppGpp	Guanosine 3'-diphosphate 5'-triphosphate
RBOHD	Respiratory burst oxidase homolog protein D
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
SAR	Systemic acquired resistance
STS	Stilbene synthase
TF	Transcription factor
Up ₂ U	Uridine 5',5'''-P ¹ ,P ² -diphosphate
UTP, pppU	Uridine 5'-triphosphate

References

1. Edel, K.H.; Marchadier, E.; Brownlee, C.; Kudla, J.; Hetherington, A.M. The evolution of calcium-based signalling in plants. *Curr. Biol.* **2017**, *27*, 667–679. [[CrossRef](#)]

2. Del Rio, L.A. ROS and RNS in plant physiology: An overview. *J. Exp. Bot.* **2015**, *66*, 2827–2837. [[CrossRef](#)]
3. Pietrowska-Borek, M.; Chadzinikolau, T.; Borek, S. Cyclic nucleotides and nucleotide cyclases in plants under stress. In *Improvement of Crops in the Era of Climatic Changes*; Ahmad, P., Wani, M.R., Azooz, M.M., Tran, L.S.P., Eds.; Springer Science and Business Media: New York, NY, USA, 2014; pp. 119–151.
4. Isner, J.C.; Maathuis, F.J.M. cGMP signalling in plants: From enigma to main stream. *Funct. Plant Biol.* **2016**, *45*, 93–101. [[CrossRef](#)]
5. Świeżawska, B.; Duszyn, M.; Jaworski, K.; Szmidt-Jaworska, A. Downstream targets of cyclic nucleotides in plants. *Front. Plant Sci.* **2018**, *9*, 1428. [[CrossRef](#)]
6. Gehring, C.; Turek, I.S. Cyclic nucleotide monophosphates and their cyclases in plant signaling. *Front. Plant Sci.* **2017**, *8*, 1704. [[CrossRef](#)]
7. Tripathi, D.; Tanaka, K. A crosstalk between extracellular ATP and jasmonate signaling pathways for plant defense. *Plant Signal. Behav.* **2018**, *13*, e1432229. [[CrossRef](#)] [[PubMed](#)]
8. Burnstock, G. Purinergic nerves. *Pharmacol. Rev.* **1972**, *24*, 509–581.
9. Drury, A.N.; Szent-Gyorgy, A. The physiological activity of adenine compounds with special reference to their action upon the mammalian heart. *J. Physiol.* **1929**, *68*, 213–237. [[CrossRef](#)] [[PubMed](#)]
10. Cheung, K.K.; Chan, W.Y.; Burnstock, G. Expression of P2X purinoceptors during rat brain development and their inhibitory role on motor axon outgrowth in neural tube explant cultures. *Neuroscience* **2005**, *133*, 937–945. [[CrossRef](#)]
11. Choi, J.; Tanaka, K.; Cao, Y.; Qi, Y.; Qiu, J.; Liang, Y.; Lee, S.Y.; Stacey, G. Identification of a plant receptor for extracellular ATP. *Science* **2014**, *343*, 290–294. [[CrossRef](#)]
12. Kim, S.Y.; Sivaguru, M.; Stacey, G. Extracellular ATP in plants. Visualization, localization, and analysis of physiological significance in growth and signaling. *Plant Physiol.* **2006**, *142*, 984–992. [[CrossRef](#)] [[PubMed](#)]
13. Wu, S.J.; Liu, Y.S.; Wu, J.Y. The signaling role of extracellular ATP and its dependence on Ca²⁺ flux in elicitation of *Salvia miltiorrhiza* hairy root cultures. *Plant Cell Physiol.* **2008**, *49*, 617–624. [[CrossRef](#)] [[PubMed](#)]
14. Weerasinghe, R.R.; Swanson, S.J.; Okada, S.F.; Garrett, M.B.; Kim, S.Y.; Stacey, G.; Boucher, R.C.; Gilroy, S.; Jones, A.M. Touch induces ATP release in *Arabidopsis* roots that is modulated by the heterotrimeric G-protein complex. *FEBS Lett.* **2009**, *583*, 2521–2526. [[CrossRef](#)] [[PubMed](#)]
15. Dark, A.; Demidchik, V.; Richards, S.L.; Shabala, S.; Davies, J.M. Release of extracellular purines from plant roots and effect on ion fluxes. *Plant Signal. Behav.* **2011**, *6*, 1855–1857. [[CrossRef](#)] [[PubMed](#)]
16. Song, C.J.; Steinebrunner, I.; Wang, X.; Stout, S.C.; Roux, S.J. Extracellular ATP induces the accumulation of superoxide via NADPH oxidases in *Arabidopsis*. *Plant Physiol.* **2006**, *140*, 1222–1232. [[CrossRef](#)] [[PubMed](#)]
17. Thomas, C.; Rajagopal, A.; Windsor, B.; Dudler, R.; Lloyd, A.; Roux, S.J. A role for ectophosphatase in xenobiotic resistance. *Plant Cell* **2000**, *12*, 519–533. [[CrossRef](#)] [[PubMed](#)]
18. Rieder, B.; Neuhaus, H.E. Identification of an *Arabidopsis* plasma membrane-located ATP transporter important for anther development. *Plant Cell* **2011**, *23*, 1932–1944. [[CrossRef](#)]
19. Lim, M.H.; Wu, J.; Clark, G.; Roux, S.J. Apyrase suppression raises extracellular ATP levels and induces gene expression and cell wall changes characteristic of stress responses. *Plant Physiol.* **2014**, *164*, 2054–2067. [[CrossRef](#)]
20. Clark, G.; Roux, S.J. Apyrases, extracellular ATP and the regulation of growth. *Curr. Opin. Plant Biol.* **2011**, *14*, 700–706. [[CrossRef](#)]
21. Parsons, H.T.; Christiansen, K.; Knierim, B.; Carroll, A.; Ito, J.; Batth, T.S.; Smith-Moritz, A.M.; Morrison, S.; McNerney, P.; Hadi, M.Z.; et al. Isolation and proteomic characterization of the *Arabidopsis* Golgi defines functional and novel components involved in plant cell wall biosynthesis. *Plant Physiol.* **2012**, *159*, 12–26. [[CrossRef](#)]
22. Chiu, T.Y.; Christiansen, K.; Moreno, I.; Lao, J.; Loqué, D.; Orellana, A.; Heazlewood, J.L.; Clark, G.; Roux, S.J. AtAPY1 and AtAPY2 function as Golgi-localized nucleoside diphosphatases in *Arabidopsis thaliana*. *Plant Cell Physiol.* **2012**, *53*, 1913–1925. [[CrossRef](#)] [[PubMed](#)]
23. Schiller, M.; Massalski, C.; Kurth, T.; Steinebrunner, I. The *Arabidopsis* apyrase AtAPY1 is localized in the Golgi instead of the extracellular space. *BMC Plant Biol.* **2012**, *12*, 123. [[CrossRef](#)] [[PubMed](#)]
24. Wu, J.; Steinebrunner, I.; Sun, Y.; Butterfield, T.; Torres, J.; Arnold, D.; Gonzalez, A.; Jacob, F.; Reichler, S.; Roux, S.J. Apyrases (nucleoside triphosphate-diphosphohydrolases) play a key role in growth control in *Arabidopsis*. *Plant Physiol.* **2007**, *144*, 961–975. [[CrossRef](#)] [[PubMed](#)]

25. Liu, X.; Wu, J.; Clark, G.; Lundy, S.; Lim, M.; Arnold, D.; Chan, J.; Tang, W.Q.; Muday, G.K.; Gardner, G.; et al. Role for apyrases in polar auxin transport in *Arabidopsis*. *Plant Physiol.* **2012**, *160*, 1985–1995. [[CrossRef](#)] [[PubMed](#)]
26. Day, R.B.; McAlvin, C.B.; Loh, J.T.; Denny, R.L.; Wood, T.C.; Young, N.D.; Stacey, G. Differential expression of two soybean apyrases, one of which is an early nodulin. *Mol. Plant Microbe Interact.* **2000**, *13*, 1053–1070. [[CrossRef](#)] [[PubMed](#)]
27. Kiba, A.; Toyoda, K.; Yoshioka, K.; Tsujimura, K.; Takahashi, H.; Ichinose, Y.; Takeda, T.; Kato, T.; Shiraishi, T. A pea NTPase, PsAPY1, recognizes signal molecules from microorganisms. *J. Gen. Plant Pathol.* **2006**, *72*, 238–246. [[CrossRef](#)]
28. Riewe, D.; Grosman, L.; Fernie, A.R.; Wucke, C.; Geigenberger, P. The potato-specific apyrase is apoplastically localized and has influence on gene expression, growth, and development. *Plant Physiol.* **2008**, *147*, 1092–1109. [[CrossRef](#)]
29. Chiu, T.Y.; Lao, J.; Manalansan, B.; Loque, D.; Roux, S.J.; Heazlewood, J.L. Biochemical characterization of *Arabidopsis* APYRASE family reveals their roles in regulating endomembrane NDP/NMP homeostasis. *Biochem. J.* **2015**, *472*, 43–54. [[CrossRef](#)]
30. Burnstock, G. Purinergic receptors. *J. Theor. Biol.* **1976**, *62*, 491–503. [[CrossRef](#)]
31. Burnstock, G. Purine and pyrimidine receptors. *Cell Mol. Life Sci.* **2007**, *64*, 1471–1483. [[CrossRef](#)]
32. Burnstock, G. A basis for distinguishing two types of purinergic receptor. In *Cell Membrane Receptors for Drugs and Hormones: A Multidisciplinary Approach*; Bolis, L., Straub, R.C.O., Eds.; Raven Press: New York, NY, USA, 1978; pp. 107–118.
33. Möhlmann, T.; Steinebrunner, I.; Neuhaus, E. Nucleotides and nucleosides: Transport, metabolism, and signaling function of extracellular ATP. In *Progress in Botany*; Lüttge, U., Beyschlag, W., Cushman, J., Eds.; Springer: Berlin/Heidelberg, Germany, 2014; Volume 75, pp. 119–144.
34. Tanaka, K.; Gilroy, S.; Jones, A.M.; Stacey, G. Extracellular ATP signaling in plants. *Trends Cell Biol.* **2010**, *20*, 601–608. [[CrossRef](#)] [[PubMed](#)]
35. Jewell, J.B.; Tanaka, K. Transcriptomic perspective on extracellular ATP signaling: A few curious trifles. *Plant Signal. Behav.* **2019**, *14*, 1659079. [[CrossRef](#)] [[PubMed](#)]
36. Liu, T.; Shang, Z. Extracellular ATP: An essential apoplastic messenger in plants. In *Progress in Botany*; Cánovas, F.M., Lüttge, U., Matyssek, R., Eds.; Springer: Cham, Switzerland, 2016; pp. 121–144.
37. Cho, S.H.; Nguyen, C.T.; Choi, J.; Stacey, G. Molecular mechanism of plant recognition of extracellular ATP. *Adv. Exp. Med. Biol.* **2017**, *1051*, 233–253.
38. Jewell, J.B.; Sowders, J.M.; He, R.; Willis, M.A.; Gang, D.R.; Tanaka, K. Extracellular ATP shapes a defense-related transcriptome both independently and along with other defense signaling pathways. *Plant Physiol.* **2019**, *179*, 1144–1158. [[CrossRef](#)] [[PubMed](#)]
39. Zhu, R.J.; Dong, X.X.; Hao, W.W.; Gao, W.; Zhang, W.Z.; Xia, S.Y.; Liu, T.; Shang, Z.L. Heterotrimeric G protein-regulated Ca²⁺ influx and PIN2 asymmetric distribution are involved in *Arabidopsis thaliana* roots' avoidance response to extracellular ATP. *Front. Plant Sci.* **2017**, *8*, 1522. [[CrossRef](#)] [[PubMed](#)]
40. Matthus, E.; Sun, J.; Wang, L.; Bhat, M.G.; Mohammad-Sidik, A.B.; Wilkins, K.A.; Leblanc-Fournier, N.; Legué, V.; Moulia, B.; Stacey, G.; et al. DORN1/P2K1 and purino-calcium signalling in plants: Making waves with extracellular ATP. *Ann. Bot.* **2020**, *124*, 1227–1242. [[CrossRef](#)] [[PubMed](#)]
41. Tanaka, K.; Choi, J.; Cao, Y.; Stacey, G. Extracellular ATP acts as a damage-associated molecular pattern (DAMP) signal in plants. *Front. Plant Sci.* **2014**, *5*, 446. [[CrossRef](#)]
42. Demidchik, V.; Nichols, C.; Oliynyk, M.; Dark, A.; Glover, B.J.; Davies, J.M. Is ATP a signaling agent in plants? *Plant Physiol.* **2003**, *133*, 456–461. [[CrossRef](#)]
43. Wu, S.; Wu, J. Extracellular ATP-induced NO production and its dependence on membrane Ca²⁺ flux in *Salvia miltiorrhiza* hairy roots. *J. Exp. Bot.* **2008**, *59*, 4007–4016. [[CrossRef](#)]
44. Rodrigues, O.; Reshetnyak, G.; Grondin, A.; Saijo, Y.; Leonhardt, N.; Maurel, C.; Verdoucq, L. Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 9200–9205. [[CrossRef](#)] [[PubMed](#)]
45. Sun, J.; Zhang, C.L.; Deng, S.R.; Lu, C.F.; Shen, X.; Zhou, X.Y.; Zheng, X.J.; Hu, Z.M.; Chen, S.L. An ATP signalling pathway in plant cells: Extracellular ATP triggers programmed cell death in *Populus euphratica*. *Plant Cell Environ.* **2012**, *35*, 893–916. [[CrossRef](#)] [[PubMed](#)]

46. Yang, Y.X.; Ahammed, G.J.; Wu, C.; Fan, S.Y.; Zhou, Y.H. Crosstalk among jasmonate, salicylate and ethylene signaling pathways in plant disease and immune responses. *Curr. Protein Pept. Sci.* **2015**, *16*, 450–461. [[CrossRef](#)] [[PubMed](#)]
47. Bouwmeester, K.; de Sain, M.; Weide, R.; Gouget, A.; Klamer, S.; Canut, H.; Govers, F. The lectin receptor kinase LecRK-I.9 is a novel *Phytophthora* resistance component and a potential host target for a RXLR effector. *PLoS Pathog.* **2011**, *7*, e1001327. [[CrossRef](#)] [[PubMed](#)]
48. Balague, C.; Gouget, A.; Bouchez, O.; Souriac, C.; Haget, N.; Boutet-Mercey, S.; Govers, F.; Roby, D.; Canut, H. The *Arabidopsis thaliana* lectin receptor kinase LecRK-I.9 is required for full resistance to *Pseudomonas syringae* and affects jasmonate signalling. *Mol. Plant Pathol.* **2017**, *18*, 937–948. [[CrossRef](#)] [[PubMed](#)]
49. Vahabi, K.; Sherameti, I.; Bakshi, M.; Mrozinska, A.; Ludwig, A.; Reichelt, M.; Oelmüller, R. The interaction of *Arabidopsis* with *Piriformospora indica* shifts from initial transient stress induced by fungus-released chemical mediators to a mutualistic interaction after physical contact of the two symbionts. *BMC Plant Biol.* **2015**, *15*, 58. [[CrossRef](#)]
50. Nizam, S.; Qiang, X.; Wawra, S.; Nostadt, R.; Getzke, F.; Schwanke, F.; Dreyer, I.; Langen, G.; Zuccaro, A. *Serendipita indica* E5'NT modulates extracellular nucleotide levels in the plant apoplast and affects fungal colonisation. *EMBO Rep.* **2019**, *20*, e47430. [[CrossRef](#)]
51. Tanaka, K.; Tóth, K.; Stacey, G. Role of ectoapyrases in nodulation. In *Biological Nitrogen Fixation*, 1st ed.; de Bruijn Frans, J., Ed.; John Wiley & Sons, Inc.: New York, NY, USA, 2015; pp. 517–524.
52. Clark, G.; Fraley, D.; Steinebrunner, I.; Cervantes, A.; Onyirimba, J.; Liu, A.; Torres, J.; Tang, W.; Kim, J.; Roux, S.J. Extracellular nucleotides and apyrases regulate stomatal aperture in *Arabidopsis*. *Plant Physiol.* **2011**, *156*, 1740–1753. [[CrossRef](#)]
53. Jeter, C.R.; Tang, W.; Henaff, E.; Butterfield, T.; Roux, S.J. Evidence of a novel cell signaling role for extracellular adenosine triphosphates and diphosphates in *Arabidopsis*. *Plant Cell* **2004**, *16*, 2652–2664. [[CrossRef](#)]
54. Hou, Q.Z.; Ye, G.J.; Wang, R.F.; Jia, L.Y.; Liang, J.Y.; Feng, H.Q.; Wen, J.; Shi, D.L.; Wang, Q.W. Changes by cadmium stress in lipid peroxidation and activities of lipoxigenase and antioxidant enzymes in *Arabidopsis* are associated with extracellular ATP. *Biologia* **2017**, *72*, 1467–1474. [[CrossRef](#)]
55. Chen, D.; Cao, Y.; Li, H.; Kim, D.; Ahsan, N.; Thelen, J.; Stacey, G. Extracellular ATP elicits DORN1-mediated RBOHD phosphorylation to regulate stomatal aperture. *Nat. Commun.* **2017**, *8*, 2265. [[CrossRef](#)] [[PubMed](#)]
56. Kim, S.H.; Yang, S.H.; Kim, T.J.; Han, J.S.; Suh, J.W. Hypertonic stress increased extracellular ATP levels and the expression of stress-responsive genes in *Arabidopsis thaliana* seedlings. *Biosci. Biotechnol. Biochem.* **2009**, *73*, 1252–1256. [[CrossRef](#)] [[PubMed](#)]
57. Wasternack, C.; Strnad, M. Jasmonates: News on occurrence, biosynthesis, metabolism and action of an ancient group of signaling compounds. *Int. J. Mol. Sci.* **2018**, *19*, 2539. [[CrossRef](#)] [[PubMed](#)]
58. Rizwan, M.; Ali, S.; Adrees, M.; Ibrahim, M.; Tsang, D.; Zia-ur-Rehman, M.; Zahir, Z.; Rinklebe, J.; Tack, F.; Ok, Y. A critical review on effects, tolerance mechanisms and management of cadmium in vegetables. *Chemosphere* **2017**, *182*, 90–105. [[CrossRef](#)]
59. Fargašová, A.; Pastierová, J.; Svetková, K. Effect of Se-metal pair combinations (Cd, Zn, Cu, Pb) on photosynthetic pigments production and metal accumulation in *Sinapis alba* L. seedlings. *Plant Soil Environ.* **2006**, *52*, 8–15. [[CrossRef](#)]
60. Sarry, J.E.; Kuhn, L.; Ducruix, C.; Lafaye, A. The early responses of *Arabidopsis thaliana* cells to cadmium exposure explored by protein and metabolite profiling analyses. *Proteomics* **2006**, *6*, 2180–2198. [[CrossRef](#)]
61. Herbette, S.; Taconnat, L.; Hugouvieux, V.; Piette, L.; Magniette, M.L.M.; Cuine, S. Genome-wide transcriptome profiling of the early cadmium response of *Arabidopsis* roots and shoots. *Biochimie* **2006**, *88*, 1751–1765. [[CrossRef](#)]
62. Sanità di Toppi, L.; Gabbriellini, R. Response to cadmium in higher plants. *Environ. Exp. Bot.* **1999**, *41*, 105–130. [[CrossRef](#)]
63. Berger, F.; Ramirez-Hernandez, M.H.; Ziegler, M. The new life of a centenarian: Signalling functions of NAD(P). *Trends Biochem. Sci.* **2004**, *29*, 111–118. [[CrossRef](#)]
64. Noctor, G. Metabolic signalling in defence and stress: The central roles of soluble redox couples. *Plant Cell Environ.* **2006**, *29*, 409–425. [[CrossRef](#)]
65. Billington, R.A.; Bruzzone, S.; De Flora, A.; Genazzani, A.A.; Koch-Nolte, F.; Ziegler, M.; Zocchi, E. Emerging functions of extracellular pyridine nucleotides. *Mol. Med.* **2006**, *12*, 324–327. [[CrossRef](#)]
66. Zhang, X.; Mou, Z. Extracellular pyridine nucleotides induce *PR* gene expression and disease resistance in *Arabidopsis*. *Plant J.* **2009**, *57*, 302–312. [[CrossRef](#)] [[PubMed](#)]

67. Zhang, X.; Mou, Z. Expression of the human NAD(P)-metabolizing ectoenzyme CD38 compromises systemic acquired resistance in *Arabidopsis*. *Mol Plant Microbe Interact.* **2012**, *25*, 1209–1218. [[CrossRef](#)]
68. Wang, C.; Zhou, M.; Zhang, X.; Yao, J.; Zhang, Y.; Mou, Z. A lectin receptor kinase as a potential sensor for extracellular nicotinamide adenine dinucleotide in *Arabidopsis thaliana*. *eLife* **2017**, *6*, e25474. [[CrossRef](#)]
69. Wang, C.; Huang, X.; Li, Q.; Zhang, Y.; Li, J.L.; Mou, Z. Extracellular pyridine nucleotides trigger plant systemic immunity through a lectin receptor kinase/BAK1 complex. *Nat. Commun.* **2019**, *10*, 4810. [[CrossRef](#)]
70. Marrian, D.H. A new adenine nucleotide. *Biochim. Biophys. Acta* **1953**, *12*, 492. [[CrossRef](#)]
71. Marrian, D.H. A new adenine nucleotide. *Biochim. Biophys. Acta* **1954**, *13*, 278–281. [[CrossRef](#)]
72. Lieberman, I. Identification of adenosine tetraphosphate from horse muscle. *J. Am. Chem. Soc.* **1955**, *77*, 3373–3375. [[CrossRef](#)]
73. Sacks, J. Adenosine pentaphosphate from commercial ATP. *Biochim. Biophys. Acta* **1955**, *16*, 436. [[CrossRef](#)]
74. Gardner, J.A.A.; Hoagland, M.B. The isolation of guanosine tetraphosphate from commercially available preparations of guanosine triphosphate. *J. Biol. Chem.* **1965**, *240*, 1244–1246.
75. Moreno, A.; Vallejo, C.G.; Sillero, A.; Sillero, M.A.G. Wide occurrence of guanosine 5'-tetraphosphate in commercial preparations of GTP. *Anal. Biochem.* **1975**, *68*, 648–650. [[CrossRef](#)]
76. Moreno, A.; Lobatón, C.D.; Günther Sillero, M.A.; Sillero, A. Dinucleosidetetraphosphatase from Ehrlich ascites tumourcells: Inhibition by adenosine, guanosine and uridine 5'-tetraphosphates. *Int. J. Biochem.* **1982**, *14*, 629–634. [[CrossRef](#)]
77. Costas, M.J.; Cameselle, J.C.; Günther Sillero, M.A.; Sillero, A. Presence of cytidine 5'-tetraphosphate in commercial samples of cytidine 5'-triphosphate. *Anal. Biochem.* **1983**, *134*, 455–458. [[CrossRef](#)]
78. Heldt, H.W.; Klingenberg, M. Endogenous nucleotides of mitochondria participating in phosphate transfer reactions as studied with ³²P labelled orthophosphate and ultramicroscale ion exchange chromatography. *Biochem. Z.* **1965**, *343*, 433–451. [[PubMed](#)]
79. Zamecnik, P.C.; Stephenson, M.L. A possible regulatory site located at the gateway to protein synthesis. In *Regulatory Mechanisms for Protein Synthesis in Mammalian Cells*; Lamborg, M.R., Kenney, P.T., Eds.; Academic Press Inc.: San Pietro, NY, USA, 1968; pp. 3–16.
80. Small, G.; Cooper, C. Studies on the occurrence and biosynthesis of adenosine tetraphosphate. *Biochemistry* **1966**, *5*, 26–33. [[CrossRef](#)]
81. Van Dyke, K.; Robinson, R.; Urquilla, P.; Smith, D.; Taylor, M.; Trush, M.; Wilson, M. An analysis of nucleotides and catecholamines in bovine medullary granules by anion exchange high pressure liquid chromatography and fluorescence. Evidence that most of catecholamines in chromaffin granules are stored without associated ATP. *Pharmacology* **1977**, *15*, 377–391. [[CrossRef](#)]
82. Sillero, M.A.; Del Valle, M.; Zaera, E.; Michelena, P.; García, A.G.; Sillero, A. Diadenosine 5',5'''-P₁,P₄-tetraphosphate (Ap₄A), ATP and catecholamine content in bovine adrenal medulla, chromaffin granules and chromaffin cells. *Biochimie* **1994**, *76*, 404–409. [[CrossRef](#)]
83. Gualix, J.; Abal, M.; Pintor, J.; Miras-Portugal, M.T. Presence of ε-adenosine tetraphosphate in chromaffin granules after transport of ε-ATP. *FEBS Lett.* **1996**, *391*, 195–198. [[CrossRef](#)]
84. Lee, J.W.; Jeon, S.J.; Kong, I.D.; Jeong, S.W. Identification of adenosine 5'-tetraphosphate in rabbit platelets and its metabolism in blood. *Korean J. Physiol.* **1995**, *29*, 217–223.
85. Jakubowski, H. Sporulation of the yeast *Saccharomyces cerevisiae* is accompanied by synthesis of adenosine 5'-tetraphosphate and adenosine 5'-pentaphosphate. *Proc. Natl. Acad. Sci. USA* **1986**, *83*, 2378–2382. [[CrossRef](#)]
86. Fraga, H.; Fontes, R. Enzymatic synthesis of mono and dinucleoside polyphosphates. *Biochim. Biophys. Acta* **2011**, *1810*, 1195–1204. [[CrossRef](#)] [[PubMed](#)]
87. Zamecnik, P.C.; Stephenson, M.L. Nucleoside pyrophosphate compounds related to the first step in protein synthesis. In *Alfred Benzon Symposium I: The Role of Nucleotides for the Function and Conformation of Enzymes*; Kalckar, H.M., Klenow, H., Munch-Peterson, G., Ottesen, M., Thuysen, J.M., Eds.; Munksgaard: Copenhagen, Denmark, 1969; pp. 276–291.
88. Theoclitou, M.E.; El-Thaher, T.S.H.; Miller, A.D. Enzymatic synthesis of diadenosine 5',5'''-P₁,P₄-tetraphosphate (Ap₄A) analogues by stress protein LysU. *J. Chem. Soc. Chem. Commun.* **1994**, *5*, 659–661. [[CrossRef](#)]
89. Zamecnik, P.C.; Stephenson, M.L.; Janeway, C.M.; Randerath, K. Enzymatic synthesis of diadenosine tetraphosphate and diadenosine triphosphate with a purified lysyl-sRNA synthetase. *Biochem. Biophys. Res. Commun.* **1966**, *24*, 91–97. [[CrossRef](#)]

90. Guranowski, A. Studies on dinucleoside polyphosphates: Some intriguing biochemical, physiological, and medical aspects. *J. Clin. Biochem.* **2000**, *28*, 177–189. [[CrossRef](#)]
91. McLennan, A.G.; Zamecnik, P.C. Dinucleoside polyphosphates—An introduction. In *Ap₄A and Other Dinucleoside Polyphosphates*; McLennan, A.G., Ed.; CRC Press: Boca Raton, FL, USA, 1992; pp. 1–7.
92. Goerlich, O.; Foeckler, R.; Holler, E. Mechanism of synthesis of adenosine(5′)tetraphospho(5′)adenosine (AppppA) by aminoacyl-tRNA synthetases. *Eur. J. Biochem.* **1982**, *126*, 135–142. [[CrossRef](#)]
93. Blanquet, S.; Plateau, P.; Brevet, A. The role of zinc in 5′,5′-diadenosine tetraphosphate production by aminoacyl-transfer RNA synthetases. *Mol. Cell. Biochem.* **1983**, *52*, 3–11. [[CrossRef](#)]
94. Biryukov, A.I.; Zhukov, Y.N.; Lavrik, O.I.; Khomutov, R.M. Influence of the aminoacyl-tRNA synthetase inhibitors and the diadenosine-5′-tetraphosphate phosphonate analogues on the catalysis of diadenosyl oligophosphates formation. *FEBS Lett.* **1990**, *273*, 208–210. [[CrossRef](#)]
95. Charlier, J.; Sanchez, R. Lysyl-tRNA synthetase from *Escherichia coli* K12. Chromatographic heterogeneity and the lysU-gene product. *Biochem. J.* **1987**, *248*, 43–51. [[CrossRef](#)]
96. Lazewska, D.; Guranowski, A. P alpha-chiral phosphorothioate analogues of bis (5′-adenosyl)tetraphosphate (Ap₄A); their enzymatic synthesis and degradation. *Nucleic Acid Res.* **1990**, *18*, 6083–6088. [[CrossRef](#)]
97. Brevet, A.; Chen, J.; Lévêque, F.; Blanquet, S.; Plateau, P. Comparison of the enzymatic properties of the two *Escherichia coli* Lysyl-tRNA synthetase species. *J. Biol. Chem.* **1995**, *270*, 14439–14444. [[CrossRef](#)]
98. Wright, M.; Boonyalai, N.; Tanner, J.A.; Hindley, A.D.; Miller, A.D. The duality of LysU, a catalyst for both Ap₄A and Ap₃A formation. *FEBS J.* **2006**, *273*, 3534–3544. [[CrossRef](#)] [[PubMed](#)]
99. Ortiz, B.; Sillero, A.; Sillero, M.A.G. Specific synthesis of adenosine(5′)tetraphospho(5′)nucleoside and adenosine(5′)oligophospho(5′)adenosine (n greater than 4) catalyzed by firefly luciferase. *Eur. J. Biochem.* **1993**, *215*, 263–270. [[CrossRef](#)] [[PubMed](#)]
100. Guranowski, A.; de Diego, A.; Sillero, A.; Sillero, M.A.G. Uridine 5′-polyphosphates (p₄U and p₅U) and uridine(5′)polyphospho(5′)nucleosides (Up_nNs) can be synthesized by UTP:glucose-1-phosphate uridylyltransferase from *Saccharomyces cerevisiae*. *FEBS Lett.* **2004**, *561*, 83–88. [[CrossRef](#)]
101. García-Díaz, M.; Canales, J.; Günther Sillero, M.A.; Sillero, A.; Cameselle, J.C. Phosphoglycerate kinase from yeast synthesizes guanosine 5′-tetraphosphate. *Biochem. Int.* **1989**, *19*, 1253–1264.
102. Kupriyanov, V.V.; Ferretti, J.A.; Balaban, R.S. Muscle adenylate kinase catalyzes adenosine 5′-tetraphosphate synthesis from ATP and ADP. *Biochim. Biophys. Acta* **1986**, *869*, 107–111. [[CrossRef](#)]
103. Luo, G.; Nishimura, J.S. Adenosine 5′-tetraphosphate is synthesized by the histidine a142—asparagine mutant of *Escherichia coli* succinyl-CoA synthetase. *J. Biol. Chem.* **1992**, *267*, 9516–9520.
104. Guranowski, A.; Günther Sillero, M.A.; Sillero, A. Adenosine 5′-tetraphosphate and adenosine 5′-pentaphosphate are synthesized by yeast acetyl coenzyme A synthetase. *J. Bacteriol.* **1994**, *176*, 2986–2990. [[CrossRef](#)]
105. Sillero, M.A.; de Diego, A.; Pérez-Zúñiga, F.; Sillero, A. Synthesis of biphosphonate derivatives of ATP by T4 DNA ligase, ubiquitin activating enzyme (E1) and other ligases. *Biochem. Pharmacol.* **2008**, *75*, 1959–1965. [[CrossRef](#)]
106. Fontes, R.; Sillero, M.A.; Sillero, A. Acyl coenzyme A synthetase from *Pseudomonas fragi* catalyzes the synthesis of adenosine 5′-polyphosphates and dinucleoside polyphosphates. *J. Bacteriol.* **1998**, *180*, 3152–3158. [[CrossRef](#)]
107. Fontes, R.; Günther Sillero, M.A.; Sillero, A. Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap₆A). *Biochimie* **1999**, *81*, 229–233. [[CrossRef](#)]
108. Madrid, O.; Martín, D.; Atencia, E.A.; Sillero, A.; Sillero, M.A.G. T4 DNA ligase synthesizes dinucleoside polyphosphates. *FEBS Lett.* **1998**, *433*, 283–286. [[CrossRef](#)]
109. Günther Sillero, M.A.; Montes, M.; de Diego, A.; del Valle, M.; Atencia, E.; Sillero, A. Thermostable *Pyrococcus furiosus* DNA ligase catalyzes the synthesis of (di)nucleoside polyphosphates. *Extremophiles* **2002**, *6*, 45–50. [[CrossRef](#)] [[PubMed](#)]
110. Sillero, M.A.; de Diego, A.; Tavares, J.E.F.; Silva, C.; Pérés-Zúñiga, J.; Sillero, A. Synthesis of ATP derivatives of compounds of the mevalonate pathway (isopentenyl di- and triphosphate; geranyl di- and triphosphate, farnesyl di- and triphosphate, and dimethylallyl diphosphate) catalyzed by T4 RNA ligase, T4 DNA ligase and other ligases potential relationship with the effect of bisphosphonates on osteoclasts. *Biochem. Pharmacol.* **2009**, *78*, 335–343. [[PubMed](#)]
111. Atencia, E.A.; Madrid, O.; Sillero, M.A.G.; Sillero, A. T4 RNA ligase catalyzes the synthesis of dinucleoside polyphosphates. *Eur. J. Biochem.* **1999**, *261*, 802–811. [[CrossRef](#)] [[PubMed](#)]

112. Bouhss, A.; Dementin, S.; van Heijenoort, J.; Parquet, C.; Blanot, D. Formation of adenosine 5'-tetrphosphate from the acyl phosphate intermediate: A difference between the MurC and MurD synthetases of *Escherichia coli*. *FEBS Lett.* **1999**, *453*, 15–19. [[CrossRef](#)]
113. Guranowski, A.; Starzyńska, E.; Rataj-Guranowska, M. Purification of apyrase from yellow lupin cotyledons after extraction with perchloric acid. *Protein. Expr. Purif.* **1991**, *4*, 235–239. [[CrossRef](#)]
114. Guranowski, A.; Starzyńska, E.; Sillero, M.A.; Sillero, A. Conversion of adenosine(5') oligophospho(5') adenosines into inosine(5')oligophospho(5')inosines by non-specific adenylate deaminase from the snail *Helix pomatia*. *Biochim. Biophys. Acta* **1995**, *1243*, 78–84. [[CrossRef](#)]
115. Mateo, J.; Miras-Portugal, M.T.; Rotllán, P. Ecto-enzymatic hydrolysis of diadenosine polyphosphates by cultured adrenomedullary vascular endothelial cells. *Am. J. Physiol.* **1997**, *273*, 6918–6927. [[CrossRef](#)]
116. Mateo, J.; Rotllán, P.; Marti, E.; Gomez de Aranda, I.; Solsona, C.; Miras-Portugal, M.T. Diadenosine polyphosphate hydrolase from presynaptic plasma membranes of Torpedo electric organ. *Biochem. J.* **1997**, *323*, 677–684. [[CrossRef](#)]
117. Plateau, P.; Fromant, M.; Brevet, A.; Gesquière, A.; Blanquet, S. Catabolism of bis(5'-nucleosidyl) oligophosphates in *Escherichia coli*; metal requirements and substrate specificity of homogeneous diadenosine-5',5'''-tetrphosphate (symmetrical) pyrophosphohydrolase. *Biochemistry* **1985**, *24*, 914–922. [[CrossRef](#)]
118. Guranowski, A.; Blanquet, S. Phosphorolytic cleavage of diadenosine 5',5'''-P1,P4-tetrphosphate; properties of homogeneous diadenosine 5',5'''-P1,P4-tetrphosphate α,β -phosphorylase from *Saccharomyces cerevisiae*. *J. Biol. Chem.* **1985**, *260*, 3542–3547. [[PubMed](#)]
119. Guranowski, A.; Just, G.; Holler, E.; Jakubowski, H. Synthesis of diadenosine 5', 5'''-P1,P4-tetrphosphate (AppppA) from adenosine 5'- phosphosulfate and adenosine 5'-triphosphate catalyzed by yeast AppppA phosphorylase. *Biochemistry* **1988**, *27*, 2959–2964. [[CrossRef](#)]
120. Guranowski, A.; Starzyńska, E.; Brown, P.; Blackburn, G.M. Adenosine 5'-tetrphosphate phosphohydrolase from yellow lupin seeds: Purification to homogeneity and some properties. *Biochem. J.* **1997**, *328*, 257–262. [[CrossRef](#)]
121. Pietrowska-Borek, M.; Stuble, H.P.; Kombrink, E.; Guranowski, A. 4-Coumarate:coenzyme A ligase has the catalytic capacity to synthesize and reuse various (di)adenosine polyphosphates. *Plant Physiol.* **2003**, *131*, 1401–1410. [[CrossRef](#)] [[PubMed](#)]
122. Dixon, R.A.; Paiva, N.L. Stress-induced phenylpropanoid metabolism. *Plant Cell.* **1995**, *7*, 1085–1097. [[CrossRef](#)] [[PubMed](#)]
123. Guranowski, A.; Miersch, O.; Staswick, P.E.; Suza, W.; Wasternack, C. Substrate specificity and products of side-reactions catalyzed by jasmonate:amino acid synthetase (JAR1). *FEBS Lett.* **2007**, *581*, 815–820. [[CrossRef](#)] [[PubMed](#)]
124. Staswick, P.E.; Tiryaki, I.; Rowe, M.L. Jasmonate response locus JAR1 and several related *Arabidopsis* genes encode enzymes of the firefly luciferase superfamily that show activity on jasmonic, salicylic, and indole 3-acetic acids in an assay for adenylation. *Plant Cell* **2002**, *14*, 1405–1415. [[CrossRef](#)]
125. Frankhauser, H.; Berkovitz, G.A.; Schiff, J.A. A nucleotide with the properties of adenosine 5'-phosphoramidate from *Chlorella* cells. *Biochem. Biophys. Res. Commun.* **1981**, *101*, 524–532. [[CrossRef](#)]
126. Frankhauser, H.; Schiff, J.A.; Garber, L.J. Purification and properties of adenylyl sulfate: Ammonia adenylyltransferase from *Chlorella* catalyzing the formation of adenosine 5'-phosphoramidate from adenosine 5'-phosphosulfate and ammonia. *Biochem. J.* **1981**, *195*, 545–560. [[CrossRef](#)]
127. Wojdyła-Mamoń, A.M.; Guranowski, A. Adenylylsulfate:ammonia adenylyltransferase activity is another inherent property of Fhit proteins. *Bioscience Rep.* **2015**, *35*, e00235. [[CrossRef](#)]
128. Kuba, M.; Okizaki, T.; Ohmori, H.; Kumon, A. Nucleoside monophosphoramidate hydrolase from rat liver: Purification and characterization. *Int. J. Biochem. Cell Biol.* **1994**, *26*, 235–245.
129. Bieganski, P.; Garrison, P.N.; Hodawadekar, S.C.; Faye, G.; Barnes, L.D.; Brenner, C. Adenosine monophosphoramidase activity of Hint and Hnt1 supports function of Kin28, Ccl1, and Tfb3. *J. Biol. Chem.* **2002**, *13*, 10852–10860. [[CrossRef](#)] [[PubMed](#)]
130. Guranowski, A.; Wojdyła, A.M.; Zimny, J.; Wypijewska, A.; Kowalska, J.; Jemielity, J.; Davis, R.E.; Bieganski, P. Dual activity of certain HIT-proteins: *A. thaliana* Hint4 and *C. elegans* DcpS act on adenosine 5'-phosphosulfate as hydrolases (forming AMP) and as phosphorylases (forming ADP). *FEBS Lett.* **2010**, *584*, 93–98. [[CrossRef](#)] [[PubMed](#)]

131. Guranowski, A.; Wojdyła, A.M.; Zimny, J.; Wypijewska, A.; Kowalska, J.; Łukaszewicz, M.; Jemielity, J.; Darzynkiewicz, E.; Jagiełło, A.; Bieganowski, P. Recognition of different nucleotidyl-derivatives as substrates of re-actions catalyzed by various HIT-proteins. *New J. Chem.* **2010**, *34*, 888–893. [[CrossRef](#)]
132. Guranowski, A.; Wojdyła, A.M.; Rydzik, A.M.; Stępiński, J.; Jemielity, J. Plant nucleoside 5'-phosphoramidate hydrolase; simple purification from yellow lupin (*Lupinus luteus*) seeds and properties of homogeneous enzyme. *Acta Biochim. Pol.* **2011**, *58*, 131–136. [[CrossRef](#)]
133. Bretes, E.; Wojdyła-Mamoń, A.M.; Kowalska, J.; Jemielity, J.; Kaczmarek, R.; Baraniak, J.; Guranowski, A. Hint2, the mitochondrial nucleoside 5'-phosphoramidate hydrolase; properties of the homogeneous protein from sheep (*Ovis aries*) liver. *Acta Biochim. Pol.* **2013**, *60*, 249–254. [[CrossRef](#)]
134. Huang, K.; Arabshahi, A.; Wei, Y.; Frey, P.A. The mechanism of action of the fragile histidine triad, Fhit: Isolation of a covalent adenylyl enzyme and chemical rescue of H96G-Fhit. *Biochemistry* **2004**, *43*, 7637–7642. [[CrossRef](#)]
135. Brenner, C. Hint, Fhit, and GalT: Function, structure, evolution, and mechanism of three branches of the histidine triad superfamily of nucleotide hydrolases and transferases. *Biochemistry* **2002**, *41*, 9003–9014. [[CrossRef](#)]
136. Guranowski, A.; Wojdyła, A.M.; Pietrowska-Borek, M.; Bieganowski, P.; Khurs, E.N.; Cliff, M.J.; Blackburn, G.M.; Błaziak, D.; Stec, W.J. Fhit proteins can also recognize substrates other than dinucleoside polyphosphates. *FEBS Lett.* **2008**, *582*, 3152–3158. [[CrossRef](#)]
137. Christie, S.M.H.; Elmore, D.T.; Kenner, G.W.; Todd, A.R.; Weymouth, F.J. Nucleotides, Part XXII. Syntheses of P¹,P²-diadenosine-5' and P¹,P²-diuridine-5'-pyrophosphates. *J. Chem. Soc.* **1953**, *3*, 2947–2953. [[CrossRef](#)]
138. Finamore, F.J.; Warner, A.H. The occurrence of P¹,P⁴-diguanosine 5'-tetraphosphate in brine shrimp eggs. *J. Biol. Chem.* **1963**, *238*, 344–348. [[PubMed](#)]
139. Warner, A.H.; Finamore, F.J. Isolation, purification, and characterization of P¹,P⁴-diguanosine 5'-tetraphosphate *asymmetrical*-pyrophosphohydrolase from brine shrimp eggs. *Biochemistry* **1965**, *4*, 1568–1575. [[CrossRef](#)] [[PubMed](#)]
140. Garrison, P.N.; Barnes, L.D. Determination of dinucleoside polyphosphates. In *Ap4A and Other Dinucleoside Polyphosphates*; McLennan, A.G., Ed.; CRC Press: Boca Raton, FL, USA, 1992; pp. 29–61.
141. Plateau, P.; Blanquet, S. Dinucleoside oligophosphates in micro-organisms. *Adv. Microb. Physiol.* **1994**, *36*, 81–109. [[PubMed](#)]
142. Lee, P.C.; Barry, R.; Ames, B.N. Diadenosine 5',5'''-P¹,P⁴-tetraphosphate and related adenylylated nucleotides in *Salmonella typhimurium*. *J. Biol. Chem.* **1983**, *258*, 6827–6834. [[PubMed](#)]
143. Coste, H.; Brevet, A.; Plateau, P.; Blanquet, S. Non-adenylylated bis(5'-nucleosidyl) tetraphosphates occur in *Saccharomyces cerevisiae* and in *Escherichia coli* and accumulate upon temperature shift or exposure to cadmium. *J. Biol. Chem.* **1987**, *262*, 12096–12103. [[PubMed](#)]
144. Baltzinger, M.; Ebel, J.P.; Remy, P. Accumulation of dinucleoside polyphosphates in *Saccharomyces cerevisiae* under stress conditions. High levels are associated with cell death. *Biochimie* **1986**, *68*, 1231–1236. [[CrossRef](#)]
145. Palfi, Z.; Suranyi, G.; Borbely, G. Alterations in the accumulation of adenylylated nucleotides in heavy-metal-ion-stressed and heat-stressed *Synechococcus* sp. strain PCC 6301, cyanobacterium, in light and dark. *Biochem. J.* **1991**, *276*, 487–491. [[CrossRef](#)]
146. Lütthje, J.; Olivie, A. The presence of diadenosine 5', 5'''-P¹,P³- triphosphate (Ap₃A) in human platelets. *Biochem. Biophys. Res. Commun.* **1983**, *115*, 253–260. [[CrossRef](#)]
147. Flodgaard, H.; Klenow, H. Abundant amounts of diadenosine 5', 5'''-P¹,P⁴-tetraphosphate are present and releasable, but metabolically inactive, in human platelets. *Biochem. J.* **1982**, *208*, 737–742. [[CrossRef](#)]
148. Morioka, M.; Shimada, H. Synthesis of diadenosine 5', 5'''-P¹,P⁴-tetraphosphate (Ap₄A) in sea urchin embryos. *Cell Differ.* **1984**, *14*, 53–58. [[CrossRef](#)]
149. Miller, D.; McLennan, A.G. Changes in intracellular levels of Ap₃A and Ap₄A in cysts and larvae of *Artemia* do not correlate with changes in protein synthesis after heat-shock. *Nucleic Acids Res.* **1986**, *14*, 6031–6040. [[CrossRef](#)] [[PubMed](#)]
150. Brevet, A.; Plateau, P.; Best-Belpomme, M.; Blanquet, S. Variation of Ap₄A and other dinucleoside polyphosphates in stressed *Drosophila* cells. *J. Biol. Chem.* **1985**, *260*, 15566–15570. [[PubMed](#)]
151. Plateau, P.; Mayaux, J.F.; Blanquet, S. Zinc (II)-dependent synthesis of diadenosine 5',5'''-P¹,P⁴-tetraphosphate by *Escherichia coli* and yeast phenylalanyl transfer ribonucleic acid synthetases. *Biochemistry* **1981**, *20*, 4654–4662. [[CrossRef](#)] [[PubMed](#)]

152. Jakubowski, H. Synthesis of diadenosine 5',5'''-P¹,P⁴-tetrphosphate and related compounds by plant (*Lupinus luteus*) seryl-tRNA and phenylalanyl-tRNA synthetases. *Acta Biochim. Pol.* **1983**, *30*, 51–69.
153. Guranowski, A.; Günther Sillero, M.A.; Sillero, A. Firefly luciferase synthesizes P¹,P⁴-bis(5'-adenosyl) tetrphosphate (Ap₄A) and other dinucleoside polyphosphates. *FEBS Lett.* **1990**, *271*, 215–218. [[CrossRef](#)]
154. Wang, D.; Shatkin, A.J. Synthesis of Gp₄N and Gp₃N compounds by guanylyltransferase purified from yeast. *Nucleic Acids Res.* **1984**, *12*, 2303–2315. [[CrossRef](#)]
155. Jin, Z.; Leveque, V.; Ma, H.; Johnson, K.A.; Klumpp, K. NTP-mediated nucleotide excision activity of hepatitis C virus RNA-dependent RNA polymerase. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 348–357. [[CrossRef](#)]
156. Guranowski, A. Specific and non-specific enzymes involved in the catabolism of mononucleoside and dinucleoside polyphosphates. *Pharma Ther.* **2000**, *87*, 117–139. [[CrossRef](#)]
157. Jakubowski, H.; Guranowski, A. Enzymes hydrolyzing ApppA and/or AppppA in higher plants purification and some properties of diadenosine triphosphatase, diadenosine tetrphosphatase, and phosphodiesterase from yellow lupin (*Lupinus luteus*) seeds. *J. Biol. Chem.* **1983**, *258*, 9982–9989.
158. Guranowski, A. Fluoride is a strong and specific inhibitor of (asymmetrical) Ap₄A hydrolases. *FEBS Lett.* **1990**, *262*, 205–208. [[CrossRef](#)]
159. Feussner, K.; Guranowski, A.; Kostka, S.; Wasternack, C. Diadenosine 5',5'''-P¹,P⁴-tetrphosphate (Ap₄A) hydrolase from tomato (*Lycopersicon esculentum* cv. Lukullus)—Purification, biochemical Ap₄A and related nucleotides in plants properties and behaviour during stress. *Z. Naturforsch. C J. Biosci.* **1996**, *51*, 477–486. [[CrossRef](#)] [[PubMed](#)]
160. Maksel, D.; Guranowski, A.; Ilgoutz, S.C.; Moir, A.; Blackburn, M.G.; Gayler, K.R. Cloning and expression of diadenosine 5',5'''-P¹,P⁴-tetrphosphate hydrolase from *Lupinus angustifolius* L. *Biochem. J.* **1998**, *329*, 313–319. [[CrossRef](#)] [[PubMed](#)]
161. Churin, J.; Hause, B.; Feussner, I.; Maucher, H.P.; Feussner, K.; Börner, T.; Wasternack, C. Cloning and expression of a new cDNA from monocotyledonous plants coding for a diadenosine 5',5'''-P¹,P⁴-tetrphosphate hydrolase from barley (*Hordeum vulgare*). *FEBS Lett.* **1998**, *431*, 481–485. [[CrossRef](#)]
162. Bartkiewicz, M.; Sierakowska, H.; Shugar, D. Nucleotide pyrophosphatase from potato tubers; purification and properties. *Eur. J. Biochem.* **1984**, *143*, 419–426. [[CrossRef](#)] [[PubMed](#)]
163. Zamecnik, P. Diadenosine 5',5'''-P¹,P⁴-tetrphosphate (Ap₄A): Its role in cellular metabolism. *Anal. Biochem.* **1983**, *134*, 1–10. [[CrossRef](#)]
164. McLennan, A.G. Dinucleoside polyphosphates - friend or foe? *Pharmacol. Ther.* **2000**, *87*, 73–89. [[CrossRef](#)]
165. Rapaport, E.; Zamecnik, P.C. Presence of diadenosine 5',5'''-P¹,P⁴-tetrphosphate (Ap₄A) in mamalian cells in levels varying widely with proliferative activity of the tissue: A possible positive “pleiotypic activator”. *Proc. Natl. Acad. Sci. USA* **1976**, *73*, 3984–3988. [[CrossRef](#)]
166. Weinmann-Dorsch, C.; Pierron, G.; Wick, R.; Sauer, H.; Grummt, F. High diadenosine tetrphosphate (Ap₄A) level at initiation of S phase in the naturally synchronous mitotic cycle of *Physarum polycephalum*. *Exp. Cell Res.* **1984**, *155*, 171–177. [[CrossRef](#)]
167. Lee, P.C.; Bochner, B.R.; Ames, B.N. AppppA, heat-shock stress, and cell oxidation. *Proc. Natl. Acad. Sci. USA* **1983**, *80*, 7496–7500. [[CrossRef](#)]
168. Bochner, B.R.; Lee, P.C.; Wilson, S.W.; Cutler, C.W.; Ames, B.N. Ap₄A and related nucleotides are synthesized as consequence of oxidation stress. *Cell* **1984**, *37*, 225–232. [[CrossRef](#)]
169. Hauryliuk, V.; Atkinson, G.C.; Murakami, K.S.; Tenson, T.; Gerdes, K. Recent functional insights into the role of (p)ppGpp in bacterial physiology. *Nat. Rev. Microbiol.* **2015**, *13*, 298–309. [[CrossRef](#)] [[PubMed](#)]
170. Perotti, V.E.; Del Vecchio, H.A.; Sansevich, A.; Meier, G.; Bello, F.; Cocco, M.; Garrán, S.M.; Anderson, C.; Vázquez, D.; Podestá, F.E. Proteomic, metabolomic, and biochemical analysis of heat treated Valencia oranges during storage. *Postharv. Biol. Technol.* **2011**, *62*, 97–114. [[CrossRef](#)]
171. Jambunathan, N.; Penaganti, A.; Tang, Y.; Mahalingam, R. Modulation of redox homeostasis under suboptimal conditions by *Arabidopsis* nudix hydrolase 7. *BMC Plant Biol.* **2010**, *10*, 173. [[CrossRef](#)] [[PubMed](#)]
172. Muthuramalingam, M.; Zeng, X.; Iyer, N.J.; Klein, P.; Mahalingam, R. A GCC-box motif in the promoter of nudix hydrolase 7 (AtNUDT7) gene plays a role in ozone response of *Arabidopsis* ecotypes. *Genomics* **2015**, *105*, 31–38. [[CrossRef](#)]
173. Maruta, T.; Ogawa, T.; Tsujimura, M.; Ikemoto, K.; Yoshida, T.; Takahashi, H.; Yoshimura, K.; Shigeoka, S. Loss-of-function of an *Arabidopsis* NADPH pyrophosphohydrolase, AtNUDX19, impacts on the pyridine nucleotides status and confers photooxidative stress tolerance. *Sci. Rep.* **2016**, *6*, 37432. [[CrossRef](#)]

174. Tanaka, S.; Kihara, M.; Sugimoto, M. Structure and molecular characterization of barley nudix hydrolase genes. *Biosci. Biotechnol. Biochem.* **2015**, *79*, 394–401. [[CrossRef](#)]
175. Xu, J.; Yang, J.Y.; Niu, Q.W.; Chua, N.H. *Arabidopsis* DCP2, DCP1, and VARICOSE form a decapping complex required for postembryonic development. *Plant Cell* **2006**, *18*, 3386–3398. [[CrossRef](#)]
176. Gunawardana, D.; Cheng, H.C.; Gayler, K.R. Identification of functional domains in *Arabidopsis thaliana* mRNA decapping enzyme (AtDcp2). *Nucleic Acids Res.* **2008**, *36*, 203–216. [[CrossRef](#)]
177. Song, M.G.; Bail, S.; Kiledjian, M. Multiple Nudix family proteins possess mRNA decapping activity. *RNA* **2013**, *19*, 390–399.
178. Despotović, D.; Brandis, A.; Savidor, A.; Levin, Y.; Fumagalli, L.; Tawfik, D.S. Diadenosine tetraphosphate (Ap₄A) – an *E. coli* alarmone or a damage metabolite? *FEBS J.* **2017**, *284*, 2194–2215. [[CrossRef](#)]
179. Ji, X.; Zou, J.; Peng, H.; Stolle, A.S.; Xie, R.; Zhang, H.; Peng, B.; Mekalanos, J.J.; Zheng, J. Alarmone Ap₄A is elevated by aminoglycoside antibiotics and enhances their bactericidal activity. *Proc. Nat. Acad. Sci. USA* **2019**, *116*, 9578–9585. [[CrossRef](#)]
180. Lee, Y.-N.; Nechushtan, H.; Figov, N.; Razin, E. The function of lysyl-tRNA synthetase and Ap₄A as signaling regulators of MITF activity in FcεRI-activated mast cells. *Immunity* **2004**, *20*, 145–151. [[CrossRef](#)]
181. Carmi-Levy, I.; Yannay-Cohen, N.; Kay, G.; Razin, E.; Nechushtan, H. Diadenosine tetraphosphate hydrolase is part of the transcriptional regulation network in immunologically activated mast cells. *Mol. Cell. Biol.* **2008**, *28*, 5777–5784. [[CrossRef](#)]
182. Marriott, A.S.; Vasieva, O.; Fang, Y.; Copeland, N.A.; McLennan, A.G.; Jones, N.J. NUDT2 disruption elevates diadenosine tetraphosphate (Ap₄A) and down-regulates immune response and cancer promotion genes. *PLoS ONE* **2016**, *11*, e0154674. [[CrossRef](#)]
183. Bobert, P.; Schlüter, H.; Schultheiss, H.P.; Reusch, H.P. Diadenosine polyphosphates Ap₃A and Ap₄A, but not Ap₅A or Ap₆A, induce proliferation of vascular smooth muscle cells. *Biochem. Pharmacol.* **2008**, *75*, 1966–1973. [[CrossRef](#)]
184. Marriott, A.S.; Copeland, N.A.; Cunningham, R.; Wilkinson, M.C.; McLennan, A.G.; Jones, N.J. Diadenosine 5', 5''-P₁,P₄-tetraphosphate (Ap₄A) is synthesized in response to DNA damage and inhibits the initiation of DNA replication. *DNA Repair.* **2015**, *33*, 90–100. [[CrossRef](#)]
185. Pietrowska-Borek, M.; Nuc, K.; Zielezińska, M.; Guranowski, A. Diadenosine polyphosphates (Ap₃A and Ap₄A) behave as alarmones triggering the synthesis of enzymes of the phenylpropanoid pathway in *Arabidopsis thaliana*. *FEBS Open Bio* **2011**, *1*, 1–6. [[CrossRef](#)]
186. Pietrowska-Borek, M.; Czekala, Ł.; Belchí-Navarro, S.; Pedreño, M.A.; Guranowski, A. Diadenosine triphosphate is a novel factor which in combination with cyclodextrins synergistically enhances the biosynthesis of trans-resveratrol in *Vitis vinifera* cv. Monastrell suspension cultured cells. *Plant Physiol. Biochem.* **2014**, *84*, 271–276. [[CrossRef](#)]
187. Pietrowska-Borek, M.; Nuc, K.; Guranowski, A. Exogenous adenosine 5'-phosphoramidate behaves as a signal molecule in plants; it augments metabolism of phenylpropanoids and salicylic acid in *Arabidopsis thaliana* seedlings. *Plant Physiol. Biochem.* **2015**, *94*, 144–152. [[CrossRef](#)]
188. Pietrowska-Borek, M.; Wojdyła-Mamoń, A.; Dobrogojski, J.; Młynarska-Cieślak, A.; Baranowski, M.R.; Dąbrowski, J.M.; Kowalska, J.; Jemielity, J.; Borek, S.; Pedreño, M.A.; et al. Purine and pyrimidine dinucleoside polyphosphates differentially affect the phenylpropanoid pathway in *Vitis vinifera* L. cv. Monastrell suspension cultured cells. *Plant Physiol. Biochem.* **2020**, *147*, 125–132. [[CrossRef](#)]
189. Ehrling, J.; Büttner, D.; Wang, Q.; Douglas, C.J.; Somssich, I.E.; Kombrink, E. Three 4-coumarate:coenzyme A ligases in *A. thaliana* represent two evolutionarily divergent classes in angiosperms. *Plant J.* **1999**, *19*, 9–20. [[CrossRef](#)] [[PubMed](#)]
190. Bru, R.; Sellés, S.; Casado-Vela, J.; Belchí-Navarro, S.; Pedreño, M.A. Modified cyclodextrins are chemically defined glucan inducers of defense responses in grapevine cell cultures. *J. Agric. Food Chem.* **2006**, *54*, 66–71. [[CrossRef](#)]
191. Winkel-Shirley, B. Biosynthesis of flavonoids and effects of stress. *Curr. Opin. Plant Biol.* **2002**, *5*, 218–223. [[CrossRef](#)]
192. Olsen, M.K.; Lea, U.S.; Slimestad, R.; Verheul, M.; Lillo, C. Differential expression of four *Arabidopsis* PAL genes; PAL1 and PAL2 have functional specialization in abiotic environmental-triggered flavonoid synthesis. *J. Plant Physiol.* **2008**, *165*, 1491–1499. [[CrossRef](#)] [[PubMed](#)]

193. Moura, J.C.M.S.; Bonine, C.A.V.; Viana, J.O.F.; Dornelas, M.C.; Mazzafera, P. Abiotic and biotic stresses and changes in the lignin content and composition in plants. *J. Integr. Plant Biol.* **2010**, *52*, 360–376. [[CrossRef](#)]
194. Pietrowska-Borek, M.; Nuc, K. Stimulating effect of cadmium on phenylpropanoid pathway in *Arabidopsis thaliana* seedlings. *Acta Soc. Bot. Polon.* **2010**, *79* (Suppl. 1), 90.
195. Steyn, W.J.; Wand, S.J.E.; Holcroft, D.M.; Jacobs, G. Anthocyanins in vegetative tissues: A proposed unified function in photoprotection. *New Phytol.* **2002**, *55*, 349–361. [[CrossRef](#)]
196. Bandurska, H.; Pietrowska-Borek, M.; Cieślak, M. Response of barley seedlings to water deficit and enhanced UV-B irradiation acting alone and in combination. *Acta Physiol. Plant.* **2012**, *34*, 161–171. [[CrossRef](#)]
197. Hayat, Q.; Hayat, S.; Irfan, M.; Ahmad, A. Effect of exogenous salicylic acid under changing environment: A review. *Environ. Exp. Bot.* **2010**, *68*, 14–25. [[CrossRef](#)]
198. Gakière, B.; Hao, J.; de Bont, L.; Pétriacq, P.; Nunes-Nesi, A.; Fernie, A.R. NAD⁺ biosynthesis and signaling in plants. *Crit. Rev. Plant Sci.* **2018**, *3*, 259–307. [[CrossRef](#)]
199. Navazio, L.; Bewell, M.A.; Siddiqua, A.; Dickinson, G.D.; Galione, A.; Sanders, D. Calcium release from the endoplasmic reticulum of higher plants elicited by the NADP metabolite nicotinic acid adenine dinucleotide phosphate. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 8693–8698. [[CrossRef](#)] [[PubMed](#)]
200. Navazio, L.; Mariani, P.; Sanders, D. Mobilization of Ca²⁺ by cyclic ADP-ribose from the endoplasmic reticulum of cauliflower florets. *Plant Physiol.* **2001**, *125*, 2129–2138. [[CrossRef](#)] [[PubMed](#)]
201. Guse, A.H.; Lee, H.C. NAADP: A universal Ca²⁺ trigger. *Sci. Signal.* **2008**, *1*, re10. [[CrossRef](#)]
202. Opendakker, K.; Remans, T.; Vangronsveld, J.; Cuyper, A. Mitogen-activated protein (MAP) kinases in plant metal stress: Regulation and responses in comparison to other biotic and abiotic stresses. *Int. J. Mol. Sci.* **2012**, *13*, 7828–7853. [[CrossRef](#)]
203. Devoto, A.; Turner, J.G. Regulation of jasmonate-mediated plant responses in *Arabidopsis*. *Ann. Bot.* **2003**, *92*, 329–337. [[CrossRef](#)]
204. Mine, A.; Berens, M.L.; Nobori, T.; Anver, S.; Fukumoto, K.; Winkel Müller, T.M.; Takeda, A.; Becker, D.; Tsuda, K. Pathogen exploitation of an abscisic acid- and jasmonate-inducible MAPK phosphatase and its interception by *Arabidopsis* immunity. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 7456–7461. [[CrossRef](#)]
205. Vian, A.; Stankovic, B.; Davies, E. Signalomics: Diversity and methods of analysis of systemic signals in plants. In *Plant Omics: The Omics of Plant Science*; Barh, D., Khan, M.S., Davies, E., Eds.; Springer: New Delhi, India, 2015; pp. 459–489.
206. Marcec, M.J.; Gilroy, S.; Poovaiah, B.W.; Tanaka, K. Mutual interplay of Ca²⁺ and ROS signaling in plant immune response. *Plant Sci.* **2019**, *283*, 343–354. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).