

CORRECTION

Correction: Dynamical Hurst analysis identifies EEG channel differences between PTSD and healthy controls

Bahareh Rahmani, Chung Ki Wong, Payam Norouzzadeh, Jerzy Bodurka, Brett McKinney

In [Table 1](#), there was an error when importing the data that made the summary calculation wrong. Please see the corrected summary statistics and [Table 1](#) here.


Table 1. Summary of EEG statistics for PTSD and healthy subjects for channel F3.

| Subject | Mean | Std. | Skewness | Kurtosis |
|---------|---------|---------|----------|----------|
| Healthy | 29.2511 | 15.7567 | -0.4091 | 4.8623 |
| PTSD | 32.4275 | 8.8147 | 0.2089 | 2.5751 |

<https://doi.org/10.1371/journal.pone.0214527.t001>

Reference

1. Rahmani B, Wong CK, Norouzzadeh P, Bodurka J, McKinney B (2018) Dynamical Hurst analysis identifies EEG channel differences between PTSD and healthy controls. *PLoS ONE* 13(7): e0199144. <https://doi.org/10.1371/journal.pone.0199144> PMID: 29969467



OPEN ACCESS

Citation: Rahmani B, Wong CK, Norouzzadeh P, Bodurka J, McKinney B (2019) Correction: Dynamical Hurst analysis identifies EEG channel differences between PTSD and healthy controls. *PLoS ONE* 14(3): e0214527. <https://doi.org/10.1371/journal.pone.0214527>

Published: March 21, 2019

Copyright: © 2019 Rahmani et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.