

Cost-Minimization Analysis of Dexmedetomidine Compared to Other Sedatives for Short-Term Sedation During Mechanical Ventilation in the United States

This article was published in the following Dove Press journal:
ClinicoEconomics and Outcomes Research

Jyoti Aggarwal¹
Jacqueline Lustrino¹
Jennifer Stephens²
Diana Morgenstern³
Wing Yu Tang⁴

¹Pharmerit International, Newton, MA, USA; ²Pharmerit International, Bethesda, MD, USA; ³Pfizer, Collegeville, PA, USA; ⁴Pfizer, New York, NY, USA

Purpose: Mechanical ventilation (MV) remains a substantial cost driver in intensive care units (ICU) in the United States (US). Evaluations of standard sedation treatments used to relieve pain and discomfort in this setting have found varying impacts on ICU length of stay. This cost analysis examines both length-of-stay costs and the total cost implications among MV patients receiving common sedative treatments (dexmedetomidine, propofol, or midazolam) in short-term sedation settings (<24 hours).

Methods: A cost-minimization model was conducted from the hospital provider perspective. Clinical outcomes were obtained from published literature and included ICU length of stay, MV duration, prescription of sedatives and pain medication, and the occurrence of adverse events. Outcomes costs were obtained from previously conducted ICU cost studies and Medicare payment fee schedules. All costs were estimated in 2018 US Dollars.

Results: The per patient costs associated with dexmedetomidine, propofol, and midazolam were estimated to be \$21,115, \$27,073, and \$27,603, respectively. Dexmedetomidine was associated with a savings of \$5958 per patient compared to propofol and a saving of \$6487 compared to midazolam. These savings were primarily driven by a reduction in ICU length of stay and the degree of monitoring and management.

Conclusion: Dexmedetomidine was associated with reduced costs when compared to propofol or midazolam used for short-term sedation during MV in the ICU, suggesting sedative choice can have a potential impact on overall cost per episode.

Keywords: sedation, dexmedetomidine, midazolam, propofol, costs, model

Introduction

Inpatient hospital stays requiring time in the intensive care unit (ICU) are approximately 2.5 times more expensive than non-ICU inpatient stays in the United States (US).¹ In 2011, approximately 27% of the hospital stays across the US involved time in the ICU, which accounted for 48% of aggregate charges from the hospital.¹ Across the 29 states providing data on inpatient and ICU discharges, these charges amounted to over \$280 million.¹

At any given time in the ICU, approximately one-third of beds are used by patients requiring mechanical ventilation (MV).^{2,3,5} Based on an analysis of ICU costs for stays with and without MV, the mean incremental costs associated with

Correspondence: Wing Yu Tang
Pfizer, 235 E. 42nd St, New York, NY
10017, USA
Email wingyu.tang@pfizer.com

MV per day in the ICU have previously been estimated at \$1,522 per day, and thus can be a significant cost burden to the institution.^{2,3}

Medications for sedation are routinely administered to critically ill patients treated in the ICU for a number of reasons, including to improve synchrony with or tolerance of MV.^{6–9} The 2018 clinical practice guidelines from the Society of Critical Care Medicine (SCCM) recommend that light sedation be used in critically ill, mechanically ventilated adults.⁷ Guidelines also generally recommend the use of nonbenzodiazepine sedatives (eg, propofol or dexmedetomidine) vs the use of benzodiazepine sedatives (eg, midazolam or lorazepam) in critically ill, mechanically ventilated patients because of the potential for improved short-term outcomes such as ICU length of stay (LOS), duration of MV, and delirium.^{7,10} Other literature have suggested benzodiazepines use in ICUs.^{11,12} Improvements in MV technologies and guideline support for the minimization of depth and duration of sedation suggest that use of short-acting agents, such as propofol or dexmedetomidine, may increase in the future.^{4,7}

Several clinical trials have compared dexmedetomidine directly with midazolam and propofol for use in the ICU for short-term sedation (< 24 hours).^{9,13–23} These studies suggest that dexmedetomidine is associated with shorter ICU stays and decreased MV time, potentially contributing to decreased costs and improved patient satisfaction.^{9,13–16,18,20–22,24,25} A larger clinical trial conducted in the US comparing dexmedetomidine to both propofol and midazolam also reported a reduction in intubation time and overall ICU/hospital LOS when dexmedetomidine was used.¹⁸

To better understand the potential cost implication of these reported outcomes, we conducted a cost-minimization analysis from the hospital perspective of the impact of sedative choice for patients requiring MV for short-term sedation.

Methods

The cost minimization analysis was conducted from the hospital perspective to reflect costs incurred by an institution for a single ICU stay requiring MV and short-term sedation (<24 hours) of dexmedetomidine, propofol, or midazolam. The model was developed using Microsoft Excel[®] 365 (2016) and Visual Basic for Applications.

Base Model Inputs

Clinical Inputs: A targeted literature search of clinical trials, systematic reviews, and meta-analyses was completed to identify clinical trials comparing dexmedetomidine to either propofol or midazolam. The 16 meta-analyses and 48 clinical trials identified were reviewed for sample size, comparators, short-term sedation focus and inclusion of key outcomes of interest (ie, ICU LOS, duration of MV, sedative duration, and use of rescue sedation and pain medications). A trial conducted and published by Maldonado et al (2009)¹⁸ (N=90) best met key criteria and was selected to supply the clinical inputs for the base case analysis. This trial enrolled and randomized patients in a large, US tertiary-care medical center to receive dexmedetomidine, propofol, or midazolam following the successful weaning from cardio-pulmonary bypass.¹⁸ It was one of the few studies that included all three sedatives within one study, had substantial sample size compared to other identified trials, and measured all key outcomes of interest. Other trials were more limited (eg, geography, sample size) but were included in the sensitivity analysis to test the robustness of results.

Resource Utilization Inputs: Resource utilization inputs included ICU LOS, MV duration, sedative duration, use of rescue sedation and analgesic medications, and the treatment of adverse events (ie, bradycardia, delirium, hypertension, infection, and low blood pressure). Drug and toxicology testing (for benzodiazepines), routine monitoring (eg, respiratory monitoring, physician visits, and intubation), and time spent preparing medications were included as well. [Tables 1](#) and [2](#) summarize the resource utilization inputs and assumptions on frequency for monitoring and management activities for the base-case analysis.

Cost Inputs: Unit cost inputs for the model included medical resource utilization unit costs (including ICU LOS, MV duration, monitoring, and management), drug costs, and sedative preparation time. Cost inputs and cost-related assumptions included in the model were based on published data and are summarized in [Table 2](#). Medical care resource unit costs as well as costs associated with the preparation of sedatives were obtained from the published literature,^{3,26,27} and estimated based on Current Procedural Terminology (CPT[®]) codes and the corresponding physician and clinical laboratory fee schedules.^{28,29} Costs for adverse events were based on the treatments described in [Table 2](#) and assumed that only a single round of treatment was required per adverse event episode. Wholesale acquisition cost was used for any

Table 1 Key Model Input Parameters

Parameter	Dexmedetomidine	Propofol	Midazolam	Data Source
Patient weight (kg)	82.5	82.5	82.5	Average weight of an adult in the US, Centers for Disease Control and Prevention ²⁰
Length of stay in ICU (days)	1.9	3.0	3.0	Maldonado et al ¹⁶
Duration of MV (days) Sedative	0.496	0.463	0.529	Intubation time, Maldonado et al ¹⁶
Treatment duration (hours)	13	11	10	Maldonado et al ¹⁶
% that receive loading dose (%)	100	0	0	Assumption based on study protocol, Maldonado et al ¹⁶
Loading dose (µg/kg)	0.4	–	–	
Duration of loading dose (min)	10	–	–	Assumption based on prescribing information ²²
Maintenance dose (µg/kg/hr or mg/kg/hr) ^a	0.35	1.578	0.018	Calculated dose for midazolam based on average patient weight used in model, Maldonado et al ¹⁶
Time to prepare one bag of sedative (seconds) ^b	35	–	35	Assumption based on time to prepare one emergency syringe, Fraind et al ²⁵ and Jelacic et al ³⁸
Pain medication				
Total dose of morphine (mg)	50.3	51.6	122.5	Maldonado et al ¹⁶
Adverse events				
Occurrence of delirium (%)	10.0	44.4	42.5	ITT population, Maldonado et al ¹⁶

Notes: ^aDexmedetomidine: µg/kg/hr; propofol and midazolam: mg/kg/hr; ^b200 µg/bag of dexmedetomidine; 50 mg/bag of midazolam.

Abbreviations: ICU, intensive care unit; ITT, intention-to-treat; LOS, length of stay; MV, mechanical ventilation; US, United States.

medication-related resource utilization in the model (eg, sedatives, analgesics, treatments for adverse events).³⁰ All costs were adjusted to 2018 US Dollars using the medical component of the Consumer Price Index.³¹

Sensitivity Analysis Inputs

To estimate the robustness of the baseline results, a univariate sensitivity analysis was performed using the clinical outcomes from other published clinical trials comparing dexmedetomidine to propofol or midazolam.^{13–16,19,32} Since not all trials compared dexmedetomidine to both propofol and midazolam, the sensitivity analysis was performed separately for the comparison of dexmedetomidine to propofol and dexmedetomidine to midazolam. The range of inputs used in the sensitivity analysis for each of the input parameters is summarized in Tables 3 and 4. Each clinical input was varied to reflect the minimum and maximum values described in the published literature. If alternative values were not identified in the published literature (eg, cost inputs), we modified the input by 10% in the univariate sensitivity analysis.

Results

The total cost per patient/per ICU stay was \$21,115 for dexmedetomidine, \$27,073 for propofol, and \$27,603 for midazolam treatment in the base-case analysis (Table 5,

Figure 1). Dexmedetomidine was associated with per patient/per ICU stay cost savings compared to both propofol and midazolam. Treatment costs for dexmedetomidine were \$5,958 lower compared to propofol and \$6,487 lower compared to midazolam.

Analysis by cost component (Table 5) showed that the greatest contributor to cost savings was the reduction in ICU LOS with dexmedetomidine. Additional cost reductions with dexmedetomidine compared to propofol and midazolam were observed for monitoring and management activities as well as for occurrence of adverse events. Compared to midazolam only, dexmedetomidine also contributed to a reduction of costs associated with MV.

Sensitivity Analysis

When compared to midazolam, dexmedetomidine was consistently associated with a cost savings in the sensitivity analysis of the base model. These cost savings ranged from \$1,653 to \$13,702. The minimum cost savings was associated with a reduction of 0.83 days in the ICU LOS for patients receiving dexmedetomidine (1.90 days in the base-case analysis vs 2.73 days in the sensitivity analysis). The maximum cost savings was associated with a reduction of 1.23 days in the ICU LOS for patients receiving midazolam (3.00 vs 4.23 days).

Table 2 Cost Inputs

Variables	Cost (2018)	Data Source
ICU room and board, cost per day		
Day 1	\$11,421.91	Dasta et al ³
Day 2	\$5,989.35	Dasta et al ³
Day 3+	\$5,454.83	Dasta et al ³
MV, cost per day		
Day 1	\$7,070.38	Dasta et al ³
Day 2	\$2,227.16	Dasta et al ³
Day 3+	\$1,343.15	Dasta et al ³
Sedative preparation		
Pharmacist hourly rate	\$55.23	Society of Critical Care Medicine ²⁴
Respiratory monitoring costs		
Arterial blood gases, per ICU day	\$26.07	CPT 82,803, Blood gases any combination, CMS ²⁶
Physician consultation cost, per ICU day	\$226.80	CPT 99,291, Critical care first 30–74 minutes, CMS ²⁷
Tracheal intubation cost, per ICU stay	\$155.52	CPT 31,730, Introduction of indwelling tube for oxygen therapy ²⁷
Toxicology testing		
Benzodiazepine testing cost, per ICU treatment day if primary sedative was midazolam, per ICU stay if midazolam is administered for rescue sedation only	\$71.83	CPT 80,307, Drug test by chemistry analyzers ²⁶
Medication costs (wholesale acquisition cost per vial)		
Dexmedetomidine	\$42.00	200 µg vial, Truven Health Analytics ²⁸
Midazolam	\$1.92	50 mg vial, Truven Health Analytics ²⁸
Midazolam	\$0.55	5 mg vial for rescue sedation, Truven Health Analytics ²⁸
Propofol	\$3.88	500 mg vial, Truven Health Analytics ²⁸
Morphine	\$7.05	250 mg vial, Truven Health Analytics ²⁸
Atropine, 0.5 mg per bradycardia event	\$1.61	1 mg vial, Truven Health Analytics ²⁸
Haloperidol, 10 mg per delirium event	\$0.88	5 mg vial, Truven Health Analytics ²⁸
Labetalol, 20 mg per hypertension event	\$3.35	100 mg vial, Truven Health Analytics ²⁸
Norepinephrine, 4 mg per hypotension event	\$4.60	4 mg vial, Truven Health Analytics ²⁸
Cefazolin, 1 g per infection event	\$6.33	2 g vial, Truven Health Analytics ²⁸

Abbreviations: CMS, Centers for Medicare and Medicaid Services; CPT, current procedural code; ICU, intensive care unit; MV, mechanical ventilation.

The sensitivity testing of propofol found similar conclusions. Dexmedetomidine compared to propofol was associated with a maximum cost savings of \$12,081 when the ICU LOS for patients receiving dexmedetomidine was reduced by 0.94 days compared to the base case (1.90 vs 0.96 days). When LOS for patients receiving propofol was reduced to 2.04 days (3.00 vs 0.96 days), the sensitivity analysis at this lower range did find dexmedetomidine was associated with an increased cost.

Discussion

In this model, dexmedetomidine was generally found to be associated with significant cost savings of ~\$6,000 compared to propofol and midazolam in mechanically

ventilated, adult patients undergoing short-term sedation (< 24 hours). The cost savings associated with reduced ICU LOS and required monitoring and management reflects potential economic value of dexmedetomidine.

These results are directionally consistent with multiple cost studies completed in the US for sedated mechanically ventilated patients.^{18,24,33} Previous cost studies have demonstrated that clinical outcomes associated with sedative choice can impact the overall cost of an ICU stay.^{18,24,30,33–37} The drivers of cost savings in this analysis were also consistent with previous studies, with the key driver of high costs being the fixed room and board costs associated with the ICU stay.^{18,24,33,37} The absolute costs and the magnitude of the cost differences, however,

Table 3 Sensitivity Analysis Clinical Inputs for Propofol Comparison

Parameters	Dexmedetomidine Arm		Propofol Arm		Data Source
	Lower Value	Upper Value	Lower Value	Upper Value	
Length of stay in ICU (days)	0.96	1.90	0.96	3.00	Corbett et al, ³⁰ Maldonado et al ¹⁶
Duration of MV (days)	0.23	0.50	0.25	0.54	Djaiani et al, ¹² Srivastava et al ¹³
Sedative					
Treatment duration relative to MV duration (\pm hours) ^a	-0.59	6.00	-0.44	-0.10	Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
% that receive loading dose (%)	90	100	0	100	Dexmedetomidine inputs did not vary in published literature, varied by 10% for analysis. Corbett et al, ³⁰ Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Loading dose (μ g/kg or mg/kg/hr) ^a	0.40	1.00	0.00	4.00	Corbett et al, ³⁰ Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Duration of loading dose (minutes) ^a	10.00	20.00	0.00	15.00	Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Maintenance dose (μ g/kg/hr or mg/kg/hr) ^b	0.31	0.55	0.04	2.00	Corbett et al, ³⁰ Djaiani et al, ¹² Srivastava et al ¹³
Time to prepare bag (200 μ g) of dexmedetomidine or vial of propofol (seconds)	25	115	0	59	Fraind et al, ²⁵ Jelacic et al, ³⁸ and van der Linden ³⁹
Rescue sedation, midazolam					
% that receive rescue sedation (%)	0	100	0	100	Corbett et al, ³⁰ Maldonado et al, ¹⁶ Srivastava et al ¹³
Rescue sedative dose (mg)	0	1.5	0	1	Corbett et al, ³⁰ Maldonado et al, ¹⁶ Srivastava et al ¹³
Pain medication					
% treated with pain medication (%)	90	100	90	100	Inputs did not vary in published literature, varied by 10% for analysis. Corbett et al, Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Total dose of morphine (mg)	2.95	50.30	6.00	51.60	Corbett et al, ³⁰ Herr et al, ¹⁷ Maldonado et al ¹⁶
Adverse events					
Occurrence of bradycardia (%)	0	3	0	1	Corbett et al, ³⁰ Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Occurrence of delirium (%)	0	12	0	44	Corbett et al, ³⁰ Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Occurrence of hypertension (%)	0	12	0	4	Corbett et al, ³⁰ Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³
Occurrence of hypotension (%)	0	81	0	67	Corbett et al, ³⁰ Djaiani et al, ¹² Maldonado et al, ¹⁶ Srivastava et al ¹³
Occurrence of infection (%)	0	10	0	10	Inputs did not vary in published literature, varied by 10% for analysis. Corbett et al, ³⁰ Djaiani et al, ¹² Herr et al, ¹⁷ Maldonado et al, ¹⁶ Srivastava et al ¹³

Notes: ^aDexmedetomidine: μ g/kg; propofol: mg/kg/hr; ^bDexmedetomidine: μ g/kg/hr; propofol: mg/kg/hr.

Abbreviations: ICU, intensive care unit; MV, mechanical ventilation.

differed between our analysis and previous cost studies which included mixed patient populations requiring both long-term and short-term sedation. Selection of inputs, and overall study design differences likely also contributed.

Of note, one published cost analysis in 2009 by Patanwala et al³⁷ did report higher costs associated with

dexmedetomidine compared to propofol for sedation in the ICU. In that analysis, total hospital costs when ICU patients were treated with dexmedetomidine were \$46,716 compared to \$31,041 in the propofol group.³⁷ However, the increased LOS that drove the cost differential was not adjusted for potential unmeasured

Table 4 Sensitivity Analysis Clinical Inputs for Midazolam Comparison

Parameters	Dexmedetomidine Arm		Midazolam Arm		Data Source
	Lower Value	Upper Value	Lower Value	Upper Value	
Length of stay in ICU (days)	1.90	2.73	3.00	4.23	Azeem et al, ¹¹ Maldonado et al ¹⁶
Duration of MV (days)	0.19	0.50	0.52	0.54	Azeem et al, ¹¹ Srivastava et al, ¹³ Wan et al ¹⁴
Sedative					
Treatment duration relative to MV duration (± hours)	-0.59	2.80	-2.70	-0.80	Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
% that receive loading dose (%)	0%	100%	0%	100%	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Loading dose (µg/kg or mg/kg) ^a	0.00	1.00	0.00	0.04	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Duration of loading dose (min)	0.00	15.00	0.00	15.00	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Maintenance dose (µg/kg/hr or mg/kg/hr) ^b	0.35	0.72	0.02	0.44	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Wan et al ¹⁴
Time to prepare one bag of sedative (sec) ^c	25	115	25	115	Fraind et al, ²⁵ Jelacic et al, ³⁸ and van der Linden ³⁹
Rescue sedation, midazolam					
% that receive rescue sedation (%)	0	100	-	-	For the midazolam arm, considered in sensitivity analysis for the primary sedative Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Rescue sedative dose (mg)	0	6.7	-	-	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Pain medication					
% treated with pain medication (%)	90	100	90	100	Inputs did not vary in published literature, varied by 10% for analysis. Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Total dose of morphine (mg)	17.61	50.30	37.21	125.06	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Wan et al ¹⁴
Adverse events					
Occurrence of bradycardia (%)	0	25	0	10	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Occurrence of delirium (%)	0	10	0	43	Maldonado et al, ¹⁶ Srivastava et al ¹³
Occurrence of hypertension (%)	0	10	0	10	Inputs did not vary in published literature, varied by 10% for analysis. Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Occurrence of hypotension (%)	0	28	0	11	Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴
Occurrence of infection (%)	0	10	0	10	Inputs did not vary in published literature, varied by 10% for analysis. Azeem et al, ¹¹ Maldonado et al, ¹⁶ Srivastava et al, ¹³ Wan et al ¹⁴

Notes: ^aDexmedetomidine: µg/kg; midazolam: mg/kg; ^bDexmedetomidine: µg/kg/hr; midazolam: mg/kg/hr; ^c200 µg/bag of dexmedetomidine; 50 mg/bag of midazolam.

Abbreviations: ICU, intensive care unit; MV, mechanical ventilation.

confounders, such as dosing, level of sedation, delirium screening, or rescue sedation. Additionally, there were several observed differences between the two patient populations (eg, age, sex, race, admission type, service category, and severity of illness) that may suggest selection bias for treatment with each sedative.³⁷

Nonetheless, there are important limitations to keep in mind regarding the current model. First, the base case analysis utilized clinical inputs from an open label trial. While an unblinded study design was reasonable due to the distinct physical and pharmacological properties of the chosen sedatives, the possibility of investigator bias could

Table 5 Base Case Results: Estimated ICU Cost Associated with Each Sedative, per Patient

	Cost per Sedative			Difference between Dexmedetomidine and Comparator	
	Dexmedetomidine	Propofol	Midazolam	v Propofol	vs Midazolam
ICU Days	\$16,812.33	\$22,866.09	\$22,866.09	-\$6,053.77	-\$6,053.77
MV Days	\$3,506.91	\$3,273.59	\$3,740.23	\$233.32	-\$233.32
Monitoring/Management	\$661.26	\$914.13	\$985.96	-\$252.87	-\$324.70
Sedative	\$127.61	\$11.64	\$2.46	\$115.97	\$125.15
Rescue Sedation	\$0.00	\$0.00	\$0.00	\$0.00	\$0.00
Pain Medication	\$7.05	\$7.05	\$7.05	\$0.00	\$0.00
Adverse Event	\$0.18	\$0.78	\$0.75	-\$0.61	-\$0.57
Total	\$21,115.33	\$27,073.28	\$27,602.54	-\$5,957.95	-\$6,487.21

Abbreviations: ICU, intensive care unit; MV, mechanical ventilation; vs, versus

not be eliminated. Further, while the results of our cost-minimization analysis were informed by national costing information and tested with the range of results found within the literature via sensitivity analysis, the analysis was performed for a specific population and caution should be employed in generalizing to all MV patients. The model was fit for purpose for one country and may not be reflective of different markets, given differences in indications, utilization, and reimbursement systems. Future explorations specific to European and Asian market dynamics could further a broader global understanding. There were also cases where a few unit costs were not readily available. In lieu of this, we applied reimbursement amounts pulled from US reimbursement fee schedules. None of these cost elements were the main drivers of overall cost estimates in the model, so the impact on conclusions drawn is likely minimal.




Figure 1 Base case results: Essential ICU cost associated with each schedule, per patient. *Other costs include cost of sedative, reduce sedative, pain medication, monitoring/management, and cost of adverse event treatment.

Abbreviations: ICU, intensive care unit; MV, mechanical ventilation.

Lastly, it is important to reflect that this model offers one perspective in the ongoing understanding and discussion of short-term sedation in the ICU. After the development of this model, new studies, such as the Spice III study have been published.³⁸ While SPICE III is out of scope geographically for this current US specific model (ie, the study was conducted in eight ex-US countries that might have variations in sedative indications, utilization, and reimbursement systems), it is still important to reflect that the understanding of these dynamics within the ICU remain an evolving conversation. The results from the current analysis and their interpretation are meant to contribute and further open value and cost perspective discussions that have in the past been limited in literature.

Conclusion

With US national healthcare spending projected to reach \$5.7 trillion by 2026³⁹ and critical care accounting for about 4.1%⁴⁰ of this spending, it is important to understand potential cost reduction opportunities in the ICU. This cost-minimization analysis suggests that choice of short-term sedation during MV may contribute to ICU LOS and associated costs. Critical care practitioners may want to consider the results of this analysis when selecting short-term sedatives for their mechanically-ventilated ICU patients.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Funding

This study was sponsored by Pfizer. Kafi Sanders of Pfizer contributed to initial core model research and construct.

Disclosure

Jennifer Stephens, Jyoti Aggarwal, and Jacqueline Lustrino are employees of Pharmerit International who were paid consultants to Pfizer in connection with this study and development of the manuscript. Wing Yu Tang and Diana Morgenstern are employees and stockholders of Pfizer. The authors report no other conflicts of interest in this work.

References

- Barrett ML, Smith MW, Elixhauser A, Leah S, Honigman, Pines JM. *Statistical Brief #185: Utilization of Intensive Care Services, 2011*. Healthcare Cost and Utilization Project; 2014.
- Kramer AA, Dasta JF, Kane-Gill SL. The impact of mortality on total costs within the ICU. *Crit Care Med*. 2017;45(9):1457–1463. doi:10.1097/CCM.0000000000002563
- Dasta JF, McLaughlin TP, Mody SH, Piech CT. Daily cost of an intensive care unit day: the contribution of mechanical ventilation. *Crit Care Med*. 2005;33(6):1266–1271. doi:10.1097/01.CCM.000164543.14619.00
- Reade MC, Finfer S. Sedation and delirium in the intensive care unit. *N Engl J Med*. 2014;370(5):444–454. doi:10.1056/NEJMra1208705
- Wunsch H, Wagner J, Herlim M, Chong DH, Kramer AA, Halpern SD. ICU occupancy and mechanical ventilator use in the United States. *Crit Care Med*. 2013;41(12):2712–2719. doi:10.1097/CCM.0b013e318298a139
- Hughes CG, McGrane S, Pandharipande PP. Sedation in the intensive care setting. *Clin Pharmacol*. 2012;4:53–63. doi:10.2147/CPAA.S26582
- Devlin JW, Skrobik Y, Gelinas C, et al. Clinical practice guidelines for the prevention and management of pain, agitation/ sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med*. 2018;46(9):e825e873. doi:10.1097/CCM.0000000000003299
- Sydow M, Neumann P. Sedation for the critically ill. *Intensive Care Med*. 1999;25(6):634–636. doi:10.1007/s001340050917
- Zhang Z, Chen K, Ni H, Zhang X, Fan H. Sedation of mechanically ventilated adults in intensive care unit: a network meta-analysis. *Sci Rep*. 2017;7:44979. doi:10.1038/srep44979
- Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med*. 2013;41(1):263–306. doi:10.1097/CCM.0b013e3182783b72
- Kotfis K, Zegan-Barańska M, Żukowski M, Kusza K, Kaczmarczyk M, Ely EW. Multicenter assessment of sedation and delirium practices in the intensive care units in Poland—is this common practice in Eastern Europe? *BMC Anesthesiol*. 2017;17(1):120. doi:10.1186/s12871-017-0415-2
- Morandi A, Piva S, Ely EW, et al. Worldwide Survey of the “Assessing pain, both spontaneous awakening and breathing trials, choice of drugs, delirium monitoring/management, early exercise/mobility, and family empowerment” (ABCDEF) Bundle. *Crit Care Med*. 2017;45(11):e1111. doi:10.1097/CCM.0000000000002640
- Azeem TMA, Yosif NE, Alansary AM, Esmat IM, Mohamed AK. Dexmedetomidine vs morphine and midazolam in the prevention and treatment of delirium after adult cardiac surgery; a randomized, double-blinded clinical trial. *Saudi J Anaesth*. 2018;12(2):190–197. doi:10.4103/sja.SJA_303_17
- Djaiani G, Silverton N, Fedorko L, et al. Dexmedetomidine versus propofol sedation reduces delirium after cardiac surgery: a randomized controlled trial. *Anesthesiology*. 2016;124(2):362–368. doi:10.1097/ALN.0000000000000951
- Srivastava VK, Agrawal S, Kumar S, Mishra A, Sharma S, Kumar R. Comparison of dexmedetomidine, propofol and midazolam for short-term sedation in postoperatively mechanically ventilated neurosurgical patients. *J Clin Diagn Res*. 2014;8(9):GC0407. doi:10.7860/JCDR/2014/8797.4817
- Wan LJ, Huang QQ, Yue JX, Lin L, Li SH. Comparison of sedative effect of dexmedetomidine and midazolam for post-operative patients undergoing mechanical ventilation in surgical intensive care unit. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue*. 2011;23(9):543–546. Chinese.
- Martin E, Ramsay G, Mantz J, Sum-Ping ST. The role of the alpha2-adrenoceptor agonist dexmedetomidine in postsurgical sedation in the intensive care unit. *J Intensive Care Med*. 2003;18(1):29–41. doi:10.1177/0885066602239122
- Maldonado JR, Wysong A, van der Starre PJ, Block T, Miller C, Reitz BA. Dexmedetomidine and the reduction of postoperative delirium after cardiac surgery. *Psychosomatics*. 2009;50(3):206–217. doi:10.1176/appi.psy.50.3.206
- Herr DL, Sum-Ping ST, England M. ICU sedation after coronary artery bypass graft surgery: dexmedetomidine-based versus propofol-based sedation regimens. *J Cardiothorac Vasc Anesth*. 2003;17(5):576–584. doi:10.1016/S1053-0770(03)00200-3
- Fan H, Zhao Y, Sun M, Ye JH, Chen GD, Zhu JH. Dexmedetomidine based sedation for post-surgery critically ill adults: a meta-analysis of randomized controlled trials. *Iran J Public Health*. 2017;46(12):1611–1622.
- Liu H, Ji F, Peng K, Applegate RL 2nd, Fleming N. Sedation after cardiac surgery: is one drug better than another? *Anesth Analg*. 2017;124(4):1061–1070. doi:10.1213/ANE.0000000000001588
- Cruickshank M, Henderson L, MacLennan G, et al. Alpha-2 agonists for sedation of mechanically ventilated adults in intensive care units: a systematic review. *Health Technol Assess*. 2016;20(25):v–xx, 1–117. doi:10.3310/hta20250
- Riker RR, Fraser GL. Altering intensive care sedation paradigms to improve patient outcomes. *Crit Care Clin*. 2009;25(3):527–538, viii–ix. doi:10.1016/j.ccc.2009.05.004
- Thoma BN, Li J, McDaniel CM, Wordell CJ, Cavarocchi N, Pizzi LT. Clinical and economic impact of substituting dexmedetomidine for propofol due to a US drug shortage: examination of coronary artery bypass graft patients at an urban medical centre. *Pharmacoeconomics*. 2014;32(2):149–157. doi:10.1007/s40273-013-0116-8
- Nguyen J, Nacpil N. Effectiveness of dexmedetomidine versus propofol on extubation times, length of stay and mortality rates in adult cardiac surgery patients: a systematic review and meta-analysis. *Joanna Briggs Inst Database Sys Rev Impl Rep*. 2018;16(5):1220–1239. doi:10.11124/JBISRIR-2017-003488
- Critical Care Statistics. Society of critical care medicine. Available from: <http://www.sccm.org/Communications/Critical-Care-Statistics>. Accessed October 20, 2018.
- Fraind DB, Slagle JM, Tubbesing VA, Hughes SA, Weinger MB. Reengineering intravenous drug and fluid administration processes in the operating room: step one: task analysis of existing processes. *Anesthesiology*. 2002;97(1):139–147. doi:10.1097/0000542-200207000-00020
- Centers for medicare & medicaid services. *Clin Lab Fee Schedule*. 2018.
- Centers for medicare & medicaid services. *Phy Fee Schedule*. 2018.
- Park JH, Jeong JH. A cost-minimization analysis comparing dexmedetomidine with midazolam and propofol for sedation in mechanically ventilated intensive care unit patients. *J Health Technol Assess*. 2016;4(2):75–81. doi:10.34161/johta.2016.4.2.006
- Bureau of Labor Statistics. *Consumer Price Index-All Urban Consumers (Current Series)*. 2018.
- Corbett SM, Rebeck JA, Greene CM, et al. Dexmedetomidine does not improve patient satisfaction when compared with propofol during mechanical ventilation. *Crit Care Med*. 2005;33(5):940–945. doi:10.1097/01.CCM.0000162565.18193.E5

33. Dasta JF, Kane-Gill SL, Pencina M, et al. A cost-minimization analysis of dexmedetomidine compared with midazolam for long-term sedation in the intensive care unit. *Crit Care Med.* 2010;38(2):497–503. doi:10.1097/CCM.0b013e3181bc81c9
34. Athanzio R, Maldini P, Roa S, Turunen H, Silva C. Cost-minimisation analysis of dexmedetomidine versus propofol in mechanical ventilated patients at ICU. *Value Health.* 2015;18(7):A842843. doi:10.1016/j.jval.2015.09.385
35. Turunen H, Jakob SM, Ruokonen E, et al. Dexmedetomidine versus standard care sedation with propofol or midazolam in intensive care: an economic evaluation. *Critical Care.* 2015;19:67. doi:10.1186/s13054-015-0787-y
36. Lachaine J, Beauchemin C. Economic evaluation of dexmedetomidine relative to midazolam for sedation in the intensive care unit. *Can J Hosp Pharm.* 2012;65(2):103–110. doi:10.4212/cjhp.v65i2.1116
37. Patanwala AE, Erstad BL. Comparison of dexmedetomidine versus propofol on hospital costs and length of stay. *J Intensive Care Med.* 2016;31(7):466–470. doi:10.1177/0885066614544452
38. Shehabi Y, Howe BD, Bellomo R, et al. Early sedation with dexmedetomidine in critically ill patients. *New Eng J Med.* 2019;380(26):2506–2517. doi:10.1056/NEJMoa1904710
39. *National Health Expenditure Projections.* Centers for Medicare & Medicaid Services; 2017.
40. Halpern NA, Pastores SM. Critical care medicine in the United States 2000-2005: an analysis of bed numbers, occupancy rates, payer mix, and costs. *Crit Care Med.* 2010;38(1):65–71. doi:10.1097/CCM.0b013e3181b090d0

ClinicoEconomics and Outcomes Research

Dovepress

Publish your work in this journal

ClinicoEconomics and Outcomes Research is an international, peer-reviewed open-access journal focusing on Health Technology Assessment, Pharmacoeconomics and Outcomes Research in the areas of diagnosis, medical devices, and clinical, surgical and pharmacological intervention. The economic impact of health policy and health systems

organization also constitute important areas of coverage. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinicoeconomics-and-outcomes-research-journal>