

Research article

Open Access

Evidence for calcium-mediated perception of plant symbiotic signals in aequorin-expressing *Mesorhizobium loti*

Roberto Moscatiello¹, Sara Alberghini², Andrea Squartini², Paola Mariani¹ and Lorella Navazio*¹

Address: ¹Dipartimento di Biologia, Università di Padova, Via U. Bassi 58/B, 35131 Padova, Italy and ²Dipartimento di Biotecnologie Agrarie, Università di Padova, Viale dell'Università 16, 35020 Legnaro, Padova, Italy

Email: Roberto Moscatiello - roberto.moscatiello@unipd.it; Sara Alberghini - sara.alberghini@unipd.it; Andrea Squartini - squart@unipd.it; Paola Mariani - mariani@bio.unipd.it; Lorella Navazio* - lorella.navazio@unipd.it

* Corresponding author

Published: 23 September 2009

Received: 7 May 2009

BMC Microbiology 2009, 9:206 doi:10.1186/1471-2180-9-206

Accepted: 23 September 2009

This article is available from: <http://www.biomedcentral.com/1471-2180/9/206>

© 2009 Moscatiello et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: During the interaction between rhizobia and leguminous plants the two partners engage in a molecular conversation that leads to reciprocal recognition and ensures the beginning of a successful symbiotic integration. In host plants, intracellular Ca^{2+} changes are an integral part of the signalling mechanism. In rhizobia it is not yet known whether Ca^{2+} can act as a transducer of symbiotic signals.

Results: A plasmid encoding the bioluminescent Ca^{2+} probe aequorin was introduced into *Mesorhizobium loti* USDA 3147^T strain to investigate whether a Ca^{2+} response is activated in rhizobia upon perception of plant root exudates. We find that *M. loti* cells respond to environmental and symbiotic cues through transient elevations in intracellular free Ca^{2+} concentration. Only root exudates from the homologous host *Lotus japonicus* induce Ca^{2+} signalling and downstream activation of nodulation genes. The extracellular Ca^{2+} chelator EGTA inhibits both transient intracellular Ca^{2+} increase and inducible *nod* gene expression, while not affecting the expression of other genes, either constitutively expressed or inducible.

Conclusion: These findings indicate a newly described early event in the molecular dialogue between plants and rhizobia and highlight the use of aequorin-expressing bacterial strains as a promising novel approach for research in legume symbiosis.

Background

Rhizobia are Gram-negative soil bacteria which can engage in a mutualistic association with leguminous plants. Under nitrogen-limiting conditions, rhizobia colonize plant roots and highly specialized plant organs, the nodules, are generated *de novo* on host roots (for a recent review see [1]). When living symbiotically, rhizobia are able to fix atmospheric nitrogen into forms usable by the

plant. In return, they receive dicarboxylic acids as a carbon and energy source for their metabolism. Nitrogen is the most frequent limiting macronutrient in many soils, and it is generally supplied as fertilizer. The rhizobium-legume mutualistic association can reduce or eliminate nitrogen fertilizer requirements, resulting also in a benefit to the environment [2].

A successful symbiosis is the result of an elaborate developmental program, regulated by the exchange of molecular signals between the two partners [3]. During growth in the rhizosphere of the host plant, rhizobia sense compounds secreted by the host root and respond by inducing bacterial nodulation (*nod*) genes which are required for the synthesis of rhizobial signal molecules of lipo-chitooligosaccharide nature, the Nod factors. In the host plant, the generation of intracellular Ca^{2+} oscillations triggered by Nod factors has been firmly established as one of the earliest crucial events in symbiosis signalling; these oscillations are transduced into downstream physiological and developmental responses [1]. It is not known whether there is a parallel key role for Ca^{2+} in rhizobia.

As in eukaryotic cells, Ca^{2+} is postulated to play essential functions in the regulation of a number of cellular processes in bacteria, including the cell cycle, differentiation, chemotaxis and pathogenicity [4,5]. Homeostatic machinery that is able to regulate intracellular free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) tightly is a prerequisite for a Ca^{2+} -based signalling system, and is known to be present in bacteria [6]. Ca^{2+} transport systems have been demonstrated in bacteria, with the identification of primary pumps and secondary exchangers, as well as putative Ca^{2+} -permeable channels [5,7]. Other Ca^{2+} regulatory components such as Ca^{2+} -binding proteins, including several EF-hand proteins, have been detected and have been putatively identified from genomic sequences [8,9].

In order to establish precisely when and how Ca^{2+} regulates processes in bacteria it is essential to measure $[\text{Ca}^{2+}]_i$ and its changes in live cells. This has proven difficult because of problems in loading fluorescent Ca^{2+} indicator dyes, such as fura-2, into bacterial cells. However, the recombinant expression of the Ca^{2+} -sensitive photoprotein aequorin, which has been demonstrated to be a suitable method to monitor $[\text{Ca}^{2+}]_i$ changes accurately in eukaryotes [10-12], has been successfully applied also to bacteria. Challenge of *E.coli* [13-17] and the cyanobacterium *Anabaena* sp. PCC7120 [18-21] expressing aequorin with different stimuli resulted in the induction of transient variations of $[\text{Ca}^{2+}]_i$ with specific Ca^{2+} signatures.

Here we report the introduction of a plasmid encoding apoaquorin in *Mesorhizobium loti*, the specific symbiont of the model legume *Lotus japonicus*, and the use of this reporter to examine the Ca^{2+} response of rhizobia to abiotic and biotic stimuli. The results obtained highlight the occurrence in *M. loti* of Ca^{2+} -based mechanisms for sensing and responding to cues originating in the rhizosphere.

Results

Construction of an inducible reporter system for Ca^{2+} measurements in rhizobia

The apoaquorin gene was cloned in the broad host-range expression vector pDB1 [22] under the control of the strong synthetic promoter P_{syn} , regulated by the *lacI*^q repressor (see Additional file 1). The pAEQ80 plasmid was mobilized by conjugation into the type strain of *M. loti* (USDA 3147^T).

Validation of the experimental system

The functioning in *M. loti* of the pAEQ80 plasmid containing the apoaquorin gene was verified by evaluating the level of aequorin expression in an *in vitro* reconstitution assay. Light emitted by total soluble protein contained in the lysates from wild-type and aequorin-expressing *M. loti* cells was monitored after reconstitution of the apoprotein with coelenterazine. The strong luminescence signal detected in protein extracts from *M. loti* cells containing the apoaquorin construct and induced with IPTG confirmed the efficient level of aequorin expression (see Additional file 2).

We analysed whether the introduced pAEQ80 plasmid (10.5 kb) encoding apoaquorin or the expressed protein could affect bacterial cell growth and the symbiotic performance of *M. loti* cells. There is no significant effect on bacterial growth kinetics exerted either by the introduced plasmid or apoaquorin expression. Nodulation efficiency of *M. loti* pAEQ80 cells on the specific plant host *Lotus japonicus* was checked 4 weeks after bacterial inoculation on roots of seedlings grown on nitrogen-free medium. *L. japonicus* roots were found to be effectively nodulated by the transformed bacterial strain, with no differences in nodule number (5 ± 1) and morphological parameters in comparison to seedlings inoculated with wild-type *M. loti*. The presence of bacteria inside nodules was verified by light microscopy (see Additional file 2). Green foliage was indicative of functional symbiosis.

The occurrence in *M. loti* cells of homeostatic control of the internal Ca^{2+} activity was then verified by preliminary Ca^{2+} measurement assays in a luminometer after *in vivo* reconstitution of apoaquorin. Unperturbed exponentially growing rhizobial cells showed a steady-state intracellular free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) residing in the submicromolar range (around 500 nM) (see Additional file 2), demonstrating a tight regulation of $[\text{Ca}^{2+}]_i$. No luminescence was detected either in cultures of the non-recombinant strain incubated with coelenterazine or in recombinant cells that had not been exposed to coelenterazine (data not shown), confirming that the recorded signal was due only to Ca^{2+} -dependent light emission from aequorin.

Environmental stimuli are sensed through transient $[Ca^{2+}]_i$ elevations by *M. loti*

To further validate the experimental system, abiotic stimuli which are known to trigger $[Ca^{2+}]_i$ changes in both plants [23] and cyanobacteria [18,19] were applied to apoaequorin-expressing *M. loti* cells. A mechanical perturbation, simulated by the injection of isoosmotic cell culture medium, resulted in a rapid Ca^{2+} transient increase ($1.08 \pm 0.24 \mu M$) that decayed within 30 sec (Fig. 1A). This Ca^{2+} trace, which is frequently referred to as a "touch

response", is often observed after the hand-operated injection of any stimulus [24]. A similar Ca^{2+} response characterized by an enhanced Ca^{2+} peak of $2.14 \pm 0.46 \mu M$ was triggered by a simple injection of air into the cell suspension with a needle (Fig. 1A).

Cold and hypoosmotic shocks, caused by supplying three volumes of ice-cold medium and distilled water, respectively, induced Ca^{2+} traces with distinct kinetics, e.g. different height of the Ca^{2+} peak ($1.36 \pm 0.13 \mu M$ and $4.41 \pm$




Figure 1
 Ca^{2+} measurements in *M. loti* cells stimulated with different physico-chemical signals. Bacteria were challenged (arrow) with: **A**, mechanical perturbation, represented by injection of an equal volume of culture medium (black trace) or 10 volumes of air (grey trace); **B**, cold shock, given by 3 volumes of ice-cold culture medium (black trace); control cells were stimulated with 3 volumes of growth medium kept at room temperature (grey trace); **C**, hypoosmotic stress, given by injection of 3 volumes of distilled water (black trace); salinity stress, represented by 200 mM NaCl (grey trace); **D**, different external Ca^{2+} concentrations. These and the following traces have been chosen to best represent the average results of at least three independent experiments.

0.51 μM , respectively) and rate of dissipation of the Ca^{2+} signal (Fig. 1B and 1C). As a control, cells were stimulated with three volumes of growth medium at room temperature, (Fig. 1B) resulting in a Ca^{2+} trace superimposable on that of the touch response (Fig. 1A). These findings eliminate the possible effect of bacterial dilution on changes in Ca^{2+} homeostasis.

Challenge of *M. loti* with a salinity stress, which has recently been shown to affect symbiosis-related events in *Rhizobium tropici* [25], resulted in a $[\text{Ca}^{2+}]_i$ elevation of large amplitude ($3.36 \pm 0.24 \mu\text{M}$) and a specific signature (Fig. 1C).

Variations in the extracellular Ca^{2+} concentration determined the induction of transient Ca^{2+} elevations whose magnitude was dependent on the level of external Ca^{2+} . After a rapidly induced increase in $[\text{Ca}^{2+}]_i$, the basal Ca^{2+} level was gradually restored with all the applied external Ca^{2+} concentrations (Fig. 1D), confirming a tight internal

homeostatic Ca^{2+} control, as previously shown for other bacteria [14,18].

All the above results indicate that aequorin-expressing *M. loti* cells comprise a functionally valid system with which to investigate the involvement of Ca^{2+} in intracellular transduction of environmental stimuli.

Host plant root exudates induce in *M. loti* a Ca^{2+} signal required for activation of nodulation genes

Root exudates from the symbiotically compatible legume *L. japonicus* were collected from 3-week-old seedlings axenically grown in water and applied to *M. loti* cells. The dose used for Ca^{2+} measurements was in the range that induced significant expression of *nodA*, *nodB*, *nodC* genes in *M. loti* (Fig. 2A). This concentration was found to trigger a transient $[\text{Ca}^{2+}]_i$ change characterized by a very rapid increase ($1.38 \pm 0.23 \mu\text{M} \text{Ca}^{2+}$) followed by a second sustained major Ca^{2+} peak ($2.01 \pm 0.24 \mu\text{M}$) at about 10 min (Fig. 2B), with a slow decay within the considered time




Figure 2
Effect of plant root exudates and tetronic acid on $[\text{Ca}^{2+}]_i$ and *nod* gene expression in *M. loti*. **A**, Analysis of gene expression by semi-quantitative RT-PCR during control conditions (lane 1, white bars) and after 1 h treatment with *L. japonicus* root exudates (lane 2, black bars) or 1.5 mM tetronic acid (lane 2, striped bars). Relative transcript abundance was normalized against 16S rRNA. Data are the means \pm SEM of three independent experiments. **B**, Monitoring of $[\text{Ca}^{2+}]_i$ changes in *M. loti* cells challenged (arrow) with *L. japonicus* root exudates (black trace) or 1.5 mM tetronic acid (grey trace).

interval (30 min). The observed induction of transient $[Ca^{2+}]_i$ changes in *M. loti* cells suggests a Ca^{2+} -mediated perception of signalling molecules contained in host plant root exudates.

Flavonoids are components of root exudates that play a prominent role as inducers of structural *nod* genes in rhizobia. Although flavonoids have been detected in *L. japonicus* seeds [26], those that specifically activate the expression of *nod* genes in *M. loti* have not yet been identified [27,28]. The most common flavonoids, known as *nod* gene inducers in other rhizobia (10 μ M naringenin, luteolin, daidzein, kaempferol, quercetin dehydrate) were not able to trigger transient Ca^{2+} elevations in *M. loti* (data not shown). Tetronic acid, an aldonic acid previously reported to promote Nod factor biosynthesis in *M. loti* [29], was found to induce a detectable Ca^{2+} response (Fig. 2B). The kinetics of the Ca^{2+} trace was similar to that induced by crude root exudates, with a prompt Ca^{2+} spike ($1.36 \pm 0.16 \mu$ M Ca^{2+}) and a subsequent flattened dome (maximal Ca^{2+} value of $1.29 \pm 0.08 \mu$ M reached around 15 min after the elicitor application). Notably, this second phase of the Ca^{2+} transient induced by tetronic acid only partially accounted for the larger Ca^{2+} increase recorded with the whole *L. japonicus* root exudates (Fig. 2B). Likewise, the level of *nod* gene expression induced by tetronic acid was found to be lower (though significantly different from the control, $P < 0.05$) than that generated by total root exudates (Fig. 2A).

Pretreatment of rhizobial cells with the extracellular Ca^{2+} chelator EGTA for 10 min effectively inhibited both the transient Ca^{2+} elevation (Fig. 3A) and *nod* gene activation (Fig. 3B) induced by *L. japonicus* root exudates. This indicates that the main source of the observed Ca^{2+} response is the extracellular medium, and that the elevation in $[Ca^{2+}]_i$ is required for *nod* gene induction. Cell viability, monitored by the BacLight Bacterial viability assay, was not altered by incubation with the Ca^{2+} chelator (Fig. 3C). The expression of both constitutive (glutamine synthetase II and 16S rRNA) and inducible (aequorin) genes was not significantly affected by EGTA treatment (Fig. 3D and 3E), ruling out possible general effects of extracellular Ca^{2+} chelation on gene induction.

To check host specificity of the Ca^{2+} signal, metabolite mixtures exuded by the non-host legumes soybean and *Vicia sativa* subsp. *nigra* were tested. After an initial rapid and steep Ca^{2+} rise ($1.77 \pm 0.34 \mu$ M), shared also by the response to *L. japonicus* root exudates, the Ca^{2+} transients triggered by non-host exudates show very different kinetics, such as a slow rate of decay of the Ca^{2+} level (Fig. 4A versus Fig. 2B). Pretreatment with EGTA also blocked these transient Ca^{2+} elevations (data not shown). The distinct Ca^{2+} signature activated by non-host legumes,

together with the lack of activation of *nod* genes (Fig. 4B), suggests the possibility of Ca^{2+} -mediated perception by *M. loti* of molecules other than *nod* gene inducers, such as non-specific chemoattractants or other signalling molecules, e.g. proteins [30,31] or plant cell wall fragments released during the detachment of border cells from the root tip [32], activating a different Ca^{2+} signalling pathway. Further confirmation of the specificity of the host plant-induced Ca^{2+} signalling comes from the complete absence of any detectable Ca^{2+} change and *nod* gene transcriptional activation by root exudates from a non-legume (tomato) (Fig. 4A and 4B).

Discussion

Even though Ca^{2+} -based signal transduction processes are well-established to underpin plant cell responses to rhizobial informational molecules, a possible involvement of Ca^{2+} as a messenger in rhizobia in response to plant symbiotic signals has not hitherto been considered. We approached this issue by constructing a *M. loti* strain expressing the bioluminescent Ca^{2+} indicator aequorin. The highly sensitive and reliable aequorin-based method is widely used to monitor the dynamic changes of $[Ca^{2+}]_i$ in both eukaryotic [33] and bacterial [18,16] living cells and represents to date the tool of choice for monitoring Ca^{2+} changes in cell populations [11]. The effectiveness of this recombinant technique has been verified at more than one level, and the results obtained demonstrate the utility of aequorin as a probe to study the early recognition events in rhizobium-legume interactions from the bacterial perspective.

The generation of a well-defined and reproducible Ca^{2+} transient in *M. loti* cells in response to root exudates of the host plant *L. japonicus* containing *nod* gene inducers is indicative of Ca^{2+} participation in sensing and transducing diffusible host-specific signals. It cannot be ruled out that the biphasic pattern of the Ca^{2+} trace (Fig. 2B), monitored by the aequorin method, may be due to an instantaneous synchronized Ca^{2+} increase in cells immediately after stimulation, followed by a sustained Ca^{2+} response probably due to the sum of asynchronous oscillations occurring in single cells. Ca^{2+} oscillations, considered as a universal mode of signalling in eukaryotic cells [34-36] have been proposed to occur in bacteria as well [37].

The significant inhibition of *nod* gene expression obtained when the Ca^{2+} elevation is blocked indicates that an upstream Ca^{2+} signal is required for *nod* gene activation. The Ca^{2+} dependence of *nod* gene expression strongly suggests that the $[Ca^{2+}]_i$ change, evoked by *L. japonicus* exudates, represents an essential prerequisite to convey the plant symbiotic message into rhizobia. All the above results fulfil the criteria required to demonstrate that a Ca^{2+} transient is a crucial intermediate in a stimulus-




Figure 3 (see legend on next page)

Figure 3 (see previous page)

Effect of EGTA on the Ca²⁺ response and nod gene expression induced by *L. japonicus* exudates. **A**, *M. loti* cells were treated with *L. japonicus* root exudates (black trace) or pretreated with 5 mM EGTA 10 min before adding *L. japonicus* root exudates (grey trace). **B**, Top: RT-PCR analysis of control cells (lane 1), cells treated for 1 h with *L. japonicus* root exudates (lane 2) and cells pretreated with 5 mM EGTA 10 min before treatment with *L. japonicus* exudates (lane 3). Bottom: Relative percentage of *nod* gene induction in response to *L. japonicus* exudates in *M. loti* cells pretreated (striped bars) or not (black bars) with 5 mM EGTA. Normalization of transcript abundance was done against 16S rRNA. Data are the means ± SEM of three independent experiments. **C**, Viability, monitored with the BacLight Bacterial Viability kit, of *M. loti* cells in control conditions or incubated with 5 mM EGTA for 1 h 10 min. As positive control, cells were treated with 70% isopropanol. Live cells fluoresce green, dead cells fluoresce red. Bar = 10 μm. **D**, Top: RT-PCR analysis of the expression of the housekeeping gene glutamine synthetase II (*GSII*) in *M. loti* cells in the absence (-) or presence (+) of 5 mM EGTA. Bottom: Relative transcript abundance of *GSII* was normalized against 16S rRNA. Bars represent SEM. **E**, Top: RT-PCR analysis of the inducible aequorin (*aeq*) gene in *M. loti* cells in the absence (-) or presence (+) of 5 mM EGTA and 1 mM IPTG. Bottom: Relative transcript abundance of *aeq* was normalized against 16S rRNA. Bars represent SEM.

response coupling [23] and confirm that Ca²⁺ signalling is operating in bacteria [5].

The inability of root exudates from non-host legumes and non legumes to duplicate the response induced by *L. japonicus* exudates (encoded in a distinct Ca²⁺ transient and downstream gene expression) further supports the symbiotic specificity of the host legume-induced Ca²⁺ signature. The possible relatedness to legume-rhizobium symbiosis of the signals contained in non-host legume exudates is supported by the absence of any Ca²⁺ response to non-legume exudates. In non-host legume root exudates *M. loti* cells may sense signalling molecules related

to the symbiotic process but not strictly specific to the compatible host-microsymbiont pair, which may enable rhizobia to distinguish non-host from compatible plants.

Plant root exudates contain a pool of molecules, both stimulatory and inhibitory, of potential relevance to the molecular signal exchange between the two partners [3]. The use of entire natural mixtures secreted by plant roots represents the first step in the evaluation of rhizobium reactions to plant factors, providing information on the global Ca²⁺ responses occurring in the bacterial partner early in the symbiosis, even before a physical contact between the two interacting organisms. Further insights





Figure 4

Monitoring [Ca²⁺]_i and nod gene expression in response to non-host legume and non-legume root exudates. Bacteria were challenged with root exudates from soybean (**A**, black trace; **B**, lane 2), *V. sativa* subsp. *nigra* (**A**, grey trace; **B**, lane 2) and tomato (**A**, light grey trace; **B**, lane 2). Control cells were treated with cell culture medium only (**B**, lane 1).

into the dynamics of the activated Ca²⁺ change may come from the comparison with the Ca²⁺ responses obtained by using fractionated root exudates or purified molecules. This would enable to assess the possible placement of the Ca²⁺ signal within the NodD-flavonoid gene expression paradigm [38] in different species of rhizobia.

Conclusion

The above results demonstrate that *M. loti* cells sense host plant symbiotic cues through Ca²⁺ and indicate that activation of *nod* genes requires an upstream Ca²⁺ signal. Transgenic rhizobium strains expressing aequorin can be used as a novel approach to the dissection of early events in legume-rhizobium symbiosis, that may shed light on a previously uninvestigated facet - bacterial Ca²⁺ signalling - of the two-way partner signal exchange and transduction.

Methods

Chemicals

Native coelenterazine was purchased from Molecular Probes (Leiden, The Netherlands). Molecular biology reagents were purchased from Promega Co. (Madison, WI, USA), Qiagen (Hilden, Germany) Clontech (Mountain View, CA, USA) and Invitrogen (Paisley, UK). Tetrionic acid was obtained from Titolchimica (Rovigo, Italy). Flavonoids (naringenin, luteolin, daidzein, quercetin dehydrate) and all other reagents were obtained from Sigma-Aldrich (St. Louis, MO, USA).

Bacterial strains and growth conditions

Mesorhizobium loti strain USDA 3147^T was kindly provided by Peter Van Berkum (USDA, Beltsville MD) and was grown in minimal BIII medium [39] with or without 30 µg/ml kanamycin, as appropriate, at 28 °C with shaking (170 rpm). *E. coli* was grown in LB medium at 37 °C.

Cloning of the apoaquorin gene and introduction into *M. loti*

The terms aequorin and apoaquorin refer to the bioluminescent protein with and without, respectively, the prosthetic group coelenterazine. The apoaquorin cassette, given by the apoaquorin cDNA fused to the first 27 nucleotides encoding hemoagglutinin (HA1-AEQ) [40] was amplified by PCR with primers designed to obtain a 5' *Xba*I site and to leave out the ATG start codon, already present into the P_{syn} promoter of the expression vector pDB1 [22]. The correct translation frame was maintained by adding a nucleotide between the 5' *Xba*I site and the apoaquorin gene. The primers used to obtain the apoaquorin cassette were: 5'-CCTACTCTAGATAAGCTT-TATGATGTTCCCT-3' and 5'TGATAGCATGCCAATTCAT-CAGTGTITTTAT-3'. PCR was run with the following parameters: 5 min at 94 °C as start step; 30 s at 94 °C, 30 s at 58 °C, 1 s at 72 °C for 30 cycle and 5 s at 72 °C as a final step using PLATINUM[®] Taq DNA polymerase (Invitro-

gen). To obtain a 3' *Xba*I site, the amplicon was then cloned into the pCR 2.1 plasmid by using TA Cloning[®] technology (Invitrogen), originating p2.1AEQ. Digestion with *Xba*I of this intermediate plasmid released the HA1-AEQ coding region, which was then ligated into the *Xba*I site of pDB1 under the control of the strong isopropylβ-D-thiogalactoside (IPTG)-inducible synthetic promoter P_{syn}. The apoaquorin gene containing construct (pAEQ80, see Additional file 1) was mobilized to *M. loti* 3147^T from *E. coli* by triparental conjugation using plasmid pRK2013 as helper [41]. Transconjugants were selected on BIII agar containing 50 µg/ml kanamycin.

Growth kinetics of the recombinant strain

To determine the effect of the plasmid presence and of apoaquorin expression on bacterial cell growth, *M. loti* wild-type or containing pAEQ80 (plus or minus IPTG) were grown in 30 ml of BIII medium (supplemented or not with 30 µg/ml kanamycin, as appropriate) as described above. Growth was determined by monitoring turbidity at 600 nm.

In vitro *L. japonicus* nodulation tests

In vitro nodulation studies were carried out as described by [42]. Briefly, seeds of *L. japonicus* B-129 GIFU were transferred after sterilization on 0.1% Jensen medium solidified with 1% agar. Inoculation with bacterial suspensions of *M. loti* wild-type or containing pAEQ80 (5 · 10⁷ cells/root) was carried out 4 days after seed germination. *Lotus* seedlings, before and after infection, were grown at 24 °C with 16 h light and 8 h dark. Growth and nodulation pattern were monitored for 4 weeks after inoculation. Microscopy observations were carried out with a Leica MZ16 stereomicroscope equipped with a DFC 480 photcamera. To check the actual occurrence of bacteria inside the nodules, they were squeezed and the content stained with 5 µg/ml 4',6-diamino-2-phenylindole (DAPI). Samples were observed with a Leica DMR fluorescence microscope. Images were acquired with a Leica IM500 digital camera.

Expression of apoaquorin

A loopful of *M. loti* USDA 3147^T pAEQ80 grown on BIII plates was used to inoculate 30 ml of BIII medium supplemented with 30 µg/ml kanamycin and 1 mM IPTG and grown at 28 °C overnight, until an absorbance at 600 nm of approximately 0.25 was reached (after about 18 h).

In vitro reconstitution of apoaquorin to aequorin

M. loti suspension cultures (300 ml) were grown to mid-exponential phase (A_{600 nm} = 0.25), pelleted by centrifugation at 3000 g for 10 min at 4 °C, washed twice with fresh medium, and finally resuspended in 2 ml reconstitution buffer (Tris-HCl 150 mM, EGTA 4 mM, supplemented with 0.8 mM phenylmethylsulfonyl fluoride, pH

8.0). Bacteria were lysed by 3 cycles (30 s each) of sonication at 35 Hz (Fisher Sonic, Artek Farmingdale, NY, USA), each followed by 30 s on ice. Non lysed bacteria were pelleted and discarded by centrifugation (1600 g for 15 min at 4°C). Protein concentration in the supernatant was estimated using the Bio-Rad (Hercules, CA) protein assay according to manufacturer's instructions. Total soluble proteins were resuspended at 1 µg/µl in reconstitution buffer and incubated with 1 mM β-mercaptoethanol and 5 µM coelenterazine for 4 h in the dark at 4°C. Aequorin luminescence was detected from 50 µl of the *in vitro* aequorin reconstitution mixture, containing 25 µg of total soluble protein diluted 1:2 with the same buffer and integrated for a 200 s time interval after the addition of an equal volume of 100 mM CaCl₂.

In vivo reconstitution of apoaequorin to aequorin

Mid-exponential phase cells (30 ml) were harvested by centrifugation at 2300 g for 15 min at room temperature and the cell pellet was washed twice in 5 ml BIII medium with intermediate centrifugation as described above. Cells were then incubated in BIII medium containing 5 µM coelenterazine in the dark for 1 h 30 min under shaking. After two washes as above, cells were resuspended in BIII medium and allowed to recover for 10 min prior to Ca²⁺ measurement experiments.

Root exudate production

Seeds of *Lotus japonicus* GIFU ecotype, soybean, *Vicia sativa* subsp. *nigra* and tomato were surface sterilized and allowed to germinate for three days on moistened filter paper at 24°C in the dark. Subsequently, seedlings were transferred aseptically on polystyrene grids covered with nylon meshes in sterile plastic containers containing different volumes of sterile H₂O, depending on the seed and seedling size (on average 5 ml of H₂O per seedling). After 3 weeks of germination crude root exudates were collected, filtered and lyophilized. The pellet was resuspended in BIII medium (50 µl per single root exudate) for cell treatments.

Ca²⁺ measurements with recombinant aequorin

Aequorin light emission was measured in a purpose-built luminometer. Bacteria (50 µl) were placed, after aequorin reconstitution, in the luminometer chamber in close proximity to a low-noise photomultiplier, with a built-in amplifier discriminator. The output of the discriminator was captured by a THORN-EMI photon counting board (Electron Tubes Limited, Middlesex, UK) and the luminescence data were converted off-line into Ca²⁺ concentration values by using a computer algorithm based on the Ca²⁺ response curve of aequorin [40]. All stimuli were administered to cells by using a light-tight syringe through the luminometer port. The experiments were terminated by lysing the cells with 15% ethanol in a Ca²⁺-rich solu-

tion (0.5 M CaCl₂ in H₂O) to discharge the remaining aequorin pool. For experiments performed in the presence of different external Ca²⁺ concentrations, cells were extensively washed and resuspended in buffer A (25 mM Hepes, 125 mM NaCl, 1 mM MgCl₂, pH 7.5), as described by [16]. When needed, cells were pretreated for 10 min with 5 mM EGTA.

Bacterial cell viability assay

Bacterial cell viability was monitored by the LIVE/DEAD® BacLight™ Bacterial Viability kit (Molecular Probes), according to manufacturer's instructions. This fluorescence-based assay uses a mixture of SYTO 9 and propidium iodide stains to distinguish live and dead bacteria. Bacteria with intact cell membranes stain fluorescent green, whereas bacteria with damaged membranes stain fluorescent red. Samples were observed with a Leica 5000B fluorescence microscope. Images were acquired with a Leica 300F digital camera using the Leica Application Suite (LAS) software.

Semi-quantitative RT-PCR experiments

M. loti cells grown to mid-exponential phase and treated as for Ca²⁺ measurement experiments (see above) were incubated for 1 h with plant root exudates, tetrionic acid or cell culture medium only (as control). To stabilize RNA, bacteria were treated with the RNA protect Bacteria Reagent (Qiagen). Bacterial cell wall was then lysed with 1 µg/ml lysozyme (Sigma) in TE buffer. Total RNA was first extracted using RNeasy Mini kit (Qiagen) and, after DNase I treatment (Promega), quantified. RNA (5 µg) was primed with Random Decamers (Ambion), reverse transcribed with PowerScript Reverse Transcriptase (Clontech) and diluted 1:5. 5 µl of diluted first-strand cDNA were used as a template in a 50 µl PCR reaction solution. Reverse transcription (RT)-PCR was performed with 5 µl diluted first-strand cDNA. The oligonucleotide primers were designed against *nodA*, *nodB*, *nodC* and glutamine synthetase II (*GSI*) sequences from *M. loti* [43] and the aequorin gene (*aeq*) from *Aequorea victoria* [44], using Primer 3 software. To amplify 16S rRNA gene, Y1 and Y2 primers were used [45].

The thermal cycler was programmed with the following parameters: 20 s at 94°C, 30 s at 68°C and Advantage 2 Polymerase mix (Clontech) was used as Taq polymerase. PCR reactions were allowed to proceed for different number of cycles to determine the exponential phase of amplification. Densitometric analysis of ethidium bromide-stained agarose gels (0.5 µg/ml) was performed using QuantityOne software (Bio-Rad). RT-PCR experiments were conducted in triplicate on three independent experiments. The primer sequences used to obtain amplicons were: 5'-TATGAGCCGACCGGAGCCTTTAAT-3' and 5'-CCGTATAGACCGAGTTCAGCGACAA-3' for *nodA*, 5'-

ATACTCGATGTGCTGGCGCAAAAT-3' and 5'-GCCTGGTTCGCTCAAATACTTCAC-3' for *nodB*, 5'-CCACCTTACGATCCTGATGCTGAAA-3' and 5'-CAATATTCTGGCCAATCACGTCCAA-3' for *nodC*, 5'-ACCGAGACTTACGGCATCGACATC-3' and 5'-GCGACGCCATAGCTAAACTTGTTC-3' for *GSII*, 5'-TAACCTTGGAGCAACACCTGAGCAA-3'

5'-ATACGGATGAGCGTTGGTTTCGTTTT-3' for aequorin, Y1 (5'-TGGCTCAGAACGAACGCTGGCGGC-3') and Y2 (5'-CCCACTGCTGCCTCCCGTAGGAGT-3') for 16S rRNA. Amplicons were sequenced by BMR Genomics (Padova, Italy).

Authors' contributions

RM cloned the apoaquorin gene, carried out the RT-PCR experiments and participated in the Ca²⁺ measurement experiments. SA and AS introduced the apoaquorin gene into *E. coli* and *M. loti*. LN performed the nodulation studies, prepared the plant root exudates and was involved in acquisition and interpretation of Ca²⁺ measurement data. MP and LN conceived of the study, designed the experiments and wrote the paper. AS helped with manuscript discussion and participated in its editing. All authors read and approved the final manuscript.

Additional material

Additional file 1

Map of the apoaquorin-expressing plasmid pAEQ80. Abbreviations: P, IPTG-inducible synthetic promoter (*P_{syn}*); HA1-AEQ, cloned apoaquorin cDNA with hemoagglutinin epitope; Km^R, kanamycin resistance gene; *lacI^q*, constitutive *lac* repressor gene. Relevant restriction endonuclease sites are also shown.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-206-S1.tiff>]

Additional file 2

Validation of the aequorin-expressing *M. loti* experimental system. A, Analysis of aequorin expression in *M. loti* based on an in vitro reconstitution assay. Data are the means \pm SEM of three experiments. B, Effect of pAEQ80 plasmid and expressed recombinant apoaquorin on *M. loti* cell growth. Data are the means of two independent experiments. C, Nodulated root of *L. japonicus* 4 weeks after inoculation with the recombinant *M. loti* strain. Bar = 2 mm. D, DAPI staining of *M. loti* cells USDA 3147^T pAEQ80 squeezed from a young nodule. Bar = 10 μ m. E, Monitoring of intracellular Ca²⁺ concentration ([Ca²⁺]_i) in resting *M. loti* cells grown to mid-exponential phase.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-206-S2.tiff>]

Acknowledgements

This work was supported by Progetti di Ricerca di Ateneo 2006 from the University of Padova (prot. CPDA063434) and Ricerca Scientifica fondi quota EX 60% 2007-2009 (prot. 60A06-9994/07, 921/08 and 5430/09) to

LN. We are grateful to M. Brini (Padova, Italy) for kindly providing the apoaquorin cassette and for helpful discussion on Ca²⁺ measurement experiments, and to D. Sanders (York, UK) for critical reading of the manuscript and insightful comments. We also thank J. Stougaard (Aarhus, Denmark), S. Varotto (Padova, Italy) and Vergerio Mangimi S.R.L. (Padova, Italy) for the kind gift of *L. japonicus*, soybean and *V. sativa* seeds, respectively.

References

- Oldroyd GED, Downie JA: **Coordinating nodule morphogenesis with rhizobial infection in legumes.** *Annu Rev Plant Biol* 2008, **59**:519-546.
- Garg N, Geetanjali: **Symbiotic nitrogen fixation in legume nodules: process and signaling. A review.** *Agron Sustain Dev* 2007, **27**:59-68.
- Cooper JE: **Early interactions between legumes and rhizobia: disclosing complexity in a molecular dialogue.** *J Appl Microbiol* 2007, **103**:1355-1365.
- Norris V, Grant S, Freestone P, Canvin J, Sheikh FN, Toth I, Trinei M, Modha K, Norman RL: **Calcium signalling in bacteria.** *J Bacteriol* 1996, **178**:3677-3682.
- Dominguez DC: **Calcium signalling in bacteria.** *Mol Microbiol* 2004, **54**:291-297.
- Case RM, Eisner D, Gurney A, Jones O, Muallem S, Verkhatsky A: **Evolution of calcium homeostasis: from birth of the first cell to an omnipresent signalling system.** *Cell Calcium* 2007, **42**:345-350.
- Kung C, Blount P: **Channels in microbes: so many holes to fill.** *Mol Microbiol* 2004, **53**:373-380.
- Yang K: **Prokaryotic calmodulins: recent developments and evolutionary implications.** *J Mol Microbiol Biotechnol* 2001, **3**:457-459.
- Michiels J, Xi C, Verhaert J, Vanderleyden J: **The functions of Ca²⁺ in bacteria: a role for EF-hand proteins?** *Trends Microbiol* 2002, **10**:87-93.
- Mithöfer A, Mazars C: **Aequorin-based measurements of intracellular Ca²⁺-signatures in plant cells.** *Biol Proced Online* 2002, **4**:105-118.
- Rudolf R, Mongillo M, Rizzuto R, Pozzan T: **Looking forward to seeing calcium.** *Nat Rev Mol Cell Biol* 2003, **4**:579-586.
- Nelson G, Kozlova-Zwinderman O, Collis A, Knight MR, Fincham JRS, Stanger CP, Renwich A, Hessing JGM, Punt PJ, Hondel CAMJJ van den, Read ND: **Calcium measurement in living filamentous fungi expressing codon-optimized aequorin.** *Mol Microbiol* 2004, **52**:1437-1450.
- Watkins NJ, Knight MR, Trewavas AJ, Campbell AK: **Free calcium transients in chemotactic and non-chemotactic strains of Escherichia coli determined by using recombinant aequorin.** *Biochem J* 1995, **306**:865-869.
- Jones HE, Holland IB, Baker HL, Campbell AK: **Slow changes in cytosolic free Ca²⁺ in Escherichia coli highlight two putative influx mechanisms in response to changes in extracellular calcium.** *Cell Calcium* 1999, **25**:265-274.
- Jones HE, Holland IB, Campbell AK: **Direct measurements of free Ca²⁺ shows different regulation of Ca²⁺ between the periplasm and the cytosol of Escherichia coli.** *Cell Calcium* 2002, **32**:183-192.
- Campbell AK, Naseem R, Wann K, Holland IB, Matthews SB: **Fermentation product butane 2,3-diol induces Ca²⁺ transients in E. coli. through activation of lanthanum-sensitive Ca²⁺ channels.** *Cell Calcium* 2007, **41**:97-106.
- Campbell AK, Naseem R, Holland IB, Matthews SB, Wann KT: **Methylglyoxal and other carbohydrate metabolites induce lanthanum-sensitive Ca²⁺ transients and inhibit growth in E. coli.** *Arch Biochem Biophys* 2007, **468**:107-113.
- Torreccilla I, Leganés F, Bonilla I, Fernández-Piñas F: **Use of recombinant aequorin to study calcium homeostasis and monitor calcium transients in response to heat and cold shock in cyanobacteria.** *Plant Physiol* 2000, **123**:161-175.
- Torreccilla I, Leganés F, Bonilla I, Fernández-Piñas F: **Calcium transients in response to salinity and osmotic stress in the nitrogen-fixing cyanobacterium Anabaena sp. PCC 120, expressing cytosolic aequorin.** *Plant Cell Environ* 2001, **24**:641-648.

20. Torrecilla I, Leganés F, Bonilla I, Fernández-Piñas F: **A calcium signal is involved in heterocyst differentiation in the cyanobacterium *Anabaena* sp. PCC7120.** *Microbiology* 2004, **150**:3731-3739.
21. Torrecilla I, Leganés F, Bonilla I, Fernández-Piñas F: **Light-to-dark transitions trigger a transient increase in intracellular Ca^{2+} modulated by the redox state of the photosynthetic electron transport chain in the cyanobacterium *Anabaena* sp. PCC7120.** *Plant Cell Environ* 2004, **27**:810-819.
22. Alberghini S, Filippini R, Marchetti E, Dindo ML, Shevelev AB, Battisti A, Squartini A: **Construction of a *Pseudomonas* sp. derivative carrying the *cry9Aa* gene from *Bacillus thuringiensis* and a proposal for new standard criteria to assess entomocidal properties of bacteria.** *Res Microbiol* 2005, **156**:690-699.
23. Sanders D, Brownlee C, Harper JF: **Communicating with calcium.** *Plant Cell* 1999, **11**:691-706.
24. Falciatore A, d'Alcalá MR, Croot P, Bowler C: **Perception of environmental signals by a marine diatom.** *Science* 2000, **288**:2363-2366.
25. Estévez J, Soria-Díaz ME, de Córdoba FF, Morón B, Manyani H, Gil A, Thomas-Oates J, van Brussel AA, Dardanelli MS, Sousa C, Megías M: **Different and new Nod factors produced by *Rhizobium tropici* CIAT899 following Na^+ stress.** *FEMS Microbiol Lett* 2009, **293**:220-231.
26. Suzuki H, Sasaki R, Ogata Y, Nakamura Y, Sakurai N, Kitajima H, Kanaya S, Aoki K, Shibata D, Saito K: **Metabolic profiling of flavonoids in *Lotus japonicus* using liquid chromatography Fourier transform ion cyclotron resonance mass spectrometry.** *Phytochemistry* 2008, **69**:99-111.
27. López-Lara IM, Berg JDJ van der, Thomas-Oates JE, Glushka J, Lugtenberg BJJ, Spaink HP: **Structural identification of the lipo-chitin oligosaccharide nodulation signals of *Rhizobium loti*.** *Mol Microbiol* 2004, **15**:627-638.
28. Saeki K, Kouchi H: **The *Lotus* symbiont, *Mesorhizobium loti*: molecular genetic techniques and application.** *J Plant Res* 2000, **113**:457-465.
29. Gagnon H, Ibrahim RK: **Aldonic acids: a novel family of nod gene inducers of *Mesorhizobium loti*, *Rhizobium lupini*, and *Sinorhizobium meliloti*.** *Mol Plant-Microbe Interact* 1998, **11**:988-998.
30. De-la-Peña C, Lei Z, Watson BS, Summer LW, Vivanco JM: **Root-microbe communication through protein secretion.** *J Biol Chem* 2008, **283**:25247-25255.
31. Wen F, VanEtten HD, Tsapraillis G, Hawes MC: **Extracellular proteins in pea root tip and border cell exudates.** *Plant Physiol* 2007, **143**:773-783.
32. Zhu Y, Pierson LS III, Hawes MC: **Induction of microbial genes for pathogenesis and symbiosis by chemicals from root border cells.** *Plant Physiol* 1997, **115**:1691-1698.
33. Brini M, Pinton P, Pozzan T, Rizzuto R: **Targeted recombinant aequorins: tools for monitoring $[Ca^{2+}]$ in the various compartments of a living cell.** *Microsc Res Tech* 1999, **46**:380-389.
34. Berridge MJ: **Calcium oscillations.** *J Biol Chem* 1990, **265**:9583-9586.
35. Dodd AN, Jakobsen MK, Baker AJ, Telzerow A, Sui-Wen Hou SW, Laplaze L, Barrot L, Poethig RS, Haseloff J, Webb AAR: **Time of day modulates low-temperature Ca^{2+} signals in *Arabidopsis*.** *Plant J* 2006, **48**:962-973.
36. McAinsh MR, Pittman JK: **Shaping the calcium signature.** *New Phytol* 2009, **181**:275-294.
37. Wolf DM, Arkin AP: **Motifs, modules and games in bacteria.** *Curr Opin Microbiol* 2003, **6**:125-134.
38. Schlaman HRM, Okker RJH, Lugtenberg BJJ: **Regulation of nodulation gene expression by NodD in rhizobia.** *J Bacteriol* 1992, **174**:5177-5182.
39. Dazzo FB: **Leguminous root nodules.** In *Experimental Microbial Ecology* Edited by: Burns R, Slater J. Oxford: Blackwell Scientific Publications; 1982:431-446.
40. Brini M, Marsault R, Bastianutto C, Alvarez J, Pozzan T, Rizzuto R: **Transfected aequorin in the measurement of cytosolic Ca^{2+} concentration ($[Ca^{2+}]_i$).** *J Biol Chem* 1995, **270**:9896-9903.
41. Figurski DH, Helinski DR: **Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans.** *Proc Natl Acad Sci USA* 1979, **76**:1648-52.
42. Barbulova A, Chiurazzi M: **A procedure for *Lotus japonicus* in vitro nodulation studies.** In *Lotus japonicus Handbook* Edited by: Márquez AJ, Stougaard J, Udvardi M, Parniske M, Spaink H, Saalbach G, Webb J, Chiurazzi M. Berlin, Springer; 2005:83-86.
43. Kaneko T, Nakamura Y, Sato S, Asamizu E, Kato T, Sasamoto S, Watanabe A, Idesawa K, Ishikawa A, Kawashima K, Kimura T, Kishida Y, Kiyokawa C, Kohara M, Matsumoto M, Matsuno A, Mochizuki Y, Nakayama S, Nakazaki N, Shimpo S, Sugimoto M, Takeuchi C, Yamada M, Tabata S: **Complete genome structure of the nitrogen-fixing symbiotic bacterium *Mesorhizobium loti*.** *DNA Res* 2000, **7**:331-338.
44. Inouye S, Noguchi M, Sakaki Y, Takagi Y, Miyata T, Iwanaga S, Miyata T, Tsuji FI: **Cloning and sequence analysis of cDNA for the luminescent protein aequorin.** *Proc Natl Acad Sci USA* 1985, **82**:3154-3158.
45. Young JPW, Downer HL, Eardly BD: **Phylogeny of the phototrophic rhizobium strain BTAi1 by polymerase chain reaction-based sequencing of a 16S rRNA gene segment.** *J Bacteriol* 1991, **173**:2271-2277.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

