

The Effect of Olea Ointment on Post-Episiotomy Pain Severity in Primiparous Women: A Paralleled Randomized Controlled Clinical Trial

Abstract

Background: Episiotomy is the most commonly performed surgical procedure during delivery, and its associated pain is a major problem in obstetrics. This study aimed to determine the effect of Olea ointment on the severity of post-episiotomy pain in primiparous women. **Materials and Methods:** This paralleled randomized controlled clinical trial was performed on 73 primiparous women in Al-Zahra hospital in Rasht, Iran in 2017-2018. Women were randomly allocated into intervention ($n = 39$) and control ($n = 34$) groups. The first intervention was performed 4 h after the episiotomy. This continued every 8 h for 10 days. The pain intensity of episiotomy was assessed by the Visual Analogue Scale (VAS) before intervention, 2 and 24 h after the beginning of intervention, and 5 and 10 days after childbirth. Descriptive and inferential statistics (Mann-Whitney, Fisher exact test, Independent t -test, Friedman test, and Chi-square) were performed for statistical analysis. **Results:** There were no significant differences among two groups in terms of demographic and obstetrics characteristics. The severity of pain in intervention and control group before the intervention was not statistically significant, but the variable depicted a meaningful difference between the groups 2 h after the intervention ($U = 483.50, p = 0.021$), 24 h after the intervention ($U = 489.50, p = 0.019$), as well as the 5th day ($U = 112.50, p < 0.001$) and 10th day postpartum ($U = 136.50, p < 0.001$). **Conclusions:** Based on the findings, Olea ointment could be used effectively for reducing of episiotomy pain. Similar studies are recommended.

Keywords: *Episiotomy, Iran, Olea, pain, postpartum period*

Introduction

Episiotomy is defined as perineal incisions in the second stage of labor which is performed for facilitation and shortening of the duration of the second stage of delivery, preventing possible damages to perineum, and pelvic organs, sexual dysfunctions, and severe perineal tears.^[1-6] In fact, episiotomy is counted as the second most common surgical experiment during delivery after cutting down of the umbilical cord.^[7]

Episiotomy is of different prevalence rates in different countries, under 10% in countries such as Sweden and Denmark, more than 50% in countries such as Netherlands and Portugal,^[8] and 100% in the Asian country (Taiwan).^[4] According to some studies, the second stage of delivery is longer in Asian women, and thus, perineal tears occur with a greater frequency.^[9,10] Even though no exact statistics is available on prevalence of episiotomy in Iran, its

distribution is assessed by cross-sectional studies as 88.32% in Mashhad, and 97.3% in Tehran.^[11]

The benefits of episiotomy is still subject to doubt.^[2-4] Thus, as recommended by World Health Organization (WHO), episiotomy should be reduced to 10%,^[12] and its application should be limited to high-risk cases.^[2]

Many women suffer from short-term side-effects of episiotomy, and more than 20% of women experience its long-term side-effects.^[13] Perineal pain is the most common complication in episiotomy, and it ranges from 96.4% on the first 24 h after the labor to 63% on the second day and 25% at the end of puerperium.^[10] This pain can be accompanied by some morbidities and causes trouble in sitting, breastfeeding, and formation of relationship between mother and infant.^[4,5] Thus, relieving the perineal pain is of great importance and, to this end, non-medical methods as well as medical

Fatemeh Jafarzadeh-Kenarsari¹, Shokoufeh Torkashvand², Bahare Gholami-Chaboki³, Yalda Donyaiei-Mobarrez⁴

¹Department of Midwifery and Reproductive Health, Social Determinants of Health Research Center (SDHRC), School of Nursing and Midwifery, Guilan University of Medical Sciences, Rasht, Iran, ²Department of Midwifery, Alborz Hospital, Karaj, Iran, ³Department of Biostatistics, School of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran, ⁴Department of Midwifery, School of Nursing and Midwifery, Guilan University of Medical Sciences, Rasht, Iran

Address for correspondence: Ms. Shokoufeh Torkashvand, Department of Midwifery, Alborz Hospital, Karaj, Iran. E-mail: shokufetorkashvand@gmail.com

Access this article online

Website: www.ijnmrjournal.net

DOI: 10.4103/ijnmr.IJNMR_151_18

Quick Response Code:



How to cite this article: Jafarzadeh-Kenarsari F, Torkashvand S, Gholami-Chaboki B, Donyaiei-Mobarrez Y. The effect of olea ointment on post-episiotomy pain severity in primiparous women: A paralleled randomized controlled clinical trial. Iranian J Nursing Midwifery Res 2019;24:348-54.

Received: October, 2018. **Accepted:** June, 2019.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

therapies such as application of lidocaine, phenytoin, normal saline, and iodine, and also oral medications or herbal remedies such as olives, honey, saffron, turmeric, cinnamon, spring flowers, chamomile, aloe vera, bromillin, and lavender can be employed.^[10,14,15] Sesame oil also has at least seven pain-relieving substances.^[16]

Olea ointment is an herbal ointment that is a combination of honey (33.4%), olive oil (33.3%), and sesame oil (33.3%). This ointment prevents infection, and by reducing the pain and inflammation, the healing process accelerates the formation of new tissue and facilitates the debridement of the necrotic tissue.^[17] For instance, a study showed that the extra virgin olive oil has been effective in the treatment of primary dysmenorrhea.^[18] Also, another study showed that the use of honey cream has not been associated with the reduction of pain in the episiotomy region.^[19]

The findings of the studies on herbal medicine are contradictory, and confirmation of the therapeutic effects of herbs requires more laboratory studies and clinical trials. Considering the importance of the reduction of pain and faster return of the mothers to daily activities, the researchers aimed to determine the effect of Olea ointment on the severity of episiotomy pain in primiparous women.

Materials and Methods

This paralleled randomized clinical trial was performed on primiparous women attending Al-Zahra hospital in

Rasht (North of Iran), who had normal vaginal delivery with episiotomy. IRCT code number was 2017020332374N1. The sample size was obtained by reference to Jahdi *et al.* study,^[20] by a statistical formula with a 95% confidence interval and a 90% test power as at least 36 individuals in each group. Taking into account 15% drop-out, 41 women were considered in each group. Thus, 82 primiparous women were selected in LDR ward (labour-delivery room) by convenient sampling and randomly allocated into intervention ($n = 41$) and control ($n = 41$) groups (by simple random sampling method and selecting cards A or B, representing intervention and control groups respectively). A total of four individuals were excluded from the study due to the opening of the wound and five patients for not referring back to the hospital; two of them were in the Olea intervening group (one due to not referring and one due to wound opening) and seven people in control group (four due to not referring and three due to wound opening). At the end of the intervention, the intervention and control groups consisted of 39 and 34 participants, respectively [Figure 1]. Sampling was conducted between December 2017 and May 2018. The inclusion criteria were Iranian ethnicity, being primiparous and aged 18-35 years old, ability to read and write, willingness to participate, live term singleton pregnancy with cephalic presentation and mediolateral episiotomy without laceration, lack of infection, and visible lesions in the perineum, and constipation, no prolonged premature rupture of the membrane, the first, second, and third stage of labor less than 14 h, 2 h, and half an hour,




Figure 1: Consort diagram

respectively, the absence of manual removal of placenta, the absence of postpartum hemorrhage, lack of perineal hematoma, and the new-born weight between 2,500 and 4,000 g. The exclusion criteria included unwillingness to continue participating in the study, use of other remedies affecting wound healing during the study, inappropriate use of the drug (more than 2 days), being allergic to Olea ointment, no referral to the hospital on the 5th and 10th days after childbirth, sexual intercourse in the first 5 days after childbirth, and resectioning of the perineum after repair of episiotomy.

Data collection scales included: 1. Demographic information form (age, education, occupation, economic status). 2. Obstetrics and delivery information form (gestational age, the number of pregnancies, the number of abortions, the gender and weight of new-borns, new-born's circumference of head and duration of the first, second, and third stages of labor). 3. The Visual Analogue Scale (VAS), which is a 10-cm paper ruler ranging from 0 to 10 where 0 indicates lack of pain, scores 1-3 indicate mild pain, scores 4-6 indicate moderate pain, and 7-10 indicate severe pain symptoms. VAS is a standard tool and its validity and reliability have been confirmed (correlation = 0.71-0.78, the intra-class coefficient = 0.95, and concurrent validity, $r > 0.8$; $p < 0.01$) in various studies.^[20-23] Data were collected through interviews, patients' files, and study, observation, and examination which were directly conducted by the researchers. Education was provided on the perineal and suture care, personal health, nutrition, and physical activity in face-to-face method and with pamphlets, alike for both groups. Before intervention, the pain intensity of the episiotomy was measured and recorded for each group based on the VAS.

Olea ointment was ordered in 30-g tubes containing equal ratios of olive oil, sesame oil, and honey from Farateb Yazd Company in Iran and were delivered by post. The first intervention was performed 4 h after the episiotomy. The samples of intervening group were asked to place a specific amount of Olea ointment (one-third of a finger) on the sutures after washing and drying the perineum area, covering it, and using a clean sanitary pad after 1-2 min. This procedure was continued every 8 h for 10 days after childbirth. It should be noted that the first intervention was carried out by the researchers and the subsequent interventions by the trained mothers. Also, the subjects in the intervention group were emphasized to use a physiologic serum (2 times a day for 10 days) to wash the perineal area along with the use of Olea ointment at home. The control group received only physiologic serum twice a day until the 10th day after delivery. We used a bottle of normal saline 0.9%, 1000cc, and a tube of Olea ointment containing 30 g effective materials for each person in intervention group. Also, we used a bottle of normal saline 0.9%, 1000cc for each participant in control group. The severity of the pain was measured and recorded

by the researchers 2 h after the first intervention, 24 h after the first intervention, then, on days 5 and 10 after delivery, and if the wound was not repaired, on the 14th day after the delivery. It should be noted that we checked women by phone to assure they used the drug properly and regularly. The patients were also asked to contact the researchers if any problems and complications would occur.

After data collection, statistical analysis was performed by SPSS version 16 (SPSS, Inc Sep 13, 2007); then descriptive statistics (quantitative variables were described by mean and standard deviation and qualitative variables with percentages and numbers), and inferential statistics (Mann-Whitney, Fisher exact test, Independent *t*-test, Friedman test, and Chi-square) were performed. To check the effect of time and compare the values, obtained over the 5 recorded times, since our variable was to be rated, we used the Friedman test separately in each group. In cases in which the data had no normal distribution, nonparametric statistical tests were used to analyse the data. A significance level of 0.05 was considered for all tests.

Ethical considerations

The study was approved in the Ethics Committee of Guilan University of Medical Sciences (IR.GUMS.REC.1396.191), Rasht, Iran, and also, it was registered at the Iranian Centre for Clinical Trials. All participants in this study were selected after obtaining their written informed consents, explaining the study objectives, as well as assuring the confidentiality of their research data.

Results

The demographic and obstetrics characteristics of the subjects were presented in Tables 1 and 2. There were no significant differences among the control and intervention groups in terms of other demographic characteristics ($p > 0.005$), except for the length of the first stage of labor ($Z = -2.81$, $p = 0.005$). Mann-Whitney test was used to compare the groups in terms of the pain severity scores before and after the intervention. The results presented in Table 3 show that the severity of pain in two groups of Olea ointment and physiologic serum before the intervention was not statistically significant ($p = 0.667$), but the variable depicted a meaningful difference between the two groups 2 h after the intervention ($U = 483.50$, $p = 0.021$), 24 h after the intervention ($U = 489.50$, $p = 0.019$), as well as on the 5th day ($U = 112.50$, $p < 0.001$) and 10th day postpartum ($U = 136.5$, $p < 0.001$). Effect of group was statistically significant, indicating that the odds of feeling mild pain would be higher [$\exp(0.62) = 1.87$] among women who received Olea ointment, compared to these who received physiologic serum. Effects of time and length of the first stage of labor were not statistically significant [Table 4]. In other words, administration of Olea ointment decreased the episiotomy pain severity.

Table 1: Qualitative demographic information and obstetrics characteristics of the study subjects in two groups

Variables	Properties	Olea ointment N (%)	Physiologic serum N (%)	Tests		
				Chi-square		
				χ^2	df	p
Age	20-25	19 (48.72)	10 (29.41)	3.52	2	0.172
	26-30	10 (25.64)	15 (44.12)			
	31-35	10 (25.64)	9 (26.47)			
Fisher's Exact Test						
Education	Primary school	6 (15.38)	3 (8.82)	1.29		0.770
	Elementary school	10 (25.64)	7 (20.59)			
	High school	17 (43.60)	17 (50.00)			
	University	6 (15.38)	7 (20.59)			
Fisher's Exact Test						
Job	Housewife	31 (79.49)	29 (85.30)	0.51		0.902
	Home employed	3 (7.69)	2 (5.88)			
	Working outside the home	5 (12.82)	3 (8.82)			
Fisher's Exact Test						
Economic Condition	Weak	6 (15.38)	3 (8.82)	1.46		0.555
	Moderate	26 (66.67)	27 (79.41)			
	Good	7 (17.95)	4 (11.76)			
Chi-square						
Fetal sex	Female	21 (53.85)	16 (47.06)	0.33	1	0.642
	Male	18 (46.15)	18 (52.94)			

Table 2: Quantitative demographic information and obstetrics characteristics of the study subjects in two groups

Variables	Olea Ointment Mean (SD)	Physiologic serum Mean (SD)	Tests		
			Mann-whitney		
			U	p	
Gestational Age	39.08 (0.84)	39.21 (0.88)	588.50	0.381	
Fisher's Exact Test					
Gravida					
One	39 (100.00)	32 (94.12)		0.213	
Two	0 (0.00)	2 (5.88)			
Fisher's Exact Test					
Abortion					
Zero	39 (100.00)	33 (97.10)			
One	0 (0.00)	1 (2.90)		0.466	
Mann-whitney					
			U	p	
Duration of the first stage of labor (h)	3.98 (1.23)	3.29 (0.42)	410	0.005*	
Duration of the second stage of labor (min)	1.00 (0.44)	0.91 (0.29)	534.50	0.148	
Duration of the third stage of labor (min)	0.07 (0.03)	0.07 (0.03)	638.50	0.745	
Newborn's head circumference	34.91 (0.84)	35.34 (1.13)	528	0.118	
Independent t-student					
			t	df	p
Newborn's Weight	3375.13 (353.72)	3328.82 (278.78)	0.61	71	0.541

*Statistically significant in confidence level $\alpha=0.05$

Discussion

The present study was the first randomized controlled clinical trial in an Iranian context aiming to determine the effect of Olea ointment on the severity of pain after episiotomy in primiparous women. Based on the results obtained by this study, Olea ointment was effective in

reducing pain at 2 h and 24 h after the beginning of intervention, as well as on the days 5 and 10 after delivery. A study found that when olive oil sitz bath was used in perineal injury after delivery, there was a significant difference between the intervention and control groups regarding the intensity of pain on the 5th and 10th day after delivery ($p < 0.05$), and olive oil was effective in

Table 3: Comparison of pain severity scores before and after the intervention in two groups

Time	Olea Ointment Mean (SD)	Physiologic serum Mean (SD)	Tests Mann-Whitney	
			U	p
Before intervention	2.33 (0.48)	2.26 (0.58)	630.50	0.667
2 h after intervention	2.08 (0.53)	2.38 (0.60)	483.50	0.021
24 h after intervention	1.92 (0.58)	2.82 (3.42)	489.50	0.019
5 days after delivery	1.03 (0.16)	1.97 (0.52)	112.50	<0.001
10 days after delivery	1.00 (0.0001)	1.79 (0.41)	136.50	<0.001

Table 4: The results of Generalized Estimating Equations (GEE) tests to assess the effects of groups, time, and demographic variables on pain severity

Parameter	Parameter estimate	Standard error	95% confidence interval		p
Pain					
Low	-2.17	0.67	-3.49	-0.85	0.001
Moderate	-0.72	0.58	-1.87	0.41	0.212
Group					
Olea ointment	0.62	0.3	0.02	1.22	0.042
Physiologic serum	ref	-	-	-	-
Time					
Length of the first stage of labor	0.078	0.09	-0.11	0.27	0.430
	-0.01	0.13	-0.26	0.24	0.939

reducing the pain.^[24] The results of another study aiming at “comparing the effect of pure and gross honey on the severity of primary dysmenorrhea, the amount, duration and interval between two menstrual cycles in female students with primary dysmenorrhea” showed that the consumption of pure honey has resulted in a significant reduction in the severity of pain and the amount of menstrual bleeding in women with primary dysmenorrhea ($p = 0.002$).^[25] According to a study aiming at “assessing the effect of back massage with sesame oil on the pain and length of delivery in primiparous women,” massage with sesame oil resulted in pain reduction at all stages of labor, especially the third stage ($p < 0.001$).^[16] It seems that fatty acids in sesame oil increase analgesia by decreasing neurotransmission, and by inhibiting the prostaglandin synthesise E2 enzyme, it increases the threshold for pain perception.^[26] Olea ointment is a mixture of olive oil, honey, and sesame oil. So, it contains the useful effects of each of these substances^[17] and can be helpful to decrease pain severity; however, in a study aiming at “comparing the effect of olive oil, aloe vera extract, and breast milk on healing of breast fissure in lactating mothers,” olive oil, aloe vera extract, and breast milk decreased pain severity and breast fissure in mothers. But, the aloe vera extract was more effective than olive oil and breast milk.^[27] According to a study aiming at “comparing the effect of honey and mefenamic acid on severe pain in women with primary dysmenorrhea,” there was no significant difference between the severity of pain in the first and second months of treatment, and mefenamic acid and honey had similar effects on reducing the pain

in women with primary dysmenorrhea.^[28] Also, based on the results of a study aiming at “comparing the effect of honey cream and Phenytoin cream on the pain severity of episiotomy in nulliparous women” in a double-blind, triplicate clinical trial, the severity of pain on the 7th and 14th days postpartum were not significantly different in the three groups.^[19] The possible reason for that was consumption of pain killers for 14 days after delivery by most of the subjects of the study in all 3 groups.

In the present study, opening of the episiotomy wound occurred in three participants of the control group, whereas this occurrence was observed in one participant of the intervention group. This may be due to the possible effect of Olea ointment on episiotomy wound healing. However, further similar studies in this area are recommended with the use of placebo.

When interpreting the results of the present study, a few weaknesses and limitations need to be considered. Lack of using the placebo in the control group might be considered as a weakness because the observed effect of Olea ointment may be unreal, and replication of similar studies with placebo is recommended. Given that the threshold for pain tolerance varies from person to person due to individual and social differences, pain intensity may be less or more than the true extent, and this is one of the limitations of the present study. Lack of complete blindness of the all researchers and study subjects was the other limitation of the present study although the statistical analyst did not know the type of treatment received by the groups.

Conclusion

Based on the findings of this study, the use of Olea ointment has been associated with the reduction of episiotomy pain. Therefore, more clinical trials are needed to support our findings. The findings of this study could encourage and stimulate researchers to design and conduct similar clinical trials with the use of Olea ointment on a wider scale.

Acknowledgement

This study is part of the findings of the research project approved by the Vice-Chancellor for Research and Technology of Guilan University of Medical Sciences (IR.GUMS.REC.1396.191), Rasht, Iran. The researchers of this study are sincerely thankful to the university research and technology department for the financial and spiritual support required for the research. The researchers would also like to acknowledge the primiparous women in this study, without whose sincere help and assistance, this research was impossible.

Financial support and sponsorship

Guilan University of Medical Sciences

Conflict of interest

There are no conflicts of interest.

References

- Malik J, Das S, Das A, Rai P. Judicial use of episiotomy in singleton normal vaginal deliveries at term: The changing trend. *Int J Reprod Contracept Obstet Gynecol* 2016;5:882-5.
- Venus D, Rao PS, Prajwal S. Comparison of use of restrictive episiotomy versus routine episiotomy in primigravidae undergoing vaginal birth at a tertiary care hospital. *Int J Reprod Contracept Obstet Gynecol* 2017;6:1770-6.
- Muhleman MA, Aly I, Walters A, Topale N, Tubbs RS, Loukas M. To cut or not to cut, that is the question: A review of the anatomy, the technique, risks and benefits of an episiotomy. *Clin Anat* 2017;30:362-72.
- Kartal B, Kızırlırmak A, Calpbıncı P, Demir G. Retrospective analysis of episiotomy prevalence. *J Turk Ger Gynecol Assoc* 2017;18:190-4.
- Shahrahmani H, Kariman N, Jannesari S, Rafieian-Kopaei M, Mirzaei M, Ghalandari S, et al. The effect of green tea ointment on episiotomy pain and wound healing in primiparous women: A randomized, double-blind, placebo-controlled clinical trial. *Phytother Res* 2018;32:522-30.
- Rezaeyan M, Geranmayeh M, Direkvand-Moghadam A. Comparison of transcutaneous electrical nerve stimulation and lidocaine on episiotomy complication in primiparous women: A randomized clinical trial. *Iran J Nurs Midwifery Res* 2017;22:26-30.
- Navi Nezhad M, Abedin Z, Asili J, Esmaeili H, Vatanchi A. Effect of alpha ointment (fundermol) on episiotomy wound healing in primiparous women. *IJOGI* 2017;20:58-65.
- Blondel B, Alexander S, Bjarnadóttir RI, Gissler M, Langhoff-Roos J, Novak-Antolič Ž, et al. Variations in rates of severe perineal tears and episiotomies in 20 European countries: A study based on routine national data in Euro-Peristat Project. *Acta Obstet Gynecol Scand* 2016;95:746-54.
- Dahlen H, Homer C. Perineal trauma and postpartum perineal morbidity in Asian and non-Asian primiparous women giving birth in Australia. *J Obstet Gynecol Neonatal Nurs* 2008;37:455-63.
- Shahrahmani H, Kariman N, Jannesari S, Ghalandari S, Asaadi N. Assessing the type of treatment methods on the intensity of episiotomy pain and wound healing in Iran: A systematic review. *IJOGI* 2016;19:17-31.
- Hajhashemi M, Ghanbari Z, Movahedi M, Rafieian M, Keivani A, Haghollahi F. The effect of *Achillea millefolium* and *Hypericum perforatum* ointments on episiotomy wound healing in primiparous women. *J Materl Fetal Neonatal Med* 2018;3:63-9.
- Roy P, Sujatha M, Biswas B, Chatterjee A, Roy P. A comparative study of perineal morbidity in vaginal delivery with and without episiotomy. *Int J Reprod Contracept Obstet Gynecol* 2015;4:1442-5.
- Ziaie T, Porhidary M, Nazifkar H, Kazemnejad E. Comparison of wound repair outcomes of chromic catgut versus vicryl sutures for episiotomy repair: A clinical trial. *Adv Nurs Midwifery* 2016;26:1-5.
- Delaram M, Dadhkah NKH. Comparing the effects of lidocaine cream and mefenamic acid on post episiotomy pain. *IJOGI* 2014;17:6-11.
- Ahmadi Z. Review of effective methods for reducing perineal damage during labor and improving it. *IJOGI* 2015;18:19-30.
- Askari M, Alavi A, Dashtinezhad E. Effect of back massage with sesame oil on pain and length of delivery in primiparous women. *IJOGI* 2016;18:12-9.
- Zahmatkesh M, Manesh MJ, Babashahabi R. Effect of Olea ointment and Acetate Mafenide on burn wounds—A randomized clinical trial. *Iran J Nurs Midwifery Res* 2015;20:599-603.
- Rezaeyan M, Abdali N, Araban M. Comparing analgesic effects of extra virgin olive oil and Ibuprofen on the intensity of primary dysmenorrhea: A crossover clinical trial. *Iran J Nutr Sci Food Technol* 2014;9:67-74.
- Lavaf M, Simbar M, Mojab F, Alavi majd H, Samimi M. Comparison of honey and phenytoin cream (PHT) effects on intensity of pain and episiotomy healing in nulliparous women. *J Complement Integr Med* 2017;5:1-12.
- Jahdi F, Sheikhan F, Marghati Khoei ES, Haghani H. The effect of lavender essence on the post-episiotomy pain intensity of perineum. *Sabzevar Univ Med Sci J* 2009;16:127-33.
- Hawker Gillian A, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res* 2011;63:240-52.
- Brokelman RBG, Haverkamp D, Van Loon C, Hol A, Kampen Albert V, Veth R. The validation of the visual analogue scale for patient satisfaction after total hip arthroplasty. *Eur Orthop Traumatol* 2012;3:101-5.
- Phan, Ngoc Quan, Blome Ch, Fritz F, Gerss J, Reich A, Ebata T, et al. Assessment of pruritus intensity: Prospective study on validity and reliability of the visual analogue scale, numerical rating scale and verbal rating scale in 471 patients with chronic pruritus. *Acta Derm Venereol* 2012;92:502-7.
- Behmanesh F, Aghamohammadi A, Zeinalzadeh M, Khafri S.

- Effects of olive oil sitz bath on improvement of perineal injury after delivery. *Koomesh* 2013;14:309-15.
25. Mirbagher-Ajorpaz N, Hafezi M, Salehi S, Tayebi A, Shenasa F, Zahtabchi S. Comparing the effect of pure and impure honey on severity of pain, amount of bleeding, and duration and interval of menstrual cycles in female students with primary dysmenorrhea. *Evidence Based Care* 2012;2:23-33.
 26. Mohammadpour-Kargar H, Shafahi M, Kesmati M. Evaluation of analgesic effect of sesame oil and lecithin diets in hot plate and formalin test on aged male rats. *J Medicin Plant* 2012;2:216-24.
 27. Eshgizade M, Basiri-Moghaddam M, Mohammadzadeh-Moghaddam H, Mahmoudian A, Mesbah M. Comparison of the effect of olive oil, aloe vera extract and breast milk on healing of breast fissure in lactating mothers: A randomized clinical trial. *Qom Univ Med Sci J* 2016;10:19-27.
 28. AmiriFarahani EL, Hasanpoor-Azghdy SB, Kasraei H, Heidari T. Comparison of the effect of honey and mefenamic acid on the severity of pain in women with primary dysmenorrhea. *Arch Gynecol Obstet* 2017;296:277-83.