



Functional Characterization of the GATA-Type Transcription Factor PaNsdD in the Filamentous Fungus *Podospira anserina* and Its Interplay with the Sterigmatocystin Pathway

Ling Shen,^{a,b} Thomas Gaslonde,^c Catherine Roullier,^d Huijuan Wang,^a Jennifer Bodin,^c François-Hugues Porée,^e
Gwenaël Ruprich-Robert,^a Florence Chapeland-Leclerc^a

^aUniversité de Paris, CNRS, Laboratoire Interdisciplinaire des Energies de Demain, Paris, France

^bShenzhen University, Shenzhen Key Laboratory of Microbial Genetic Engineering, College of Life Sciences and Oceanography, Shenzhen, China

^cUniversité de Paris, CNRS, Cibles Thérapeutiques et Conception de Médicaments, Paris, France

^dUniversité de Nantes, Mer Molécules Santé, Nantes, France

^eUniversité de Rennes 1, Rennes Institute of Chemical Sciences, UMR 6226 CNRS, Laboratoire de Chimie Thérapeutique, Faculté de Pharmacie, Rennes, France

Gwenaël Ruprich-Robert and Florence Chapeland-Leclerc contributed equally to this article. Author order was determined in order of decreasing seniority.

ABSTRACT The model ascomycete *Podospira anserina*, distinguished by its strict sexual development, is a prolific but yet unexploited reservoir of natural products. The GATA-type transcription factor NsdD has been characterized by the role in balancing asexual and sexual reproduction and governing secondary metabolism in filamentous fungi. In the present study, we functionally investigated the NsdD ortholog PaNsdD in *P. anserina*. Compared to the wild-type strain, vegetative growth, ageing processes, sexual reproduction, stress tolerance, and interspecific confrontations in the mutant were drastically impaired, owing to the loss of function of PaNsdD. In addition, the production of 3-acetyl-4-methylpyrrole, a new metabolite identified in *P. anserina* in this study, was significantly inhibited in the $\Delta PaNsdD$ mutant. We also demonstrated the interplay of PaNsdD with the sterigmatocystin biosynthetic gene pathway, especially as the deletion of *PaNsdD* triggered the enhanced red-pink pigment biosynthesis that occurs only in the presence of the core polyketide synthase-encoding gene *PaStcA* of the sterigmatocystin pathway. Taken together, these results contribute to a better understanding of the global regulation mediated by PaNsdD in *P. anserina*, especially with regard to its unexpected involvement in the fungal ageing process and its interplay with the sterigmatocystin pathway.

IMPORTANCE Fungal transcription factors play an essential role in coordinating multiple physiological processes. However, little is known about the functional characterization of transcription factors in the filamentous fungus *Podospira anserina*. In this study, a GATA-type regulator PaNsdD was investigated in *P. anserina*. The results showed that PaNsdD was a key factor that can control the fungal ageing process, vegetative growth, pigmentation, stress response, and interspecific confrontations and positively regulate the production of 3-acetyl-4-methylpyrrole. Meanwhile, a molecular interaction was implied between PaNsdD and the sterigmatocystin pathway. Overall, loss of function of PaNsdD seems to be highly disadvantageous for *P. anserina*, which relies on pure sexual reproduction in a limited life span. Therefore, PaNsdD is clearly indispensable for the survival and propagation of *P. anserina* in its complex ecological niches.

KEYWORDS transcription factor PaNsdD, sterigmatocystin, sexual development, 3-acetyl-4-methylpyrrole, *Podospira anserina*

Editor Irina S. Druzhinina, Nanjing Agricultural University

Copyright © 2022 Shen et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Gwenaël Ruprich-Robert, gwenael.ruprich-robert@u-paris.fr.

The authors declare no conflict of interest.

Received 3 December 2021

Accepted 16 January 2022

Accepted manuscript posted online

26 January 2022

Published 22 March 2022

The model fungus *Podospora anserina*, recently described as *Triangularia anserina* (Sordariales, Ascomycota) (1), has long been used in the laboratory to study various biological processes, such as senescence, meiosis, prions, sexual development, signal transduction, cell fusion, hyphal interference, vegetative incompatibility, mitochondrial physiology, and plant biomass degradation (2–5). *P. anserina* is frequently recovered from herbivore dung and is therefore regarded as a typically coprophilous fungus (4, 6). Dung-inhabiting fungi are predominately known to produce various secondary metabolites that act as chemical weapons to enhance competitiveness and ecological fitness (7, 8). These bioactive products are of intense interest to humankind because of their potential pharmaceutical properties (9, 10). Notably, mining of the *P. anserina* genome revealed a large number of putative biosynthetic gene clusters (BGCs) for secondary metabolites (11). However, to date, only very few products that exhibited antibacterial, antifungal, and larvicidal activities have been characterized in this species during past decades (12–14), probably because most BGCs are silent under standard culture conditions (10, 15). Genomics-driven BGC detection and preliminary chemical investigations so far imply that *P. anserina* is a prolific, as-yet unexploited natural product reservoir.

In fungi, secondary metabolism is often governed by intricate regulatory mechanisms that also control other processes (e.g., asexual and sexual development) (16, 17). For example, the global transcription factor LaeA (loss of *afIR* expression) simultaneously regulates numerous metabolic BGCs and fungal morphogenesis in *Aspergillus* spp., as well as in *Fusarium verticillioides* (Hypocreales, Ascomycota), *Monascus pilosus* (Eurotiales, Ascomycota), and *Penicillium dipodomyis* (Eurotiales, Ascomycota) (18–22). As core regulatory elements, transcription factors indeed play an essential role in multilevel gene expression coordination (23). Our previous work characterized the mycotoxin sterigmatocystin (ST) pathway-specific transcription factor PaAfIR, which mediates biological processes, including metabolite production, sexual development, pigmentation, and interspecific competition in *P. anserina* (14). Nevertheless, plenty of transcription factors in *P. anserina* still remain to be elucidated (24).

The *NsdD* gene (never undergo sexual development) encodes a GATA-type zinc finger transcription factor, with highly conserved DNA-binding domain in many filamentous fungi but not in plants or animals (25). NsdD was initially identified as a key activator of sexual reproduction in *Aspergillus nidulans* (Eurotiales, Ascomycota) due to the complete loss of fertility in the corresponding mutant (25). From then on, NsdD and its orthologs (SUB-1/Pro44/Ltf1/Csm1/Nsd1) were characterized according to their involvements in biological processes in filamentous fungi. The roles of NsdD and its orthologs are summarized in Table 1. For example, NsdD affected morphogenesis and aflatoxin (AF) production in *Aspergillus flavus* (Eurotiales, Ascomycota) (26) and was identified as a repressor of asexual development and negatively affected ST production in *A. nidulans* (27). Deletion of *Pro44* caused developmental defects and sterility in *Sordaria macrospora* (Sordariales, Ascomycota) (28). Deletion of *Csm1* resulted in enhanced microconidia formation, reduced stress tolerance, and deregulated expression of BGCs in *Fusarium fujikuroi* (Hypocreales, Ascomycota) (29). Loss of SsNsd1 function affected morphological transition, appressoria formation, and pathogenicity in the plant pathogen *Sclerotinia sclerotiorum* (Helotiales, Ascomycota) (30). However, the role of NsdD in the model fungus *P. anserina*, which lacks asexual reproduction, has not been studied.

In the present study, we functionally investigated the global regulator PaNsdD in *P. anserina*. Our results revealed that PaNsdD was required for vegetative growth, sexual reproduction, stress tolerance, interspecific confrontation, and production of 3-acetyl-4-methylpyrrole, a new secondary metabolite identified in *P. anserina* in this study. We also demonstrated the interplay of PaNsdD with the ST pathway. Indeed, the deletion of *PaNsdD* triggers the enhanced red-pink pigment biosynthesis in the presence of the core polyketide synthase (PKS)-encoding gene *PaStcA* of the ST pathway.

RESULTS

Identification of PaNsdD and generation of mutants. In *P. anserina*, a total of 507 transcription factors classified into 41 families can be retrieved from the Fungal Transcription Factor Database (24). Among them, the *P. anserina* genome contains a putative *NsdD* ortholog, namely *PaNsdD* (*Pa_2_1880*). The complete CDS of *PaNsdD*

TABLE 1 Role of NsdD and orthologs in some filamentous fungi

Fungi	Protein	Role	Reference
<i>Aspergillus nidulans</i>	NsdD	Fertility and sterigmatocystin production	25, 27
<i>Neurospora crassa</i>	SUB-1	Perithecia production	56
<i>Aspergillus flavus</i>	NsdD	Morphogenesis and production of aflatoxin	26
<i>Sordaria macrospora</i>	Pro44	Developmental cycle and fertility	28
<i>Trichoderma reesei</i>	SUB1	Regulation in a nutrient- and light-dependent manner of female fertility, pheromone system, and secondary metabolism	36
<i>Botrytis cinerea</i>	BcLtf1	Regulation of light-dependent stress responses and expression of secondary metabolism-related genes	45
<i>Fusarium fujikuroi</i>	Csm1	Microconidia formation, stress tolerance, and expression of biosynthetic gene clusters	29
<i>Sclerotinia sclerotinium</i>	SsNsd1	Morphological transition, appressoria formation, and pathogenicity	30
<i>Penicillium oxalicum</i>	PoxNsdD	Regulation of major genes involved in plant biomass degradation, conidiation, and pigment biosynthesis	37
<i>Metarhizium rileyi</i>	MrNsdD	Dimorphic transition and microsclerotia formation	46
<i>Podospora anserina</i>	PaNsdD	Vegetative growth, sexual reproduction, stress tolerance, interspecific confrontation, and production of 3-acetyl-4-methylpyrrole	This study

consists of 1,659 bp and potentially encodes a GATA-type transcription factor, namely PaNsdD, of 411 amino acids. PaNsdD shares 99.32, 57.43, and 56.57% identities with NsdD orthologs from *Podospora comata*, *Neurospora crassa*, and *S. macrospora*, respectively. In order to study the biological functions of PaNsdD in *P. anserina*, a deletion mutant lacking *PaNsdD* ($\Delta PaNsdD$) and complemented strain ($\Delta PaNsdD^C$) were constructed, verified (Fig. S1), and subjected to further analysis.

Impact of PaNsdD on vegetative growth and colony morphology. To evaluate the impact of PaNsdD on vegetative growth and colony morphogenesis, wild-type and mutant strains were cultured on M2 solid medium for 12 days. In the first stage, the $\Delta PaNsdD$ mutant displayed a reduction in growth rate compared to the wild-type strain (Fig. 1). Moreover, the $\Delta PaNsdD$ strain formed a flat colony due to sparse aerial




FIG 1 Colony morphology of *P. anserina* strains. (A) The wild-type (WT), $\Delta PaNsdD$, and $\Delta PaNsdD^C$ strains were grown on M2 solid medium at 27°C under dark conditions. The pictures were taken after several days of incubation as indicated. Inoculum was positioned on the edge of the petri dish in order to observe the growth of the strains on a large surface, necessary for 12 days of growth. (B) Colony diameter of tested strains after 5 days of growth. As the inoculation point was made on the edge of the plate, the radius of the colony was measured, the diameter is then deduced by multiplying this length by two. The data represent the means \pm standard deviation (SD) from three independent experiments. Asterisks indicate significant differences ($P < 0.01$) of each strain relative to the WT, followed by Student's *t* test.

hyphae formation. Intriguingly, in the second stage from the 6th day to the 9th day, growth of $\Delta PaNsdD$ appeared to suddenly cease since culture began to slow down, and hyphal prolongation at the edge of the colony was clearly blocked to a large extent. Meanwhile, accelerated and excessive accumulation of pigment started to appear on the surface of colony in a short period, usually less than 3 days (Fig. 1). Nevertheless, the remaining hyphae that were not arrested in partial section could escape the front of growth and persist with outward extension. In the third stage from the 9th to the 12th day, active hyphae that went beyond the colony margin restarted a series of processes, including vegetative proliferation, cessation, and melanization (Fig. 1). It should be noted that more and more mycelia were gradually arrested during this periodic process; only some hyphae can avoid the melanized fate and continue their weak growth until final growth cessation. As shown in Fig. 1, deletion of *PaNsdD* eventually led to the irregular peripheral growth, accompanied by hyphal staling and enhanced pigmentation. In contrast, the $\Delta PaNsdD^C$ strain showed wild-type-like growth characteristics, such as normal growth rate, fluffy colony with extensive aerial hyphae, and normal pigmentation. These results indicated that *PaNsdD* was crucially required for the correct growth and morphogenesis in *P. anserina*; otherwise, a cycled stop-start growth pattern might be triggered, leading to an affected ageing process.

Role of *PaNsdD* in sexual reproduction. To study the role of *PaNsdD* in sexual development, we first examined the formation of reproductive organs through microscopic observation. Both opposite mating type strains (*mat+* and *mat-*) of *P. anserina* were able to differentiate male gametes (spermatia) and female gametes (ascogonia). Obviously, the $\Delta PaNsdD$ mutant exhibited enhanced production of spermatia on M2 solid medium following 3 days of cultivation (Fig. 2A). Subsequently, quantification analysis demonstrated that spermatia yield of $\Delta PaNsdD$ was significantly increased (4.5-fold, $P < 0.01$), compared with wild-type strain (Fig. 2B). The complemented strain restored normal spermatia production. Therefore, these data suggested *PaNsdD* functions as a negative regulator of spermatia differentiation in *P. anserina*.

In view of the aberrant formation of spermatia caused by *PaNsdD* deletion, we further determined the fertility of wild-type and mutant strains. As expected, the wild-type strain produced a ring of perithecia after a fertilization event. In contrast, loss of *PaNsdD* led to the radial growth, but no perithecia were produced even if it was cultured for up to 1 month. Genetically complemented strain regained the wild-type-like fertility (Fig. 2C). This result indicated that the lack of *PaNsdD* completely abolished sexual reproductive ability. Due to the sterile characteristic, we then dissected the potential defects that could be present in the process of gamete differentiation, fertilization, or perithecial maturation. We showed that the $\Delta PaNsdD$ mutant acts as a maternal partner that failed to be fertilized by the spermatia of the wild-type strain, which acts as paternal partner (Fig. 2D). It implies that the absence of *PaNsdD* triggered potential defects in ascogonia. We finally confirmed that the female sterile phenotype in $\Delta PaNsdD$ was not dependent on mating type by reciprocal crosses. Taken together, largely consistent with the initial nomenclature, the data above revealed that the *PaNsdD* mutant indeed never undergo sexual development, which may be mainly due to the overproduction of spermatia and the dysfunction of ascogonia.

Function of *PaNsdD* in multiple stress responses. To clarify the function of *PaNsdD* in response to various stressors, we determined the fungal growth on M2 plates supplemented with stress-inducing agents. Strikingly, compared to the wild-type strain, the $\Delta PaNsdD$ mutant showed significantly reduced tolerance to different oxidative stressors, as H_2O_2 (500 μM), menadione (Mena, 25 μM), and *t*-butyl hydroxyperoxide (TBY, 50 μM) (Fig. 3). Especially, the growth of $\Delta PaNsdD$ was almost completely inhibited in the presence of methylglyoxal (MG) at 5 mM, which can induce reactive oxygen species (ROS) production by depleting intracellular glutathione (31). Moreover, growth of $\Delta PaNsdD$ was highly affected in the presence of sorbitol at 1.5 M. Additionally, to a different extent, growth of mutants was also inhibited under cell wall stress conditions triggered by exposure to Congo red (CR, 100 μM) and Calcofluor white (CFW 10 μM), which impair assembly of β -1,4-glucans and chitin in the cell wall,




FIG 2 Sexual development of *P. anserina* strains. (A) Microscopic observation of spermatia formation. The pictures were taken after 3 days of growth on M2 plate. The arrows indicate typical spermatia. (B) Quantification analysis of spermatia. The tested strains were cultured on M2 medium for 3 days and then used for the counting and observation of spermatia. The data represent the means \pm SD from three independent experiments. Asterisks indicate significant differences ($P < 0.01$) of each strain relative to the WT, followed by Student's *t* test. (C) Fertility assays. Fragmented mycelia from opposite mating type strains were mixed and then inoculated in the center of plate. The plates were grown at 27°C under constant illumination. The photos were taken after 8 days of incubation. (D) Spreading experiments. Crosses were made by inoculating both strains 1.5 cm apart. After 3 days of incubation, 1.5 mL of water was added and spread all over the plate. Mature perithecia were visible as small dots when the fertilization event was completed.

respectively (Fig. 3). Taken together, increased sensitivity of $\Delta PaNsdD$ to various stressors indicated that PaNsdD was required for the maintenance of stress tolerance and cell wall integrity in *P. anserina*.

Contribution of PaNsdD in interspecific confrontations. To elucidate the contribution of PaNsdD in interspecific interactions, the $\Delta PaNsdD$ mutant and the wild-type strain were coconfronted with three other fungal challengers, as previously described (14, 32). In the cell death assay, we found that *Penicillium chrysogenum* (Eurotiales, Ascomycota) and *Botrytis cinerea* that have been described to be usually killed by *P. anserina* displayed a lower level of cell death in the contact zone when confronted to $\Delta PaNsdD$ compared to wild type (Fig. 4A). In contrast, *Trametes versicolor* (Polyporales, Basidiomycota), which usually can kill *P. anserina* (14, 32), aggravated cell death specifically on the thallus of $\Delta PaNsdD$, compared to confrontation with wild type. These results implied that loss of PaNsdD triggered the declining competitiveness of *P. anserina* during interspecific antagonistic processes. Then, we detected the level of peroxide, which is usually restricted to the contact zone, and superoxide, that is accumulated more or less uniformly on fungal thallus, without any necessary contact with the competitor (32). No obvious difference in peroxide accumulation is observed between wild type and $\Delta PaNsdD$ at the confrontation with contestant strains (Fig. 4B). Moreover,




FIG 3 Growth of *P. anserina* strains on M2 plates exposed to abiotic stressors. The tested strains were grown on M2 medium supplemented with different stress agents under certain concentrations as indicated. (A) Colony morphology in the presence of stress-inducing agents. The pictures were taken after 3 days of incubation at 27°C. (B) Colony diameter (cm) of tested strains after 3 days of incubation at 27°C. The corresponding strains that grown on M2 medium without supplements were used for normalization. CFW, Calcofluor white; CR, Congo red; Mena, menadione; TBY, *t*-butyl hydroxyperoxide; MG, methylglyoxal; Sorb, sorbitol.

peroxide accumulation was not detected on wild-type and mutant thalli, in the absence of competitor (Fig. S2). Finally, nitroblue tetrazolium (NBT) detection assay revealed that Δ PaNsdD produced a lower level of superoxide over the thallus than that of the wild-type strain (Fig. 4C), indicating that PaNsdD was probably involved in the production of ROS, mainly superoxide. Combined with the decreased tolerance of Δ PaNsdD to abiotic stresses, we suggested that PaNsdD might contribute to a positive role in protecting *P. anserina* from hostile environment by mediating ROS production.

Effects on fungal pigmentation and interplay of PaNsdD with the ST pathway.

To investigate the impact of PaNsdD on fungal pigmentation, all of the strains were cultured on M2 and LB media, liquid and solid in both cases. As shown in Fig. 5, *P. anserina* wild-type strain was normally greenish on M2 medium, due to accumulation of melanin, and presented a slight red-pink pigmentation in M2 liquid medium and on LB plate. Nevertheless, we first observed that the coloration is more intense under all conditions with Δ PaNsdD compared to wild type, suggesting an excessive accumulation of pigments in this mutant. Then, one might expect that PaNsdD could downregulate the pigment production or affect its accumulation and could be involved in the melanin biosynthesis in particular. It should be noted here that when strains were grown on G medium and G + yeast extract medium (culture media usually used to promote ascospore germination), the mycelium remained everywhere very poorly pigmented, and the culture medium remained unstained (Fig. S3). Taken together, all these results are in accordance with the fact that fungal coloration highly depends on the medium composition (5).

Nevertheless, in a previous work, we observed (i) that the absence of the red-pink pigmentation when the Δ PaStcA mutant lacking the ST-core polyketide synthase PaStcA was cultured on M2 and (ii) the accumulation of red-pink pigments in the ST-overproducing strain OE-PaAflR (14), which seem to be identical to those produced by




FIG 4 Confrontation of *P. anserina* strains against other fungal species. Cell death assay (A), peroxide detection (B), and superoxide detection (C) were conducted as previously described (14, 32). (A) In the contact zone, the intensity difference of blue color indicates the Trypan blue-stained dead cells triggered by the contestant strains. Asterisks indicate more serious cell death than that indicated by arrows. (B) Peroxide detection of *P. anserina* strains when contestants were involved. (C) Superoxide detection in WT, $\Delta PaNsdD$, and $\Delta PaNsdD^C$ in the absence of challengers.

the $\Delta PaNsdD$ strain. Similar results were obtained here on M2 but also on LB medium (Fig. 5B to D). In order to better understand the putative interplay of PaNsdD with the ST pathway and the occurrence of a similar red-pink pigmentation phenotype, the double mutant $\Delta PaNsdD\Delta PaStcA$ was constructed.

This mutant exhibited a colorless colony phenotype identical to $\Delta PaStcA$ on M2 liquid and LB media (Fig. 5B to D). Therefore, these results suggested that the presence of PaStcA was a prerequisite for the biosynthesis of red-pink pigments and that this gene interplays with pigment production. Our findings here are consistent with those previously described; i.e., that such a red-pink pigmentation requires the presence of functional ST pathway (14). It is worth noting that cultures of $\Delta PaStcA$ were always colorless even when they were subjected to 1 month of incubation under the same conditions (data not shown).

Involvement of PaNsdD in the biosynthesis of a new metabolite in *P. anserina*.

To examine secondary metabolites regulated by the global transcription factor PaNsdD in *P. anserina*, we first conducted a qualitative and semiquantitative analysis by high-performance liquid chromatography UV (HPLC-UV) analysis in all extracts after 14-day cultivation, as previously described (14). We found that the secondary metabolite profile was significantly altered owing to the *PaNsdD* mutation. First, under M2 liquid condition, the profile of wild-type strain evidenced two major peaks at 16.05 and 27.3 min, respectively (Fig. 6). The second peak at 27.3 min could unambiguously be correlated with ST, as previously seen (14). We may note that the HPLC-UV profiling on LB culture extracts showed only one peak, clearly correlated with ST, in wild-type and $\Delta PaNsdD$ strains, meaning that the second compound is absent under these culture conditions (Fig. S4).




FIG 5 Pigmentation of *P. anserina* on various media. Tested strains were cultured on M2 solid medium (A), M2 liquid medium (B), LB solid medium (C) and LB liquid medium (D), respectively. After 14 days of growth at 27°C, altered pigmentation were clearly observed, and photos were therefore taken.

In order to identify the compound corresponding to the peak at 16.05 min, it was necessary to proceed to its isolation by silica chromatography and preparative HPLC successively. Characterization of this isolated product was conducted by liquid chromatography-mass spectrometry (LC-MS) in electron spray ionization (ESI)-positive and -negative modes. Our results showed that the corresponding signals of this product are two ions at m/z 124 for $[M+H]^+$ and m/z 122 for $[M-H]^-$, respectively (Fig. S5), indicating that the molecular formula of this compound could be C_7H_9NO . 1H and ^{13}C nuclear magnetic resonance (NMR) along with 2D spectra (Fig. S6) allowed identification of 3-acetyl-4-methylpyrrole for the compound at 16.05 min. All of these data are in accordance with the literature (33–35). Thus, deletion of *PaNsdD* in both wild-type and $\Delta PaStcA$ background almost completely blocked the production of 3-acetyl-4-methylpyrrole (at 16.05 min), implying that *PaNsdD* positively regulates its biosynthesis. In contrast, the amount of ST, determined by the UV signal intensity (mAU), was similar in both wild type and $\Delta PaNsdD$, indicating that the production of ST was not affected by the deletion of *PaNsdD* (Fig. 6).

DISCUSSION

PaNsdD is an activator of sexual reproduction and a repressor of male gamete production in *P. anserina*. Given the critical role of *NsdD* in balancing sexual and asexual development in several fungi (29, 30, 36–38), we wondered about the function of *PaNsdD* in *P. anserina*, which has only a pure sexual life cycle without any asexual phase (4). Our data suggested that *PaNsdD* is a positive regulator of sexual reproduction and a repressor of male gametes production in *P. anserina*. Unlike other asexual spores, spermatia produced by *P. anserina* were only used to fertilize ascogonia and had almost no germination (38). Here, we show that the erected sterile hyphae of




FIG 6 High-performance liquid chromatography UV (HPLC-UV) detection of culture extracts of *P. anserina*. The HPLC-UV profiling displayed the peak intensity (mAU) according to the retention time (rt). Sterigmatocystin (ST) presents a rt of 27.3 min (14). The other major peak presented a rt of 16.05 min. The x axis (min) shows the retention time. The y axis (mAU) shows the signal intensity.

$\Delta PaNsdD$ mutant was able to form anamorphic-like structures containing dozens of spermatia, which strikingly resemble those of *Cladorrhinum/Bahupaathra* species (5).

PaNsdD is involved in the vegetative growth pattern of *P. anserina*. Remarkably, $\Delta PaNsdD$ mutant was distinguished by an unusual growth characteristic that involved an alternation of irregular periods of growth and nongrowth on M2 solid medium. In this study, we presented the infrequent alternative growth pattern in *P. anserina*, however, that resembled the ones observed in *Neurospora* and *Aspergillus* species. For instance, unlike the continuous outward extension of hyphae in the wild-type strain, the “stopper” mutants of *N. crassa* had an aberrant phenotype in which growth of the mycelium started, stopped, and then started again (39). The “vegetative death” mutants of *Aspergillus glaucus* (Eurotiales, Ascomycota) showed many features of senescence similar to those described for *P. anserina*. However, a partial section of *A. glaucus* mutant culture sometimes can escape the dying front of mycelia and continue the patchy extension until ultimate growth cessation (40). Taken together, compared to the wild-type strain of *P. anserina*, the $\Delta PaNsdD$ mutant showed persistent cycles of stopping and impaired starting, with a concomitant excessive accumulation of pigment, leading to a clear disturbance in the ageing process of the thallus. As we know, the ageing model *P. anserina* clearly did not propagate indefinitely and underwent

senescence during its limited vegetative growth period (4, 41). In *P. anserina*, strain-specific senescent phenotypes were characterized by some common features, such as darker pigmentation, growth stop at the hyphal tips, and formation of “barrier of senescence” (5). Noticeably, visible morphological changes in $\Delta PaNsdD$ mutant, such as sparse aerial hyphae, overproduction of dark pigment, accelerated cessation of hyphal growth, rapid transition from juvenile phase to senescent phase, and ragged elongation of partial peripheral hyphae, indicated the premature senescence syndrome caused by *PaNsdD* mutation.

PaNsdD is required for fungal pigmentation and stress tolerance. Pigmentation is generally considered the output of secondary protective mechanisms that protect the producer from adverse environmental stresses, including oxidative stress, light, UV radiation, and temperature variation (42–44). Moreover, given the essential role of NsdD and its orthologs in maintaining the redox homeostasis in other fungi (29, 45), we hypothesized that the overpigmented $\Delta PaNsdD$ mutant would respond to multiple stressors. However, we found that $\Delta PaNsdD$ mutant was hypersensitive in the presence of various oxidant and osmotic stressors, suggesting that here, enhanced accumulation of pigment did not confer a protective function onto *P. anserina*. Moreover, a likely scenario could be that defects in cell morphogenesis might lead to internal stress signals caused by aberrant cell wall formation, which in turn could be the reason for the observed sensitivity toward several stresses or the enhanced melanization. Our findings were in accordance with the functions of BcLtf1 in *B. cinerea* and MrNsdD in *Metarhizium releyi* in mediating ROS homeostasis and stress response (45, 46).

PaNsdD affects secondary metabolism in *P. anserina*. Apart from the altered growth pattern and differentiation affected by the loss of function of PaNsdD, pigment variation of mutant cultures in both solid and liquid medium indicated that PaNsdD played a vital role in governing some secondary metabolism pathways in *P. anserina*. Indeed, previous studies revealed that NsdD and its orthologs function as global regulators in secondary metabolism processes in several fungi, and in particular, NsdD influenced ST biosynthesis in *A. nidulans* (27). In this study, we have highlighted a dark green pigment and a red-pink pigment that are overproduced in $\Delta PaNsdD$. As the DHN-melanin has been described to be the major pigment that contributes to the greenish colony morphology of *P. anserina* in standard conditions (47), we hypothesize here that the dark green pigment is linked to the biosynthesis of melanin, but further experiments are necessary to clarify this point. Intriguingly, deletion of *PaNsdD* in both wild-type and $\Delta PaStcA$ backgrounds triggered a very dark pigmentation of the whole thalli, which is probably also linked to an excessive accumulation of melanin. We also found that the single mutant $\Delta PaStcA$ and the double mutant $\Delta PaNsdD\Delta PaStcA$ lost their abilities to produce the red-pink pigment, which is visible in both wild-type and $\Delta PaNsdD$ strains on M2 liquid medium and on LB plates, implying that the $\Delta PaStcA$ phenotype predominated in the double mutant under these conditions. In contrast, it seems that growth of *P. anserina* on LB medium was not able to induce fungal melanin production but led to the formation of a red-pink pigment, indicating that at least two pigment biosynthetic pathways exist in *P. anserina*. We do not know yet which chemical compound is responsible for the pigmentation, but we have demonstrated here that the presence of PaStcA was a prerequisite for the biosynthesis of this red-pink pigment and that PaNsdD negatively regulated the two pigmentation pathways.

Finally, secondary metabolites may serve as defense mechanisms during fungal colonization and the adaptation processes (48, 49). The interspecific competitiveness of $\Delta PaNsdD$ was suppressed when confronted with other fungal antagonists. Accordingly, in addition to the reduction of ROS production, the role of the new compound, identified here as 3-acetyl-4-methylpyrrole, needs to be further investigated. To this day, this compound was identified only in *Eupenicillium hirayamae* (Eurotiales, Ascomycota) under the name of desoxyverrucarin E (35) and in *Aspergillus terreus* (Eurotiales, Ascomycota) (34), but its involvement in fungal physiology has not yet been investigated.

To conclude, in this study, we demonstrated that deletion of *PaNsdD* led to a huge change in vegetative growth, pigmentation, stress response, and secondary metabolism

regulation and that a molecular interaction was observed between PaNsdD and PaStcA. Overall, the loss of PaNsdD leads to premature ageing, sterility, and sensitivity to environmental stresses. Such features seemed to be highly disadvantageous for this species whose propagation before senescence and death exclusively relies on sexual reproduction. The presence of PaNsdD is clearly indispensable for the survival and propagation of *P. anserina* in its complex ecological niches.

MATERIALS AND METHODS

Strains and culture conditions. The *P. anserina* strains used in this study were derived from the “S” (big S) wild-type strain, which was used for the sequencing of the *P. anserina* genome (2, 11, 50). The two culture media used for this study are the most commonly used M2 medium (0.25 g/L KH_2PO_4 , 0.3 g/L K_2HPO_4 , 0.25 g/L MgSO_4 , 0.5 g/L urea, 0.05 mg/L thiamine, 0.05 $\mu\text{g/L}$ biotine, 5 mg/L citric acid, 5 mg/L ZnSO_4 , 0.25 mg/L CuSO_4 , 50 $\mu\text{g/L}$ MnSO_4 , 50 $\mu\text{g/L}$ boric acid, 50 $\mu\text{g/L}$ natrium molybdate, 1 mg/L iron alum, 5.5 g/L dextrin, and 10 g/L agar) with pH maintenance at 7 by a phosphate buffer and the G (germination) medium (25 g/L corn flour, 25 g/L corn cream, 6 g/L ammonium acetate, and 12g/L agar).

All the protocols, including standard culture conditions and genetic manipulation for this microorganism, are described by Silar (4) and can be accessed at <http://podospora.i2bc.paris-saclay.fr>. The $\Delta\text{mus51}::\text{nourR}$ strain lacking a *mus51* gene was used for deletion mutant construction, which resulted in an increased frequency of targeted gene replacement (51). The $\Delta\text{PaStcA}::\text{genetR}$ strain harboring a mutation in the PKS-encoding gene *PaStcA*, which acts at the first step of ST biosynthesis, was previously constructed by Shen et al. (14).

Deletion and complementation of PaNsdD. The *PaNsdD* (*Pa_2_1880*) gene sequence was retrieved from the *P. anserina* genome database by BLAST, using the *N. crassa* protein SUB-1 (NCU01154) as query (11). To investigate the function of PaNsdD, targeted gene deletion was carried out according to the “split marker” method, as previously described (4, 52). Briefly, a phleomycin resistance cassette was used as selective marker. The flanking regions (approximately 800 bp) of the open reading frame of *PaNsdD* were amplified and then were fused with the resistance cassette, respectively. The fusion products were then used to transform protoplasts of the $\Delta\text{mus51}::\text{nourR}$ strain. All transformants were screened on minimal medium containing phleomycin at 10 $\mu\text{g/mL}$. Putative transformants were then diagnosed by PCR. The primers used in this study are listed in Table 2. To eliminate the untransformed nuclei and to segregate the Δmus51 mutation, two primary transformants $\Delta\text{PaNsdD}\Delta\text{mus51}$ (PhleoR HygroR) were genetically purified by crossing with the wild-type strain to generate both *mat+* and *mat-* homokaryotic mutant strains ΔPaNsdD (PhleoR HygroS) that contain only the *PaNsdD* deletion but lack Δmus51 . Correct gene replacement was further confirmed by Southern blotting (Fig. S1). The ΔPaNsdD mutant was genetically crossed with ΔPaStcA to generate the double mutant $\Delta\text{PaNsdD}\Delta\text{PaStcA}$, which was resistant to both phleomycin and Geneticin.

Complemented strains were generated by ectopically introducing a wild-type allele of *PaNsdD* into corresponding mutant, as described by Shen et al. (14). To obtain the plasmid pBC-hygro-PaNsd-Comp, a wild-type *PaNsdD* gene containing native promoter and terminator was amplified with primers PaNsdD_cF and PaNsdD_cR (Table S1). Fragment was inserted into EcoRV-linearized pBC-hygro plasmid. The reconstructed vector was validated by sequencing (Genewiz, Germany) and then was transformed into the ΔPaNsdD mutant strain. All hygromycin-resistant transformants were verified by PCR and phenotypic analysis; three out of putative transformants that restored the wild-type-like phenotype were genetically purified, resulting in the complemented strain ΔPaNsdD^c , which was selected for detailed phenotypic characterization.

Phenotypic analysis. To determine the role of PaNsdD in fungal physiology, wild-type and mutant strains were cultured on M2 medium at 27°C. Aliquots (5 μL) of standardized fragmented mycelial suspension were spotted on the plates. For vegetative growth observation, colony morphology, size, and pigmentation were documented each day. For fertility assessment, perithecia formation, ascospore production, and dispersal were examined during their sexual cycle. In order to observe the fungal pigmentation of thallus and the coloration of culture filtrate, M2 and LB liquid cultivation were also similarly conducted. All experiments were performed in triplicate.

Microscopy analysis and quantitative measurement of spermatia. Microscopic observation was made on fresh mycelia growing on solid M2 medium. Quantitative measurement of spermatia was conducted as described previously (53). Briefly, the strains were cultured for 3 days, and then the spermatia were harvested by flooding the plates with 1.5 mL of 0.05% Tween 20 in sterile water. The suspension was counted through a hemacytometer.

Sensitivity to various abiotic stresses. To assess the sensitivity of fungal cells to environmental stresses, fungal growth was determined on M2 medium supplemented with various chemical stressors at a certain concentration: (i) H_2O_2 (500 μM), menadione (Mena, 25 μM), t-butyl hydroxylperoxide (TBH, 50 μM), or methylglyoxal (MG, 5 mM) for oxidative stresses; (ii) Sorbitol (1.5 M) for osmotic stress, and (iii) Congo red (CR, 100 μM) or Calcofluor white (CFW, 10 μM) for cell wall-perturbing stresses. Aliquots (5 μL) of standardized fragmented mycelial suspension were spotted on the plates. Colony diameters were measured daily from three replicates. Quantification of relative growth inhibition of strains exposed to different chemicals was calculated as D_i/D_c (where D_c is diameter of control colony without treatment, and D_i is the diameter of colony treated with stressors).

Interspecific confrontation assay. Detection of cell death and oxidative burst were conducted as previously described (14, 32). Briefly, *P. anserina* wild-type and mutant strains were inoculated on M2 plates with three neighboring *P. chrysogenum* strains and then incubated for 3 days. *T. versicolor* and *B. cinerea* strains were centrally precultured on M2 plates for 4 days, prior to the inoculation of *P. anserina* wild-type and mutant strains. Cocultivation can be ceased until the central fungus was completely in

TABLE 2 Primers used in this study

Primers	Sequence (5' to 3')	Application
PaNsdD_1F	tccagattggcaggacagt	Deletion of <i>Pa_2_1880</i>
PaNsdD_2R	ctatttaacgacctgacctgaaccgggactggaccgttttg	
PaNsdD_2F	ccaaaaacggctccaagtccgggtcagggcagggtcgtaaatag	
PaNsdD_3R	ggcactactagctccggctccatcgaactggatctcaacagcggaag	
PaNsdD_3F	cttaccgctgttgagatccagttcagtgaggaccggagctagtattgacc	Junction verification for <i>Pa_2_1880</i> deletion
PaNsdD_4R	actaaagggtgcccacaaactgc	
PaNsdD_verify_F	cgaattgttccgcttttcg	
Valid5'	tgagaagcacacggtcac	
Valid3'	tcggggcgaaaactctc	Complementation construct of <i>Pa_2_1880</i>
PaNsdD_verify_R	tgaatccgagcctctgttgc	
PaNsdD_cF	attggcaggacagtgcacatc	
PaNsdD_cR	gtcttcaggcgagaacagc	
PaNsdD_verify_cF	ccagtcctcaacaaatgcc	Verification for <i>Pa_2_1880</i> complementation
PaNsdD_verify_cR	gctgtttcctcacatagc	

contact with competing fungal thalli (14). Trypan blue staining was used to reveal the cell death in the contact zone (32). Diaminobenzidine (DAB) and nitroblue tetrazolium (NBT) staining were used to detect peroxide and superoxide, respectively (54, 55).

Metabolite extraction and analysis. To monitor the production of putative compounds regulated by PaNsdD, *P. anserina* wild-type and mutant strains were stationarily cultured on M2 and LB liquid medium at 27°C for 14 days. Secondary metabolite extraction was conducted as described previously (14). The extracts were dissolved in 2 mL of methanol, filtered over a 0.2- μ m filter, and subjected to reversed-phase high performance liquid chromatography coupled to a diode-array detector (DAD) analysis. HPLC analysis was performed on Dionex UltiMate 3000 HPLC Systems using a column (X-bridge C18, 4.6 \times 250 mm, 5 μ m, Waters, Ireland) with a flow rate of 1 mL/min. Fresh extracts of all strains were detected with a gradient of acetonitrile in H₂O: 0% for 5 min, 0 to 100% in 35 min, and 100% for 10 min.

In order to identify peaks detected on HPLC-UV chromatogram, ST standard previously isolated was used (14). Concerning the peak at 16.05 min, which disappeared in the Δ PaNsdD strain, its purification was conducted as follows. The extract (125 mg) was purified by chromatography on silica (20 to 45 μ m) with a gradient of ethyl acetate in cyclohexane (0 to 100%) to give a fraction of 4.5 mg containing the unknown compound. A subsequent purification was made on a preparative HPLC system (Shimadzu, Kyoto, Japan) consisting of an LC-40 delivery system, Rheodyne manual injection valve, FRC-40 fraction collector, and SPD-M40 detector connected to a SCL-40 control unit. The fraction was dissolved in 0.5 mL of MeOH and purified on a Nucleodur C18 Htec (250 mm \times 32 mm, 5 μ m) column (Macherey-Nagel) with a gradient of acetonitrile in water (10 to 80% for 30 min) at a flow rate of 45 mL/min. Fractions containing the products were pooled, and the mobile phase was evaporated under reduced pressure to furnish 1.9 mg of the desired pure compound. NMR and MS data of this metabolite were in accordance with the literature (33, 34) and were permitted to identify the product as 3-acetyl-4-methylpyrrole.

Data availability. All of the data generated in this study are included in the main text and its supplementary information files.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 2.3 MB.

ACKNOWLEDGMENTS

We sincerely thank Sylvie Cangemi for her expert technical assistance.

H. Wang and L. Shen were funded by the China Scholarship Council (CSC). This study was financially supported by Université de Paris.

We declare no conflict of interest.

REFERENCES

- Wang XW, Bai FY, Bensch K, Meijer M, Sun BD, Han YF, Crous PW, Samson RA, Yang FY, Houbraken J. 2019. Phylogenetic re-evaluation of *Thielavia* with the introduction of a new family Podosporaceae. *Stud Mycol* 93: 155–252. <https://doi.org/10.1016/j.simyco.2019.08.002>.
- Rizet G. 1952. Les phénomènes de barrage chez *Podospore anserina*. I. Analyse génétique des barrages entre souches S and s. *Rev Cytol Biol Veg* 13:51–92.
- Osiewacz HD. 1990. Molecular analysis of aging processes in fungi. *Mutat Res* 237:1–8. [https://doi.org/10.1016/0921-8734\(90\)90026-n](https://doi.org/10.1016/0921-8734(90)90026-n).
- Silar P. 2013. *Podospore anserina*: from laboratory to biotechnology, p 283–309. In Horwitz BA, Mukherjee PK, Mukherjee M, Kubicek CP (ed), *Genomics of soil- and plant-associated fungi*. Springer, Berlin, Germany.
- Silar P. 2020. *Podospore anserina*. <https://hal.archives-ouvertes.fr/hal-02475488>.
- Richardson MJ. 2001. Diversity and occurrence of coprophilous fungi. *Mycological Res* 105:387–402. <https://doi.org/10.1017/S0953756201003884>.
- Bills GF, Gloer JB, An Z. 2013. Coprophilous fungi: antibiotic discovery and functions in an underexplored arena of microbial defensive

- mutualism. *Curr Opin Microbiol* 16:549–565. <https://doi.org/10.1016/j.mib.2013.08.001>.
8. Sarrocco S. 2016. Dung-inhabiting fungi: a potential reservoir of novel secondary metabolites for the control of plant pathogens. *Pest Manag Sci* 72:643–652. <https://doi.org/10.1002/ps.4206>.
 9. Keller NP. 2019. Fungal secondary metabolism: regulation, function and drug discovery. *Nat Rev Microbiol* 17:167–180. <https://doi.org/10.1038/s41579-018-0121-1>.
 10. Caesar LK, Kelleher NL, Keller NP. 2020. In the fungus where it happens: history and future propelling *Aspergillus nidulans* as the archetype of natural products research. *Fungal Genet Biol* 144:103477. <https://doi.org/10.1016/j.fgb.2020.103477>.
 11. Espagne E, Lespinet O, Malagnac F, Da Silva C, Jaillon O, Porcel BM, Couloux A, Aury J-M, Ségurens B, Poulain J, Anthouard V, Grossetete S, Khalili H, Coppin E, Déquard-Chablat M, Picard M, Contamine V, Arnaise S, Bourdais A, Berteaux-Lecellier V, Gautheret D, de Vries RP, Battaglia E, Coutinho PM, Danchin EG, Henrissat B, Khoury RE, Sainsard-Chanet A, Boivin A, Pinan-Lucarré B, Sellem CH, Debuchy R, Wincker P, Weissenbach J, Silar P. 2008. The genome sequence of the model ascomycete fungus *Podospira anserina*. *Genome Biol* 9:R77. <https://doi.org/10.1186/gb-2008-9-5-r77>.
 12. Wang H, Gloer KB, Gloer JB, Scott JA, Malloch D. 1997. Anserinones A and B: new antifungal and antibacterial benzoquinones from the coprophilous fungus *Podospira anserina*. *J Nat Prod* 60:629–631. <https://doi.org/10.1021/np970071k>.
 13. Matasyoh JC, Dittrich B, Schueffler A, Laatsch H. 2011. Larvicidal activity of metabolites from the endophytic *Podospira* sp. against the malaria vector *Anopheles gambiae*. *Parasitol Res* 108:561–566. <https://doi.org/10.1007/s00436-010-2098-1>.
 14. Shen L, Porée F, Gaslonde T, Lalucque H, Chapeland-Leclerc F, Ruprich-Robert G. 2019. Functional characterization of the sterigmatocystin secondary metabolite gene cluster in the filamentous fungus *Podospira anserina*: involvement in oxidative stress response, sexual development, pigmentation and interspecific competitions. *Environ Microbiol* 108:3011–3026.
 15. Brakhage AA. 2013. Regulation of fungal secondary metabolism. *Nat Rev Microbiol* 11:21–32. <https://doi.org/10.1038/nrmicro2916>.
 16. Calvo AM, Wilson RA, Bok JW, Keller NP. 2002. Relationship between secondary metabolism and fungal development. *Microbiol Mol Biol Rev* 66:447–459. <https://doi.org/10.1128/MMBR.66.3.447-459.2002>.
 17. Bills GF, Gloer JB. 2016. Biologically active secondary metabolites from the fungi. *Microbiol Spectr* 4:e0009. <https://doi.org/10.1128/microbiolspec.FUNK-0009-2016>.
 18. Bok JW, Keller NP. 2004. LaeA, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot Cell* 3:527–535. <https://doi.org/10.1128/EC.3.2.527-535.2004>.
 19. Butchko RAE, Brown DW, Busman M, Tudzynski B, Wiemann P. 2012. Lae1 regulates expression of multiple secondary metabolite gene clusters in *Fusarium verticillioides*. *Fungal Genet Biol* 49:602–612. <https://doi.org/10.1016/j.fgb.2012.06.003>.
 20. Lee SS, Lee JH, Lee I. 2013. Strain improvement by overexpression of the laeA gene in *Monascus pilosus* for the production of monascus-fermented rice. *J Microbiol Biotechnol* 23:959–965. <https://doi.org/10.4014/jmb.1303.03026>.
 21. Grau MF, Entwistle R, Oakley CE, Wang CCC, Oakley BR. 2019. Overexpression of an LaeA-like methyltransferase upregulates secondary metabolite production in *Aspergillus nidulans*. *ACS Chem Biol* 14:1643–1651. <https://doi.org/10.1021/acscchembio.9b00380>.
 22. Yu J, Han H, Zhang X, Ma C, Sun C, Che Q, Gu Q, Zhu T, Zhang G, Li D. 2019. Discovery of two new sorbicillinoids by overexpression of the global regulator LaeA in a marine-derived fungus *Penicillium dipodomys* YJ-11. *Mar Drugs* 17:446. <https://doi.org/10.3390/md17080446>.
 23. Shelest E. 2017. Transcription factors in fungi: TFome dynamics, three major families, and dual-specificity TFs. *Front Genet* 8:53. <https://doi.org/10.3389/fgene.2017.00053>.
 24. Park J, Park J, Jang S, Kim S, Kong S, Choi J, Ahn K, Kim J, Lee S, Kim S, Park B, Jung K, Kim S, Kang S, Lee Y-H. 2008. FTFD: an informatics pipeline supporting phylogenomic analysis of fungal transcription factors. *Bioinformatics* 24:1024–1025. <https://doi.org/10.1093/bioinformatics/btn058>.
 25. Han K-H, Han K-Y, Yu J-H, Chae K-S, Jahng K-Y, Han D-M. 2001. The *nsdD* gene encodes a putative GATA-type transcription factor necessary for sexual development of *Aspergillus nidulans*. *Mol Microbiol* 41:299–309. <https://doi.org/10.1046/j.1365-2958.2001.02472.x>.
 26. Cary JW, Harris-Coward PY, Ehrlich KC, Mack BM, Kale SP, Larey C, Calvo AM. 2012. NsdC and NsdD affect *Aspergillus flavus* morphogenesis and aflatoxin production. *Eukaryot Cell* 11:1104–1111. <https://doi.org/10.1128/EC.00069-12>.
 27. Lee M-K, Kwon N-J, Choi JM, Lee I-S, Jung S, Yu J-H. 2014. NsdD is a key repressor of asexual development in *Aspergillus nidulans*. *Genetics* 197:159–173. <https://doi.org/10.1534/genetics.114.161430>.
 28. Nowrousian M, Teichert I, Masloff S, Kück U. 2012. Whole-genome sequencing of *Sordaria macrospora* mutants identifies developmental genes. *G3* 2:261–270. <https://doi.org/10.1534/g3.111.001479>.
 29. Niehaus E-M, Schumacher J, Burkhardt I, Rabe P, Münsterkötter M, Güldener U, Sieber CMK, Dickschat JS, Tudzynski B. 2017. The GATA-type transcription factor Csm1 regulates conidiation and secondary metabolism in *Fusarium fujikuroi*. *Front Microbiol* 8:1175. <https://doi.org/10.3389/fmicb.2017.01175>.
 30. Li J, Mu W, Veluchamy S, Liu Y, Zhang Y, Pan H, Rollins JA. 2018. The GATA-type IVb zinc-finger transcription factor SsNsd1 regulates asexual development and appressoria formation in *Sclerotinia sclerotiorum*. *Mol Plant Pathol* 19:1679–1689. <https://doi.org/10.1111/mpp.12651>.
 31. Desai KM, Chang T, Wang H, Banigesh A, Dhar A, Liu J, Untereiner A, Wu L. 2010. Oxidative stress and aging: is methylglyoxal the hidden enemy? *Can J Physiol Pharmacol* 88:273–284. <https://doi.org/10.1139/Y10-001>.
 32. Silar P. 2005. Peroxide accumulation and cell death in filamentous fungi induced by contact with a contestant. *Mycol Res* 109:137–149. <https://doi.org/10.1017/s0953756204002230>.
 33. Wang P, Chau Nguyen K, Lindsey JS. 2019. Synthesis of the ring C pyrrole of native chlorophylls and bacteriochlorophylls. *J Org Chem* 84:11286–11293. <https://doi.org/10.1021/acs.joc.9b01650>.
 34. Arai K, Yoshimura T, Itatani Y, Yamamoto Y. 1983. Metabolic products of *Aspergillus terreus*. VIII. Astepyrone: a novel metabolite of the strain IFO 4100. *Chem Pharm Bull* 31:925–933. <https://doi.org/10.1248/cpb.31.925>.
 35. Arndt RR, Holzapfel CW, Ferreira NP, Marsh JJ. 1974. The structure and biogenesis of desoxyverrucarin E, a metabolite of *Eupenicillium hirayamae*. *Phytochemistry* 13:1865–1870. [https://doi.org/10.1016/0031-9422\(74\)85103-4](https://doi.org/10.1016/0031-9422(74)85103-4).
 36. Bazaifan H, Beier S, Stappler E, Böhmendorfer S, Oberlerchner JT, Suljok M, Schmol M. 2017. SUB1 has photoreceptor dependent and independent functions in sexual development and secondary metabolism in *Trichoderma reesei*. *Mol Microbiol* 106:742–759. <https://doi.org/10.1111/mmi.13842>.
 37. He Q-P, Zhao S, Wang J-X, Li C-X, Yan Y-S, Wang L, Liao L-S, Feng J-X. 2018. Transcription factor NsdD regulates the expression of genes involved in plant biomass-degrading enzymes, conidiation, and pigment biosynthesis in *Penicillium oxalicum*. *Appl Environ Microbiol* 84:e01039-18. <https://doi.org/10.1128/AEM.01039-18>.
 38. Grognet P, Timpano H, Carlier F, Ait-Benkhalil J, Berteaux-Lecellier V, Debuchy R, Bidard F, Malagnac F. 2019. A RID-like putative cytosine methyltransferase homologue controls sexual development in the fungus *Podospira anserina*. *PLoS Genet* 15:e1008086. <https://doi.org/10.1371/journal.pgen.1008086>.
 39. Bertrand H, Collins RA, Stohl LL, Goewert RR, Lambowitz AM. 1980. Deletion mutants of *Neurospora crassa* mitochondrial DNA and their relationship to the “stop-start” growth phenotype. *Proc Natl Acad Sci U S A* 77:6032–6036. <https://doi.org/10.1073/pnas.77.10.6032>.
 40. Jinks JLY. 1959. Lethal suppressive cytoplasm in aged clones of *Aspergillus glaucus*. *Microbiology* 21:397–409. <https://doi.org/10.1099/00221287-21-2-397>.
 41. Osiewacz HD, Hamann A, Zintel S. 2013. Assessing organismal aging in the filamentous fungus *Podospira anserina*, p 439–462. In Galluzzi L, Vitale I, Kepp O, Kroemer G (ed), *Cell senescence: methods and protocols*. Humana Press, Totowa, NJ.
 42. Gmoser R, Ferreira JA, Lennartsson PR, Taherzadeh MJ. 2017. Filamentous ascomycetes fungi as a source of natural pigments. *Fungal Biol Biotechnol* 4:4. <https://doi.org/10.1186/s40694-017-0033-2>.
 43. Li J, Zhou L, Yin C, Zhang D, Klosterman SJ, Wang B, Song J, Wang D, Hu X, Subbarao KV, Chen J, Dai X. 2019. The *Verticillium dahliae* Sho1-MAPK pathway regulates melanin biosynthesis and is required for cotton infection. *Environ Microbiol* 21:4852–4874. <https://doi.org/10.1111/1462-2920.14846>.
 44. Zhou S, Zhang P, Zhou H, Liu X, Li S-M, Guo L, Li K, Yin W-B. 2019. A new regulator RsdA mediating fungal secondary metabolism has a detrimental impact on asexual development in *Pestalotiopsis fici*. *Environ Microbiol* 21:416–426. <https://doi.org/10.1111/1462-2920.14473>.
 45. Schumacher J, Simon A, Cohrs KC, Viaud M, Tudzynski P. 2014. The transcription factor BcLTF1 regulates virulence and light responses in the

- necrotrophic plant pathogen *Botrytis cinerea*. PLoS Genet 10:e1004040. <https://doi.org/10.1371/journal.pgen.1004040>.
46. Xin C, Yang J, Mao Y, Chen W, Wang Z, Song Z. 2020. GATA-type transcription factor MrNsdD regulates dimorphic transition, conidiation, virulence and microsclerotium formation in the entomopathogenic fungus *Metarhizium rileyi*. Microb Biotechnol 13:1489–1501. <https://doi.org/10.1111/1751-7915.13581>.
 47. Coppin E, Silar P. 2007. Identification of PaPKS1, a polyketide synthase involved in melanin formation and its use as a genetic tool in *Podospira anserina*. Mycol Res 111:901–908. <https://doi.org/10.1016/j.mycres.2007.05.011>.
 48. Künzler M. 2018. How fungi defend themselves against microbial competitors and animal predators. PLoS Pathog 14:e1007184. <https://doi.org/10.1371/journal.ppat.1007184>.
 49. Pang G, Sun T, Yu Z, Yuan T, Liu W, Zhu H, Gao Q, Yang D, Kubicek CP, Zhang J, Shen Q. 2020. Azaphilones biosynthesis complements the defence mechanism of *Trichoderma guizhouense* against oxidative stress. Environ Microbiol 22:4808–4824. <https://doi.org/10.1111/1462-2920.15246>.
 50. Rizet G, Esser K. 1953. Sur des phénomènes d'incompatibilité entre souches d'origines différentes chez *Podospira anserina*. C R Hebd Seances Acad Sci Paris 237:760–761.
 51. El-Khoury R, Sellem CH, Coppin E, Boivin A, Maas MFPM, Debuchy R, Sainsard-Chanet A. 2008. Gene deletion and allelic replacement in the filamentous fungus *Podospira anserina*. Curr Genet 53:249–258. <https://doi.org/10.1007/s00294-008-0180-3>.
 52. Xie N, Chapeland-Leclerc F, Silar P, Ruprich-Robert G. 2014. Systematic gene deletions evidences that laccases are involved in several stages of wood degradation in the filamentous fungus *Podospira anserina*. Environ Microbiol 16:141–161. <https://doi.org/10.1111/1462-2920.12253>.
 53. Benkhali JA, Coppin E, Brun S, Peraza-Reyes L, Martin T, Dixelius C, Lazar N, van Tilbeurgh H, Debuchy R. 2013. A network of HMG-box transcription factors regulates sexual cycle in the fungus *Podospira anserina*. PLoS Genet 9:e1003642. <https://doi.org/10.1371/journal.pgen.1003642>.
 54. Munkres KD. 1990. Histochemical detection of the secretion of superoxide radicals and hydrogen peroxide by *age-1* mutants of *Neurospora*. Fungal Genetics Rep 37:12. <https://doi.org/10.4148/1941-4765.1477>.
 55. Malagnac F, Lalucque H, Lepère G, Silar P. 2004. Two NADPH oxidase isoforms are required for sexual reproduction and ascospore germination in the filamentous fungus *Podospira anserina*. Fungal Genet Biol 41:982–997. <https://doi.org/10.1016/j.fgb.2004.07.008>.
 56. Colot HV, Park G, Turner GE, Ringelberg C, Crew CM, Litvinkova L, Weiss RL, Borkovich KA, Dunlap JC. 2006. A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. Proc Natl Acad Sci U S A 103:10352–10357. <https://doi.org/10.1073/pnas.0601456103>.