

Towards a unifying, systems biology understanding of large-scale cellular death and destruction caused by poorly liganded iron: Parkinson's, Huntington's, Alzheimer's, prions, bactericides, chemical toxicology and others as examples

Douglas B. Kell

Received: 14 June 2010 / Accepted: 14 July 2010 / Published online: 17 August 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract Exposure to a variety of toxins and/or infectious agents leads to disease, degeneration and death, often characterised by circumstances in which cells or tissues do not merely die and cease to function but may be more or less entirely obliterated. It is then legitimate to ask the question as to whether, despite the many kinds of agent involved, there may be at least some **unifying mechanisms** of such cell death and destruction. I summarise the evidence that in a great many cases, one underlying mechanism, providing major stresses of this type, entails *continuing and autocatalytic production* (based on positive feedback mechanisms) of hydroxyl radicals via Fenton chemistry involving **poorly liganded iron**, leading to cell death via apoptosis (probably including via pathways induced by changes in the NF- κ B system). While every pathway is in some sense connected to every other one, I highlight the literature evidence suggesting that the degenerative effects of **many diseases and toxicological insults converge on iron dysregulation**. This highlights specifically the role of iron metabolism, and the detailed speciation of iron, in chemical and other toxicology, and has significant implications for the use of iron chelating substances (probably in *partnership* with appropriate antioxidants) as nutritional or therapeutic agents in inhibiting both the progression of these mainly degenerative diseases and the sequelae of both chronic and acute toxin exposure. The complexity of biochemical networks, especially those involving autocatalytic behaviour and positive feedbacks,

means that multiple interventions (e.g. of iron chelators plus antioxidants) are likely to prove most effective. A variety of systems biology approaches, that I summarise, can predict both the mechanisms involved in these cell death pathways and the optimal sites of action for nutritional or pharmacological interventions.

Keywords Antioxidants · Apoptosis · Atherosclerosis · Cell death · Chelation · Chemical toxicology · Iron · Neurodegeneration · Phlebotomy · Polyphenols · Sepsis · SIRS · Stroke · Systems biology · Toxicity

Abbreviations


A β	β -amyloid
AD	Alzheimer's disease
ALS	Amyotrophic lateral sclerosis (Lou Gehrig's disease)
BSE	Bovine spongiform encephalopathy
CJD	Creutzfeldt-Jakob disease
CNS	Central nervous system
EAE	Experimental autoimmune encephalomyelitis
GBA	Glucocerebrosidase
HD	Huntington's disease
Htt	Huntingtin protein
MCMC	Markov chain Monte Carlo
MIRIAM	Minimum information requested in the annotation of biochemical models
MPTP	1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine
MS	Multiple sclerosis
PD	Parkinson's disease
RNS(s)	Reactive nitrogen species
ROS(s)	Reactive oxygen species
SBML	Systems biology markup language

D. B. Kell (✉)
School of Chemistry and the Manchester Interdisciplinary
Biocentre, The University of Manchester,
Manchester M1 7DN, UK
e-mail: dbk@manchester.ac.uk
URL: <http://dbkgroup.org/>; <http://twitter.com/dbkell>


SBRML	Systems biology results markup language
SIRS	Systemic inflammatory response syndrome
TCDD, 'dioxin'	2,3,7,8-tetrachlorodibenzo- <i>p</i> -dioxin
TSEs	Transmissible spongiform encephalopathies
2,4-D	2,4-dichlorophenoxyacetic acid
2,4,5-T	2,4,5-trichlorophenoxyacetic acid

Introduction

As a transition metal that can exist in several valencies, and that can bind up to six ligands, iron is an important component of industrial catalysts in the chemical industry (Hagen 2006), especially for redox reactions. Its catalysis of specific reactions requires rather exact architectures at the catalytic centre, and indeed much of the art and science of catalyst production involves determining and synthesising them. Nearly half of all enzymes are metalloproteins (Waldron et al. 2009), and iron is also of considerable importance in biology as a component of all kinds of metalloproteins (Andreini et al. 2008, 2009) from haemoglobin to cytochromes, as well as in the directed evolution of *novel* enzyme activities (Pordea and Ward 2008; Que and Tolman 2008; Turner 2009). When serving in enzymes, the iron is normally safely liganded, and any reactions catalysed are usually fairly specific. However, as is widely recognised, iron can also have a dark side (Kell 2009a), in that when it is not properly liganded (Graf et al. 1984), and in the ferrous form, it can react with hydrogen peroxide (produced by mitochondria (e.g. Brennan and Kantorow 2009; Fato et al. 2008; Orrenius et al. 2007) or (per)oxidases (Bedard and Krause 2007; Cave et al. 2006) via the Fenton reaction (Goldstein et al. 1993; Kruszewski 2003; Toyokuni 2002; Wardman and Candeias 1996; Winterbourn 1995), leading to the very reactive and damaging hydroxyl radical (OH[•])



Superoxide (also produced by mitochondria) can react with ferric iron in the Haber-Weiss reaction (Kehrer 2000) to produce Fe(II) again, thereby effecting redox cycling of the iron (Fig. 1):



Ascorbate (vitamin C) can also replace O₂^{•-} for reducing the Fe(III) to Fe(II) (Hershko and Weatherall 1988), as can other reducing agents, and indeed too low a redox poise leads to DNA damage (e.g. Li and Marbán 2010; Seifried et al. 2007).

The hydroxyl radical is exceptionally reactive and damaging to cellular components, and, for instance, can liberate




Fig. 1 The Haber-Weiss and Fenton reactions combine using poorly liganded iron in a **catalytic cycle** to produce the very damaging hydroxyl radical. Poorly liganded iron can also be liberated via the destruction of haem and other iron-containing substances. Peroxynitrite anion (ONOO⁻) is produced by the reaction of superoxide and nitric oxide (NO[•]) which when protonated (pH ca 6.5–6.8) decomposes to OH[•] and NO₂

further Fe(II) from iron-sulphur centres and other iron-containing compounds such as ferritin (Arosio et al. 2009), thereby driving reaction (1) in an autocatalytic, runaway kind of reaction. This kind of phenomenon has the potential to overwhelm any kinds of attempts at repair, and inflammation and oxidative stress are the hallmarks of each of the conditions I summarise. Related reactions include peroxynitrite production (from the reaction of NO and superoxide) (Babior 2000; Beckman et al. 1990; Beckman and Koppenol 1996; Goldstein and Merényi 2008; Koppenol et al. 1992; Murphy et al. 1998; Pacher et al. 2007; Pavlovic and Santaniello 2007; Pryor and Squadrito 1995; Radi et al. 2001, 2002; Rubbo and O'Donnell 2005; Rubbo et al. 2009; Smith et al. 1997b; Squadrito and Pryor 1998; Szabo 1996; Szabó et al. 2007; Torreilles et al. 1999; White et al. 1994; Zimmet and Hare 2006). These can lead to nitrotyrosine (Beckman 1996; Goldstein and Merényi 2008; Herce-Pagliai et al. 1998) (a reaction catalysed by poorly liganded iron, Beckman et al. 1992), or nitro-fatty acid (Aslan et al. 2001;

Some small molecules derived from macromolecule oxidation

- DNA → 8-OH-2'-deoxyguanosine
- Proteins → nitrotyrosine, methionine sulphoxime
- Lipids → 4-hydroxy-non-2-enal

Fig. 2 Some small molecules that are derived from the oxidative attack of hydroxyl and other radicals on cellular macromolecules and that can act as biomarkers of oxidative stress, including that mediated by iron

O'Donnell and Freeman 2001) production or protein cysteine nitrosylation (Lancaster 2008; Landino 2008; Vaz and Augusto 2008) that can provide a means of their detection downstream. Some of these are shown in Fig. 2. A key point here is that despite the widespread and uncritical use of the term ROS to describe any 'Reactive Oxygen Species', most such as superoxide and peroxide are not terribly reactive, in contrast to the hydroxyl radical (and peroxyxynitrite) which is, and **unliganded iron is required for hydroxyl radical production in the Fenton reaction**. Hence the focus on unliganded iron rather than the more nebulous ROSs, albeit that (su)peroxide is necessarily involved.

Clearly the occurrence of these activities leads to the (more or less irreversible) formation of a variety of substances that can act as biomarkers of these activities (Kell 2009a), and such oxidised compounds are represented by both small and macro-molecules. For instance, 8-hydroxy-2'-deoxyguanosine (8-OHdG) (ChEBI 40304) also referred to as its tautomer 8-oxo-7,8-dihydro-2'-deoxyguanosine (8-oxodG) (HMDB03333) is produced when the hydroxyl radical reacts with DNA and damages it (e.g. Bal and Kasprzak 2002; Burrows and Muller 1998; Cooke et al. 2003, 2008; Lloyd et al. 1998; Loft and Poulsen 1996; Orhan et al. 2004; Shi et al. 2003; Toyokuni and Sagripanti 1996; Valavanidis et al. 2009; Valko et al. 2005), and **body-iron status correlates rather strongly with its production or urinary excretion** (Agarwal et al. 2004; Broedbaek et al. 2009; Fujita et al. 2007a, 2009; Gackowski et al. 2002; Hori et al. 2010; Kang et al. 1998; Kuo et al. 2008; Maruyama et al. 2007; Nakano et al. 2003; Toyokuni and Sagripanti 1996; Tuomainen et al. 2007; Valavanidis et al. 2005). The data in (Hori et al. 2010)

relating urinary 8-OHdG levels to serum ferritin concentrations are particularly striking, and I have digitised them (cf. Pettifer et al. 2009 for pointers to automated ways of doing this in the future) and replotted some of them in Fig. 3.

Ferritin levels (a common measure of body iron stores) are also widely associated with disease development/severity/ poor outcomes (e.g. Alissa et al. 2007; Armand et al. 2007; Bartzokis et al. 2007; Braun et al. 2004; Bugianesi et al. 2004; Chen et al. 2006; Choi et al. 2005; Double et al. 2000; Fargion et al. 2001; Fernández-Real et al. 2002; Ford and Cogswell 1999; Forouhi et al. 2007; He et al. 2007; Hubel et al. 2004; Ishizaka et al. 2005; Jehn et al. 2004, 2007; Kaur et al. 2007; Kiechl et al. 1997; Lee et al. 2006f; Lim et al. 2001; Rayman et al. 2002; Rouault 2006; Salonen et al. 1992; Sheth and Brittenham 2000; Tuomainen et al. 1998; Valenti et al. 2007; Wilson et al. 2003; You et al. 2003; You and Wang 2005) (Adamkiewicz et al. 2009; Baune et al. 2010; Busca et al. 2010; DePalma et al. 2010; Ferrara et al. 2009; Ferrucci et al. 2010; Gamberini et al. 2008; Goodall et al. 2008; Inati et al. 2010; Kaysen 2009; Knovich et al. 2009; Kolberg et al. 2009; Lecube et al. 2008; Lim et al. 2010; Mateo-Gallego et al. 2010; McNeill et al. 2008; Menke et al. 2009; Qureshi et al. 2008; Rajpathak et al. 2009; Sharifi et al. 2008; Skinner et al. 2010; Song et al. 2009; Storey et al. 2009; Sun et al. 2008; Tsimikas et al. 2009; Valenti et al. 2010; Walker et al. 2010; Wang et al. 2010b; Yoneda et al. 2010; Zandman-Goddard and Shoenfeld 2008), though note that most assays are for the protein itself, and not for the full molecule including any iron that it may sequester effectively (Balla et al. 1992; Hintze and Theil 2006; Theil 2007) or otherwise.

Although iron is most commonly bivalent, Fe(II), or trivalent, Fe(III), this simple statement does not remotely cover the relevant chemistry and speciation that are necessary to recognise what forms of 'iron' may be safe and which are likely to catalyse damaging reactions. The first issue is that at neutral pH Fe(III) is more or less insoluble, and in aerobic environments it is necessary to chelate the otherwise 'free' Fe(III) with appropriate chelators or siderophores (especially for microbes, e.g. Andrews et al. 2003; Barry and Challis 2009; Cornelis and Andrews 2010; de Carvalho and Fernandes 2010; Haas et al. 2008; Johnson 2008; Miethke and Marahiel 2007; Raymond et al. 2003; Sandy and Butler 2009; Wandersman and Delepelaire 2004; Winkelmann 2002). These siderophores—secreted as are other bacterial pheromones (Kell et al. 1995)—typically have extremely tight binding constants ($K_f > 10^{30}$, e.g. Clifton et al. 2009; Loomis and Raymond 1991) and can solubilise and sequester iron such that it can be internalised via suitable transporter molecules within the plasma membrane (Stintzi et al. 2000).




Fig. 3 Very strong relationship between serum ferritin concentrations and urinary concentrations of the DNA damage/oxidative stress marker 8-hydroxy-2'-deoxyguanosine. Data are replotted from Fig. 1 of Hori et al. (2010)

Iron contains up to six individual chelation sites, arranged octahedrally, and ligands can typically satisfy them partially (i.e. some ligands are bidentate or tridentate and need three or two molecules for full liganding/activity) or fully i.e. are hexadentate. **Since iron cannot be transmuted into any other substance, the only way to stop the damaging activity of free or partially liganded 'iron' is to ensure that all of its six possible liganding sites are satisfied, whether by endogenous chelators or those added from the diet or as pharmaceuticals.** Put another way, it is not simply enough to know that 'iron' is present at an adequate level but that it is available *in a suitably liganded form*. **Anaemia can be caused by poor liganding as well as by an actual shortage of 'iron' itself.** Note too that *partial* chelation in the presence of an antioxidant agent such as ascorbate (vitamin C) can in fact make ascorbate (or other reducing agent) act as a *pro-oxidant* and

thus actually *promote* the production of OH^\bullet radicals in the presence of inappropriately or inadequately liganded Fe(II) (Allen and Cornforth 2009; Fábíán and Csordás 2003; Halliwell 2009; Hininger et al. 2005; Lachili et al. 2001; Long et al. 2000; Miller et al. 1990; Reif 1992; Sugihara et al. 1999). This very likely explains the often and indeed surprisingly disappointing clinical results obtained when using antioxidants alone (Bjelakovic et al. 2008; Giustarini et al. 2009; Kell 2009a; Miller et al. 2005).

The above facts are well known (e.g. Halliwell and Gutteridge 2006; Weinberg 2004), and I discussed them at length in a recent and wide-ranging review (Kell 2009a). My purpose here is to look more closely at the evidence that they are part of the sequelae of a number of predispositions to cellular and organismal death (whether by necrosis or apoptosis) that follow from a large variety of initial 'insults' or distal causes, whether of genetic or




Fig. 4 A mind map (Buzan 2002) setting out the structure of this review. To read this start at “1 o’clock” and move outwards and clockwise

environmental origin (or both). While, as a systems property, many other cellular processes contribute measurably to any specific activity, **I suggest, in all the disparate cases I review, that it is the binding of poorly liganded iron (mainly bivalent) to inappropriate cellular structures that serves to generate this catalytic activity, albeit that the overall manifestations may differ at a physiological level, and I here review what evidence is available. This basic suggestion can be tested explicitly more or less easily.** I suggest further that it is this ongoing, autocatalytic activity based on positive feedback that is responsible for the really large-scale damage, leading to organismal death, that can occur in affected cells, tissues, organs and organisms. **This analysis also suggests an important role for iron chelation as part of combination approaches in the acute and chronic therapy of these conditions.** An overview of this article is given as a Mind Map (Buzan 2002) in Fig. 4.

Note that I do recognise that other transition and polyvalent metal ions (Al^{n+} , Cr^{n+} , Cu^{n+} , Mn^{n+} , Se^{n+} , Zn^{n+} etc.) may also contribute to the kinds of process I describe. However, for reasons of simplicity, focus, and because of the natural abundance of this metal in biological systems, it is 'iron' on which I shall concentrate. As previously (Kell 2009a), **I use the term 'iron' to include iron of any valencies or speciation**, unless specified otherwise.

Since the role of iron is obvious in cases of primary diseases of iron overload, such as hereditary haemochromatosis (e.g. Camaschella and Merlini 2005; Ellervik et al. 2007; Gan et al. 2010; Limdi and Crampton 2004; Mair and Weiss 2009; Marx 2002; McLaren and Gordeuk 2009; Pantopoulos 2008; Papanikolaou and Pantopoulos 2005; Pietrangelo 2006; Weiss 2010), thalassaemias (e.g. Borgna-Pignatti et al. 2005; Camaschella and Merlini 2005; Cao and Galanello 2010; Cohen et al. 2004; Gattermann 2009; Lam et al. 2008; Li et al. 2010; Mohkam et al. 2008; Peng et al. 2008; Taher et al. 2010; Vichinsky et al. 2005) and myelodysplastic syndrome (e.g. Cazzola et al. 2008; Cuijpers et al. 2010; Dreyfus 2008; Gattermann 2008; Greenberg 2006; Greenberg et al. 2008; Jabbour et al. 2009; Jädersten and Hellström-Lindberg 2010; Leitch 2007; Mahesh et al. 2008; Malcovati 2009; Porter et al. 2008; Wimazal et al. 2009), I largely ignore this literature. However, I recognise that the sequelae of iron overload, especially various kinds of organ failure, share many similarities to those I describe below, consistent with the role of iron in these other cases where its involvement has been less widely recognised. Clearly the commonality of any specific effect with those of known iron overload (Dever et al. 2010) might give strong hints for the involvement of iron in specific toxicological processes.

The literature survey at the time of initial submission for review extended to 14 June, 2010.

To start our survey, we look at the consequences of an acute cerebral infarction or stroke.

Stroke

Stroke is a term used to describe the destruction of brain cells, typically following a temporary blockage in blood flow (ischaemia-reperfusion injury) or an intra-cranial haemorrhage. The extent of damage varies considerably, and can be exacerbated because affected cells can release inflammatory cytokines that activate other cells in a similar way, providing a positive autocatalytic effect and leading to spreading of the damage. It is this secondary spreading that is especially damaging, but because it is slower (4–7 days), it also affords the opportunity for therapeutic intervention (Qureshi et al. 2009). It is now clear that iron is intimately involved (Armengou and Davalos 2002; Bailey et al. 2006; Bishop and Robinson 2001; Chang et al. 2005; Dávalos, et al. 1994; Demougeot et al. 2004; Ferro and Dávalos 2006; Garoufi et al. 2006; Gillum et al. 1996; Lee et al. 2006e; Marniemi et al. 2005; Mascitelli and Pezzetta 2006; Mehta et al. 2004; Mu et al. 2005; Nakamura et al. 2006; Saxena et al. 2005; Selim and Ratan 2004; Switzer et al. 2006; Wagner et al. 2003; Wu et al. 2003; Zuliani et al. 2006; Adams 2007; Altamura et al. 2009; Assenza et al. 2009; Basak et al. 2008; Bosomtwi et al. 2008; Carbonell and Rama 2007; Cho et al. 2007; Ekblom et al. 2007; Hanson et al. 2009; Heckl 2007; Helal 2008; Jolkkonen et al. 2007; Justicia et al. 2008; Kaushal and Schlichter 2008; Kim et al. 2008a; Kobayashi et al. 2008; Lou et al. 2009; Maguire et al. 2007; Mazumdar et al. 2007; Mehdiratta et al. 2008; Millan et al. 2007, 2008; Millerot-Serruot et al. 2008; Nighoghossian et al. 2008; O'Rourke et al. 2008; Ratan et al. 2008; Ross and Meschia 2009; Saleh et al. 2007; Santhosh et al. 2009; Verduzco and Nathan 2009; Walters and Rye 2009; Weng et al. 2008), whether by release from (ferritin in) cells or from the haem of haemoglobin. Iron chelators have thus shown promise in decreasing the sequelae of an initial stroke-inducing event (Demougeot et al. 2004; Ferro and Dávalos 2006; Hurn et al. 1995; Kompala et al. 1986; Mu et al. 2005; Patt et al. 1990; Prass et al. 2002; Selim and Ratan 2004; Soloniuk et al. 1992; White et al. 1988; Chen-Roetling et al. 2009; Gu et al. 2009; Hanson et al. 2009; Hua et al. 2008; Mazumdar et al. 2007; Méthy et al. 2008; Millerot-Serruot et al. 2008; Mirre et al. 2010; Okauchi et al. 2009, 2010; Ratan et al. 2008; Robinson et al. 2009; Selim 2009; Verduzco and Nathan 2009), providing further evidence for the primary importance of iron in causing injury.

While stroke is a sudden occurrence, albeit with secondary consequences, a number of neurodegenerative diseases are rather more long term in their development, and

we now look at several, each of which is seen to involve iron intimately.

Huntington's disease

The pathology of HD reveals striking neurodegeneration in the corpus striatum and shrinkage of the brain, leading to its most prominent manifestation, viz. movement disorders or chorea (Bhidayasiri and Truong 2004). Huntington's disease occurs via the addition of trinucleotide CAG repeats within exon 1 of the relevant gene, encoding (poly)glutamine (polyQ) repeats in the huntingtin protein (htt) and leading to a gain of (toxic) function (Bauer and Nukina 2009; Gusella and MacDonald 2000; Imarisio et al. 2008; Quintanilla and Johnson 2009). The number of these repeats determines both the time of onset of observable disease (more repeats meaning earlier onset) (Perutz and Windle 2001; Ross 1995; Walters and Murphy 2009), accounting for 70% of the variance (Imarisio et al. 2008), and the disease severity, implying that it is indeed the polyglutamines *themselves* that are the chief culprits. Some of the polyQ-containing huntingtin proteins can also aggregate to form inclusion bodies, and aggregation and neurodegeneration can in part be related (Chopra et al. 2007; Cowan et al. 2003; Michalik and Van Broeckhoven 2003; Nagai et al. 2003; Wang et al. 2005, 2009; Zhang et al. 2005a). Now the evidence for the involvement of poorly liganded iron in a wide variety of neurodegenerative diseases is overwhelming (e.g. Benarroch 2009; Bishop et al. 2010a; Brown 2009a; Friedman et al. 2007; Jellinger 1999; Ke and Qian 2003; Kell 2009a; Lee et al. 2006a; Perez and Franz 2010; Thompson et al. 2001; Youdim et al. 2004b; Zecca et al. 2004), and the question arises as to whether huntingtin (or, more likely, its degradation fragments), containing these polyQ tracts, can catalyse oxidative stress and/or hydroxyl formation directly. In an important paper, Firdaus et al. (Firdaus et al. 2006) show that that huntingtin inclusion bodies act as centres of oxidative stress and that partially purified inclusion bodies contain large amounts of oxidised proteins. Iron metabolism is deranged in gene knockdown models of Huntington's (Henshall et al. 2009; Lumsden et al. 2007), and experiments with the iron chelator deferoxamine revealed that the oxidation, localisation and structural organisation of the inclusion bodies formed by mutant htt were indeed iron-dependent. They did not demonstrate the ability of these inclusion bodies to catalyse hydroxyl formation, but this could easily be shown using specific assays, such as the hydroxylation of suitable aromatics (e.g. Grootveld and Halliwell 1986; Halliwell and Gutteridge 2006; Thomas et al. 2009a). The iron chelator clioquinol is also protective (Nguyen et al. 2005).

Parkinson's disease

Parkinson's disease (PD) results primarily from the death of dopaminergic neurons in the substantia nigra part of the brain and is characterised in particular by the presence of intracytoplasmic inclusions from protein aggregates called Lewy bodies (LB) and at the physiological level by a variety of movement disorders (e.g. Singh et al. 2007). It is strongly linked with oxidative stress (e.g. Alam et al. 1997; Büeler 2009; Jenner 2003; Jenner and Olanow 1996; Kidd 2000; Mandel et al. 2003; Olivares et al. 2009; Seet et al. 2010; Van Laar and Berman 2009; Zhang et al. 1999). Again (e.g. Berg et al. 2001; Bharath et al. 2002; Buchanan et al. 2002; Bush 2000; Castellani et al. 2002; Double et al. 2000; Good et al. 1998; Gotz et al. 2004; Hirsch and Faucheux 1998; Jellinger 1999; Johnson 2000; Olanow and Arendash 1994; Ostrerova-Golts et al. 2000; Sofic et al. 1991; Wolozin and Golts 2002; Altamura and Muckenthaler 2009; Andersen 2004; Barapatre et al. 2010; Baudrexel et al. 2010; Becker 2010; Becker et al. 2010a, b; Berg 2006, 2007; Berg and Hochstrasser 2006; Berg et al. 2006, 2008; Brar et al. 2009; Brown 2009b; Crichton and Ward 2006; Fasano et al. 2006; Friedman et al. 2007, 2009; Ghosh et al. 2010; Hirsch 2009; Jellinger 2009; Jimenez-Del-Rio et al. 2010; Kaur and Andersen 2004; Lee and Andersen 2010; Levenson 2003; Mandemakers et al. 2007; Matusch et al. 2010; Oakley et al. 2007; Salvador 2010; Shi et al. 2010; Wayne Martin 2009; Wypijewska et al. 2010; Yeager and Coleman 2010; Zecca et al. 2004; Zhang et al. 2005, 2010b)—there is overwhelming evidence for the involvement of iron in this neurodegeneration. Why these dopaminergic neurons are especially susceptible is not entirely clear, although dopamine is capable of reacting with iron directly to form a toxic complex (Arreguin et al. 2009; Paris et al. 2005) that probably itself catalyses hydroxyl formation. Iron chelators inhibit these inimical processes (Gal et al. 2006; Perez et al. 2008; Xu et al. 2008; Youdim et al. 2004a, b; Zheng et al. 2005), including (Kaur et al. 2003; Reznichenko et al. 2010; Youdim 2003) in the case of a Parkinson-like disease induced (Blum et al. 2001; Choi et al. 2009; Gal et al. 2009; Yokoyama et al. 2008) by 1-methyl-4-phenyl-1,2,3,6-tetrahydro-pyridine (MPTP) (ChEBI 17963). The Lewy bodies contain lipid and a variety of proteins, including ubiquitin, neurofilament, various proteasomal elements and α -synuclein, which may be oxidatively modified (Double et al. 2008). Ferric iron may itself catalyse the formation of α -synuclein oligomers (Brown 2009a; Hillmer et al. 2009; Peng et al. 2010), and copper may also be involved (Wang et al. 2010c).

Melanins are polymers of polyphenols, especially of L-dopamine, although neuromelanin, which also contains oxidatively polymerised dopamine or noradrenaline with the possible involvement of cysteinyl derivatives (Fedorow

et al. 2005), is of special interest here. The substantia nigra also contains a substantial amount of the dark, insoluble polymeric pigment neuromelanin, and this has been shown directly to bind (and release) iron in high amounts (Double et al. 2003a, b; Fasano et al. 2006; Faucheux et al. 2003; Gerlach et al. 2003, 2008; Shamoto-Nagai et al. 2006), and thereby to produce an iron-loaded form that seems to serve as a reservoir catalysing (hydroxyl) radical formation and whose amounts correlate with PD. In this case, the in vitro experiments have been done, and incubation of human neuromelanin with iron in vitro stimulates oxidative tissue damage (Ben-Shachar et al. 1991; Double et al. 2003a; Gerlach et al. 2008).

Another dark pigment commonly observed in PD patients (and other neurodegenerative diseases) is lipofuscin (Double et al. 2008; Jung et al. 2007; Terman and Brunk 2004). Lipofuscin too consists of lipids and oxidised proteins, often crosslinked by oxidised lipid derivatives such as (E)-4-hydroxy-non-2-enal (ChEBI 58968) and can adsorb high contents of metals (including iron) of up to 2% by weight (Jung et al. 2007). It is typically formed in lysosomes from degrading mitochondria (Terman et al. 2006a), themselves rich in metalloproteins, and the inability of cells to degrade it means that (purportedly) non-dividing cells (such as brain tissue) simply accumulate it as cells age (Terman and Brunk 2004, 2006). This arguably explains why non- or rarely-dividing tissues such as brain tissues are particularly susceptible, though note the important and increasing evidence (e.g. Abrous et al. 2005; Christie and Cameron 2006; Demir et al. 2009; Fuchs and Gould 2000; Götz and Huttner 2005; Gould 2007; Imayoshi et al. 2009; Leuner et al. 2007; Lledo et al. 2006; Ohnuma and Harris 2003; Taupin 2007; Zhao et al. 2008) for considerable turnover—neurogenesis—in at least some regions of the brain. The **autocatalytic nature** of the process—lipofuscin loaded with iron catalyses more lipofuscin production—is especially dangerous and can (as one would predict) be inhibited using appropriate iron chelators (Persson et al. 2003). Iron chelation also assists neurogenesis (Nowicki et al. 2009).

Thus, as with Huntington's, Parkinson's disease is clearly characterised by all the hallmarks of hyperactive hydroxyl radical generation, leading to cell death and destruction, in this case mainly, it would seem (Chen et al. 2007; Chiueh et al. 2000; Ekshyyan and Aw 2004; Jellinger 2002; Jenner 2003; Jenner and Olanow 1996; Kermer et al. 2004; Levenson 2005; Loh et al. 2006; Mandel et al. 2005; Mattson 2006; Okouchi et al. 2007; Xu et al. 2008a; Yasuda and Mochizuki 2010), by apoptosis.

Gaucher's disease

Gaucher's disease is an autosomally recessive inborn error of metabolism due to deficiency of a lysosomal enzyme,

glucocerebrosidase (GBA), resulting in the accumulation of glucocerebroside in large macrophages throughout the reticuloendothelial system, leading to various neuronopathies. Given that the lysosome is the site of most labile iron in the cell (Fakih et al. 2008; Gorria et al. 2008; Kurz et al. 2004, 2008a, b; Persson 2005; Tenopoulou et al. 2007; Terman et al. 2006b; Yu et al. 2003), it is also of interest that there is an increased frequency of mutations in the gene encoding GBA among patients with Parkinson's disease (e.g. Aharon-Peretz et al. 2004; Clark et al. 2007; Gan-Or et al. 2008; Lwin et al. 2004), and vice versa (Sidransky et al. 2009). Iron dysregulation is also well established in Gaucher's disease (Finch et al. 1986; Lee et al. 1967, 1977; Lorber 1960, 1970; Morgan et al. 1983; Schiano et al. 1993; Weisberger et al. 2004). This recognition of the role of iron dysregulation in Gaucher's disease may offer novel therapeutic approaches.

Alzheimer's disease

Alzheimer's disease (AD), the commonest of the neurodegenerative disease of ageing, shares many similarities with Huntington's and Parkinson's diseases, not least the extensive evidence for the role of oxidative stress (e.g. Butterfield et al. 2007; Christen 2000; DiMauro and Schon 2008; Good et al. 1996; Milton 2004; Miranda et al. 2000; Moreira et al. 2009; Nunomura et al. 2001, 2006; Reddy and Beal 2008; Reddy et al. 2009; Rottkamp et al. 2000; Smith et al. 1996, 2000; Zhu et al. 2007) and of iron (e.g. Adlard and Bush 2006; Avramovich-Tirosh et al. 2007a; Becker et al. 2010b; Bishop et al. 2002; Blázquez et al. 2007; Bolognin et al. 2009b; Brar et al. 2009; Bush 2000, 2003, 2008; Casadesus et al. 2004; Castellani et al. 2007; Collingwood and Dobson 2006; Collingwood et al. 2008; Connor and Lee 2006; Ding et al. 2009; Doraiswamy and Finefrock 2004; Gerlach et al. 1994; Good et al. 1996; Hegde et al. 2009; Honda et al. 2004; Jellinger et al. 1990; Jellinger 2009; Kala et al. 1996; Lehmann et al. 2006; LeVine 1997; Lovell et al. 1998; Malecki and Connor 2002; Mandel et al. 2007; Markesbery 1997; Markesbery and Lovell 1998; Mascitelli et al. 2009; Olanow and Arendash 1994; Ong and Farooqui 2005; Quintana et al. 2006; Rival et al. 2009; Robson et al. 2004; Silvestri and Camaschella 2008; Smith et al. 1997a, 2010; Tabner et al. 2005; Thomas and Jankovic 2004; Thompson et al. 2001; Valko et al. 2005; Zatta et al. 2009; Zecca et al. 2004; Zheng et al. 2005). It is especially noteworthy that iron correlates with disease severity as measured by cognitive ability (Ding et al. 2009; Gómez Ravetti et al. 2010; Grossi et al. 2009; Lavados et al. 2008; Perez et al. 2010; Smith et al. 2010). AD is characterized by the loss of neurons in the cognitive centres of the brain and by the presence of

two separate pathological lesions, extracellular β -amyloid (A β) plaques (e.g. Dong et al. 2003; Scott and Orvig 2009) and neurofibrillary tangles within neurons (Shcherbatykh and Carpenter 2007).

Again, there is evidence that insoluble polymers (fibrillary tangles) of proteins such as the β -amyloid can bind iron (Castellani et al. 2007; Dickens and Franz 2010; Exley 2006; Good et al. 1992; Jiang et al. 2009; Mancino et al. 2009; Rival et al. 2009; Sayre et al. 2000b; Smith et al. 2007) and thereby act in an autocatalytic manner to promote further radical production and oxidative stress. There is also a relationship between the ability to bind haem and neurotoxicity (Atamna 2009; Atamna et al. 2009), and (HFE) mutations that cause haemochromatosis increase the susceptibility to AD (Bartzokis et al. 2010; Candore et al. 2003; Combarros et al. 2003; Connor and Lee 2006; Haacke et al. 2005; Kauwe et al. 2010; Lleó et al. 2002; Moalem et al. 2000; Pulliam et al. 2003; Robson et al. 2004; Sampietro et al. 2001), especially when in the presence of the APOE4 allele (Pulliam et al. 2003). (HFE mutations also increase susceptibility to PD (Biasotto et al. 2008; Dekker et al. 2003)—but cf. (Aamodt et al. 2007)—and to ALS (Ellervik et al. 2007)).

Importantly, iron chelators have been shown to ameliorate the development of fibril formation/neurodegeneration/dementia/AD (Amit et al. 2008; Avramovich-Tirosh et al. 2007a; Bandyopadhyay et al. 2006; Barnham et al. 2004; Biran et al. 2009; Bolognin et al. 2009a; Crapper McLachlan et al. 1991; Cuajungco et al. 2000; Faux et al. 2010; Finefrock et al. 2003; Liu et al. 2009b, c, 2010; Mancino et al. 2009; Mandel 2007, 2008; Reznichenko et al. 2006; Scott and Orvig 2009; Weinreb et al. 2009a; Zheng et al. 2005). Although this fact rarely appears in papers setting out therapeutic options for preventing (or at least ameliorating the progress of) AD, this already considerable literature suggests that trials using modern iron chelators would be of worth.

Amyotrophic lateral sclerosis (ALS, Lou Gehrig's disease)

ALS is another neurodegenerative disease in which iron has been strongly implicated (and thus the same kinds of mechanism as described previously) (Bush 2000; Carri et al. 2003; Cozzolino et al. 2008; Goodall et al. 2008; Jeong et al. 2009; Kasarskis et al. 1995; Mattson 2004; Migliore et al. 2005; Molfino et al. 2009; Qureshi et al. 2008; Reynolds et al. 2007; Sadrzadeh and Saffari 2004; Sayre et al. 2000a; Scott and Orvig 2009; Spasojević et al. 2010; Sutudja et al. 2007; Wang et al. 2004). Again, the clear benefits of iron chelators in mouse models (Jeong et al. 2009; Petri et al. 2007) and elsewhere (Avramovich-

Tirosh et al. 2007b; Bolognin et al. 2009a) would seem to merit trials in humans known to be at risk. Long-standing associations of some forms of ALS with mutations in genes coding for superoxide dismutase 1 (e.g. Dalle-Donne 2007; Pasinelli and Brown 2006; Rosen et al. 1993; Vucic and Kiernan 2009; Wijesekera and Leigh 2009) are consistent with this, and recent genetic associations (e.g. Kwiatkowski Jr. et al. 2009; Valdmanis et al. 2009; Vance et al. 2009) include one with optineurin (Maruyama et al. 2010) that together with other evidence implies an interaction with NF- κ B (see later) leading to the downstream effects of apoptotic neuronal cell death.

Friedreich's ataxia

Friedreich's ataxia is another neurodegenerative disease caused by the insertion of a trinucleotide repeat (or occasionally a missense mutation) in the gene encoding a protein called frataxin (Adibhatla and Hatcher 2010; Babady et al. 2007; Becker and Richardson 2001; Bencze et al. 2006; Chakravarty 2003; González-Cabo et al. 2009; Palau 2001; Pandolfo 2009; Patel and Isaya 2001; Puccio 2009; Rötig et al. 2002; Santos et al. 2010; Sheftel et al. 2010). Driven not least by studies in bacteria and lower eukaryotes (Ayala-Castro et al. 2008; Becker and Richardson 2001; Bulteau et al. 2007; Cavadini et al. 2000; Cook et al. 2006; De Freitas et al. 2003; Foury 1997; Foury and Cazzalini 1997; Foury and Talibi 2001; González-Cabo et al. 2005; Irazusta et al. 2006, 2008; Knight et al. 1999; Pastore et al. 2007; Vázquez-Manrique et al. 2006; Wang and Craig 2008; cf. Seguin, et al. 2010), it is now understood that frataxin is a mitochondrial iron chaperone protein (Atkinson and Winge 2009; Ba et al. 2009; Babady et al. 2007; Bencze et al. 2007; Cavadini et al. 2002; Correia et al. 2008; Gakh et al. 2006; Haugen et al. 2010; Huang et al. 2008a; Huang et al. 2009; Karlberg et al. 2006; Kondapalli et al. 2008; Lill 2009; Lill and Muhlenhoff 2008; Lu and Cortopassi 2007; MacKenzie et al. 2008; Mandemakers et al. 2007; Marmolino and Acquaviva 2009; Napier et al. 2005; Napoli et al. 2006; O'Neill et al. 2005; Pandolfo 2009; Pandolfo and Pastore 2009; Pastore et al. 2007; Reddy 2008; Schagerlöf et al. 2008; Sparaco et al. 2009; Wilson 2006; Zanella et al. 2008) involved in the safe insertion of Fe(II) during the production of Fe-S centres in the mitochondrial respiratory chain (Anderson et al. 2005). The defect thus leads to the liberation of mitochondrial iron (Huang et al. 2009; Popescu et al. 2007; Richardson et al. 2009a). Fenton chemistry is strongly implicated (Park et al. 2002; Sturm et al. 2005), as the attenuation of H₂O₂ production (but not of superoxide) (Anderson et al. 2005) ameliorates the disease (Anderson et al. 2008). As would be expected, appropriate iron chelation may have

therapeutic benefits (Boddaert et al. 2007; Campanella et al. 2009; Goncalves et al. 2008; Kakhlon et al. 2010; Lim et al. 2008; Marmolino and Acquaviva 2009; Santos et al. 2010; Schulz et al. 2009; Sohn et al. 2008). This very clear molecular picture underlines strongly the problems of iron release leading to cell death and destruction.

Multiple sclerosis

MS is an inflammatory demyelinating autoimmune disease affecting the CNS. While its underlying causes remain uncertain, and do not seem to have a significant genetic component (Baranzini et al. 2010; Oksenberg et al. 2008; Ramagopalan et al. 2008), ROS are certainly implicated (Adibhatla and Hatcher 2010), and antioxidant therapies may be useful (Mirshafiey and Mohsenzadegan 2009). It is at least plausible that part of the demyelination is caused by ROS and in particular hydroxyl radicals, and it is thus of much interest that iron dysregulation and/or deposition have indeed been implicated directly (Adams 1988; Bakshi et al. 2000, 2001, 2002, 2008; Bermel et al. 2005; Brass et al. 2006a, b; Ceccarelli et al. 2009, 2010; Chard and Miller 2009; Craelius et al. 1982; Drayer et al. 1987; Eissa et al. 2009; Exley et al. 2006; Filippi and Agosta 2009; Ge et al. 2007; Grimaud et al. 1995; Haacke et al. 2005, 2009, 2010; Hammond et al. 2008; He and Yablonskiy 2009; Khalil et al. 2009; LeVine and Chakrabarty 2004; Neema et al. 2007a, b, 2009a, b; Pirko et al. 2009; Sayre et al. 2005; Schenck and Zimmerman 2004; Simka and Rybak 2008; Singh and Zamboni 2009; Stankiewicz et al. 2007; Tjoa et al. 2005; Walter et al. 2009; Zamboni 2006; Zivadinov and Bakshi 2004; Zivadinov and Minagar 2009). While the extent of causality of the iron dysregulation in MS is not yet established, it is highly pertinent that iron chelators have been shown to suppress disease progression in experimental autoimmune encephalomyelitis (EAE), an animal model of multiple sclerosis (Bowern et al. 1984; Mitchell et al. 2007; Pedchenko and LeVine 1998). To say the least, the question of the involvement of iron in the human disease would seem to need pursuing with some urgency.

Age-related macular degeneration

While this review is intended to be much more circumscribed than the previous one (Kell 2009a), there are some pertinent diseases in which my understanding has improved significantly, and age-related macular degeneration is one such. Age-related macular degeneration (AMD) (Rattner and Nathans 2006) is recognised as the leading cause of blindness and visual disability in the elderly in developed

countries (Fine 2005; Harvey 2003; Lotery and Trump 2007; Tielsch et al. 1995). Interestingly, despite the relatively modest impact of genome-wide association studies to date (Maher 2008; Moore et al. 2010), and notwithstanding their promise as their statistical power improves, 80% of the variation in susceptibility to AMD is said to be explainable (Collins 2010) (and see e.g. Chu et al. 2008) though cf. e.g. (Chen et al. 2010; Hadley et al. 2010; Katta et al. 2009; Neale et al. 2010; Patel et al. 2008; Sobrin et al. 2010) by just two genes concerned with inflammation combined with two environmental factors (smoking and obesity). The environmental factors can certainly be related to iron dysregulation. Indeed, as previously discussed, iron is significantly implicated in AMD (and other related diseases such as glaucoma) (Blasiak et al. 2009; Cai et al. 2000; Charkoudian et al. 2008; Chen et al. 2009; Chowers et al. 2006; Dentchev et al. 2005; Dunaief 2006; García-Castiñeiras 2010; Goralska et al. 2009a, b, c; Hahn et al. 2003, 2004; He et al. 2007; Loh et al. 2009; Richer et al. 2002; Wong et al. 2007), iron chelation may help to reverse the process (Voloboueva et al. 2007), and dietary and other antioxidants are also protective (see Tan et al. 2008; Wang et al. 2009).

Prion diseases: including transmissible spongiform encephalopathies and Creutzfeldt–Jakob disease

A particularly striking set of diseases, where the downstream events manifest as extreme cellular destruction leading to large areas of (brain) tissue disappearing completely, are represented by the prion diseases, in particular the TSEs including bovine spongiform encephalopathy (BSE) (e.g. Aguzzi et al. 2008a; Aguzzi and Heikenwälder 2006; Caughey and Baron 2006; Caughey and Lansbury 2003; Linden et al. 2008) and scrapie (e.g. Bulgin and Melson 2007; Hur et al. 2002; Roostae et al. 2010; Wisniewski and Sigurdsson 2007), kuru (Gajdusek 2008; McLean 2008) and CJD (e.g. McLean 2008; Peden and Ironside 2004) (note too the co-location of amyloid plaques and spongiform degeneration in CJD, Ghoshal et al. 2009).

Now it is well known that prion diseases are caused in part (or at least accompanied by) the conversion of the prion protein PrP from its normal form PrP^C to a conformationally distinct version PrP^{Sc} and that the latter can autocatalyse this conversion (Aguzzi and Heikenwälder 2006; Aguzzi et al. 2008b; Prusiner 1998, 2001; Tamgüney et al. 2008; Watts and Westaway 2007). However, PrP^{Sc} levels are in many cases poorly correlated with prion disease progression, suggesting that it is necessary but not sufficient; ‘something else’ is required (Caughey and Baron 2006). Equally, it is far from clear how a simple conformational change of a protein leads to the vacuolation or

spongiosis (e.g. Aguzzi 2006; Aguzzi et al. 2007; Armstrong et al. 2001; Chung et al. 1999; Crozet et al. 2008; Diedrich et al. 1991; Foster et al. 2001; Julius et al. 2008; Kourie 2002; Mallucci et al. 2003; Mallucci et al. 2007; Miele et al. 2001; Sakudo and Ikuta 2009a, b; Westaway et al. 1994; Williams et al. 1997) (formation of holes) in brain tissue characteristic of late-stage prion disease (and indeed of Alzheimer's (Erkinjuntti et al. 1996; Sakudo and Ikuta 2009a). The question then arises as to whether it is possible that PrP^{Sc} modifies iron metabolism in an unfavourable way (and/or vice versa), not least since prion proteins bear phylogenetic relationships to the ZIP family of metal ion transporters (Schmitt-Ulms et al. 2009). Building on earlier work relating prion biology with (especially) iron-induced oxidative stress (e.g. Bareggi et al. 2009; Barnham et al. 2006; Basu et al. 2007; Choi et al. 2006; Feraeus et al. 2005a, b; Feraeus and Land 2005; Iwamaru et al. 2008; Kim et al. 2000, 2001, 2007; Lee et al. 2007; Lehmann 2002; Milhavet and Lehmann 2002; Pamplona et al. 2008; Park et al. 2008; Petersen et al. 2005; Pollera et al. 2005; Rana et al. 2009; Stegmann and Grohmann 2003; Westergard et al. 2007; Wong et al. 2001), this is exactly what Singh and colleagues (Singh et al. 2009a, b, c, 2010a, b) have now demonstrated, which is that the 'something else' (Kell 2009a, b) is (as so often) an imbalance in iron metabolism. They start by showing (Singh et al. 2009c) that even normal PrP^C mediates cellular iron uptake and transport and that mutant PrP forms alter cellular iron levels differentially.

PrP^C binds iron tightly, and this helps effect its conversion to PrP^{Sc} (Singh et al. 2010a, b). Singh et al. (2009a) showed that PrP^{Sc} can sequester cellular iron in insoluble PrP^{Sc}–ferritin complexes, making it bio-unavailable, and that this leads to an upregulation of iron uptake via transferrin and its receptor, leading to an overall iron excess in brain tissue. This was true for both scrapie and sporadic Creutzfeldt–Jakob disease, and stresses that it is not only the total amount of Fe(II) and Fe(III) that matter but their *speciation*. Mice knocked out for PrP also show derangement in iron metabolism (Singh et al. 2009b), and given what we know already (Kell 2009a) this leads to many obvious predictions of other diseases (Altamura and Muckenthaler 2009; Granic et al. 2009; Reddy et al. 2009; Serra et al. 2009) to which they might be susceptible. It has not yet been shown that PrP^{Sc}–ferritin complexes catalyse OH[•] production directly, but if this is increased in diseased brains this continuous, *autocatalytic* spewing out of toxic OH[•] radicals would clearly be able to account for the massive damage observed. A systems biology approach by Hwang et al. (Hwang et al. 2009) also found many genes of iron metabolism modified in prion disease. There is little doubt that all these new findings about the role of iron (and other metals, e.g. Brown et al. 1997) open up many new

inroads into understanding the important role of iron in prion diseases, and thereby novel therapeutic options involving both anti-oxidants and cell-permeable iron chelators. Some of these new ideas and findings are illustrated in Fig. 5.

Finally, here, it is worth pointing up the evidence for the role of iron and its metabolic dysregulation in the oxidative stress-mediated aetiology of scrapie (Choi et al. 2006; Feraeus et al. 2005a; Feraeus and Land 2005; Gudmundsdóttir et al. 2006; Kim et al. 2000, 2007; Pamplona et al. 2008; Singh et al. 2009a, 2010a, b; Wong et al. 2001a, b; Yun et al. 2006), a prion disease of sheep and goats (Bulgin and Melson 2007; Hur et al. 2002) and of CJD (Choi et al. 2006; Freixes et al. 2006; Petersen et al. 2005; Singh et al. 2009a, 2010a, b). As such, the use of geostatistics relating the variation of metals present in soils to prion disease prevalence is also a powerful tool here (e.g. Gudmundsdóttir et al. 2006; Imrie et al. 2009; Purdey 2000; Stevens et al. 2009).

Iron deposition and disease: cause or effect? The case of atherosclerosis

I have implied that the role of unliganded iron is *causative* of a variety of sequelae, but sometimes this is hard to infer as the networks in which iron is involved are multiple and complex (hence the need for a systems approach—see later), and often simple one-time snapshots of covariates do not permit the inference of causality. However, one effect of the reaction of hydroxyl radicals (their formation catalysed by unliganded iron) with proteins (or lipids) is to tend to make those proteins (or lipids) insoluble or fibrotic (Davies 2005; Fig. 6), and if one believes that this is




Fig. 5 Some of the interactions between the prion protein in its two main conformations, reactive oxygen species and iron dysregulation. This diagram is based on Fig. 12 of (Singh et al. 2010b), and illustrates in particular the autocatalytic nature of the ROS- and iron-dependent conversion of PrP^C to PrP^{Sc} and the neurotoxicity of the latter




Fig. 6 Iron catalyses the formation of the hydroxyl radical (and thence other kinds of ROS) that can react with proteins and lipids to denature them, leading to insoluble plaques and other fibrotic structures that can themselves bind/entrap the iron that caused their formation. This bound iron can cause further hydroxyl radical formation such that the process is autocatalytic. Eventually this overwhelms cellular defences, leading to cell death (with further release of iron)

causative, the presence of insoluble plaques should correlate with the presence of the iron that is seen to have caused their formation and thereby became entrapped (and may yet be further reactive). Because of its biomedical importance, considerable work has been performed on atherosclerotic lesions, which provide a very clear example.

Atherosclerosis is a progressive (and inflammatory) disease (Altman 2003; Binder et al. 2002; Blake and Ridker 2001; Dwevel et al. 2010; Dwyer et al. 2004; Forrester 2002, 2004; Gieseg et al. 2009; Grainger 2007; Hansson 2001, 2005; Himmelfarb et al. 2002; Kibel et al. 2008; Kunsch and Medford 1999; Libby 2002; Libby et al. 2002; Madamanchi et al. 2005a, b; Mullenix et al. 2005; Nigro et al. 2006; Packard and Libby 2008; Paoletti et al. 2004; Popa et al. 2007; Rader and Daugherty 2008; Ridker et al. et al. 2004; Ross 1999; Schleicher and Friess 2007; Subramanian and Ferrante 2009; Sullivan 2009; Tan and Lip 2008; Tang et al. 2009; Taqueti et al. 2006; Tedgui and Mallat 2006; van Leuven et al. 2008; van Oostrom et al. 2004; Willerson and Ridker 2004; Young et al. 2002) characterized by the accumulation of both oxidised lipids and various fibrous elements in arteries, often as plaques (Lusis 2000; Stocker and Keaney 2004). Several lines of evidence point to the involvement of iron in these processes:

- Both iron and oxidised lipids (de Valk and Marx 1999; Smith et al. 1992) are found in atherosclerotic lesions (Altamura and Muckenthaler 2009; Brewer 2007; Chau 2000; Fernandes de Godoy et al. 2007; Gajda et al. 2008; Halliwell 2009; Horwitz et al. 1998; Kazi et al. 2008; Lee et al. 1998; McRae et al. 2009; Rajendran et al. 2007; Ramakrishna et al. 2003; Roijers et al. 2005; Smith et al. 1992; Stadler et al. 2004; Stanley et al. 2006; Stocker and Keaney 2005; Sullivan 2009; Watt et al. 2006; Wolff et al. 2004; Yuan and Li 2008)




Fig. 7 Large (17-fold) accumulation of EPR-detectable iron in atherosclerotic plaques relative to healthy non-atherosclerotic ('intima') controls. The difference is highly significant ($P = 0.0001$). Data are replotted from those in Fig. 1b of Stadler et al. (2004)

- Iron depletion by dietary or other means delays the formation of such lesions (Ferrara and Taylor 2005; Halliwell 2009; Lee et al. 1999; Matthews et al. 1997; Ponraj et al. 1999; Ren et al. 2005).
- There is a correlation between iron status and atherosclerosis (Ahluwalia et al. 2010; Chau 2000; Day et al. 2003; de Valk and Marx 1999; DePalma et al. 2010; Gozzelino et al. 2010; Howes et al. 2000; Kallianpur 2005; Kartikasari et al. 2009; Kiechl et al. 1997; Lapenna et al. 2007; Marx et al. 2008; McRae et al. 2009; Minqin et al. 2003; Ramakrishna et al. 2003; Ren et al. 2003; Salonen et al. 1992; Shah and Alam 2003; Sullivan 2007, 2009; You et al. 2003; You and Wang 2005; Zacharski et al. 2000; Zacharski and Gerhard 2003) (and zinc is protective only against subsequent calcification, Stadler et al. 2008)
- Exogenous ferric iron is deleterious to endothelial function (Rooyackers et al. 2002)
- Iron chelation improves endothelial function (Altamura and Muckenthaler 2009; Duffy et al. 2001; Halliwell 2009; Ishizaka et al. 2005; Saito et al. 2005; Thomas et al. 2006).
- Iron levels in plaques correlate with the amount of oxidised proteins therein (Stanley et al. 2006); indeed in one study (Stadler et al. 2004), the EPR-detectable iron (essentially Fe(III)) in atherosclerotic tissue was *seventeen times greater* than that in the equivalent healthy tissue. I redraw these data in Fig. 7.

It seems clear that the application of similar studies to other diseases in which plaque formation occurs will be of benefit. Indeed, the location of iron in plaques has also

been observed, for instance, for Alzheimer's (e.g. Collingwood et al. 2008; Connor et al. 1992; El Tannir El Tayara et al. 2006; Ghribi et al. 2006; Haacke et al. 2005; Lovell et al. 1998; Morgan et al. 2004; Quintana 2007; Quintana et al. 2006; Sayre et al. 2000b; Schrag et al. 2009), ALS (Leveugle et al. 1997), MS (Haacke et al. 2009) and Parkinson's (see above).

Sepsis, septic shock and the systemic inflammatory response syndrome

Although prions lack nucleic acids, they are seen as infectious agents. Bacteria are more conventional infective agents, and bacterial sepsis provides a very interesting example in which iron release is the key to cell death and the multiple organ failure that normally precedes organismal death. This said, the processes leading to septic shock and ultimately death (Fig. 8) are independent of the presence of viable/culturable bacteria (as defined in Kaprelyants et al. 1993; Kell et al. 1998), i.e. in the apparent absence of infection (Goris 1990; Suntharalingam et al. 2006). This is because it is the *products* of bacteria, especially lipopolysaccharide (LPS), that cause the cells themselves to initiate the progressive processes of self-destruction. The sequelae are typically referred to as a systemic inflammatory response syndrome (SIRS) (e.g. Baue et al. 1998; Beal and Cerra 1994; Bolton 1996; Bone 1996; Davies and Hagen 1997; Gutteridge and Mitchell 1999; Hotchkiss and Karl 2003; Johnson and Mayers 2001; Keel and Trentz 2005; Leithead et al. 2009; Lolis and Bucala 2003; Martin et al. 2003; Nguyen et al. 2006; Protti and Singer 2006; Rittirsch et al. 2008; Ulloa et al. 2009).




Fig. 8 The major events accompanying sepsis and the Systemic Inflammatory Response Syndrome. Note in particular the positive feedback by which the release of pro-inflammatory cytokines and ROSs leads to the release of poorly liganded iron (e.g. from ferritin and haem) that causes the release of further inflammatory cytokines and ROSs. In principle, iron chelators could interfere with this vicious cycle

It is by now well established that—while a ‘cytokine storm’ (Huang et al. 2005; Sriskandan and Altmann 2008; Stacey et al. 2009; Suntharalingam et al. 2006; Wang and Ma 2008; Woo et al. 2010) is an important part of the aetiology of sepsis/septic shock—these phenomena are associated with the hyperproduction of ROSs (Abdelrahman et al. 2005; Andrades et al. 2005; Bulger and Maier 2001; Cadenas and Cadenas 2002; Closa and Folch-Puy 2004; Crimi et al. 2006a, b; Goode and Webster 1993; Gutteridge and Mitchell 1999; Horn 1998; Javadi et al. 2005; Lagan et al. 2008; Lemineur et al. 2006; Mishra 2007; Protti and Singer 2006; Quinlan et al. 2001; Redl et al. 1993; Victor et al. 2004, 2005; Vlessis et al. 1995), and (local and circulating) free iron is raised in sepsis and related conditions (Duvigneau et al. 2008; Galley et al. 1996, 1997; Galley and Webster 1996; Ghio et al. 2003; Lagan et al. 2008), as is the iron siderophore-binding protein lipocalin-2 (Hattori et al. 2009).

The natural iron-chelating antioxidant melatonin has been found to be particularly effective in preventing septic shock (Carrillo-Vico et al. 2005; Escames et al. 2006; Reynolds et al. 2003), and a variety of suitable antioxidants have shown promise (Bekyarova et al. 2009; Cuzzocrea et al. 2001, 2004; Di Paola et al. 2006; Galley et al. 1997; Mishra 2007; Nathens et al. 2002; Sebai et al. 2009; Streck et al. 2008; Zapelini et al. 2008). Most importantly, antioxidants are especially effective when given in combination with iron chelators (Barichello et al. 2007; Ritter et al. 2004, 2006) (see also Bulucu et al. 2009; Cassol et al. 2009; Petronilho et al. 2009; Pinho et al. 2005; Teixeira et al. 2008; Valvassori et al. 2008). Erythropoietin is also protective (Abdelrahman et al. 2004; Cuzzocrea et al. 2006), as is the iron-binder lactoferrin (Tian et al. 2010) and as are anti-inflammatory statins (Durant et al. 2004; Gao et al. 2008; Kopterides and Falagas 2009). All of these observations make it very clear indeed that iron is intimately involved in the death following SIRS and sepsis, and—while not even mentioned in the guidelines (Dellinger et al. 2008)—that its effective chelation should be considered for research leading towards its inclusion as an important part of any therapy.

Bactericidal antibiotics

The classical test for if a bacterium is (or was) ‘alive’ is to analyze its ability to multiply, i.e. viability = culturability (Kell et al. 1998; Postgate 1967, 1976). Accordingly, bacterial antibiotics have long been classified into those that are bacteriostatic, i.e. stop individual cells growing but allow culturability to recover if the antibiotic is removed, and those that are bactericidal, i.e. that actually kill cells and thereby decrease the numbers of those that *are*

culturable (capable of multiplication) even if the antibiotic is removed. Since this clearly implies that causing growth to cease is not of itself bactericidal, and anyway different antibiotics have a variety of different modes of action, it has always been a slight mystery as to what it is that makes a particular antibiotic bactericidal rather than merely bacteriostatic.

A very important paper (Kohanski et al. 2007) provides much of the answer. As one may suppose from the subject matter of this review, iron is again involved, specifically via the Fenton reaction in which poorly liganded ferrous ions reduce the comparatively harmless hydrogen peroxide to the deadly hydroxyl radical.

Based on a previous paper from the same group (Dwyer et al. 2007), Kohanski et al. (2007) (see also Davies et al. 2009; Dwyer et al. 2007, 2009; Imlay 2008; Kohanski et al. 2008; Liu et al. 2009a; Wang and Zhao, 2009) reported that the three major classes of bactericidal antibiotics (exemplified by norfloxacin, ampicillin and kanamycin), regardless of drug–target interaction, utilize internal iron liberated from iron–sulphur clusters to promote Fenton-mediated hydroxyl radical formation in bacteria, that such hydroxyl radical generation contributes to the killing efficiency of these lethal drugs (probably via their known ability to damage DNA), and that in contrast antibiotics that are merely bacteriostatic do not cause the production of such hydroxyl radicals. The availability of multiple genetically marked strains allowed several predictions (such as the resistance to be expected in mutants that could not make iron-sulphur centres so effectively) to be tested successfully. Nitric oxide is correspondingly protective (Gusarov et al. 2009). This suggests, much as above, that there is a common mechanism of iron/hydroxyl-mediated cellular death underlying the action of all classes of bactericidal antibiotics and opens up many new possibilities for combination therapies based on promoting this kind of activity selectively (see Fig. 9). This said, how synergistic a combination is does depend on the exact speciation of the liganded iron; surprisingly, the iron chelator deferasirox promoted the antibacterial activity of ciprofloxacin against *V. vulnificus* (Neupane and Kim 2010), as did deferiprone with various azoles against *Aspergillus fumigatus*, while deferoxamine was antagonistic to azoles (Zarembek et al. 2009). As with any complex system, the interactions can be highly dosedependent (Hegreness et al. 2008; Kohanski et al. 2010; Yeh et al. 2009).

Viruses

A typical viral infection involves the binding of virus to the cell surface followed by internalisation and other effects. Some viruses are especially virulent, and as well as




Fig. 9 A diagram, based very loosely on the narrative in (Kohanski et al. 2007), illustrating how the autocatalytic activity of the reactions of superoxide, peroxide and the hydroxyl radical involving poorly liganded iron can exert a positive feedback leading to the death of cells

inflammation cause the death of cells that they have invaded. It is obviously of interest to establish whether iron is involved in this kind of cell death (and if so whether iron chelation might contribute to a suitable therapy during the acute phase of an infection).

The above suggestions are certainly true of infections caused by **influenza A** (Visseren et al. 2002) (see also Akaike et al. 1996; Beck et al. 2004), **herpes simplex virus** (Gennero et al. 2010; Lamey and Biagioni 1995; Romeo et al. 2001), **hepatitis C** (Alavian and Tabatabaei 2009; Alla and Bonkovsky 2005; Ameli et al. 2008; Bassett 2007; Batts 2007; Bonkovsky 2002; Bonkovsky et al. 2006; Desai et al. 2008; Di Marco et al. 2008; Drakesmith and Prentice 2008; Fargion et al. 2002; Ferrara et al. 2009; Franchini et al. 2008; Fujita et al. 2007a, b; Fujita and Takei 2007; Girelli et al. 2009; Gürkan et al. 2005; Guyader et al. 2007; Hayashi and Yano 2002; Heathcote 2004; Kaito 2007; Kaito et al. 2006; Kato et al. 2001, 2007; Ko et al. 2007; Mahmoud et al. 2008; Mueller et al. 2006; Nahon et al. 2008; Nishina et al. 2008; Otagawa et al. 2008; Price and Kowdley 2009; Rigamonti et al. 2005; Sebastiani and Walker 2007; Sikorska et al. 2010; Sugimoto et al. 2009; Tanaka et al. 2007; Tanaka and Kiyosawa 2004; Trinder et al. 2008; Tung et al. 2003; Valenti et al. 2008; Venturini et al. 2010; Won et al. 2009) and **HIV** (Boelaert et al. 1996; Butensky James et al. 2009; de Monyé et al. 1999; Debebe et al. 2007; Drakesmith and Prentice 2008; Georgiou et al. 2000; Gordeuk et al. 2001, 2006; Kagu et al. 2007; Ketonen et al. 1996; McDermid et al. 2007; McDermid and Prentice 2006; McDermid et al.

2009; Meyer 2006; Northrop-Clewes 2008; Rawat et al. 2008; Savarino et al. 1999; Schreck et al. 1992; Szajerka and Jablecki 2007; Traoré and Meyer 2004, 2007; van Asbeck et al. 2001; Weinberg 2006; Ziegler et al. 2001).

Microbial, plant and animal toxins

Before considering chemical toxicants, it is worth noting that the natural world contains a very great many molecules that are toxic to cells and hence organisms. Although the mechanisms of cell death are rather rarely pursued, there is evidence for the involvement of iron in the sequelae of exposure to toxic doses of gentamicin (Petronilho et al. 2009), ochratoxin (Hasinoff et al. 1990; Omar et al. 1990; Størmer and Hoiby 1996) and ricin (Hassoun and Wang 1999; Kumar et al. 2003; Muldoon et al. 1996), as well as to certain snake venoms (Gutierrez et al. 2010).

Chemical toxicants: aniline, asbestos, dioxins, hydrazines, methyl mercury and paraquat

The most dangerous kinds of man-made toxic chemical agents, those that kill cells or organisms, exhibit a rich diversity of chemical structures and can bind to multiple cellular targets, to the extent that a unitary mode of action is not considered likely. (This said, the same was thought to be true for the mode of action of general anaesthetics or narcotics, and they turn out to have a comparatively small number of rather specific protein targets (Dobson et al. 2009; Dobson and Kell 2008; Franks 2006, 2008; Grasshoff et al. 2006)). However, this does not mean that multiple initial events do not at some point converge to a comparatively small number of crucial events of cell death (whether through necrosis or apoptosis, Vanden Berghe et al. 2009), equivalent to the ‘bow-tie’ model of cellular networks (see e.g. Csete and Doyle 2004; Ma’ayan 2009; Ma et al. 2004, 2007; Ma and Zeng 2003; Oda and Kitano 2006; Rodríguez-Caso et al. 2009; Wang and Chen 2010; Zhao et al. 2006, 2010) and (Fig. 10), and my purpose here is to highlight the evidence that iron-based mechanisms, largely based on Fenton chemistry, are involved in a number of these cases. A recent special issue on toxicants and neurodegeneration is also of interest here (Gupta and Milatovic 2009).

Acetaminophen

Acetaminophen (paracetamol in the United Kingdom) (ChEBI 46195) is a well-known analgesic that is nevertheless highly toxic at concentrations in excess of the therapeutic dose, as it is converted to the electrophilic N-acetyl-p-benzoquinone imine (NAPQI). If glutathione is in

‘Bow-tie’ models of cellular networks




Fig. 10 ‘Bow-tie’ models of cellular networks. There are many examples of cellular networks in which a large variety of possible initial events leads to complex sequelae, but these are mediated via a comparatively small number of ‘intermediate’ reactions. While in the present case it is suggested that these in part involve complex positive feedbacks, the general ‘bow-tie’ idea does allow one to recognise that despite the many different possible inputs, a broadly unitary kind of mechanism of action—here involving iron dysregulation—can reasonably be invoked to explain the multiple causes that can lead to cell death and destruction

insufficient supply, the NAPQI cannot be conjugated harmlessly by glutathione and so it reacts covalently with proteins, especially mitochondrial proteins (Hinson et al. 2004; Jaeschke and Bajt 2006; Kon et al. 2004; Masubuchi et al. 2005; Reid et al. 2005), resulting in apoptotic cell death and organismal death via liver damage (e.g. Barker et al. 1977; Beckett et al. 1989; James et al. 2003; Martin-Murphy et al. 2010; Olaleye and Rocha 2008). One of the effects of mitochondrial damage is iron release, and it has long been known that iron chelators help prevent this toxicity (Sakaida et al. 1995; Schnellmann et al. 1999). Appropriate hydroxyl radical scavengers are also partly protective (Chandrasekaran et al. 2009), while lactoferrin is also protective, although not apparently only via its iron-binding capacity (Yin et al. 2010).

Aniline

As is well known, the adulteration with aniline of cooking oil in Spain showed that aniline so administered can be exceedingly toxic to humans (the ‘toxic oil syndrome’, Hill et al. 1987; Vázquez Roncero et al. 1983), especially in the lungs and spleen. It is now clear (Bomhard and Herbold 2005; Khan et al. 1995, 1999; Ma et al. 2008; Mellert et al. 2004; Wang et al. 2005, 2008, 2010) that iron is heavily involved in this via Fenton chemistry.

Asbestos

Many mutagens are carcinogens, typically by intercalating into DNA, so it was not a priori obvious why asbestos fibres—enormously larger than the diameter of DNA—should

be so toxic. The mystery was solved when it was realised that asbestos does in fact contain and bind iron, that the sharp surface of the iron-loaded fibres can catalyse hydroxyl radical formation and that it is *this* that provides the toxic mechanism (Aljandali et al. 2001; Burrows and Muller 1998; Chao et al. 1996; Chen et al. 1996; Dai and Churg 2001; Fubini and Arean 1999; Gilmour et al. 1997; Governa et al. 1999; Gulumian and van Wyk 1987; Hardy and Aust 1995a, b; Hippeli and Elstner 1999; Jaurand 1997; Kamp et al. 1992; Kamp and Weitzman 1997, 1999; Kelleher et al. 2000; Levrresse et al. 2000; Lund and Aust 1991, 1992; Mossman and Churg 1998; Mossman et al. 1987; Park and Aust 1998; Quinlan et al. 1994; Shatos et al. 1987; Shen et al. 1995; Weinberg 1989; Weitzman and Graceffa 1984; Xu et al. 1999; Azad et al. 2008; Bhattacharya et al. 2005; Daghino et al. 2006; Fantauzzi et al. 2010; Fattman et al. 2006; Gazzano et al. 2007; Ghio et al. 2004, 2009; Heintz et al. 2010; Kamp 2009; Knaapen et al. 2004; Liu et al. 2010; Nymark et al. 2008; Panduri et al. 2006; Poser et al. 2004; Quinlan et al. 2002; Shannahan et al.; Shukla et al. 2003a, b; Sørensen et al. 2007; Toyokuni 2009a, b; Upadhyay and Kamp 2003; Valavanidis et al. 2009).

Dioxins

Dioxins are formed typically via the thermal combustion of halogen-containing compounds, and a chief product, 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD, often referred to simply as ‘dioxin’ (ChEBI 28119), notoriously produced during the 1976 Seveso chemical plant explosion and as an impurity in Agent Orange (a roughly 1:1 mixture of the plant hormone (auxin—ChEBI 22676) analogues) 2,4-dichlorophenoxyacetic acid (2,4-D; ChEBI 28854) and 2,4,5-trichlorophenoxyacetic acid (2,4,5-T; ChEBI 27903), is both a cause of chloracne and is a hepato-carcinogen. It was found that mice low in iron are resistant to this hepatotoxicity however (Poland and Knutson 1982; Sweeny et al. 1979), strongly implicating iron. In addition, iron enhances the toxicity of dioxin (Al-Turk et al. 1988; Smith et al. 1998) while iron chelators reduce it (Al-Bayati and Stohs 1987; Wahba et al. 1990). While the exact details have still not been worked out, there is little doubt that iron is important in the toxicity of a wide variety of polyhalogenated aromatics (Smith and Chernova 2009; Smith and Elder 2010), including hexachlorobenzene (ChEBI 5692) (Franklin et al. 2005).

Hydrazines

Phenylhydrazine (ChEBI 27924) (like other hydrazines) is a reactive substance that is commonly used in toxicity studies. It is known (Agrawal et al. 2001; Atanasova et al.

2004; Ferrali et al. 1997a, b; Fraenkel et al. 2009; Karbownik et al. 2000; Kim et al. 2009; Latunde-Dada et al. 2004; Rokushima et al. 2007a, b; Sharma and Haldar 2009; Yamamoto and Kawanishi 1992) to induce haemolytic anaemia and iron release and thence cytotoxicity. Melatonin can be protective (Sharma and Haldar 2009).

Methyl mercury

Originally highlighted via ‘Minamata disease’ (a prominent symptom of which, as with many neurological disease, is ataxia), methylmercury (ChEBI 30785) is a well-known neurotoxicant (Clarkson et al. 2003). Like most if not all other xenobiotics (Dobson et al. 2009a; Dobson and Kell 2008; Dobson et al. 2009b; Kell and Dobson 2009), it requires one or more carriers for cellular uptake—in this case, when complexed to L-cysteine, the ‘large amino acid’ transporter (Kanai and Endou 2003; Simmons-Willis et al. 2002; Yin et al. 2008). There is certainly evidence that exposure to this agent causes oxidative stress (Ali et al. 1992; Aschner et al. 2007; Atchison and Hare 1994; Franco et al. 2009; Johansson et al. 2007; Sanfeliu et al. 2003; Sarafian et al. 1994; Yee and Choi 1994, 1996; Yin et al. 2007) and raises cellular iron levels (Ilbäck et al. 2000) and that anti-oxidants (Daré et al. 2000; Kaur et al. 2006; Sarafian and Verity 1991; Shanker and Aschner 2003; Usuki et al. 2001; Watanabe et al. 2009; Yamashita et al. 2004; Zurich and Monnet-Tschudi 2009), metallothioneins (West et al. 2008) and/or iron chelators (Castoldi et al. 2001; LeBel et al. 1992; Sarafian and Verity 1991; Stohs and Bagchi 1995) can be protective. Taken together, this does imply strongly a role for iron-mediated cytotoxicity of the type highlighted herein in the pathogenesis of methylmercury poisoning.

Paraquat

Paraquat (N,N'-dimethyl-4,4'-bipyridinium dichloride; ChEBI 34905) is a well-known herbicide and toxicant, whose mode of action relies on the generation of superoxide radicals. As such, its toxicity may be expected to be exacerbated by iron, and this is the case (Andersen 2003; Ayaki et al. 2005; Fernandez et al. 2000; Peng et al. 2007, 2009). In addition, again as expected from the above (and consistent with its structural similarity to the Parkinson-inducing agent MPTP-1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, ChEBI 17963), it can induce Parkinson-like symptoms (Kaur et al. 2009; Miller et al. 2009; Ossowska et al. 2006; Peng et al. 2007, 2009). The protection by molecules—such as ferritin (Kaur et al. 2009) and polyphenols (Jimenez-Del-Rio et al. 2010)—capable of chelating iron harmlessly is thus of considerable interest.

Other toxicants in which iron release has been implicated as part of the mechanisms of their toxicity

The list of known and potential toxicants is limitless, and part of the role of this review is to alert domain experts to the likelihood that iron, and dysregulation in iron metabolism, may mediate the toxicity of such molecules (and thus that its chelation may be important, under both acute and chronic conditions, in assisting resolution of the toxicity). To this end, I note the role of iron-induced oxidative stress in the toxicity of anthracyclines (including adriamycin/doxorubicin [ChEBI 28748](#)) (Ceyssens et al. 2004; Kaiserová et al. 2007; Kotamraju et al. 2002, 2004; Luft 2006; Othman et al. 2008; Šimůnek et al. 2009; Thougard et al. 2010; Xu et al. 2008), arsenic (De Vizcaya-Ruiz et al. 2009; Fujihara et al. 2009), beer brewed in cast iron pots (Kew and Asare 2007), bleomycin ([ChEBI 3139](#)) (Wu et al. 2004; Yang et al. 2010), C60 fullerenes (Usenko et al. 2008), carbon nanotubes (Murray et al. 2009) (and see Guo et al. 2007; Jurkschat et al. 2007), carbon tetrachloride (Mansour 2000; Ogeturk et al. 2004), organophosphates (Goel et al. 2006), sulphur mustard agents (Korkmaz et al. 2008; Korkmaz et al. 2006; Naghii 2002) and talc nanoparticles (Akhtar et al. 2010). Welding fumes are toxic and contain both iron and manganese (Flynn and Susi 2010; Sriram et al. 2010), and the toxicity of nano-iron itself has also been studied (Li et al. 2009).

Apoptosis or necrosis?

As is well known, cells can die by both necrosis and by a more regulated form, apoptosis. While this—nor ‘normal’ ageing (summarised e.g. by Bishop et al. 2010b; Finkel 2005; Franco et al. 2007; Gems 2009; Kenyon 2010; Kirkwood 2008; Muller et al. 2007; Vaupel 2010), in which various degradative processes including iron dysregulation can be implicated (e.g. Brewer 2010; Cass et al. 2007; Cook and Yu 1998; Facchini 2002; Galaris et al. 2008; Kell 2009a; Kujoth et al. 2005; Kurz et al. 2008b; Massie et al. 1983; Napoli et al. 2006; Pfefferbaum et al. 2009; Terman et al. 2010; Xu et al. 2010)—is not the focus of this review, it is reasonable to enquire as to what kind of evidence exists for the mechanism(s) of cell death contingent upon iron dysregulation. As it turns out, while there is a little evidence that iron overload can cause necrosis (Babatasi et al. 1996; Traoré and Meyer 2007; Vanden Berghe et al. 2009; Vanlangenakker et al. 2008), there is abundant evidence for the ability of iron to cause apoptosis (Kotamraju et al. 2004a, b; Kurz et al. 2007, 2008b; Vanden Berghe et al. 2009), and—depending presumably on the exact liganding of the iron (Kell 2009a)—this can be effected when iron is ostensibly depleted (e.g. Bergman et al. 2005; Bidle and Bender 2008; Brard et al. 2006; Buss et al. 2003; Fukuchi

et al. 1994; Greene et al. 2002; Haq et al. 1995; Hileti et al. 1995; Jiang et al. 2002; Jin et al. 2007; Koc et al. 2006; Kovář et al. 1997, 2001; Leardi et al. 1998; Lee et al. 2006b, c; Pan et al. 2004; Simonart et al. 2000; Sun et al. 2009; Truksa et al. 2003; Zhang et al. 2003; Zhao et al. 2004) or in excess (e.g. Aljandali et al. 2001; Cozzi et al. 2003, 2010; Devireddy et al. 2005; Franke et al. 2010; Jacob et al. 1997, 2000; Javadi et al. 2004; Jiménez Del Río and Vélez-Pardo 2004; Kamp et al. 2002; Kawabata et al. 1997; Kermer et al. 2004; Kooncumchoo et al. 2006; Kruman et al. 1997; Kummer et al. 2008; Kurz et al. 2007, 2008b; Lakshmi Devi and Anuradha 2009; Lee et al. 2006d; Levenson 2005; Lu et al. 2008; Ma et al. 2009; Messer et al. 2009; Murakami et al. 2006; Natoli et al. 2009; Persson 2005; Polla et al. 2003; Schlawe et al. 2004; Shin and Kim 2009; Shukla et al. 2003; Tenopoulou et al. 2005; Upadhyay and Kamp 2003; Vélez-Pardo et al. 1997; Zhao et al. 1997).

Inflammation and NF- κ B

As reviewed in the previous survey (Kell 2009a) and elsewhere (e.g. Shoelson and Goldfine 2009; Shoelson et al. 2007), inflammation lies at the heart of many of these processes, and pathways involving NF- κ B (see e.g. <http://www.nf-kb.org/>) are especially core here (e.g. Ali and Mann 2004; Circu and Aw 2010; Granic et al. 2009; Karin 2008; Karin and Greten 2005; Li et al. 2005; Maeda and Omata 2008; Simmonds and Foxwell 2008; Zhang and Rigas 2006). Of particular relevance in ROS/iron metabolism are the facts that **NF- κ B is redox-sensitive** (e.g. Aggarwal and Sung 2009; Baeuerle and Henkel 1994; Chandrasekaran and Taylor 2008; Cho et al. 1998; Dröge 2002; Fulda et al. 2010; Galaris et al. 2008; Gloire et al. 2006; Gloire and Piette 2009; Ha et al. 2010; Haddad et al. 2000; Hayashi et al. 1993; Janssen-Heininger et al. 2008; Lagan et al. 2008; Lukosz et al. 2010; Nishi et al. 2002; Phillips et al. 2010; Schreck et al. 1991, 1992; Schreck and Baeuerle 1994; Sen and Packer 1996; She et al. 2002; Surh et al. 2005; Surh and Na 2008; Thannickal and Fanburg 2000; Trachootham et al. 2008, 2009; Valko et al. 2007; Weinberg and Chandel 2009; Wondrak 2009; Xiong et al. 2003) and that there are links between NF- κ B and NGAL/ICN2 (Bu et al. 2006; Cowland et al. 2003, 2006, 2008; Karlsen et al. 2010) and/or hepcidin (Sow et al. 2009; Tacchini et al. 2008). Specificity is likely to come, in part, via the frequency encoding of NF- κ B activity (Ashall et al. 2009; Nelson et al. 2004).

Catalytic behaviour of polypeptides and proteins

I have mentioned several times the idea that different proteins can bind unliganded iron to catalyse the

production of hydroxyl radical, and it is certainly the case that the degree of liganding can affect these kinds of activities (Kell 2009a). Here it is appropriate to highlight the abilities of polypeptides and even poly(aminoacids)—whether binding metals (or not)—to exhibit catalytic activities, e.g. poly(leucine) (Carrea et al. 2005; Porter et al. 1999), and how adding metals to polypeptides to make metalloenzymes can have huge effects on their catalytic activities (Pierron et al. 2008; Pordea and Ward 2008; Turner 2009). The recent development of orthogonal ribosomes capable of incorporating novel amino acids into polypeptides (Neumann et al. 2010) opens up considerable possibilities here.

Dietary and/or pharmacological treatment of iron-mediated toxicity

Pharmaceutical


Developing novel pharmaceutical agents is a costly enterprise, and many candidate drugs fall by the wayside during the development process (see e.g. Dobson et al. 2009b; Kola and Landis 2004; Leeson and Springthorpe 2007). As most recently reviewed (Agarwal 2010; Berdoukas et al. 2009; Bolognin et al. 2009a; Cappellini and Pattoneri 2009; Cappellini et al. 2010; Clifton et al. 2009; Hider et al. 2008; Kell 2009a; Kontoghiorghes et al. 2010; Kwiatkowski

2010; Mladěnka et al. 2009; Pennell et al. 2010; Perez and Franz 2010; Richardson et al. 2009b and references therein), a number of (pharmacological drug) approved iron chelators do exist and thus clearly have the potential to prove beneficial in treating diseases and the sequelae of toxin exposure whose aetiology involves poorly liganded iron via the Fenton reaction. The benefits of administration of desferrioxamine in Alzheimer's, for instance, have been known for nearly 20 years (Crapper McLachlan et al. 1991), benefits since observed with other iron chelators (e.g. Amit et al. 2008; Bolognin et al. 2009a; Bush 2008; Liu et al. 2009b, 2010b; Perez and Franz 2010; Smith et al. 2010; Zhang et al. 2009). The implications for this and other neurodegenerative diseases—and potentially for the more acute and even chronic effects of other toxins—are obvious. Those approved clinically include Deferoxamine (Desferal; ChEBI 31460), Deferiprone (Ferriprox; PubChem CID 2972) and Deferasirox (Exjade; ChEBI 49005) (Fig. 11). Note that—unlike the other two—Deferoxamine is not orally active and must be injected, and even then may fail to cross cellular membranes to exert its action appropriately (Warkentin et al. 2010).

Dietary or nutritional/nutraceutical

In addition, many organisms make so-called 'secondary' or *natural products*, whose functions are often unclear but

Fig. 11 The three main iron chelators approved for clinical use



that clearly have evolutionary benefits for the host (Hadacek et al. 2010; Kell et al. 1995). As I also reviewed (Kell 2009a), many **iron-chelating** natural products exist in (or may be added to, Pócsi et al. 2008) foodstuffs for which no full pharmaceutical regulatory controls are required and which are classed as nutritional substances. Many are **polyphenols** (ChEBI 26195) (Amic et al. 2007; Arts and Hollman 2005; Auclair et al. 2009; Cemeli et al. 2009; Cheynier 2005; Geronikaki and Gavalas 2006; Halliwell 2007; Jovanovic and Simic 2000; Kandaswami and Middleton 1994; Kiokias et al. 2008; Korkina and Afanas'ev 1997; Linseisen and Rohrmann 2008; Loke et al. 2009; Manach et al. 2004, 2005a, b; Pérez-Jiménez et al. 2010; Perron and Brumaghim 2009; Perron et al. 2008; Petti and Scully 2009; Pietta 2000; Rice-Evans and Packer 2003; Scalbert et al. 2005; Scalbert and Williamson 2000; Seifried et al. 2007; Slemmer et al. 2008; Spencer et al. 2008; Yadav and Bhatnagar 2007a, b, 2010), including the anthocyanins that proved chemoprotective (Butelli et al. 2008) in a mouse p53 cancer model. Since the previous review (Kell 2009a), I would draw particular attention to a comprehensive overview of the subject (Perron and Brumaghim 2009), as well as new reviews and papers on the chief polyphenol in green tea, (–)**epigallocatechin-3-gallate** (ChEBI 4806) (Adhami et al. 2009; Butt and Sultan 2009; Hsieh et al. 2009; Jimenez-Del-Rio et al. 2010; Johnson et al. 2010; Pandey and Gupta 2009; Paterniti et al. 2009; Ramesh et al. 2009; Reznichenko et al. 2010; Suh et al. 2009; Weinreb et al. 2009a), on **curcumin** (ChEBI 3962) (a constituent of turmeric) (Aggarwal and Harikumar 2009; Aggarwal and Sung 2009; Anand et al. 2010; Basile et al. 2009; Epstein et al. 2009; Hegde et al. 2009; Iqbal et al. 2009; Jiao et al. 2009; Jurenka 2009; Kalpravidh et al. 2010; Majumdar et al. 2009; Messner et al. 2009; Patil et al. 2009; Ravindran et al. 2009; Rivera-Espinoza and Muriel 2009; Thephinlap et al. 2009; Thomas et al. 2009; Tuntipipipat et al. 2009; Wang et al. 2009a), on **eugenol** (ChEBI 4917) (Nagababu et al. 2010) (and cf. Allen and Cornforth 2009), on nepetoidin caffeic acid esters (Maioli et al. 2010), on **quercetin** (ChEBI 16243) (Chobot 2010; Kitagawa et al. 2009; Krukoski et al. 2009; Lorrain et al. 2010; Park et al. 2010; Pavlica and Gebhardt 2010; Smirnova et al. 2009; Terao 2009; Vlachodimitropoulou et al. 2010) and on **melatonin** (CHEBI 16796) (Borah and Mohanakumar 2009; Korkmaz et al. 2008, 2009; Reiter et al. 2009; Sener et al. 2009; Sharma and Haldar 2009; Signorini et al. 2009). Many of the protective effects observed are likely due to the iron chelating, as well as directly antioxidative (redox) properties of these molecules (e.g. Ferrali et al. 1997a; Hague et al. 2006; Halliwell 2009; Halliwell et al. 2005; Hider et al. 2001; Končić et al. 2010; Lopes et al. 1999; Mandel et al. 2004; Mandel and Youdim 2004; Morel et al. 1994; Nijveldt et al. 2001;

Rice-Evans et al. 1997; Rice-Evans and Packer 2003; Weinreb et al. 2007, 2009a; Yoshino and Murakami 1998). Other molecules that are not themselves directly antioxidant, but have iron-binding characteristics, may also be of benefit. This is certainly true of **phytic acid** (phytate; myo-inositol-1,2,3,4,5,6-hexakis phosphate; CHEBI 17401) (Aljandali et al. 2001; Allen and Cornforth 2009; Brune et al. 1989; Davidsson et al. 1994; Doria et al. 2009; Glahn et al. 2002; Graf and Eaton 1990, 1993; Graf et al. 1987; Grases et al. 2001; Hallberg et al. 1989; Han et al. 1994; Hanson et al. 2006; Hawkins et al. 1993; Kamp et al. 1995a, b, 2002; Kim et al. 2008; Kumar et al. 2010; Minihane and Rimbach 2002; Miyamoto et al. 2000; Panduri et al. 2006; Poser et al. 2004; Rao et al. 1991; Rimbach et al. 2008; Sandberg et al. 1999; Schlemmer et al. 2009; Shamsuddin 1995; Vucenik and Shamsuddin 2006; Xu et al. 2008).

Phlebotomy

In addition, as follows implicitly from Sullivan's 'iron hypothesis' (Sullivan 1981, 2001, 2003, 2004) (and see Kell 2009a), one way to decrease the amount of iron in the body is to remove it by blood-letting or phlebotomy. While phlebotomy is a very traditional nostrum, often assumed or considered to have rather dubious or at best modest scientific support, there is in fact increasing literature implying its benefits in a variety of conditions (e.g. Aigner et al. 2008; Beutler 2007; Broedbaek et al. 2009; Brudevold et al. 2008; Busca et al. 2010; DePalma et al. 2007, 2010; Dereure et al. 2008; Desai et al. 2008; Dwyer et al. 2009; Equitani et al. 2008; Facchini et al. 2002; Fargion et al. 2002; Fernández-Real et al. 2002; Fujita et al. 2009; Fujita and Takei 2007; Hayashi et al. 2006; Hayashi and Yano 2002; Heathcote 2004; Horwitz and Rosenthal 1999; Hua et al. 2001; Kaito 2007; Kaito et al. 2006; Kato et al. 2001, 2007; Kom et al. 2006; Rajpathak et al. 2009; Sullivan 2009; Sumida et al. 2009; Tanaka et al. 2007, 2009; Toyokuni 2009b; Zacharski et al. 2008). Plausibly such benefits are due to its role in decreasing iron stores.

Discussion: toxicology as systems biology

“When sorrows come, they come not single spies but in battalions.” (Shakespeare, Hamlet.)

It is certainly the case that reasons for the co-occurrence of two or more observables are that one causes the other or they have the same, separate cause. Since I have argued strongly here for the essentiality of iron dysregulation in a variety of syndromes (and have omitted many others, such as non-alcoholic (Baptista-Gonzalez et al. 2008; Kirsch

et al. 2006; Kowdley 2010) or alcoholic liver disease (Breitkopf et al. 2009; Lecube et al. 2009)), and indeed cancers of various kinds (see Kell 2009a), the implication is that iron dysregulation can manifest in multiple different ways and that these might co-occur. There is considerable and increasing evidence for that (see e.g. Adibhatla and Hatcher 2010; Baker et al. 2009; Beeri et al. 2008; Cole et al. 2007; Franco et al. 2009; Janson et al. 2004; Kell 2009a; Li and Hölscher 2007; Mascitelli et al. 2009; Neumann et al. 2008; Pandolfo 2009; Roriz-Filho et al. 2009; Schnaider Beeri et al. 2004; Starley et al. 2010; Utzschneider and Kowdley 2010; Verduzco and Nathan 2009; Xu et al. 2009), with the exact outcomes being determined by different levels of expression (see for instance <http://www.proteinatlas.org/>) of different proteins in different tissues.

In a similar vein, it is not news (Attwood et al. 2009; Evans 2008; Hull et al. 2008; Kostoff 2002) that the large floods of literature have led to something of a ‘balkanisation’ of the literature into increasingly specialised segments, and that this makes it hard to seek to bring together such disparate parts of the literature so as to draw out general truths (Dobson and Kell 2008; Kell 2009a), via the methods of inductive reasoning (Kell and Oliver 2004; King et al. 2004, 2009). As previously discussed, it is well established on theoretical grounds alone that the activity of any step in a biochemical network can influence the concentration of any variable (such as—fleetingly—the amount of hydroxyl radicals produced). Thus, while it is better to derive quantitative models (e.g. Coveney and Fowler 2005; Csikasz-Nagy 2009; Kell 2006a, b; Kell and Knowles 2006; Kell and Westerhoff 1986a, b; Milo et al. 2010; Mogilner et al. 2006; Noble 2002, 2006, 2010; Oates et al. 2009; Phillips and Milo 2009; Tomlin and Axelrod 2007), at the very minimum one should be looking qualitatively at those factors that should be increased or decreased in order to improve the organism’s ability to resist oxidative stress (Fig. 12).

Indeed, it is the very act of doing this that leads one to recognise these kinds of complex problems as problems of systems or network biology (Alon 2006; Klipp et al. 2005; Palsson 2006). What do I mean by this?

A systems biology view of toxicology: toxicology as systems biology

In the present case, I have adduced evidence that a range of distinct aetologies, which manifest as a variety of different syndromes or diseases, converge at some point on an autocatalytic series of reactions, involving imperfectly liganded iron, in the production of the very damaging hydroxyl radical, that by causing the destruction of further appropriate cellular structures and proteins liberates further




Fig. 12 The extent of cell damage caused by the unliganded iron-catalysed production of ROSs and RNSs is determined by many factors, some of which promote and some of which act against it. While the existence of hormesis (see text) means that the see-saw illustrated here is an imperfect metaphor, the diagram serves to illustrate the complexity of the problem and the need for a systems biology approach to its solution

‘free’ iron and other reactive oxygen species, thereby effecting positive feedback (also known as ‘vicious cycles’, Lin and Beal 2006; Proctor and Gray 2010; Proctor et al. 2007) that leads to the production of yet further damage. The implication of this is that to be more specific we need suitable network models of the biological systems of interest (Kell 2007), and of how toxicants of any kind can interact with their components.

Of course in the steady state, the amount of a substance is governed both by the processes that create it and those that decompose it (Fig. 13), and individual effects of particular substances can vary qualitatively depending on concentrations, as in hormesis (e.g. Calabrese 2008a, b, c; Calabrese and Baldwin 1997, 2001, 2003; Calabrese and Blain 2005, 2009; Cook and Calabrese 2006; Cox 2009; Gems and Partridge 2008; Gómez-Pinilla 2008; Hadacek et al. 2010; Hayes 2009; Hoffmann 2009; Mattson 2008a, b; Radak et al. 2008; Rattan 2004, 2008; Rattan and Demirovic 2009;




Fig. 13 The steady-state concentrations of a molecule depend on the rates of production and removal of the molecule in question. Both lowering creation and increasing removal represents a particularly effective strategy relative to doing just one of these alone. Thus if C is the hydroxyl radical we can lower its concentration by decreasing A and/or B and by increasing the rates of the reaction to D and E. The latter (decrease of concentration of C) may sometimes better be effected by increasing the activities of enzymes yet further downstream

Ristow and Zarse 2010; Sørensen et al. 2009), so any analysis should be a systems analysis. The simple fact that, for instance, iron chelators improve prospects in Alzheimer's implies that there is a background level of *natural* repair (otherwise chelators would only slow the neurodegeneration and not reverse it) and that is not necessarily apparent from measuring only net effects. Hence, the need for accurate systems biology models.

The very fact that multiple reactions typically lead to the creation (and destruction) of a given metabolite (such as the hydroxyl radical) means that to minimise its concentration it is best to modify many reactions simultaneously. This exceptionally important point from systems biology should underpin any scientific and therapeutic programs.

Systems biology methods

Systems biology methods are typically classified into 'forward' and 'inverse' (Fig. 14). For the purposes of this discussion, I shall concentrate on *metabolic* networks (in contrast to signalling networks they transfer both matter and information, Kell 2004). Metabolic and other networks are 'graphs' in that they consist of entities ('nodes', such as metabolites) that interact with each other via 'edges' (enzymatic reactions). There may of course be other interactions such those that do not involve substrates or products but are mediated via molecules (effectors) that modify the rates of reactions. The distinction between forward and inverse models is based on whether one starts with the parameters or variables of the system. In metabolic networks, the parameters are typically the starting concentrations of substances present and the kinetic constants (K_m , K_i , V_{max} , etc.) of the enzymes. The variables are typically the concentrations of metabolic intermediates, and the metabolic fluxes that change over time until a steady state (if such exists) is attained. It is important to note that only the parameters control the system. In *forward systems biology*, we thus seek to create a model of the

network based on its parameters, and this is done in four phases, the first two of which are qualitative and the second quantitative (Herrgård et al. 2008; Kell and Knowles 2006; Thiele and Palsson 2010).

Forward modelling

The first qualitative stage is one in which are listed all the reactions that are known to occur in the system or organism of interest; in the modern era, and especially for metabolic networks, these reaction lists are often derived in part from genomic annotations (Arakawa et al. 2006; Palsson 2006) with curation based on literature ('bibliomic') data (Duarte et al. 2007; Thiele and Palsson 2010). A second stage, again qualitative, adds known effectors (e.g. those involved in forming feedback loops), while the third and fourth stages—essentially amounting to molecular enzymology—include the known kinetic rate equations and the values of their parameters. Armed with such information, it is then possible to provide stochastic and ordinary differential equation models of the entire metabolic network of interest. In some cases, it is possible to make simplifying assumptions that require minimal knowledge of the kinetic parameters of the system and allow one to generate a time series of the system variables (Smallbone et al. 2007). An attractive feature of metabolism, for the purposes of modelling, is that, in contrast to signalling pathways, metabolism is subject to direct thermodynamic and (in particular) stoichiometric constraints (Palsson 2006). These constraints are extremely important in reducing the search space for the values of the kinetic parameters of the model, and indeed constrain significantly the states that the system *can* adopt (Kell 2004; Palsson 2006).

The modelling community has access to many tools, which are interoperable because they can be exchanged using community standards (Kell 2008) of which the most popular (Klipp et al. 2007) is SBML, the Systems Biology Markup Language (Hucka et al. 2003) (<http://www.sbml.org>). Models are best made, and made available, in SBML. In addition, there is recognition that it is desirable to be clear about the *meaning* (semantics) of the elements in the model (Ananiadou et al. 2006; Attwood et al. 2009; Kell and Mendes 2008), since SBML puts no constraints on the names given to entities, and we have pioneered (Herrgård et al. 2008; Pettifer et al. 2009) the use of semantic annotation of all the entities, whether by reference to persistent databases or via database-independent means (e.g. the use of SMILES (Weininger 1988) and InChI (Coles et al. 2005; Stein et al. 2003) strings for small molecules). This is consistent with the MIRIAM proposals for describing the contents of a model accurately (Le Novère et al. 2005), and we have also used (Herrgård et al. 2008) the Systems Biology Ontology (<http://www.ebi.ac.uk/sbo/>)

Forward and inverse methods in systems biology

Forward: start with the system's parameters (including its topology, kinetic constants and the 'starting' concentrations) and model the time evolution of its variables

Inverse: start with the time series of the system variables and seek to infer the topology and parameters of the system that could produce them

Fig. 14 The relationship between 'forward' and 'inverse' methods of systems biology. The comparison of two 'inverse' systems (one a control and one treated with a toxin) in terms of the estimation of which parameters have changed most allows one to infer the sites or modes of action of that toxin [In a related manner we also discriminate forward and reverse genetics, including chemical genetics (Kell 2006a, b)].

and the Gene Ontology (<http://www.geneontology.org/>) as especially pertinent and useful ontologies. Tools also exist for creating such networks using CellDesigner (<http://www.celldesigner.org/>) and SBMLMerge (Schulz et al. 2006) for merging submodels, including visualising them (via the Systems Biology Graphical Notation (Le Novère et al. 2009) (<http://www.sbgn.org/>) e.g. via Arcadia (Villéger et al. 2010) (<http://www.sourceforge.net/projects/arcadiapathways/>). Armed with such networks, it is possible to analyse their dynamics and other properties using tools/ modelling environments such as COPASI (Hoops et al. 2006) (<http://www.copasi.org/>) and to compare model and experimental data, conveniently stored using SBRML (Dada et al. 2010). Models and attendant metadata are best stored directly in databases, such as that set down in the B-Net database model (see Li et al. 2003; Mehrotra et al. 2010) as used in our yeast model (Herrgård et al. 2008) (<http://www.comp-sys-bio.org/yeastnet>).

Interrogating the model

Sensitivity analysis

Armed with a model, there are a number of important studies that are usefully performed. The first is the study of the contribution of individual steps to the control of the variables of interest. This is the domain of local and global sensitivity analysis (Saltelli et al. 2004, 2008), to which we have made significant contributions (e.g. Lüdtke et al. 2008; Sahle et al. 2008; Yue et al. 2006, 2007, 2008). Sensitivity analysis studies the quantitative contribution of individual steps to the overall control of a process. The ‘local’ version has been popularised in biochemistry as ‘metabolic control analysis’ (e.g. Fell 1996; Heinrich and Schuster 1996; Kell and Westerhoff 1986a), and, for instance COPASI (Hoops et al. 2006) calculates the control or sensitivity coefficients directly from a model. The results of these kinds of analyses are represented by a rank order of the most important reactions or parameters, which provides an important starting point for the combinatorial analysis of these complex networks. Formalisms such as Metabolic Control Analysis also explain why the modulation of individual steps singly rarely has much of an effect on an overall pathway (see e.g. Pérez et al. 2009a, b), reviewed by Gems (Gems and Doonan 2009), for a typical example.

Combinatorial analysis of parameter sensitivity

As indicated above, it is commonly the case in complex biochemical networks that while changes in individual parameters have little effect on the behaviour of biochemical pathways—indeed evolution is thought to have

selected for robustness to individual parameter changes (Bornholdt and Sneppen 2000; Kitano 2007; Lehár et al. 2008; Morohashi et al. 2002; Stelling et al. 2004; Wagner 2005)—combinations of parameter changes can and do have major effects. A straightforward corollary of this is that in order to cause or inhibit the progression of diseases—including inflammatory diseases such as asthma (Biswas and Rahman 2009; Bowler 2004; Bowler and Crapo 2002; Boyer and Liu 2004; Chodaczek et al. 2007; Ciencewicz et al. 2008; Comhair and Erzurum 2010; Galli et al. 2008; Greene 1995; Han et al. 2001; Horváth et al. 1998; Jarvis et al. 2005; Kocyigit et al. 2004; Muñoz et al. 2009; Narula et al. 2007; Nemery 1990; Pae et al. 2008; Riedl and Nel 2008; Shore 2008; Vlašić et al. 2009)—it is necessary to affect multiple steps simultaneously. The problem is to find which ones, and that is a combinatorial problem that scales poorly. It is exactly here where the ability to interrogate a suitable model of the pathway comes to the fore, since computations to determine which combinations of parameters to affect can be done enormously more rapidly than can ‘wet’ experiments. To be explicit, if manipulating (let us say knocking out) just three genes out of 1,000 is necessary to secure a desired effect (as is often the case) and the question is to choose which three, the answer involves recognizing that this is a combinatorial optimization problem involving ca 100 million combinations—a trivial number computationally given a model and an easy number (i.e. 3) to work with post hoc, but an implausible number to interrogate with present experimental methods. See (Park et al. 2007, 2008) for a nice example of the approach.

At all events, the results of running a systems biology model in the forward direction are the time-varying concentrations of variables that result from the network topology and parameters of the system, and these can then be compared with measured variables to see how accurate the model is.

Of course it is typically the case (i) that the variables predicted by the model differ from those that one has measured, and (ii) one has only measurements of the variables and not the parameters anyway. Since it is the parameters that determine the system behaviour, it is then necessary to try and estimate or infer the parameters of the systems from measurements of the variables, the domain of inverse modelling (Liu and Wang 2008; Mendes and Kell 1996) or system identification (Bongard and Lipson 2005; Ljung 1987).

Inverse modelling

Finding the sets of parameters that can reflect a set (or sets) of variables is a much harder problem. If there are just 20 parameters, each of which can take just 10 values, there are

at once 10^{20} combinations, although there may be multiple combinations that give a reasonable solution (the problem or system is then said to be under-determined). At all events, finding these combinations requires methods that can effectively search these combinatorial search spaces to find good solutions (they are not provably optimal solutions as these kinds of problems are NP-hard, Garey and Johnson 1979).

There are many kinds of approaches to attacking these kinds of combinatorial optimisation problems, and I give two examples. The first uses evolutionary methods to modify the search for these parameters, while the second is Bayesian in character.

Evolutionary computing methods (e.g. Bäck et al. 1997; Goldberg 1989, 2002) use ideas from Darwinian evolution to select and modify parameters to achieve a desired objective (or objectives—this is known as multi-objective optimisation, Handl et al. 2007). In an evolutionary algorithm, one selects a population of models encoding a different candidate set of parameters, assesses their performance, rejects those that perform poorly and modifies the parameters of the better-performing models using methods akin to recombination and mutation to create a second generation population and so on. Finally, one finds a candidate set of solutions that perform well and can indeed interrogate these solutions to determine the parameters that differ most frequently in the selected solutions from those in the starting pathway. These kinds of methods are recognised as highly effective (Banga 2008; Banga and Balsa-Canto 2008), and they have proved so for both modelling (Dada and Mendes 2009; Mendes and Kell 1998; Moles et al. 2003; Rodriguez-Fernandez et al. 2006) and other optimisation problems (e.g. Broadhurst et al. 1997; Kell et al. 2001; Knight et al. 2009; O'Hagan et al. 2005, 2007; Platt et al. 2009; Rowe et al. 2010). A related and novel approach that we have developed (Wilkinson et al. 2008) (and see Brown et al. 2010) also evolves estimates of parameters, but in a somewhat different way. Evolutionary computing methods are amenable to ('coarse-grained') parallelisation that allows them to run on many (hundreds or even thousands of) processors simultaneously, and we have used this approach with great effectiveness (Wedge and Kell 2008; Wedge et al. 2009). Overall, the result of this kind of approach is the effective determination of the multiple parameters that are required to explain the system variables.

A conceptually quite different approach, which may of course be expected to lead to the same result (and which in consequence strengthens further any conclusions from e.g. an evolutionary computing approach), is an approach based on Bayesian inference (Jayawardhana et al. 2008; Liebermeister and Klipp 2006; Vyshemirsky and Girolami 2008) and known as Markov Chain Monte Carlo or MCMC. In

this approach (which we do not rehearse in detail as those details have recently been published by us, Jayawardhana et al. 2008), we have knowledge of the variables of interest and sample candidate models that can explain those variables using various estimates of their parameters *from a distribution*. The MCMC algorithm then slowly builds up the likeliest values of the parameters and their possible distribution given the demonstrable uncertainty.

Detecting the modes or sites of action of a toxin

Clearly, if we have measurements of the variables of a system before and after the addition of a toxin, we can estimate the parameters in the two cases and see which have most plausibly changed by applying these inferencing methods to the two cases. This automatically gives the mode(s) or site(s) of action of the intervention, whether of a toxic inhibitor or by any other means. In other words these methods can detect the modes or sites of action of anything that causes a system to change its state. Taking the Bayesian approach (see also e.g. Nidhi et al. 2006; O'Hagan 2006; Vyshemirsky and Girolami 2008; Wilkinson 2007), in the work described above (Jayawardhana et al. 2008) we 'calibrated' our approach using purely simulated data in which we knew exactly which parameters had been changed (and in which direction) and showed that we could detect this accurately. We then applied our approach to an experimental system of metabolic engineering in *Lactococcus lactis* (Hoefnagel et al. 2002) where the investigators had manipulated the activity of a specific enzyme to achieve a desired metabolic effect. Our approach detected both the enzyme whose activity had been modified and suggested the site of another significant pleiotropic effect (these always occur, Featherstone and Broadie 2002). Thus, the power of this general approach is that it is free of specific hypotheses and thus guides us to the hypotheses that can best explain the data. These are classic data mining methods (e.g. Breiman 2001; Kell and King 2000; Kell and Oliver 2004; Wu et al. 2008) that allow us to discover mechanisms of action that are unknown and/or have not previously been contemplated. Note, however, that such methods need careful handling from a statistical point of view (e.g. Altman and Royston 2000; Broadhurst and Kell 2006; Hastie et al. 2001; Ioannidis 2005, 2008). A variety of other network inferencing methods also exist (e.g. Lorenz et al. 2009; Managbanag et al. 2008; Stolovitzky and Califano 2006; Stolovitzky et al. 2009).

Why most toxins will have multiple sites of action

A variety of means are available for testing these mechanisms further, including the use of modulation of the

expression of constituent parts of the proposed pathways by genetic means (e.g. Jones et al. 2007), and in particular the use of appropriate iron chelators and antioxidants, alone or (preferably) in combination. The combinatorial aspect is important, since there are theoretical grounds, as well as abundant evidence, to the effect that several steps in a network must be modulated *simultaneously* to exhibit substantial effects (see e.g. Ágoston et al. 2005; Ambesi-Impiombato and di Bernardo 2006; Bender et al. 2006; Borisy et al. 2003; Csermely et al. 2005; Dancey and Chen 2006; Fan et al. 2003, 2006; Haggarty et al. 2003; Hopkins et al. 2006; Kaelin 2005; Keith et al. 2005; Kostoff and Delafuente 2006; Millan 2006; Perlstein et al. 2006; Sharom et al. 2004; Shiomi et al. 1999; Wermuth 2004) (Brötz-Oesterhelt and Brunner 2008; Chong and Sullivan 2007; Costanzo et al. 2010; Gerber et al. 2007; Goltsov et al. 2009; Hegreness et al. 2008; Huang et al. 2008b; Jia et al. 2009; Koufaki et al. 2009; Kuhn et al. 2010; Lee et al. 2007; Lehár et al. 2007, 2008a, b; Li et al. 2007; Liby et al. 2007; Morphy and Rankovic 2007; Pachauri et al. 2009; Reznichenko et al. 2010; Sarkar and Li 2009; Sarkar et al. 2009; Sawyers 2007; Schratzenholz and Šoškić 2008; Silver et al. 2009; Silver 2007; Sporn et al. 2007; Stommel et al. 2007; Suchy et al. 2009; Ulrich-Merzenich et al. 2009; Watterson et al. 2008; Weinreb et al. 2009b; Whitehurst et al. 2007; Wong et al. 2008; Yeh et al. 2006, 2009; Ying et al. 2008; Zamin et al. 2009; Zhang 2006; Zhang et al. 2010; Zimmermann et al. 2007). The same holds true for biotechnology (Kell and Westerhoff 1986a), in which typically we seek to maximise fluxes (Park et al. 2007, 2008). Of course some molecules—both nutrients/natural products and pharmaceutical agents—can act as both antioxidants and iron chelators; these are likely to be especially useful in minimising the damage from toxicants of many kinds.

Concluding remarks

Science consists of both analysis and synthesis, and thus a substantial amount of Science involves finding regularities ('Laws') that can be extracted from observables, even when some of those observables or their presumed causes seem to have little in common (Kell and Oliver 2004; Oldroyd 1986; Westerhoff and Kell 2007). Thus, there are many kinds of toxins that can cause death or injury, but it is not necessarily obvious what they share in terms of their modes of action. What I have therefore sought to do here is to bring together the evidence for the autocatalytic involvement of poorly liganded iron (leading, *inter alia*, to hydroxyl radical formation) in a considerable number of these. Certainly the exact molecular mechanisms, cascades and networks involved in each case depend on many other factors, and

appropriate systems biology methods can tell us what they are, but I would argue that the extensive evidence for iron's involvement that I cite here and elsewhere (Kell 2009a) is very hard to ignore. Hence, the view that iron is the nexus.

Acknowledgments I thank a great many colleagues and correspondents for their contribution to the refinement of my thinking about these issues, and Julie Cowley for assistance with literature, etc. Previous work that has led to this review has been funded by the BBSRC, EPSRC and MRC.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Aamodt AH, Stovner LJ, Thorstensen K, Lydersen S, White LR, Aasly JO (2007) Prevalence of haemochromatosis gene mutations in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 78(3):315–317
- Abdelrahman M, Mazzon E, Bauer M, Bauer I, Delbosc S, Cristol JP et al (2005) Inhibitors of NADPH oxidase reduce the organ injury in hemorrhagic shock. *Shock* 23(2):107–114
- Abdelrahman M, Sharples EJ, McDonald MC, Collin M, Patel NS, Yaqoob MM et al (2004) Erythropoietin attenuates the tissue injury associated with hemorrhagic shock and myocardial ischemia. *Shock* 22(1):63–69
- Abrous DN, Koehl M, Le Moal M (2005) Adult neurogenesis: from precursors to network and physiology. *Physiol Rev* 85(2): 523–569
- Adamkiewicz TV, Abboud MR, Paley C, Olivieri N, Kirby-Allen M, Vichinsky E et al (2009) Serum ferritin level changes in children with sickle cell disease on chronic blood transfusion are nonlinear and are associated with iron load and liver injury. *Blood* 114(21):4632–4638
- Adams CWM (1988) Perivascular iron deposition and other vascular damage in multiple sclerosis. *J Neurol Neurosurg Psychiatry* 51(2):260–265
- Adams RJ (2007) Big strokes in small persons. *Arch Neurol* 64(11):1567–1574
- Adhami VM, Siddiqui IA, Sarfaraz S, Khwaja SI, Hafeez BB, Ahmad N et al (2009) Effective prostate cancer chemopreventive intervention with green tea polyphenols in the TRAMP model depends on the stage of the disease. *Clin Cancer Res* 15(6):1947–1953
- Adibhatla RM, Hatcher JF (2010) Lipid oxidation and peroxidation in CNS health and disease: from molecular mechanisms to therapeutic opportunities. *Antioxid Redox Signal* 12:125–169
- Adlard PA, Bush AI (2006) Metals and Alzheimer's disease. *J Alzheimers Dis* 10(2–3):145–163
- Agarwal MB (2010) Deferasirox: oral, once daily iron chelator—an expert opinion. *Indian J Pediatr* 77(2):185–191
- Agarwal R, Vasavada N, Sachs NG, Chase S (2004) Oxidative stress and renal injury with intravenous iron in patients with chronic kidney disease. *Kidney Int* 65(6):2279–2289
- Aggarwal BB, Harikumar KB (2009) Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *Int J Biochem Cell Biol* 41(1):40–59
- Aggarwal BB, Sung B (2009) Pharmacological basis for the role of curcumin in chronic diseases: an age-old spice with modern targets. *Trends Pharmacol Sci* 30(2):85–94

- Ágoston V, Csermely P, Pongor S (2005) Multiple weak hits confuse complex systems: a transcriptional regulatory network as an example. *Phys Rev E* 71(5):051909
- Agrawal R, Sharma PK, Rao GS (2001) Release of iron from ferritin by metabolites of benzene and superoxide radical generating agents. *Toxicology* 168(3):223–230
- Aguzzi A (2006) Prion diseases of humans and farm animals: epidemiology, genetics, and pathogenesis. *J Neurochem* 97(6):1726–1739
- Aguzzi A, Heikenwalder M (2006) Pathogenesis of prion diseases: current status and future outlook. *Nat Rev Microbiol* 4(10):765–775
- Aguzzi A, Heikenwalder M, Polymenidou M (2007) Insights into prion strains and neurotoxicity. *Nat Rev Mol Cell Biol* 8(7):552–561
- Aguzzi A, Baumann F, Bremer J (2008a) The prion's elusive reason for being. *Annu Rev Neurosci* 31:439–477
- Aguzzi A, Sigurdson C, Heikenwalder M (2008b) Molecular mechanisms of prion pathogenesis. *Annu Rev Pathol* 3:11–40
- Aharon-Peretz J, Rosenbaum H, Gershoni-Baruch R (2004) Mutations in the glucocerebrosidase gene and Parkinson's disease in Ashkenazi Jews. *N Engl J Med* 351(19):1972–1977
- Ahluwalia N, Genoux A, Ferrieres J, Perret B, Carayol M, Drouet L et al (2010) Iron status is associated with carotid atherosclerotic plaques in middle-aged adults. *J Nutr* 140(4):812–816
- Aigner E, Theurl I, Theurl M, Lederer D, Haufe H, Dietze O et al (2008) Pathways underlying iron accumulation in human nonalcoholic fatty liver disease. *Am J Clin Nutr* 87(5):1374–1383
- Akaike T, Noguchi Y, Ijiri S, Setoguchi K, Suga M, Zheng YM et al (1996) Pathogenesis of influenza virus-induced pneumonia: involvement of both nitric oxide and oxygen radicals. *Proc Natl Acad Sci USA* 93(6):2448–2453
- Akhtar MJ, Kumar S, Murthy RC, Ashquin M, Khan MI, Patil G et al. (2010) The primary role of iron-mediated lipid peroxidation in the differential cytotoxicity caused by two varieties of talc nanoparticles on A(549) cells and lipid peroxidation inhibitory effect exerted by ascorbic acid. *Toxicol In Vitro*
- Al-Bayati ZA, Stohs SJ (1987) The role of iron in 2,3,7,8-tetrachlorodibenzo-*p*-dioxin-induced lipid peroxidation by rat liver microsomes. *Toxicol Lett* 38(1–2):115–121
- Al-Turk WA, Shara MA, Mohammadpour H, Stohs SJ (1988) Dietary iron and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin induced alterations in hepatic lipid peroxidation, glutathione content and body weight. *Drug Chem Toxicol* 11(1):55–70
- Alam ZI, Jenner A, Daniel SE, Lees AJ, Cairns N, Marsden CD et al (1997) Oxidative DNA damage in the parkinsonian brain: An apparent selective increase in 8-hydroxyguanine levels in substantia nigra. *J Neurochem* 69(3):1196–1203
- Alavian SM, Tabatabaei SV (2009) Therapy of hepatitis C in thalassemia: the influence of iron on achieving sustained viral response. *Ann Hematol*
- Ali S, Mann DA (2004) Signal transduction via the NF-kappaB pathway: a targeted treatment modality for infection, inflammation and repair. *Cell Biochem Funct* 22(2):67–79
- Ali SF, LeBel CP, Bondy SC (1992) Reactive oxygen species formation as a biomarker of methylmercury and trimethyltin neurotoxicity. *Neurotoxicology* 13:637–648
- Alissa EM, Ahmed WH, Al-Ama N, Ferns GA (2007) Relationship between indices of iron status and coronary risk factors including diabetes and the metabolic syndrome in Saudi subjects without overt coronary disease. *J Trace Elem Med Biol* 21(4):242–254
- Aljandali A, Pollack H, Yeldandi A, Li Y, Weitzman SA, Kamp DW (2001) Asbestos causes apoptosis in alveolar epithelial cells: role of iron-induced free radicals. *J Lab Clin Med* 137(5):330–339
- Alla V, Bonkovsky HL (2005) Iron in nonhemochromatotic liver disorders. *Semin Liver Dis* 25(4):461–472
- Allen K, Cornforth D (2009) Effect of chelating agents and spice-derived antioxidants on myoglobin oxidation in a lipid-free model system. *J Food Sci* 74(5):C375–379
- Alon U (2006) An introduction to systems biology: design principles of biological circuits. Chapman and Hall/CRC, London
- Altamura C, Squitti R, Pasqualetti P, Gaudino C, Palazzo P, Tibuzzi F et al (2009) Ceruloplasmin/Transferrin system is related to clinical status in acute stroke. *Stroke* 40(4):1282–1288
- Altamura S, Muckenthaler MU (2009) Iron toxicity in diseases of aging: Alzheimer's disease, Parkinson's disease and atherosclerosis. *J Alzheimers Dis* 16(4):879–895
- Altman DG, Royston P (2000) What do we mean by validating a prognostic model? *Stat Med* 19(4):453–473
- Altman R (2003) Risk factors in coronary atherosclerosis athero-inflammation: the meeting point. *Thromb J* 1(1):4
- Ambesi-Impimbato A, di Bernardo D (2006) Computational biology and drug discovery: from single-target to network drugs. *Curr Bioinform* 1(1):3–13
- Ameli M, Besharati S, Nemati K, Zamani F (2008) Relationship between elevated liver enzyme with iron overload and viral hepatitis in thalassemia major patients in Northern Iran. *Saudi Med J* 29(11):1611–1615
- Amic D, Davidovic-Amic D, Beslo D, Rastija V, Lucic B, Trinajstic N (2007) SAR and QSAR of the antioxidant activity of flavonoids. *Curr Med Chem* 14(7):827–845
- Amit T, Avramovich-Tirosh Y, Youdim MB, Mandel S (2008) Targeting multiple Alzheimer's disease etiologies with multimodal neuroprotective and neurorestorative iron chelators. *FASEB J* 22(5):1296–1305
- Anand P, Nair HB, Sung B, Kunnumakkara AB, Yadav VR, Tekmal RR et al (2010) Design of curcumin-loaded PLGA nanoparticles formulation with enhanced cellular uptake, and increased bioactivity in vitro and superior bioavailability in vivo. *Biochem Pharmacol* 79(3):330–338
- Ananiadou S, Kell DB, Tsujii J-i (2006) Text mining and its potential applications in systems biology. *Trends Biotechnol* 24:571–579
- Andersen JK (2003) Paraquat and iron exposure as possible synergistic environmental risk factors in Parkinson's disease. *Neurotox Res* 5(5):307–313
- Andersen JK (2004) Iron dysregulation and Parkinson's disease. *J Alzheimers Dis* 6(6 Suppl):S47–S52
- Anderson PR, Kirby K, Hilliker AJ, Phillips JP (2005) RNAi-mediated suppression of the mitochondrial iron chaperone, frataxin, in *Drosophila*. *Hum Mol Genet* 14(22):3397–3405
- Anderson PR, Kirby K, Orr WC, Hilliker AJ, Phillips JP (2008) Hydrogen peroxide scavenging rescues frataxin deficiency in a *Drosophila* model of Friedreich's ataxia. *Proc Natl Acad Sci USA* 105(2):611–616
- Andrades M, Ritter C, Moreira JC, Dal-Pizzol F (2005) Oxidative parameters differences during non-lethal and lethal sepsis development. *J Surg Res* 125(1):68–72
- Andreini C, Bertini I, Cavallaro G, Holliday GL, Thornton JM (2008) Metal ions in biological catalysis: from enzyme databases to general principles. *J Biol Inorg Chem* 13(8):1205–1218
- Andreini C, Bertini I, Cavallaro G, Najmanovich RJ, Thornton JM (2009) Structural analysis of metal sites in proteins: non-heme iron sites as a case study. *J Mol Biol* 388(2):356–380
- Andrews SC, Robinson AK, Rodriguez-Quinones F (2003) Bacterial iron homeostasis. *FEMS Microbiol Rev* 27(2–3):215–237
- Arakawa K, Yamada Y, Shinoda K, Nakayama Y, Tomita M (2006) GEM System: automatic prototyping of cell-wide metabolic pathway models from genomes. *BMC Bioinform* 7:168
- Armand P, Kim HT, Cutler CS, Ho VT, Koreth J, Alyea EP et al (2007) Prognostic impact of elevated pretransplantation serum

- ferritin in patients undergoing myeloablative stem cell transplantation. *Blood* 109(10):4586–4588
- Armengou A, Davalos A (2002) A review of the state of research into the role of iron in stroke. *J Nutr Health Aging* 6(3):207–208
- Armstrong RA, Lantos PL, Cairns NJ (2001) Spatial correlations between the vacuolation, prion protein deposits, and surviving neurons in the cerebral cortex in sporadic Creutzfeldt-Jakob disease. *Neuropathology* 21(4):266–271
- Arosio P, Ingrassia R, Cavadini P (2009) Ferritins: a family of molecules for iron storage, antioxidation and more. *Biochim Biophys Acta* 1790:589–599
- Arreguin S, Nelson P, Padway S, Shirazi M, Pierpont C (2009) Dopamine complexes of iron in the etiology and pathogenesis of Parkinson's disease. *J Inorg Biochem* 103(1):87–93
- Arts ICW, Hollman PCH (2005) Polyphenols and disease risk in epidemiologic studies. *Am J Clin Nutr* 81(1):317S–325S
- Aschner M, Syversen T, Souza DO, Rocha JB, Farina M (2007) Involvement of glutamate and reactive oxygen species in methylmercury neurotoxicity. *Braz J Med Biol Res* 40(3):285–291
- Ashall L, Horton CA, Nelson DE, Paszek P, Ryan S, Sillitoe K et al (2009) Pulsatile stimulation determines timing and specificity of NF-kappa-B-dependent transcription. *Science* 324:242–246
- Aslan M, Ryan TM, Adler B, Townes TM, Parks DA, Thompson JA et al (2001) Oxygen radical inhibition of nitric oxide-dependent vascular function in sickle cell disease. *Proc Natl Acad Sci* 98(26):15215–15220
- Assenza G, Zappasodi F, Squitti R, Altamura C, Ventriglia M, Ercolani M et al (2009) Neuronal functionality assessed by magnetoencephalography is related to oxidative stress system in acute ischemic stroke. *Neuroimage* 44(4):1267–1273
- Atamna H (2009) Amino acids variations in amyloid-beta peptides, mitochondrial dysfunction, and new therapies for Alzheimer's disease. *J Bioenerg Biomembr* 41:457–464
- Atamna H, Frey W H 2nd, Ko N (2009) Human and rodent amyloid-beta peptides differentially bind heme: relevance to the human susceptibility to Alzheimer's disease. *Arch Biochem Biophys* 487(1):59–65
- Atanasova B, Mudway IS, Laftah AH, Latunde-Dada GO, McKie AT, Peters TJ et al (2004) Duodenal ascorbate levels are changed in mice with altered iron metabolism. *J Nutr* 134(3):501–505
- Atchison WD, Hare MF (1994) Mechanisms of methylmercury-induced neurotoxicity. *FASEB J* 8(9):622–629
- Atkinson A, Winge DR (2009) Metal acquisition and availability in the mitochondria. *Chem Rev* 109(10):4708–4721
- Attwood TK, Kell DB, McDermott P, Marsh J, Pettifer SR, Thorne D (2009) Calling international rescue: knowledge lost in literature and data landslide!. *Biochem J* 424:317–333
- Auclair S, Milenkovic D, Besson C, Chauvet S, Gueux E, Morand C et al (2009) Catechin reduces atherosclerotic lesion development in apo E-deficient mice: a transcriptomic study. *Atherosclerosis* 204(2):e21–e27
- Avramovich-Tirosh Y, Amit T, Bar-Am O, Zheng HL, Fridkin M, Youdim MBH (2007a) Therapeutic targets and potential of the novel brain-permeable multifunctional iron chelator-monoamine oxidase inhibitor drug, M-30, for the treatment of Alzheimer's disease. *J Neurochem* 100(2):490–502
- Avramovich-Tirosh Y, Reznichenko L, Mit T, Zheng H, Fridkin M, Weinreb O et al (2007b) Neurorescue activity, APP regulation and amyloid-beta peptide reduction by novel multi-functional brain permeable iron-chelating-antioxidants, M-30 and green tea polyphenol, EGCG. *Curr Alzheimer Res* 4(4):403–411
- Ayaki H, Lee MJ, Sumino K, Nishio H (2005) Different cytoprotective effect of antioxidants and change in the iron regulatory system in rodent cells exposed to paraquat or formaldehyde. *Toxicology* 208(1):73–79
- Ayala-Castro C, Saini A, Outten FW (2008) Fe-S cluster assembly pathways in bacteria. *Microbiol Mol Biol Rev* 72(1):110–125. Table of contents
- Azad N, Rojanasakul Y, Vallyathan V (2008) Inflammation and lung cancer: roles of reactive oxygen/nitrogen species. *J Toxicol Environ Health B Crit Rev* 11(1):1–15
- Ba LA, Doering M, Burkholz T, Jacob C (2009) Metal trafficking: from maintaining the metal homeostasis to future drug design. *Metallomics* 1(4):292–311
- Babady NE, Carelle N, Wells RD, Rouault TA, Hirano M, Lynch DR et al (2007) Advancements in the pathophysiology of Friedreich's Ataxia and new prospects for treatments. *Mol Genet Metab* 92(1–2):23–35
- Babatasi G, Masetti M, Galateau F, Mosquet B, Khayat A, Evrard C (1996) Bronchial necrosis induced by inhalation of an iron tablet. *J Thorac Cardiovasc Surg* 112(5):1397–1399
- Babior BM (2000) Phagocytes and oxidative stress. *Am J Med* 109(1):33–44
- Bäck T, Fogel DB, Michalewicz Z (eds) (1997) Handbook of evolutionary computation. IOP Publishing/Oxford University Press, Oxford
- Baeuerle PA, Henkel T (1994) Function and activation of NF-kappa B in the immune system. *Annu Rev Immunol* 12:141–179
- Bailey DM, Robach P, Thomsen JJ, Lundby C (2006) Erythropoietin depletes iron stores: antioxidant neuroprotection for ischemic stroke? *Stroke* 37(10):2453
- Baker ML, Wang JJ, Rogers S, Klein R, Kuller LH, Larsen EK et al (2009) Early age-related macular degeneration, cognitive function, and dementia: the cardiovascular health study. *Arch Ophthalmol* 127(5):667–673
- Bakshi R, Benedict RH, Bermel RA, Caruthers SD, Puli SR, Tjoa CW et al (2002) T2 hypointensity in the deep gray matter of patients with multiple sclerosis: a quantitative magnetic resonance imaging study. *Arch Neurol* 59(1):62–68
- Bakshi R, Dmochowski J, Shaikh ZA, Jacobs L (2001) Gray matter T2 hypointensity is related to plaques and atrophy in the brains of multiple sclerosis patients. *J Neurol Sci* 185(1):19–26
- Bakshi R, Shaikh ZA, Janardhan V (2000) MRI T2 shortening ('black T2') in multiple sclerosis: frequency, location, and clinical correlation. *Neuroreport* 11(1):15–21
- Bakshi R, Thompson AJ, Rocca MA, Pelletier D, Dousset V, Barkhof F et al (2008) MRI in multiple sclerosis: current status and future prospects. *Lancet Neurol* 7(7):615–625
- Bal W, Kasprzak KS (2002) Induction of oxidative DNA damage by carcinogenic metals. *Toxicol Lett* 127(1–3):55–62
- Balla G, Jacob HS, Balla J, Rosenberg M, Nath K, Apple F et al (1992) Ferritin: a cytoprotective antioxidant strategem of endothelium. *J Biol Chem* 267(25):18148–18153
- Bandyopadhyay S, Huang X, Cho H, Greig NH, Youdim MB, Rogers JT (2006) Metal specificity of an iron-responsive element in Alzheimer's APP mRNA 5' untranslated region, tolerance of SH-SY5Y and H4 neural cells to desferrioxamine, clioquinol, VK-28, and a piperazine chelator. *J Neural Trans Suppl* 71:237–247
- Banga JR (2008) Optimization in computational systems biology. *BMC Syst Biol* 2(1):47
- Banga JR, Balsacanto E (2008) Parameter estimation and optimal experimental design. *Essays Biochem* 45:195–209
- Baptista-Gonzalez H, Chavez-Tapia NC, Zamora-Valdes D, Uribe M, Mendez-Sanchez N (2008) Importance of iron and iron metabolism in nonalcoholic fatty liver disease. *Mini Rev Med Chem* 8(2):171–174
- Baranzini SE, Mudge J, van Velkinburgh JC, Khankhanian P, Khrebtukova I, Miller NA et al (2010) Genome, epigenome and RNA sequences of monozygotic twins discordant for multiple sclerosis. *Nature* 464(7293):1351–1356

- Barapatre N, Morawski M, Butz T, Reinert T (2010) Trace element mapping in Parkinsonian brain by quantitative ion beam microscopy. *Nucl Instrum Methods Phys Res B* 268:2156–2159
- Bareggi SR, Braida D, Pollera C, Bondiolotti G, Formentin E, Puricelli M et al (2009) Effects of clioquinol on memory impairment and the neurochemical modifications induced by scrapie infection in golden hamsters. *Brain Res* 1280:195–200
- Barichello T, Machado RA, Constantino L, Valvassori SS, Reus GZ, Martins MR et al (2007) Antioxidant treatment prevented late memory impairment in an animal model of sepsis. *Crit Care Med* 35(9):2186–2190
- Barker JD Jr, de Carle DJ, Anuras S (1977) Chronic excessive acetaminophen use and liver damage. *Ann Intern Med* 87(3):299–301
- Barnham KJ, Cappai R, Beyreuther K, Masters CL, Hill AF (2006) Delineating common molecular mechanisms in Alzheimer's and prion diseases. *Trends Biochem Sci* 31(8):465–472
- Barnham KJ, Masters CL, Bush AI (2004) Neurodegenerative diseases and oxidative stress. *Nat Rev Drug Discov* 3(3):205–214
- Barry SM, Challis GL (2009) Recent advances in siderophore biosynthesis. *Curr Opin Chem Biol* 13(2):205–215
- Bartzokis G, Lu PH, Tishler TA, Peters DG, Kosenko A, Barrall KA et al (2010) Prevalent iron metabolism gene variants associated with increased brain ferritin iron in healthy older men. *J Alzheimers Dis* 20(1):333–341
- Bartzokis G, Tishler TA, Lu PH, Villablanca P, Altshuler LL, Carter M et al (2007) Brain ferritin iron may influence age- and gender-related risks of neurodegeneration. *Neurobiol Aging* 28(3):414–423
- Basak R, Chowdhury A, Fatmi L, Saha N, Mollah A, Yasmin S (2008) Stroke in the young: relationship with iron deficiency anemia and thrombocytosis. *Mymensingh Med J* 17(1):74–77
- Basile V, Ferrari E, Lazzari S, Belluti S, Pignedoli F, Imbriano C (2009) Curcumin derivatives: molecular basis of their anti-cancer activity. *Biochem Pharmacol* 78(10):1305–1315
- Bassett ML (2007) Iron and hepatitis C: beginning to make sense. *J Gastroenterol Hepatol* 22(11):1703–1704
- Basu S, Mohan ML, Luo X, Kundu B, Kong Q, Singh N (2007) Modulation of proteinase K-resistant prion protein in cells and infectious brain homogenate by redox iron: implications for prion replication and disease pathogenesis. *Mol Biol Cell* 18(9):3302–3312
- Batts KP (2007) Iron overload syndromes and the liver. *Mod Pathol* 20(Suppl 1):S31–S39
- Baudrexel S, Nurnberger L, Rub U, Seifried C, Klein JC, Deller T et al. (2010) Quantitative mapping of T1 and T2* discloses nigral and brainstem pathology in early Parkinson's disease. *Neuroimage*
- Baue AE, Durham R, Faist E (1998) Systemic inflammatory response syndrome (SIRS), multiple organ dysfunction syndrome (MODS), multiple organ failure (MOF): are we winning the battle? *Shock* 10(2):79–89
- Bauer PO, Nukina N (2009) The pathogenic mechanisms of polyglutamine diseases and current therapeutic strategies. *J Neurochem* 110(6):1737–1765
- Baune BT, Neuhauser H, Ellert U, Berger K (2010) The role of the inflammatory markers ferritin, transferrin and fibrinogen in the relationship between major depression and cardiovascular disorders—the German health interview and examination survey. *Acta Psychiatr Scand* 121(2):135–142
- Beal AL, Cerra FB (1994) Multiple organ failure syndrome in the 1990s—systemic inflammatory response and organ dysfunction. *JAMA* 271(3):226–233
- Beck MA, Handy J, Levander OA (2004) Host nutritional status: the neglected virulence factor. *Trends Microbiol* 12(9):417–423
- Becker E, Richardson DR (2001) Frataxin: its role in iron metabolism and the pathogenesis of Friedreich's ataxia. *Int J Biochem Cell Biol* 33(1):1–10
- Becker JS (2010) Bioimaging of metals in brain tissue from micrometre to nanometre scale by laser ablation inductively coupled plasma mass spectrometry: state of the art and perspectives. *Int J Mass Spectrom* 289(2–3):65–75
- Becker JS, Matusch A, Palm C, Salber D, Morton KA, Becker S (2010a) Bioimaging of metals in brain tissue by laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) and metallomics. *Metallomics* 2(2):104–111
- Becker JS, Zoriy M, Matusch A, Wu B, Salber D, Palm C et al (2010b) Bioimaging of metals by laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS). *Mass Spectrom Rev* 29(1):156–175
- Beckett GJ, Foster GR, Hussey AJ, Oliveira DB, Donovan JW, Prescott LF et al (1989) Plasma glutathione S-transferase and F protein are more sensitive than alanine aminotransferase as markers of paracetamol (acetaminophen)-induced liver damage. *Clin Chem* 35(11):2186–2189
- Beckman JS (1996) Oxidative damage and tyrosine nitration from peroxynitrite. *Chem Res Toxicol* 9(5):836–844
- Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA (1990) Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA* 87(4):1620–1624
- Beckman JS, Ischiropoulos H, Zhu L, van der Woerd M, Smith C, Chen J et al (1992) Kinetics of superoxide dismutase- and iron-catalyzed nitration of phenolics by peroxynitrite. *Arch Biochem Biophys* 298(2):438–445
- Beckman JS, Koppenol WH (1996) Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol* 271(5 Pt 1):C1424–C1437
- Bedard K, Krause KH (2007) The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 87(1):245–313
- Beeri MS, Schmeidler J, Silverman JM, Gandy S, Wysocki M, Hannigan CM et al (2008) Insulin in combination with other diabetes medication is associated with less Alzheimer neuropathology. *Neurology* 71(10):750–757
- Bekyarova G, Tancheva S, Hristova M (2009) Protective effect of melatonin against oxidative hepatic injury after experimental thermal trauma. *Methods Find Exp Clin Pharmacol* 31(1):11–14
- Ben-Shachar D, Riederer P, Youdim MB (1991) Iron-melanin interaction and lipid peroxidation: implications for Parkinson's disease. *J Neurochem* 57(5):1609–1614
- Benarroch EE (2009) Brain iron homeostasis and neurodegenerative disease. *Neurology* 72(16):1436–1440
- Bencze KZ, Kondapalli KC, Cook JD, McMahon S, Millán-Pacheco C, Pastor N et al (2006) The structure and function of frataxin. *Crit Rev Biochem Mol Biol* 41(5):269–291
- Bencze KZ, Yoon T, Millán-Pacheco C, Bradley PB, Pastor N, Cowan JA et al (2007) Human frataxin: iron and ferrocyclase binding surface. *Chem Commun (Camb)* 18:1798–1800
- Bender A, Jenkins JL, Glick M, Deng Z, Nettles JH, Davies JW (2006) “Bayes affinity fingerprints” improve retrieval rates in virtual screening and define orthogonal bioactivity space: when are multitarget drugs a feasible concept? *J Chem Inf Model* 46(6):2445–2456
- Berdoukas V, Chouliaras G, Moraitis P, Zannikos K, Berdoussi E, Ladis V (2009) The efficacy of iron chelator regimes in reducing cardiac and hepatic iron in patients with thalassaemia major: a clinical observational study. *J Cardiovasc Magn Reson* 11(1):20
- Berg D (2006) In vivo detection of iron and neuromelanin by transcranial sonography—a new approach for early detection of substantia nigra damage. *J Neural Transm* 113(6):775–780

- Berg D (2007) Disturbance of iron metabolism as a contributing factor to SN hyperechogenicity in Parkinson's disease: implications for idiopathic and monogenetic forms. *Neurochem Res* 32(10):1646–1654
- Berg D, Gerlach M, Youdim MBH, Double KL, Zecca L, Riederer P et al (2001) Brain iron pathways and their relevance to Parkinson's disease. *J Neurochem* 79(2):225–236
- Berg D, Hochstrasser H (2006) Iron metabolism in parkinsonian syndromes. *Mov Disord* 21(9):1299–1310
- Berg D, Hochstrasser H, Schweitzer KJ, Riess O (2006) Disturbance of iron metabolism in Parkinson's disease ultrasonography as a biomarker. *Neurotox Res* 9(1):1–13
- Berg D, Riederer P, Gerlach M (2008) Contribution of disturbed iron metabolism to the pathogenesis of Parkinson's disease. *Future Med* 3(4):447–461
- Bergman M, Salman H, Pinchasi R, Straussberg R, Djaldetti M, Bessler H (2005) Phagocytic capacity and apoptosis of peripheral blood cells from patients with iron deficiency anemia. *Biomed Pharmacother* 59(6):307–311
- Bermel RA, Puli SR, Rudick RA, Weinstock-Guttman B, Fisher E, Munschauer F E 3rd et al (2005) Prediction of longitudinal brain atrophy in multiple sclerosis by gray matter magnetic resonance imaging T2 hypointensity. *Arch Neurol* 62(9):1371–1376
- Beutler E (2007) Iron storage disease: facts, fiction and progress. *Blood Cells Mol Dis* 39(2):140–147
- Bharath S, Hsu M, Kaur D, Rajagopalan S, Andersen JK (2002) Glutathione, iron and Parkinson's disease. *Biochem Pharmacol* 64(5–6):1037–1048
- Bhattacharya K, Dopp E, Kakkar P, Jaffery FN, Schiffmann D, Jaurand MC et al (2005) Biomarkers in risk assessment of asbestos exposure. *Mutat Res* 579(1–2):6–21
- Bhidayasiri R, Truong DD (2004) Chorea and related disorders. *Postgrad Med J* 80(947):527–534
- Biasiotto G, Goldwurm S, Finazzi D, Tunesi S, Zecchinelli A, Sironi F et al (2008) HFE gene mutations in a population of Italian Parkinson's disease patients. *Parkinsonism Relat Disord* 14(5):426–430
- Bidle KD, Bender SJ (2008) Iron starvation and culture age activate metacaspases and programmed cell death in the marine diatom *Thalassiosira pseudonana*. *Eukaryot Cell* 7(2):223–236
- Binder CJ, Chang MK, Shaw PX, Miller YI, Hartvigsen K, Dewan A et al (2002) Innate and acquired immunity in atherosclerosis. *Nat Med* 8(11):1218–1226
- Biran Y, Masters CL, Barnham KJ, Bush AI, Adlard PA (2009) Pharmacotherapeutic targets in Alzheimer's disease. *J Cell Mol Med* 13(1):61–86
- Bishop GM, Robinson SR (2001) Quantitative analysis of cell death and ferritin expression in response to cortical iron: implications for hypoxia-ischemia and stroke. *Brain Res* 907(1–2):175–187
- Bishop GM, Robinson SR, Liu Q, Perry G, Atwood CS, Smith MA (2002) Iron: a pathological mediator of Alzheimer disease? *Dev Neurosci* 24(2–3):184–187
- Bishop GM, Dang TN, Dringen R, Robinson SR (2010a) Accumulation of non-transferrin-bound iron by neurons, astrocytes, and microglia. *Neurotox Res* (in press)
- Bishop NA, Lu T, Yankner BA (2010b) Neural mechanisms of ageing and cognitive decline. *Nature* 464(7288):529–535
- Biswas SK, Rahman I (2009) Environmental toxicity, redox signaling and lung inflammation: the role of glutathione. *Mol Aspects Med* 30(1–2):60–76
- Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C (2008) Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database of Systematic Reviews* 2008, CD007176.
- Blake GJ, Ridker PM (2001) Novel clinical markers of vascular wall inflammation. *Circ Res* 89(9):763–771
- Blasiak J, Sklodowska A, Ulinska M, Szaflik JP (2009) Iron and age-related macular degeneration. *Klin Oczna* 111(4–6):174–177
- Blázquez L, De Juan D, Ruiz-Martínez J, Empanaza JI, Saenz A, Otaegui D et al (2007) Genes related to iron metabolism and susceptibility to Alzheimer's disease in Basque population. *Neurobiol Aging* 28(12):1941–1943
- Blum D, Torch S, Lambeng N, Nissou MF, Benabid AL, Sadoul R et al (2001) Molecular pathways involved in the neurotoxicity of 6-OHDA, dopamine and MPTP: contribution to the apoptotic theory in Parkinson's disease. *Prog Neurobiol* 65(2):135–172
- Boddaert N, Le Quan Sang KH, Rotig A, Leroy-Willig A, Gallet S, Brunelle F et al (2007) Selective iron chelation in Friedreich ataxia: biologic and clinical implications. *Blood* 110(1):401–408
- Boelaert JR, Weinberg GA, Weinberg ED (1996) Altered iron metabolism in HIV infection: mechanisms, possible consequences, and proposals for management. *Infect Agents Dis* 5(1):36–46
- Bolognin S, Drago D, Messori L, Zatta P (2009a) Chelation therapy for neurodegenerative diseases. *Med Res Rev* 29(4):547–570
- Bolognin S, Messori L, Zatta P (2009b) Metal ion physiopathology in neurodegenerative disorders. *Neuromol Med* 11(4):223–238
- Bolton CF (1996) Sepsis and the systemic inflammatory response syndrome: neuromuscular manifestations. *Crit Care Med* 24(8):1408–1416
- Bomhard EM, Herbold BA (2005) Genotoxic activities of aniline and its metabolites and their relationship to the carcinogenicity of aniline in the spleen of rats. *Crit Rev Toxicol* 35(10):783–835
- Bone RC (1996) Toward a theory regarding the pathogenesis of the systemic inflammatory response syndrome: what we do and do not know about cytokine regulation. *Crit Care Med* 24(1):163–172
- Bongard JC, Lipson H (2005) Nonlinear system identification using coevolution of models and tests. *IEEE Trans Evol Comput* 9(4):361–384
- Bonkovsky HL (2002) Iron as a comorbid factor in chronic viral hepatitis. *Am J Gastroenterol* 97(1):1–4
- Bonkovsky HL, Naishadham D, Lambrecht RW, Chung RT, Hoefs JC, Nash SR et al (2006) Roles of iron and HFE mutations on severity and response to therapy during retreatment of advanced chronic hepatitis C. *Gastroenterology* 131(5):1440–1451
- Borah A, Mohanakumar KP (2009) Melatonin inhibits 6-hydroxydopamine production in the brain to protect against experimental parkinsonism in rodents. *J Pineal Res* 47(4):293–300
- Borgna-Pignatti C, Cappellini MD, De Stefano P, Del Vecchio GC, Forni GL, Gamberini MR et al (2005) Survival and complications in thalassemia. *Ann NY Acad Sci* 1054:40–47
- Borisy AA, Elliott PJ, Hurst NW, Lee MS, Lehar J, Price ER et al (2003) Systematic discovery of multicomponent therapeutics. *Proc Natl Acad Sci USA* 100(13):7977–7982
- Bornholdt S, Sneppen K (2000) Robustness as an evolutionary principle. *Proc R Soc B Biol Sci* 267(1459):2281–2286
- Bosomtvi A, Jiang Q, Ding GL, Zhang L, Zhang ZG, Lu M et al (2008) Quantitative evaluation of microvascular density after stroke in rats using MRI. *J Cereb Blood Flow Metab* 28(12):1978–1987
- Bowern N, Ramshaw IA, Clark IA, Doherty PC (1984) Inhibition of autoimmune neuropathological process by treatment with an iron-chelating agent. *J Exp Med* 160(5):1532–1543
- Bowler RP (2004) Oxidative stress in the pathogenesis of asthma. *Curr Allergy Asthma Rep* 4(2):116–122
- Bowler RP, Crapo JD (2002) Oxidative stress in allergic respiratory diseases. *J Allergy Clin Immunol* 110(3):349–356
- Boyer J, Liu RH (2004) Apple phytochemicals and their health benefits. *Nutr J* 3:5
- Brar S, Henderson D, Schenck J, Zimmerman EA (2009) Iron accumulation in the substantia nigra of patients with Alzheimer disease and parkinsonism. *Arch Neurol* 66(3):371–374

- Brard L, Granai CO, Swamy N (2006) Iron chelators deferoxamine and diethylenetriamine pentaacetic acid induce apoptosis in ovarian carcinoma. *Gynecol Oncol* 100(1):116–127
- Brass SD, Benedict RH, Weinstock-Guttman B, Munschauer F, Bakshi R (2006a) Cognitive impairment is associated with subcortical magnetic resonance imaging grey matter T2 hypointensity in multiple sclerosis. *Mult Scler* 12(4):437–444
- Brass SD, Chen NK, Mulkern RV, Bakshi R (2006b) Magnetic resonance imaging of iron deposition in neurological disorders. *Top Magn Reson Imaging* 17(1):31–40
- Braun S, Ndrepepa G, von Beckerath N, Vogt W, Schomig A, Kastrati A (2004) Value of serum ferritin and soluble transferrin receptor for prediction of coronary artery disease and its clinical presentations. *Atherosclerosis* 174(1):105–110
- Breiman L (2001) Statistical modeling: the two cultures. *Stat Sci* 16(3):199–215
- Breitkopf K, Nagy LE, Beier JI, Mueller S, Weng H, Dooley S (2009) Current experimental perspectives on the clinical progression of alcoholic liver disease. *Alcohol Clin Exp Res* 33(10):1647–1655
- Brennan LA, Kantorow M (2009) Mitochondrial function and redox control in the aging eye: role of MsrA and other repair systems in cataract and macular degenerations. *Exp Eye Res* 88(2):195–203
- Brewer GJ (2007) Iron and copper toxicity in diseases of aging, particularly atherosclerosis and Alzheimer's disease. *Exp Biol Med* 232(2):323–335
- Brewer GJ (2010) Risks of copper and iron toxicity during aging in humans. *Chem Res Toxicol* 23(2):319–326
- Broadhurst D, Goodacre R, Jones A, Rowland JJ, Kell DB (1997) Genetic algorithms as a method for variable selection in multiple linear regression and partial least squares regression, with applications to pyrolysis mass spectrometry. *Anal Chim Acta* 348(1–3):71–86
- Broadhurst D, Kell DB (2006) Statistical strategies for avoiding false discoveries in metabolomics and related experiments. *Metabolomics* 2(4):171–196. doi:10.1007/s11306-006-0037-z
- Broedbaek K, Poulsen HE, Weimann A, Kom GD, Schwedhelm E, Nielsen P et al (2009) Urinary excretion of biomarkers of oxidatively damaged DNA and RNA in hereditary hemochromatosis. *Free Radic Biol Med* 47:1230–1233
- Brötz-Oesterhelt H, Brunner NA (2008) How many modes of action should an antibiotic have? *Curr Opin Pharmacol* 8:567–573
- Brown DR (2009a) Brain proteins that mind metals: a neurodegenerative perspective. *Dalton Trans* 21:4069–4076
- Brown DR (2009b) Role of microglia in age-related changes to the nervous system. *ScientificWorldJournal* 9:1061–1071
- Brown DR, Qin K, Herms JW, Madlung A, Manson J, Strome R et al (1997) The cellular prion protein binds copper in vivo. *Nature* 390(6661):684–687
- Brown M, He F, Wilkinson SJ (2010) Properties of the proximate parameter tuning regularization algorithm. *Bull Math Biol* 72(3):697–718
- Brudevold R, Hole T, Hammerstrøm J (2008) Hyperferritinemia is associated with insulin resistance and fatty liver in patients without iron overload. *PLoS One* 3(10):e3547
- Brune M, Rossander L, Hallberg L (1989) Iron absorption and phenolic compounds: importance of different phenolic structures. *Eur J Clin Nutr* 43(8):547–557
- Bu DX, Hemdahl AL, Gabrielsen A, Fuxe J, Zhu C, Eriksson P et al (2006) Induction of neutrophil gelatinase-associated lipocalin in vascular injury via activation of nuclear factor-kappaB. *Am J Pathol* 169(6):2245–2253
- Buchanan DD, Silburn PA, Chalk JB, Le Couteur DG, Mellick GD (2002) The Cys282Tyr polymorphism in the HFE gene in Australian Parkinson's disease patients. *Neurosci Lett* 327(2):91–94
- Büeler H (2009) Impaired mitochondrial dynamics and function in the pathogenesis of Parkinson's disease. *Exp Neurol* 218(2):235–246
- Bugianesi E, Manzini P, D'Antico S, Vanni E, Longo F, Leone N et al (2004) Relative contribution of iron burden, HFE mutations, and insulin resistance to fibrosis in nonalcoholic fatty liver. *Hepatology* 39(1):179–187
- Bulger EM, Maier RV (2001) Antioxidants in critical illness. *Arch Surg* 136(10):1201–1207
- Bulgin MS, Melson SS (2007) What veterinary practitioners should know about scrapie. *J Am Vet Med Assoc* 230(8):1158–1164, 1120.
- Bulteau AL, Dancis A, Gareil M, Montagne JJ, Camadro JM, Lesuisse E (2007) Oxidative stress and protease dysfunction in the yeast model of Friedreich ataxia. *Free Radic Biol Med* 42(10):1561–1570
- Bulucu F, Ocal R, Karadurmus N, Sahin M, Kenar L, Aydin A et al (2009) Effects of N-acetylcysteine, deferoxamine and selenium on doxorubicin-induced hepatotoxicity. *Biol Trace Elem Res* 132:184–196
- Burrows CJ, Muller JG (1998) Oxidative nucleobase modifications leading to strand scission. *Chem Rev* 98(3):1109–1152
- Busca A, Falda M, Manzini P, D'Antico S, Valfre A, Locatelli F et al (2010) Iron overload in patients receiving allogeneic hematopoietic stem cell transplantation: quantification of iron burden by a superconducting quantum interference device (SQUID) and therapeutic effectiveness of phlebotomy. *Biol Blood Marrow Transpl* 16(1):115–122
- Bush AI (2000) Metals and neuroscience. *Curr Opin Chem Biol* 4(2):184–191
- Bush AI (2003) The metallobiology of Alzheimer's disease. *Trends Neurosci* 26(4):207–214
- Bush AI (2008) Drug development based on the metals hypothesis of Alzheimer's disease. *J Alzheimers Dis* 15(2):223–240
- Buss JL, Neuzil J, Gellert N, Weber C, Ponka P (2003) Pyridoxal isonicotinoyl hydrazone analogs induce apoptosis in hematopoietic cells due to their iron-chelating properties. *Biochem Pharmacol* 65(2):161–172
- Butelli E, Titta L, Giorgio M, Mock HP, Matros A, Peterek S et al (2008) Enrichment of tomato fruit with health-promoting anthocyanins by expression of select transcription factors. *Nat Biotechnol* 26(11):1301–1308
- Butensky James E, Harmatz P, Lee M, Kennedy C, Petru A, Wara D et al (2009) Altered iron metabolism in children with human immunodeficiency virus disease. *Pediatr Hematol Oncol* 26(2):69–84
- Butt MS, Sultan MT (2009) Green tea: nature's defense against malignancies. *Crit Rev Food Sci Nutr* 49(5):463–473
- Butterfield DA, Reed T, Newman SF, Sultana R (2007) Roles of amyloid beta-peptide-associated oxidative stress and brain protein modifications in the pathogenesis of Alzheimer's disease and mild cognitive impairment. *Free Radic Biol Med* 43(5):658–677
- Buzan T (2002) How to mind map. Thorsons, London
- Cadenas S, Cadenas AM (2002) Fighting the stranger-antioxidant protection against endotoxin toxicity. *Toxicology* 180(1):45–63
- Cai J, Nelson KC, Wu M, Sternberg P Jr, Jones DP (2000) Oxidative damage and protection of the RPE. *Prog Retin Eye Res* 19(2):205–221
- Calabrese E (2008a) Hormesis: why it is important to toxicology and toxicologists. *Environ Toxicol Chem* 27:1451–1474
- Calabrese EJ (2008b) Converging concepts: adaptive response, preconditioning, and the Yerkes-Dodson Law are manifestations of hormesis. *Ageing Res Rev* 7(1):8–20
- Calabrese EJ (2008c) Hormesis and medicine. *Br J Clin Pharmacol*

- Calabrese EJ, Baldwin LA (1997) The dose determines the stimulation (and poison): development of a chemical hormesis database. *Int J Toxicol* 16(6):545–559
- Calabrese EJ, Baldwin LA (2001) Hormesis: U-shaped dose responses and their centrality in toxicology. *Trends Pharmacol Sci* 22(6):285–291
- Calabrese EJ, Baldwin LA (2003) Hormesis: the dose-response revolution. *Annu Rev Pharmacol Toxicol* 43:175–197
- Calabrese EJ, Blain R (2005) The occurrence of hormetic dose responses in the toxicological literature, the hormesis database: an overview. *Toxicol Appl Pharmacol* 202(3):289–301
- Calabrese EJ, Blain RB (2009) Hormesis and plant biology. *Environ Pollut* 157(1):42–48
- Camaschella C, Merlini R (2005) Inherited hemochromatosis: from genetics to clinics. *Minerva Med* 96(3):207–222
- Campanella A, Rovelli E, Santambrogio P, Cozzi A, Taroni F, Levi S (2009) Mitochondrial ferritin limits oxidative damage regulating mitochondrial iron availability: hypothesis for a protective role in Friedreich ataxia. *Hum Mol Genet* 18(1):1–11
- Candore G, Licastro F, Chiappelli M, Franceschi C, Lio D, Rita Balistreri C et al (2003) Association between the HFE mutations and unsuccessful ageing: a study in Alzheimer's disease patients from Northern Italy. *Mech Ageing Dev* 124(4):525–528
- Cao A, Galanello R (2010) Beta-thalassemia. *Genet Med* 12(2):61–76
- Cappellini MD, Pattoneri P (2009) Oral iron chelators. *Annu Rev Med* 60:25–38
- Cappellini MD, Porter J, El-Beshlawy A, Li CK, Seymour JF, Elalfy M et al (2010) Tailoring iron chelation by iron intake and serum ferritin: the prospective EPIC study of deferasirox in 1744 patients with transfusion-dependent anemias. *Haematologica* 95(4):557–566
- Carbonell T, Rama R (2007) Iron, oxidative stress and early neurological deterioration in ischemic stroke. *Curr Med Chem* 14(8):857–874
- Carrea G, Colonna S, Kelly DR, Lazcano A, Ottolina G, Roberts SM (2005) Polyamino acids as synthetic enzymes: mechanism, applications and relevance to prebiotic catalysis. *Trends Biotechnol* 23(10):507–513
- Carri MT, Ferri A, Cozzolino M, Calabrese L, Rotilio G (2003) Neurodegeneration in amyotrophic lateral sclerosis: the role of oxidative stress and altered homeostasis of metals. *Brain Res Bull* 61(4):365–374
- Carrillo-Vico A, Lardone PJ, Naji L, Fernández-Santos JM, Martín-Lacave I, Guerrero JM et al (2005) Beneficial pleiotropic actions of melatonin in an experimental model of septic shock in mice: regulation of pro-/anti-inflammatory cytokine network, protection against oxidative damage and anti-apoptotic effects. *J Pineal Res* 39(4):400–408
- Casadesus G, Smith MA, Zhu X, Aliev G, Cash AD, Honda K et al (2004) Alzheimer disease: evidence for a central pathogenic role of iron-mediated reactive oxygen species. *J Alzheimers Dis* 6(2):165–169
- Cass WA, Grondin R, Andersen AH, Zhang Z, Hardy PA, Hussey-Andersen LK et al (2007) Iron accumulation in the striatum predicts aging-related decline in motor function in rhesus monkeys. *Neurobiol Aging* 28(2):258–271
- Cassol OJ Jr, Rezin GT, Petronilho FC, Scaini G, Goncalves CL, Ferreira GK et al. (2009) Effects of N-acetylcysteine/deferrioxamine, taurine and RC-3095 on respiratory chain complexes and creatine kinase activities in rat brain after sepsis. *Neurochem Res*
- Castellani RJ, Moreira PI, Liu G, Dobson J, Perry G, Smith MA et al (2007) Iron: the redox-active center of oxidative stress in Alzheimer disease. *Neurochem Res* 32(10):1640–1645
- Castellani RJ, Perry G, Siedlak SL, Nunomura A, Shimohama S, Zhang J et al (2002) Hydroxynonenal adducts indicate a role for lipid peroxidation in neocortical and brainstem lewy bodies in humans. *Neurosci Lett* 319(1):25–28
- Castoldi AF, Coccini T, Ceccatelli S, Manzo L (2001) Neurotoxicity and molecular effects of methylmercury. *Brain Res Bull* 55(2):197–203
- Caughey B, Baron GS (2006) Prions and their partners in crime. *Nature* 443(7113):803–810
- Caughey B, Lansbury PT (2003) Protofibrils, pores, fibrils, and neurodegeneration: separating the responsible protein aggregates from the innocent bystanders. *Annu Rev Neurosci* 26:267–298
- Cavadini P, Gellera C, Patel PI, Isaya G (2000) Human frataxin maintains mitochondrial iron homeostasis in *Saccharomyces cerevisiae*. *Hum Mol Genet* 9(17):2523–2530
- Cavadini P, O'Neill HA, Benada O, Isaya G (2002) Assembly and iron-binding properties of human frataxin, the protein deficient in Friedreich ataxia. *Hum Mol Genet* 11(3):217–227
- Cave AC, Brewer AC, Narayanapanicker A, Ray R, Grieve DJ, Walker S et al (2006) NADPH oxidases in cardiovascular health and disease. [Review]. *Antioxid Redox Signal* 8(5–6):691–728
- Cazzola M, Della Porta MG, Malcovati L (2008) Clinical relevance of anemia and transfusion iron overload in myelodysplastic syndromes. *Hematol Am Soc Hematol Educ Program* 2008:166–175
- Ceccarelli A, Filippi M, Neema M, Arora A, Valsasina P, Rocca MA et al (2009) T2 hypointensity in the deep gray matter of patients with benign multiple sclerosis. *Mult Scler* 15(6):678–686
- Ceccarelli A, Rocca MA, Neema M, Martinelli V, Arora A, Tauhid S et al (2010) Deep gray matter T2 hypointensity is present in patients with clinically isolated syndromes suggestive of multiple sclerosis. *Mult Scler* 16(1):39–44
- Cemeli E, Baumgartner A, Anderson D (2009) Antioxidants and the comet assay. *Mutat Res* 681(1):51–67
- Ceysens B, Pauwels M, Meulemans B, Verbeelen D, Van den Branden C (2004) Increased oxidative stress in the mouse adriamycin model of glomerulosclerosis is accompanied by deposition of ferric iron and altered GGT activity in renal cortex. *Ren Fail* 26(1):21–27
- Chakravarty A (2003) Friedreich's ataxia—yesterday, today and tomorrow. *Neurol India* 51(2):176–182
- Chandrasekaran V, Taylor EW (2008) Molecular modeling of the oxidized form of nuclear factor-kappa B suggests a mechanism for redox regulation of DNA binding and transcriptional activation. *J Mol Graph Model* 26(5):861–867
- Chandrasekaran VR, Hsu DZ, Liu MY (2009) The protective effect of sesamol against mitochondrial oxidative stress and hepatic injury in acetaminophen-overdosed rats. *Shock* 32(1):89–93
- Chang EF, Claus CP, Vreman HJ, Wong RJ, Noble-Haesslein LJ (2005) Heme regulation in traumatic brain injury: relevance to the adult and developing brain. *J Cereb Blood Flow Metab* 25(11):1401–1417
- Chao CC, Park SH, Aust AE (1996) Participation of nitric oxide and iron in the oxidation of DNA in asbestos-treated human lung epithelial cells. *Arch Biochem Biophys* 326(1):152–157
- Chard DT, Miller DH (2009) What you see depends on how you look: gray matter lesions in multiple sclerosis. *Neurology* 73(12):918–919
- Charkoudian LK, Dentshev T, Lukinova N, Wolkow N, Dunaief JL, Franz KJ (2008) Iron prochelator BSIH protects retinal pigment epithelial cells against cell death induced by hydrogen peroxide. *J Inorg Biochem* 102(12):2130–2135
- Chau LY (2000) Iron and atherosclerosis. *Proc Natl Sci Councl Repub China B* 24(4):151–155
- Chen-Roetling J, Chen L, Regan RF (2009) Minocycline attenuates iron neurotoxicity in cortical cell cultures. *Biochem Biophys Res Commun*

- Chen H, Lukas TJ, Du N, Suyeoka G, Neufeld AH (2009) Dysfunction of the retinal pigment epithelium with age: increased iron decreases phagocytosis and lysosomal activity. *Invest Ophthalmol Vis Sci* 50(4):1895–1902
- Chen LW, Wang YQ, Wei LC, Shi M, Chan YS (2007) Chinese herbs and herbal extracts for neuroprotection of dopaminergic neurons and potential therapeutic treatment of Parkinson's disease. *CNS Neurol Disord Drug Targets* 6(4):273–281
- Chen Q, Marsh J, Ames B, Mossman B (1996) Detection of 8-oxo-2'-deoxyguanosine, a marker of oxidative DNA damage, in culture medium from human mesothelial cells exposed to crocidolite asbestos. *Carcinogenesis* 17(11):2525–2527
- Chen W, Stambolian D, Edwards AO, Branham KE, Othman M, Jakobsdottir J et al (2010) Genetic variants near TIMP3 and high-density lipoprotein-associated loci influence susceptibility to age-related macular degeneration. *Proc Natl Acad Sci USA*
- Chen X, Scholl TO, Stein TP (2006) Association of elevated serum ferritin levels and the risk of gestational diabetes mellitus in pregnant women: the Camden study. *Diabetes Care* 29(5):1077–1082
- Cheynier V (2005) Polyphenols in foods are more complex than often thought. *Am J Clin Nutr* 81(1 Suppl):223S–229S
- Chiueh CC, Andoh T, Lai AR, Lai E, Krishna G (2000) Neuroprotective strategies in Parkinson's disease: protection against progressive nigral damage induced by free radicals. *Neurotox Res* 2(2–3):293–310
- Cho S, Urata Y, Iida T, Goto S, Yamaguchi M, Sumikawa K et al (1998) Glutathione downregulates the phosphorylation of I kappa B: autoloop regulation of the NF-kappa B-mediated expression of NF-kappa B subunits by TNF-alpha in mouse vascular endothelial cells. *Biochem Biophys Res Commun* 253(1):104–108
- Cho TH, Nighoghossian N, Wiart M, Desestret V, Cakmak S, Berthezene Y et al (2007) USPIO-enhanced MRI of neuroinflammation at the sub-acute stage of ischemic stroke: preliminary data. *Cerebrovasc Dis* 24(6):544–546
- Chobot V (2010) Simultaneous detection of pro- and antioxidative effects in the variants of the deoxyribose degradation assay. *J Agric Food Chem* 58(4):2088–2094
- Chodaczek G, Saavedra-Molina A, Bacsí A, Kruzel ML, Sur S, Boldogh I (2007) Iron-mediated dismutation of superoxide anion augments antigen-induced allergic inflammation: effect of lactoferrin. *Postepy Hig Med Dosw (Online)* 61:268–276
- Choi CJ, Kanthasamy A, Anantharam V, Kanthasamy AG (2006) Interaction of metals with prion protein: possible role of divalent cations in the pathogenesis of prion diseases. *Neurotoxicology* 27(5):777–787
- Choi KM, Lee KW, Kim HY, Seo JA, Kim SG, Kim NH et al (2005) Association among serum ferritin, alanine aminotransferase levels, and metabolic syndrome in Korean postmenopausal women. *Metabolism* 54(11):1510–1514
- Choi YG, Park JH, Lim S (2009) Acupuncture inhibits ferric iron deposition and ferritin-heavy chain reduction in an MPTP-induced parkinsonism model. *Neurosci Lett* 450(2):92–96
- Chong CR, Sullivan D J Jr (2007) New uses for old drugs. *Nature* 448(7154):645–646
- Chopra V, Fox JH, Lieberman G, Dorsey K, Matson W, Waldmeier P et al (2007) A small-molecule therapeutic lead for Huntington's disease: preclinical pharmacology and efficacy of C2–8 in the R6/2 transgenic mouse. *Proc Natl Acad Sci USA* 104(42):16685–16689
- Chowers I, Wong R, Dentchev T, Farkas RH, Iacovelli J, Gunatilaka TL et al (2006) The iron carrier transferrin is upregulated in retinas from patients with age-related macular degeneration. *Invest Ophthalmol Vis Sci* 47(5):2135–2140
- Christen Y (2000) Oxidative stress and Alzheimer disease. *Am J Clin Nutr* 71(2):621S–629S
- Christie BR, Cameron HA (2006) Neurogenesis in the adult hippocampus. *Hippocampus* 16(3):199–207
- Chu J, Zhou CC, Lu N, Zhang X, Dong FT (2008) Genetic variants in three genes and smoking show strong associations with susceptibility to exudative age-related macular degeneration in a Chinese population. *Chin Med J (Engl)* 121(24):2525–2533
- Chung YL, Williams A, Ritchie D, Williams SC, Changani KK, Hope J et al (1999) Conflicting MRI signals from gliosis and neuronal vacuolation in prion diseases. *Neuroreport* 10(17):3471–3477
- Cienciewicki J, Trivedi S, Kleeberger SR (2008) Oxidants and the pathogenesis of lung diseases. *J Allergy Clin Immunol* 122(3):456–468 (quiz 469–470)
- Circu ML, Aw TY (2010) Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic Biol Med* 48(6):749–762
- Clark LN, Ross BM, Wang Y, Mejia-Santana H, Harris J, Louis ED et al (2007) Mutations in the glucocerebrosidase gene are associated with early-onset Parkinson disease. *Neurology* 69(12):1270–1277
- Clarkson TW, Magos L, Myers GJ (2003) The toxicology of mercury—current exposures and clinical manifestations. *N Engl J Med* 349(18):1731–1737
- Clifton MC, Corrent C, Strong RK (2009) Siderocalins: siderophore-binding proteins of the innate immune system. *Biometals* 22(4):557–564
- Closa D, Folch-Puy E (2004) Oxygen free radicals and the systemic inflammatory response. *IUBMB Life* 56(4):185–191
- Cohen AR, Galanello R, Pennell DJ, Cunningham MJ, Vichinsky E (2004) Thalassemia. *Hematology* 14–34
- Cole AR, Astell A, Green C, Sutherland C (2007) Molecular connexions between dementia and diabetes. *Neurosci Biobehav Rev* 31(7):1046–1063
- Coles SJ, Day NE, Murray-Rust P, Rzepa HS, Zhang Y (2005) Enhancement of the chemical semantic web through the use of InChI identifiers. *Organic & Biomolecular Chemistry* 3(10):1832–1834
- Collingwood J, Dobson J (2006) Mapping and characterization of iron compounds in Alzheimer's tissue. *J Alzheimers Dis* 10(2–3):215–222
- Collingwood JF, Chong RK, Kasama T, Cervera-Gontard L, Dunin-Borkowski RE, Perry G et al (2008) Three-dimensional tomographic imaging and characterization of iron compounds within Alzheimer's plaque core material. *J Alzheimers Dis* 14(2):235–245
- Collins FS (2010) *The language of life*. Profile Books, London
- Combarros O, Garcia-Roman M, Fontalba A, Fernandez-Luna JL, Llorca J, Infante J et al (2003) Interaction of the H63D mutation in the hemochromatosis gene with the apolipoprotein E epsilon 4 allele modulates age at onset of Alzheimer's disease. *Dement Geriatr Cogn Disord* 15(3):151–154
- Comhair SA, Erzurum SC (2010) Redox control of asthma: molecular mechanisms and therapeutic opportunities. *Antioxid Redox Signal* 12(1):93–124
- Connor JR, Lee SY (2006) HFE mutations and Alzheimer's disease. *J Alzheimers Dis* 10(2–3):267–276
- Connor JR, Menzies SL, St Martin SM, Mufson EJ (1992) A histochemical study of iron, transferrin, and ferritin in Alzheimer's diseased brains. *J Neurosci Res* 31(1):75–83
- Cook CI, Yu BP (1998) Iron accumulation in aging: modulation by dietary restriction. *Mech Ageing Devel* 102(1):1–13
- Cook JD, Bencze KZ, Jankovic AD, Crater AK, Busch CN, Bradley PB et al (2006) Monomeric yeast frataxin is an iron-binding protein. *Biochemistry* 45(25):7767–7777
- Cook R, Calabrese EJ (2006) The importance of hormesis to public health. *Environ Health Perspect* 114(11):1631–1635

- Cooke MS, Evans MD, Dizdaroglu M, Lunec J (2003) Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J* 17(10):1195–1214
- Cooke MS, Olinski R, Loft S (2008) Measurement and meaning of oxidatively modified DNA lesions in urine. *Cancer Epidemiol Biomarkers Prev* 17(1):3–14
- Cornelis P, Andrews SC (eds) (2010) Iron uptake and homeostasis in microorganisms. Horizon Press, New York
- Correia AR, Pastore C, Adinolfi S, Pastore A, Gomes CM (2008) Dynamics, stability and iron-binding activity of frataxin clinical mutants. *FEBS J* 275(14):3680–3690
- Costanzo M, Baryshnikova A, Bellay J, Kim Y, Spear ED, Sevier CS et al (2010) The genetic landscape of a cell. *Science* 327(5964):425–431
- Coveney PV, Fowler PW (2005) Modelling biological complexity: a physical scientist's perspective. *J R Soc Interface* 2(4):267–280
- Cowan KJ, Diamond MI, Welch WJ (2003) Polyglutamine protein aggregation and toxicity are linked to the cellular stress response. *Hum Mol Genet* 12(12):1377–1391
- Cowland JB, Muta T, Borregaard N (2006) IL-1beta-specific up-regulation of neutrophil gelatinase-associated lipocalin is controlled by IkappaB-zeta. *J Immunol* 176(9):5559–5566
- Cowland JB, Sorensen OE, Sehested M, Borregaard N (2003) Neutrophil gelatinase-associated lipocalin is up-regulated in human epithelial cells by IL-1 beta, but not by TNF-alpha. *J Immunol* 171(12):6630–6639
- Cox LA Jr (2009) Hormesis without cell killing. *Risk Anal* 29(3):393–400
- Cozzi A, Levi S, Corsi B, Santambrogio P, Campanella A, Gerardi G et al (2003) Role of iron and ferritin in TNFalpha-induced apoptosis in HeLa cells. *FEBS Lett* 537(1–3):187–192
- Cozzi A, Rovelli E, Frizzale G, Campanella A, Amendola M, Arosio P et al (2010) Oxidative stress and cell death in cells expressing L-ferritin variants causing neuroferritinopathy. *Neurobiol Dis* 37(1):77–85
- Cozzolino M, Ferri A, Carri MT (2008) Amyotrophic lateral sclerosis: from current developments in the laboratory to clinical implications. *Antioxid Redox Signal* 10(3):405–443
- Craeli W, Migdal MW, Luessenhop CP, Sugar A, Mihalakis I (1982) Iron deposits surrounding multiple sclerosis plaques. *Arch Pathol Lab Med* 106(8):397–399
- Crapper McLachlan DR, Dalton AJ, Kruck TP, Bell MY, Smith WL, Kalow W et al (1991) Intramuscular desferrioxamine in patients with Alzheimer's disease. *Lancet* 337(8753):1304–1308
- Crichton RR, Ward RJ (2006) Metal-based neurodegeneration. Wiley/Blackwell, Chichester
- Crimi E, Sica V, Slutsky AS, Zhang HB, Williams-Ignarro S, Ignarro LJ et al (2006a) Role of oxidative stress in experimental sepsis and multisystem organ dysfunction. *Free Radic Res* 40(7):665–672
- Crimi E, Sica V, Williams-Ignarro S, Zhang H, Slutsky AS, Ignarro LJ et al (2006b) The role of oxidative stress in adult critical care. *Free Radic Biol Med* 40(3):398–406
- Crozet C, Beranger F, Lehmann S (2008) Cellular pathogenesis in prion diseases. *Vet Res* 39(4):44
- Csermely P, Agoston V, Pongor S (2005) The efficiency of multi-target drugs: the network approach might help drug design. *Trends Pharmacol Sci* 26(4):178–182
- Csete M, Doyle J (2004) Bow ties, metabolism and disease. *Trends Biotechnol* 22(9):446–450
- Csikasz-Nagy A (2009) Computational systems biology of the cell cycle. *Brief Bioinform* 10(4):424–434
- Cuajungco MP, Fagét KY, Huang X, Tanzi RE, Bush AI (2000) Metal chelation as a potential therapy for Alzheimer's disease. *Ann NY Acad Sci* 920:292–304
- Cuijpers ML, Raymakers RA, Mackenzie MA, de Witte TJ, Swinkels DW (2010) Recent advances in the understanding of iron overload in sideroblastic myelodysplastic syndrome. *Br J Haematol* 149:322–333
- Cuzzocrea S, Di Paola R, Mazzon E, Patel NS, Genovese T, Muia C et al (2006) Erythropoietin reduces the development of nonseptic shock induced by zymosan in mice. *Crit Care Med* 34(4):1168–1177
- Cuzzocrea S, Riley DP, Caputi AP, Salvemini D (2001) Antioxidant therapy: a new pharmacological approach in shock, inflammation, and ischemia/reperfusion injury. *Pharmacol Rev* 53(1):135–159
- Cuzzocrea S, Thiemermann C, Salvemini D (2004) Potential therapeutic effect of antioxidant therapy in shock and inflammation. *Curr Med Chem* 11(9):1147–1162
- Dada JO, Mendes P (2009) Design and architecture of web services for simulation of biochemical systems. *Data Integr Life Sci Proc* 5647:182–195
- Dada JO, Spasic I, Paton NW, Mendes P (2010) SBRML: a markup language for associating systems biology data with models. *Bioinformatics*
- Daghino S, Turci F, Tomatis M, Favier A, Perotto S, Douki T et al (2006) Soil fungi reduce the iron content and the DNA damaging effects of asbestos fibers. *Environ Sci Technol* 40(18):5793–5798
- Dai J, Chung A (2001) Relationship of fiber surface iron and active oxygen species to expression of procollagen, PDGF-A, and TGF-beta(1) in tracheal explants exposed to amosite asbestos. *Am J Respir Cell Mol Biol* 24(4):427–435
- Dalle-Donne I (2007) Familial amyotrophic lateral sclerosis (FALS): emerging hints from redox proteomics. Highlight commentary on: "Redox proteomics analysis of oxidatively modified proteins in G93A-SOD1 transgenic mice—a model of familial amyotrophic lateral sclerosis". *Free Radic Biol Med* 43(2):157–159
- Dancey JE, Chen HX (2006) Strategies for optimizing combinations of molecularly targeted anticancer agents. *Nat Rev Drug Discov* 5(8):649–659
- Daré E, Götz ME, Zhivotovsky B, Manzo L, Ceccatelli S (2000) Antioxidants J811 and 17beta-estradiol protect cerebellar granule cells from methylmercury-induced apoptotic cell death. *J Neurosci Res* 62(4):557–565
- Dávalos A, Fernandezreal JM, Ricart W, Soler S, Molins A, Planas E et al (1994) Iron-related damage in acute ischemic stroke. *Stroke* 25(8):1543–1546
- Davidsson L, Galan P, Kastenmayer P, Cherouvrier F, Juillerat MA, Hercberg S et al (1994) Iron bioavailability studied in infants: the influence of phytic acid and ascorbic acid in infant formulas based on soy isolate. *Pediatr Res* 36(6):816–822
- Davies BW, Kohanski MA, Simmons LA, Winkler JA, Collins JJ, Walker GC (2009) Hydroxyurea induces hydroxyl radical-mediated cell death in *Escherichia coli*. *Mol Cell* 36(5):845–860
- Davies MG, Hagen PO (1997) Systemic inflammatory response syndrome. *Br J Surg* 84(7):920–935
- Davies MJ (2005) The oxidative environment and protein damage. *Biochim Biophys Acta* 1703(2):93–109
- Day SM, Duquaine D, Mundada LV, Menon RG, Khan BV, Rajagopalan S et al (2003) Chronic iron administration increases vascular oxidative stress and accelerates arterial thrombosis. *Circulation* 107(20):2601–2606
- de Carvalho CC, Fernandes P (2010) Production of metabolites as bacterial responses to the marine environment. *Mar Drugs* 8(3):705–727
- De Freitas J, Wintz H, Kim JH, Poynton H, Fox T, Vulpe C (2003) Yeast, a model organism for iron and copper metabolism studies. *Biometals* 16(1):185–197

- de Monyé C, Karcher DS, Boelaert JR, Gordeuk VR (1999) Bone marrow macrophage iron grade and survival of HIV-seropositive patients. *Aids* 13(3):375–380
- de Valk B, Marx JJM (1999) Iron, atherosclerosis, and ischemic heart disease. *Arch Internal Med* 159(14):1542–1548
- De Vizcaya-Ruiz A, Barbier O, Ruiz-Ramos R, Cebrian ME (2009) Biomarkers of oxidative stress and damage in human populations exposed to arsenic. *Mutat Res* 674(1–2):85–92
- Debebe Z, Ammosova T, Jerebtsova M, Kurantsin-Mills J, Niu X, Charles S et al (2007) Iron chelators ICL670 and 311 inhibit HIV-1 transcription. *Virology* 367(2):324–333
- Dekker MC, Giesbergen PC, Njajou OT, van Swieten JC, Hofman A, Breteler MM et al (2003) Mutations in the hemochromatosis gene (HFE), Parkinson's disease and parkinsonism. *Neurosci Lett* 348(2):117–119
- Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R et al (2008) Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 36(1):296–327
- Demir O, Singh S, Klimaschewski L, Kurnaz IA (2009) From birth till death: neurogenesis, cell cycle, and neurodegeneration. *Anat Rec (Hoboken)* 292(12):1953–1961
- Demougeot C, Van Hoecke M, Bertrand N, Prigent-Tessier A, Mossiat C, Beley A et al (2004) Cytoprotective efficacy and mechanisms of the liposoluble iron chelator 2, 2'-dipyridyl in the rat photothrombotic ischemic stroke model. *J Pharm Exp Therapeut* 311(3):1080–1087
- Dentchev T, Hahn P, Dunaief JL (2005) Strong labeling for iron and the iron-handling proteins ferritin and ferroportin in the photoreceptor layer in age-related macular degeneration. *Arch Ophthalmol* 123(12):1745–1746
- DePalma RG, Hayes VW, Chow BK, Shamayeva G, May PE, Zacharski LR (2010) Ferritin levels, inflammatory biomarkers, and mortality in peripheral arterial disease: a substudy of the iron (Fe) and atherosclerosis study (FeAST) Trial. *J Vasc Surg*
- DePalma RG, Hayes VW, Zacharski LR (2007) Bloodletting: past and present. *J Am Coll Surg* 205(1):132–144
- Dereure O, Jumez N, Bessis D, Gallix B, Guillot B (2008) Measurement of liver iron content by magnetic resonance imaging in 20 patients with overt porphyria cutanea tarda before phlebotomy therapy: a prospective study. *Acta Derm Venereol* 88(4):341–345
- Desai TK, Jamil LH, Balasubramaniam M, Koff R, Bonkovsky HL (2008) Phlebotomy improves therapeutic response to interferon in patients with chronic hepatitis C: a meta-analysis of six prospective randomized controlled trials. *Dig Dis Sci* 53(3):815–822
- Dever JB, Mallory MA, Mallory JE, Wallace D, Kowdley KV (2010) Phenotypic characteristics and diagnoses of patients referred to an iron overload clinic. *Dig Dis Sci* 55:803–807
- Devireddy LR, Gazin C, Zhu X, Green MR (2005) A cell-surface receptor for lipocalin 24p3 selectively mediates apoptosis and iron uptake. *Cell* 123(7):1293–1305
- Di Marco V, Capra M, Gagliardotto F, Borsellino Z, Cabibi D, Barbaria F et al (2008) Liver disease in chelated transfusion-dependent thalassemics: the role of iron overload and chronic hepatitis C. *Haematologica* 93(8):1243–1246
- Di Paola R, Mazzon E, Muia C, Crisafulli C, Genovese T, Di Bella P et al (2006) Green tea polyphenol extract attenuates zymosan-induced non-septic shock in mice. *Shock* 26(4):402–409
- Dickens MG, Franz KJ (2010) A prochelator activated by hydrogen peroxide prevents metal-induced amyloid beta aggregation. *Chembiochem* 11(1):59–62
- Diedrich JF, Bendheim PE, Kim YS, Carp RI, Haase AT (1991) Scrapie-associated prion protein accumulates in astrocytes during scrapie infection. *Proc Natl Acad Sci USA* 88(2):375–379
- DiMauro S, Schon EA (2008) Mitochondrial disorders in the nervous system. *Annu Rev Neurosci* 31:91–123
- Ding B, Chen KM, Ling HW, Sun F, Li X, Wan T et al (2009) Correlation of iron in the hippocampus with MMSE in patients with Alzheimer's disease. *J Magn Reson Imaging* 29(4):793–798
- Dobson P, Lanthaler K, Oliver SG, Kell DB (2009a) Implications of the dominant role of cellular transporters in drug uptake. *Curr Top Med Chem* 9:163–184
- Dobson PD, Patel Y, Kell DB (2009b) "Metabolite-likeness" as a criterion in the design and selection of pharmaceutical drug libraries. *Drug Disc Today* 14:31–40
- Dobson PD, Kell DB (2008) Carrier-mediated cellular uptake of pharmaceutical drugs: an exception or the rule? *Nat Rev Drug Discov* 7:205–220
- Dong J, Atwood CS, Anderson VE, Siedlak SL, Smith MA, Perry G et al (2003) Metal binding and oxidation of amyloid-beta within isolated senile plaque cores: Raman microscopic evidence. *Biochemistry* 42(10):2768–2773
- Doraiswamy PM, Finefrock AE (2004) Metals in our minds: therapeutic implications for neurodegenerative disorders. *Lancet Neurol* 3(7):431–434
- Doria E, Galleschi L, Calucci L, Pinzino C, Pilu R, Cassani E et al (2009) Phytic acid prevents oxidative stress in seeds: evidence from a maize (*Zea mays* L.) low phytic acid mutant. *J Exp Bot* 60(3):967–978
- Double KL, Dedov VN, Fedorow H, Kettle E, Halliday GM, Garner B et al (2008) The comparative biology of neuromelanin and lipofuscin in the human brain. *Cell Mol Life Sci* 65(11):1669–1682
- Double KL, Gerlach M, Schunemann V, Trautwein AX, Zecca L, Gallorini M et al (2003a) Iron-binding characteristics of neuromelanin of the human substantia nigra. *Biochem Pharmacol* 66(3):489–494
- Double KL, Halliday GM, Henderson J, Griffiths FM, Heinemann T, Riederer P et al (2003b) The dopamine receptor agonist lisuride attenuates iron-mediated dopaminergic neurodegeneration. *Exp Neurol* 184(1):530–535
- Double KL, Gerlach M, Youdim MB, Riederer P (2000) Impaired iron homeostasis in Parkinson's disease. *J Neural Transm Suppl* 60: 37–58
- Drakesmith H, Prentice A (2008) Viral infection and iron metabolism. *Nat Rev Microbiol* 6(7):541–552
- Drayer B, Burger P, Hurwitz B, Dawson D, Cain J (1987) Reduced signal intensity on MR images of thalamus and putamen in multiple sclerosis—increased iron content. *Am J Neuroradiol* 8(3):413–419
- Dreyfus F (2008) The deleterious effects of iron overload in patients with myelodysplastic syndromes. *Blood Rev* 22(Suppl 2):S29–S34
- Dröge W (2002) Free radicals in the physiological control of cell function. *Physiol Rev* 82(1):47–95
- Duarte NC, Becker SA, Jamshidi N, Thiele I, Mo ML, Vo TD et al (2007) Global reconstruction of the human metabolic network based on genomic and bibliomic data. *Proc Natl Acad Sci* 104(6):1777–1782
- Duwell P, Kono H, Rayner KJ, Sirois CM, Vladimer G, Bauernfeind FG et al (2010) NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 464(7293):1357–1361
- Duffy SJ, Biegelsen ES, Holbrook M, Russell JD, Gokce N, Keaney J F Jr et al (2001) Iron chelation improves endothelial function in patients with coronary artery disease. *Circulation* 103(23): 2799–2804
- Dunaief JL (2006) Iron induced oxidative damage as a potential factor in age-related macular degeneration: The Cogan Lecture. *Invest Ophthalmol Vis Sci* 47(11):4660–4664

- Durant R, Klouche K, Delbosch S, Morena M, Amigues L, Beraud JJ et al (2004) Superoxide anion overproduction in sepsis: effects of vitamin E and simvastatin. *Shock* 22(1):34–39
- Duvigneau JC, Piskernik C, Haindl S, Kloesch B, Hartl RT, Huttemann M et al (2008) A novel endotoxin-induced pathway: upregulation of heme oxygenase 1, accumulation of free iron, and free iron-mediated mitochondrial dysfunction. *Lab Invest* 88(1):70–77
- Dwyer BE, Zacharski LR, Balestra DJ, Lerner AJ, Perry G, Zhu X et al (2009a) Getting the iron out: phlebotomy for Alzheimer's disease? *Med Hypotheses* 72(5):504–509
- Dwyer DJ, Kohanski MA, Collins JJ (2009b) Role of reactive oxygen species in antibiotic action and resistance. *Curr Opin Microbiol* 12(5):482–489
- Dwyer DJ, Kohanski MA, Hayete B, Collins JJ (2007) Gyrase inhibitors induce an oxidative damage cellular death pathway in *Escherichia coli*. *Mol Syst Biol* 3:91
- Dwyer JH, Allayee H, Dwyer KM, Fan J, Wu H, Mar R et al (2004) Arachidonate 5-lipoxygenase promoter genotype, dietary arachidonic acid, and atherosclerosis. *N Engl J Med* 350(1):29–37
- Eissa A, Lebel RM, Korzan JR, Zavodni AE, Warren KG, Catz I et al (2009) Detecting lesions in multiple sclerosis at 4.7 tesla using phase susceptibility-weighting and T2-weighting. *J Magn Reson Imaging* 30(4):737–742
- Eklblom K, Hultdin J, Stegmayr B, Johansson I, Van Guelpen B, Hallmans G et al (2007) Iron stores and HFE genotypes are not related to increased risk of ischemic stroke. A prospective nested case-referent study. *Cerebrovasc Dis* 24(5):405–411
- Ekshyyan O, Aw TY (2004) Apoptosis: a key in neurodegenerative disorders. *Curr Neurovasc Res* 1(4):355–371
- El Tannir El Tayara N, Delatour B, Le Cudennec C, Guegan M, Volk A, Dhenain M (2006) Age-related evolution of amyloid burden, iron load, and MR relaxation times in a transgenic mouse model of Alzheimer's disease. *Neurobiol Dis* 22(1):199–208
- Ellervik C, Birgens H, Tybjaerg-Hansen A, Nordestgaard BG (2007) Hemochromatosis genotypes and risk of 31 disease endpoints: meta-analyses including 66, 000 cases and 226, 000 controls. *Hepatology* 46(4):1071–1080
- Epstein J, Sanderson IR, MacDonald TT (2009) Curcumin as a therapeutic agent: the evidence from in vitro, animal and human studies. *Br J Nutr*. doi:10.1017/S0007114509993667
- Equitani F, Fernandez-Real JM, Menichella G, Koch M, Calvani M, Nobili V et al (2008) Bloodletting ameliorates insulin sensitivity and secretion in parallel to reducing liver iron in carriers of HFE gene mutations. *Diabetes Care* 31(1):3–8
- Erkinjuntti T, Benavente O, Eliasziw M, Munoz DG, Sulkava R, Haltia M et al (1996) Diffuse vacuolization (spongiosis) and arteriolosclerosis in the frontal white matter occurs in vascular dementia. *Arch Neurol* 53(4):325–332
- Escames G, Acuna-Castroviejo D, López LC, Tan DX, Maldonado MD, Sanchez-Hidalgo M et al (2006) Pharmacological utility of melatonin in the treatment of septic shock: experimental and clinical evidence. *J Pharm Pharmacol* 58(9):1153–1165
- Evans JA (2008) Electronic publication and the narrowing of science and scholarship. *Science* 321(5887):395–399
- Exley C (2006) Aluminium and iron, but neither copper nor zinc, are key to the precipitation of beta-sheets of Aβ₄₂ in senile plaque cores in Alzheimer's disease. *J Alzheimers Dis* 10(2–3):173–177
- Exley C, Mamutse G, Korchazhkina O, Pye E, Strekopytov S, Polwart A et al (2006) Elevated urinary excretion of aluminium and iron in multiple sclerosis. *Mult Scler* 12(5):533–540
- Fábián I, Csordás V (2003) Metal ion catalyzed autoxidation reactions: Kinetics and mechanisms. *Adv Inorg Chem* 54:395–461
- Facchini FS (2002) The iron factor of aging: why do Americans age faster?. Fenestra Books, Tucson, AZ
- Facchini FS, Hua NW, Stoohs RA (2002) Effect of iron depletion in carbohydrate-intolerant patients with clinical evidence of non-alcoholic fatty liver disease. *Gastroenterology* 122(4):931–939
- Fakih S, Podinovskaia M, Kong X, Collins HL, Schaible UE, Hider RC (2008) Targeting the lysosome: fluorescent iron(III) chelators to selectively monitor endosomal/lysosomal labile iron pools. *J Med Chem* 51(15):4539–4552
- Fan QW, Knight ZA, Goldenberg DD, Yu W, Mostov KE, Stokoe D et al (2006) A dual PI3 kinase/mTOR inhibitor reveals emergent efficacy in glioma. *Cancer Cell* 9(5):341–349
- Fan QW, Specht KM, Zhang C, Goldenberg DD, Shokat KM, Weiss WA (2003) Combinatorial efficacy achieved through two-point blockade within a signaling pathway—a chemical genetic approach. *Cancer Res* 63(24):8930–8938
- Fantauzzi M, Pacella A, Atzei D, Gianfagna A, Andreozzi GB, Rossi A (2010) Combined use of X-ray photoelectron and Mössbauer spectroscopic techniques in the analytical characterization of iron oxidation state in amphibole asbestos. *Anal Bioanal Chem* 396(8):2889–2898
- Fargion S, Fracanzani AL, Rossini A, Borzio M, Riggio O, Belloni G et al (2002) Iron reduction and sustained response to interferon-alpha therapy in patients with chronic hepatitis C: results of an Italian multicenter randomized study. *Am J Gastroenterol* 97(5):1204–1210
- Fargion S, Mattioli M, Fracanzani AL, Sampietro M, Tavazzi D, Fociani P et al (2001) Hyperferritinemia, iron overload, and multiple metabolic alterations identify patients at risk for nonalcoholic steatohepatitis. *Am J Gastroenterol* 96(8):2448–2455
- Fasano M, Bergamasco B, Lopiano L (2006) Modifications of the iron-neuromelanin system in Parkinson's disease. *J Neurochem* 96(4):909–916
- Fato R, Bergamini C, Leoni S, Stocchi P, Lenaz G (2008) Generation of reactive oxygen species by mitochondrial complex I: implications in neurodegeneration. *Neurochem Res* 33(12):2487–2501
- Fattman CL, Tan RJ, Tobolewski JM, Oury TD (2006) Increased sensitivity to asbestos-induced lung injury in mice lacking extracellular superoxide dismutase. *Free Radic Biol Med* 40(4):601–607
- Faucheux BA, Martin ME, Beaumont C, Hauw JJ, Agid Y, Hirsch EC (2003) Neuromelanin associated redox-active iron is increased in the substantia nigra of patients with Parkinson's disease. *J Neurochem* 86(5):1142–1148
- Faux NG, Ritchie CW, Gunn A, Rembach A, Tsatsanis A, Bedo J et al (2010) PBT2 rapidly improves cognition in Alzheimer's disease: additional phase II analyses. *J Alzheimers Dis*
- Featherstone DE, Broadie K (2002) Wrestling with pleiotropy: genomic and topological analysis of the yeast gene expression network. *Bioessays* 24(3):267–274
- Fedorow H, Tribl F, Halliday G, Gerlach M, Riederer P, Double KL (2005) Neuromelanin in human dopamine neurons: comparison with peripheral melanins and relevance to Parkinson's disease. *Prog Neurobiol* 75(2):109–124
- Fell DA (1996) Understanding the control of metabolism. Portland Press, London
- Fernaes S, Halldin J, Bedecs K, Land T (2005a) Changed iron regulation in scrapie-infected neuroblastoma cells. *Brain Res Mol Brain Res* 133(2):266–273
- Fernaes S, Land T (2005) Increased iron-induced oxidative stress and toxicity in scrapie-infected neuroblastoma cells. *Neurosci Lett* 382(3):217–220
- Fernaes S, Reis K, Bedecs K, Land T (2005b) Increased susceptibility to oxidative stress in scrapie-infected neuroblastoma cells is associated with intracellular iron status. *Neurosci Lett* 389(3):133–136

- Fernandes de Godoy M, Takakura IT, Machado RD, Grassi LV, Nogueira PR (2007) Serum ferritin and obstructive coronary artery disease: angiographic correlation. *Arquivos Brasileiros De Cardiologia* 88(4):430–433
- Fernández-Real JM, Peñarroja G, Castro A, García-Bragado F, López-Bermejo A, Ricart W (2002) Blood letting in high-ferritin type 2 diabetes: effects on vascular reactivity. *Diabetes Care* 25(12):2249–2255
- Fernandez Y, Anglade F, Mitjavila S (2000) Paraquat and iron-dependent lipid peroxidation. NADPH versus NADPH-generating systems. *Biol Trace Elem Res* 74(3):191–201
- Ferrali M, Signorini C, Caciotti B, Sugherini L, Ciccoli L, Giachetti D et al (1997a) Protection against oxidative damage of erythrocyte membrane by the flavonoid quercetin and its relation to iron chelating activity. *FEBS Lett* 416(2):123–129
- Ferrali M, Signorini C, Sugherini L, Pompella A, Lodovici M, Caciotti B et al (1997b) Release of free, redox-active iron in the liver and DNA oxidative damage following phenylhydrazine intoxication. *Biochem Pharmacol* 53(11):1743–1751
- Ferrara DE, Taylor WR (2005) Iron chelation and vascular function: in search of the mechanisms. *Arterioscler Thromb Vasc Biol* 25(11):2235–2237
- Ferrara F, Ventura P, Vegetti A, Guido M, Abbati G, Corradini E et al (2009) Serum ferritin as a predictor of treatment outcome in patients with chronic hepatitis C. *Am J Gastroenterol* 104(3):605–616
- Ferro JM, Dávalos A (2006) Other neuroprotective therapies on trial in acute stroke. *Cerebrovasc Dis* 21(Suppl 2):127–130
- Ferrucci L, Semba RD, Guralnik JM, Ershler WB, Bandinelli S, Patel KV et al (2010) Proinflammatory state, hepcidin, and anemia in older persons. *Blood* 115(18):3810–3816
- Filippi M, Agosta F (2009) Closing the clinical-imaging gap in multiple sclerosis? Imaging iron deposition in deep gray matter. *J Neuroimaging* 19(1):1–2
- Finch CA, Bellotti V, Stray S, Lipschitz DA, Cook JD, Pippard MJ et al (1986) Plasma ferritin determination as a diagnostic tool. *West J Med* 145(5):657–663
- Fine SL (2005) Age-related macular degeneration 1969–2004: a 35-year personal perspective. *Am J Ophthalmol* 139(3):405–420
- Finefrock AE, Bush AI, Doraiswamy PM (2003) Current status of metals as therapeutic targets in Alzheimer's disease. *J Am Geriatr Soc* 51(8):1143–1148
- Finkel T (2005) Radical medicine: treating ageing to cure disease. *Nat Rev Mol Cell Biol* 6(12):971–976
- Firdaus WJ, Wytttenbach A, Giuliano P, Kretz-Remy C, Currie RW, Arrigo AP (2006) Huntingtin inclusion bodies are iron-dependent centers of oxidative events. *FEBS J* 273(23):5428–5441
- Flynn MR, Susi P (2010) Manganese, iron, and total particulate exposures to welders. *J Occup Environ Hyg* 7(2):115–126
- Ford ES, Cogswell ME (1999) Diabetes and serum ferritin concentration among U.S. adults. *Diabetes Care* 22(12):1978–1983
- Forouhi NG, Harding AH, Allison M, Sandhu MS, Welch A, Luben R et al (2007) Elevated serum ferritin levels predict new-onset type 2 diabetes: results from the EPIC-Norfolk prospective study. *Diabetologia* 50(5):949–956
- Forrester JS (2002) Prevention of plaque rupture: a new paradigm of therapy. *Ann Intern Med* 137(10):823–833
- Forrester JS (2004) Common ancestors: chronic progressive diseases have the same pathogenesis. *Clin Cardiol* 27(4):186–190
- Foster J, Goldmann W, Parnham D, Chong A, Hunter N (2001) Partial dissociation of PrP(Sc) deposition and vacuolation in the brains of scrapie and BSE experimentally affected goats. *J Gen Virol* 82(Pt 1):267–273
- Foury F (1997) Human genetic diseases: a cross-talk between man and yeast. *Gene* 195:1–10
- Foury F, Cazzalini O (1997) Deletion of the yeast homologue of the human gene associated with Friedreich's ataxia elicits iron accumulation in mitochondria. *FEBS Lett* 411:373–377
- Foury F, Talibi D (2001) Mitochondrial control of iron homeostasis. A genome wide analysis of gene expression in a yeast frataxin-deficient strain. *J Biol Chem* 276(11):7762–7768
- Fraenkel PG, Gibert Y, Holzheimer JL, Lattanzi VJ, Burnett SF, Dooley KA et al (2009) Transferrin-a modulates hepcidin expression in zebrafish embryos. *Blood* 113(12):2843–2850
- Franchini M, Targher G, Capra F, Montagnana M, Lippi G (2008) The effect of iron depletion on chronic hepatitis C virus infection. *Hepatol Internat* 2(3):335–340
- Franco JL, Posser T, Dunkley PR, Dickson PW, Mattos JJ, Martins R et al (2009a) Methylmercury neurotoxicity is associated with inhibition of the antioxidant enzyme glutathione peroxidase. *Free Radic Biol Med* 47(4):449–457
- Franco R, Sánchez-Olea R, Reyes-Reyes EM, Panayiotidis MI (2009b) Environmental toxicity, oxidative stress and apoptosis: manage a trois. *Mutat Res* 674(1–2):3–22
- Franco OH, Kirkwood TB, Powell JR, Catt M, Goodwin J, Ordovas JM et al (2007) Ten commandments for the future of ageing research in the UK: a vision for action. *BMC Geriatr* 7:10
- Franke JC, Plotz M, Prokop A, Geilen CC, Schmalz HG, Eberle J (2010) New caspase-independent but ROS-dependent apoptosis pathways are targeted in melanoma cells by an iron-containing cytosine analogue. *Biochem Pharmacol* 79(4):575–586
- Franklin MR, Phillips JD, Kushner JP (2005) Attenuation of polychlorinated biphenyl induced uroporphyrin by iron deprivation. *Environ Toxicol Pharmacol* 20(3):417–423
- Franks NP (2006) Molecular targets underlying general anaesthesia. *Br J Pharmacol* 147(Suppl 1):S72–S81
- Franks NP (2008) General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nat Rev Neurosci* 9(5):370–386
- Freixes M, Rodríguez A, Dalfó E, Ferrer I (2006) Oxidation, glycooxidation, lipoxidation, nitration, and responses to oxidative stress in the cerebral cortex in Creutzfeldt-Jakob disease. *Neurobiol Aging* 27(12):1807–1815
- Friedman A, Gałazka-Friedman J, Bauminger ER (2007) Iron as a trigger of neurodegeneration in Parkinson's disease. *Handb Clin Neurol* 83:493–505
- Friedman A, Galazka-Friedman J, Koziorowski D (2009) Iron as a cause of Parkinson disease—a myth or a well established hypothesis? *Parkinsonism Rel Disord* 15S3:S212–S214
- Fubini B, Arean CO (1999) Chemical aspects of the toxicity of inhaled mineral dusts. *Chem Soc Rev* 28(6):373–381
- Fuchs E, Gould E (2000) Mini-review: in vivo neurogenesis in the adult brain: regulation and functional implications. *Eur J Neurosci* 12(7):2211–2214
- Fujihara J, Agusa T, Tanaka J, Fujii Y, Moritani T, Hasegawa M et al (2009) 8-Hydroxy-2'-deoxyguanosine (8-OHdG) as a possible marker of arsenic poisoning: a clinical case study on the relationship between concentrations of 8-OHdG and each arsenic compound in urine of an acute promyelocytic leukemia patient being treated with arsenic trioxide. *Forensic Toxicol* 27(1):41–44
- Fujita N, Horiike S, Sugimoto R, Tanaka H, Iwasa M, Kobayashi Y et al (2007a) Hepatic oxidative DNA damage correlates with iron overload in chronic hepatitis C patients. *Free Radic Biol Med* 42(3):353–362
- Fujita N, Sugimoto R, Takeo M, Urawa N, Mifuji R, Tanaka H et al (2007b) Hepcidin expression in the liver: relatively low level in patients with chronic hepatitis C. *Mol Med* 13(1–2):97–104
- Fujita N, Sugimoto R, Urawa N, Tanaka H, Konishi M, Kobayashi Y et al (2007c) Influence of phlebotomy on iron-related gene

- expression levels in the livers of patients with chronic hepatitis C. *J Gastroenterol* 42(4):326–327
- Fujita N, Miyachi H, Tanaka H, Takeo M, Nakagawa N, Kobayashi Y et al (2009) Iron overload is associated with hepatic oxidative damage to DNA in nonalcoholic steatohepatitis. *Cancer Epidemiol Biomarkers Prev* 18(2):424–432
- Fujita N, Takei Y (2007) Iron, hepatitis C virus, and hepatocellular carcinoma: iron reduction preaches the gospel for chronic hepatitis C. *J Gastroenterol* 42(11):923–926
- Fukuchi K, Tomoyasu S, Tsuruoka N, Gomi K (1994) Iron deprivation-induced apoptosis in HL-60 cells. *FEBS Lett* 350(1):139–142
- Fulda S, Gorman AM, Hori O, Samali A (2010) Cellular stress responses: cell survival and cell death. *Int J Cell Biol* 2010:214074
- Gackowski D, Kruszewski M, Bartłomiejczyk T, Jawien A, Ciecierski M, Olinski R (2002) The level of 8-oxo-7, 8-dihydro-2'-deoxyguanosine is positively correlated with the size of the labile iron pool in human lymphocytes. *J Biol Inorg Chem* 7(4–5):548–550
- Gajda M, Banas K, Banas A, Jawien J, Mateuszuk L, Chlopicki S et al (2008) Distribution of selected elements in atherosclerotic plaques of apoE/LDLR-double knockout mice assessed by synchrotron radiation-induced micro-XRF spectrometry. *X-Ray Spectrom* 37(5):495–502
- Gajdusek DC (2008) Review. Kuru and its contribution to medicine. *Philos Trans R Soc Lond B Biol Sci* 363(1510):3697–3700
- Gakh O, Park S, Liu G, Macomber L, Imlay JA, Ferreira GC et al (2006) Mitochondrial iron detoxification is a primary function of frataxin that limits oxidative damage and preserves cell longevity. *Human Mol Genet* 15(3):467–479
- Gal S, Fridkin M, Amit T, Zheng H, Youdim MBH (2006) M30, a novel multifunctional neuroprotective drug with potent iron chelating and brain selective monoamine oxidase-ab inhibitory activity for Parkinson's disease. *J Neural Transm Suppl* 70:447–456
- Gal S, Zheng H, Fridkin M, Youdim MB (2009) Restoration of nigrostriatal dopamine neurons in post-MPTP treatment by the novel multifunctional brain-permeable iron chelator-monoamine oxidase inhibitor drug, M30. *Neurotox Res*
- Galaris D, Mantzaris M, Amorgianiotis C (2008a) Oxidative stress and aging: the potential role of iron. *Hormones (Athens)* 7(2):114–122
- Galaris D, Skiada V, Barbouti A (2008b) Redox signaling and cancer: the role of “labile” iron. *Cancer Lett* 266(1):21–29
- Galley HF, Davies MJ, Webster NR (1996) Ascorbyl radical formation in patients with sepsis: effect of ascorbate loading. *Free Radic Biol Med* 20(1):139–143
- Galley HF, Howdle PD, Walker BE, Webster NR (1997) The effects of intravenous antioxidants in patients with septic shock. *Free Radic Biol Med* 23(5):768–774
- Galley HF, Webster NR (1996) Elevated serum bleomycin-detectable iron concentrations in patients with sepsis syndrome. *Intensive Care Med* 22(3):226–229
- Galli SJ, Tsai M, Piliponsky AM (2008) The development of allergic inflammation. *Nature* 454(7203):445–454
- Gamberini MR, De Sanctis V, Gilli G (2008) Hypogonadism, diabetes mellitus, hypothyroidism, hypoparathyroidism: incidence and prevalence related to iron overload and chelation therapy in patients with thalassaemia major followed from 1980 to 2007 in the Ferrara Centre. *Pediatr Endocrinol Rev* 6(Suppl 1):158–169
- Gan-Or Z, Giladi N, Rozovski U, Shifrin C, Rosner S, Gurevich T et al (2008) Genotype-phenotype correlations between GBA mutations and Parkinson disease risk and onset. *Neurology* 70(24):2277–2283
- Gan EK, Ayonrinde OT, Trinder D, Olynyk JK (2010) Phenotypic expression of hereditary hemochromatosis: what have we learned from the population studies? *Curr Gastroenterol Rep* 12(1):7–12
- Gao F, Linhartova L, Johnston AM, Thickett DR (2008) Statins and sepsis. *Br J Anaesth* 100(3):288–298
- García-Castañeiras S (2010) Iron, the retina and the lens: A focused review. *Exp Eye Res* 90:664–678
- Garey M, Johnson D (1979) Computers and intractability: a guide to the theory of NP-completeness. Freeman, San Francisco
- Garoufi AJ, Prassouli AA, Attilakos AV, Voudris KA, Katsarou ES (2006) Homozygous MTHFR C677T gene mutation and recurrent stroke in an infant. *Pediatr Neurol* 35(1):49–51
- Gattermann N (2008) Overview of guidelines on iron chelation therapy in patients with myelodysplastic syndromes and transfusional iron overload. *Int J Hematol* 88(1):24–29
- Gattermann N (2009) The treatment of secondary hemochromatosis. *Dtsch Arztebl Int* 106(30): 499–504, I
- Gazzano E, Turci F, Foresti E, Putzu MG, Aldieri E, Silvagno F et al (2007) Iron-loaded synthetic chrysotile: a new model solid for studying the role of iron in asbestos toxicity. *Chem Res Toxicol* 20(3):380–387
- Ge Y, Jensen JH, Lu H, Helpem JA, Miles L, Inglese M et al (2007) Quantitative assessment of iron accumulation in the deep gray matter of multiple sclerosis by magnetic field correlation imaging. *AJNR Am J Neuroradiol* 28(9):1639–1644
- Gems D (2009) Ageing and oxidants in the nematode *Caenorhabditis elegans*. *SEB Exp Biol Ser* 62:31–56
- Gems D, Doonan R (2009) Antioxidant defense and aging in *C. elegans*: is the oxidative damage theory of aging wrong? *Cell Cycle* 8(11):1681–1687
- Gems D, Partridge L (2008) Stress-response hormesis and aging: “that which does not kill us makes us stronger”. *Cell Metab* 7(3):200–203
- Gennero L, Roos MA, D'Amelio P, Denysenko T, Morra E, Sperber K et al (2010) Iron metabolism markers and haptoglobin phenotypes in susceptibility to HSV-1 or/and HSV-2 lesion relapses. *Cell Biochem Funct* 28(2):142–148
- Georgiou NA, van der Bruggen T, Oudshoorn M, Nottet HH, Marx JJ, van Asbeck BS (2000) Combining iron chelators with the nucleoside analog didanosine in anti-HIV therapy. *Transfus Sci* 23(3):249–250
- Gerber S, Assmus H, Bakker B, Klipp E (2007) Drug-efficacy depends on the inhibitor type and the target position in a metabolic network—a systematic study. *J Theor Biol* 255:442–455
- Gerlach M, Benschachar D, Riederer P, Youdim MBH (1994) Altered brain metabolism of iron as a cause of neurodegenerative diseases. *J Neurochem* 63(3):793–807
- Gerlach M, Double KL, Ben-Shachar D, Zecca L, Youdim MB, Riederer P (2003) Neuromelanin and its interaction with iron as a potential risk factor for dopaminergic neurodegeneration underlying Parkinson's disease. *Neurotox Res* 5(1–2):35–44
- Gerlach M, Riederer P, Double KL (2008) Neuromelanin-bound ferric iron as an experimental model of dopaminergic neurodegeneration in Parkinson's disease. *Parkinsonism Relat Disord* 14(Suppl 2):S185–S188
- Geronikaki AA, Gavalas AM (2006) Antioxidants and inflammatory disease: synthetic and natural antioxidants with anti-inflammatory activity. *Comb Chem High Throughput Screen* 9(6):425–442
- Ghio A, Tan RJ, Ghio K, Fattman CL, Oury TD (2009) Iron accumulation and expression of iron-related proteins following murine exposure to crocidolite. *J Environ Pathol Toxicol Oncol* 28(2):153–162

- Ghio AJ, Carter JD, Richards JH, Richer LD, Grissom CK, Elstad MR (2003) Iron and iron-related proteins in the lower respiratory tract of patients with acute respiratory distress syndrome. *Crit Care Med* 31(2):395–400
- Ghio AJ, Churg A, Roggli VL (2004) Ferruginous bodies: implications in the mechanism of fiber and particle toxicity. *Toxicol Pathol* 32(6):643–649
- Ghosh B, Antonio T, Reith ME, Dutta AK (2010) Discovery of 4-(4-(2-((5-Hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)(propyl)amino)ethyl) piperazin-1-yl)quinolin-8-ol and its analogues as highly potent dopamine D2/D3 agonists and as iron chelator: in vivo activity indicates potential application in symptomatic and neuroprotective therapy for Parkinson's disease. *J Med Chem* 53(5):2114–2125
- Ghoshal N, Cali I, Perrin RJ, Josephson SA, Sun N, Gambetti P et al (2009) Codistribution of amyloid beta plaques and spongiform degeneration in familial Creutzfeldt-Jakob disease with the E200K–129M haplotype. *Arch Neurol* 66(10):1240–1246
- Ghribi O, Golovko MY, Larsen B, Schrag M, Murphy EJ (2006) Deposition of iron and beta-amyloid plaques is associated with cortical cellular damage in rabbits fed with long-term cholesterol-enriched diets. *J Neurochem* 99(2):438–449
- Gieseg SP, Leake DS, Flavall EM, Amit Z, Reid L, Yang YT (2009) Macrophage antioxidant protection within atherosclerotic plaques. *Frontiers Biosci* 14:1230–1246
- Gillum RF, Sempos CT, Makuc DM, Looker AC, Chien CY, Ingram DD (1996) Serum transferrin saturation, stroke incidence, and mortality in women and men. The NHANES I epidemiologic followup study. National health and nutrition examination survey. *Am J Epidemiol* 144(1):59–68
- Gilmour PS, Brown DM, Beswick PH, MacNee W, Rahman I, Donaldson K (1997) Free radical activity of industrial fibers: role of iron in oxidative stress and activation of transcription factors. *Environ Health Perspect* 105(Suppl 5):1313–1317
- Girelli D, Pasino M, Goodnough JB, Nemeth E, Guido M, Castagna A et al (2009) Reduced serum hepcidin levels in patients with chronic hepatitis C. *J Hepatol* 51(5):845–852
- Giustarini D, Dalle-Donne I, Tsikas D, Rossi R (2009) Oxidative stress and human diseases: origin, link, measurement, mechanisms, and biomarkers. *Crit Rev Clin Lab Sci* 46(5–6):241–281
- Glahn RP, Wortley GM, South PK, Miller DD (2002) Inhibition of iron uptake by phytic acid, tannic acid, and ZnCl₂: studies using an in vitro digestion/Caco-2 cell model. *J Agric Food Chem* 50(2):390–395
- Gloire G, Legrand-Poels S, Piette J (2006) NF-kappaB activation by reactive oxygen species: fifteen years later. *Biochem Pharmacol* 72(11):1493–1505
- Gloire G, Piette J (2009) Redox regulation of nuclear post-translational modifications during NF-kappaB activation. *Antioxid Redox Signal* 11(9):2209–2222
- Goel A, Dani V, Dhawan DK (2006) Role of zinc in mitigating the toxic effects of chlorpyrifos on hematological alterations and electron microscopic observations in rat blood. *Biometals* 19(5):483–492
- Goldberg DE (1989) Genetic algorithms in search, optimization and machine learning. Addison-Wesley, Reading, MA
- Goldberg DE (2002) The design of innovation: lessons from and for competent genetic algorithms. Kluwer, Boston
- Goldstein S, Merényi G (2008) The chemistry of peroxyntirite: implications for biological activity. *Methods Enzymol* 436:49–61
- Goldstein S, Meyerstein D, Czapski G (1993) The Fenton reagents. *Free Radic Biol Med* 15(4):435–445
- Goltsov A, Maryashkin A, Swat M, Kosinsky Y, Humphery-Smith I, Demin O et al (2009) Kinetic modelling of NSAID action on COX-1: focus on in vitro/in vivo aspects and drug combinations. *Eur J Pharm Sci* 36(1):122–136
- Gómez-Pinilla F (2008) The influences of diet and exercise on mental health through hormesis. *Ageing Res Rev* 7(1):49–62
- Gómez Ravetti M, Rosso OA, Berretta R, Moscato P (2010) Uncovering molecular biomarkers that correlate cognitive decline with the changes of hippocampus' gene expression profiles in Alzheimer's disease. *PLoSOne* 5(4):e10153
- Goncalves S, Paupe V, Dassa EP, Rustin P (2008) Deferiprone targets aconitase: implication for Friedreich's ataxia treatment. *BMC Neurol* 8:20
- González-Cabo P, Llorens JV, Palau F, Moltó MD (2009) Friedreich ataxia: an update on animal models, frataxin function and therapies. *Adv Exp Med Biol* 652:247–261
- González-Cabo P, Vázquez-Manrique RP, García-Gimeno MA, Sanz P, Palau F (2005) Frataxin interacts functionally with mitochondrial electron transport chain proteins. *Hum Mol Genet* 14(15):2091–2098
- Good PF, Hsu A, Werner P, Perl DP, Olanow CW (1998) Protein nitration in Parkinson's disease. *J Neuropathol Exp Neurol* 57(4):338–342
- Good PF, Perl DP, Bierer LM, Schmeidler J (1992) Selective accumulation of aluminum and iron in the neurofibrillary tangles of Alzheimer's disease: a laser microprobe (LAMMA) study. *Ann Neurol* 31(3):286–292
- Good PF, Werner P, Hsu A, Olanow CW, Perl DP (1996) Evidence of neuronal oxidative damage in Alzheimer's disease. *Am J Pathol* 149(1):21–28
- Goodall EF, Haque MS, Morrison KE (2008) Increased serum ferritin levels in amyotrophic lateral sclerosis (ALS) patients. *J Neurol* 255(11):1652–1656
- Goode HF, Webster NR (1993) Free radicals and antioxidants in sepsis. *Crit Care Med* 21(11):1770–1776
- Goralska M, Ferrell J, Harned J, Lall M, Nagar S, Fleisher LN et al (2009a) Iron metabolism in the eye: a review. *Exp Eye Res* 88(2):204–215
- Goralska M, Nagar S, Colitz CM, Fleisher LN, McGahan MC (2009b) Changes in ferritin H- and L-chains in canine lenses with age-related nuclear cataract. *Invest Ophthalmol Vis Sci* 50(1):305–310
- Goralska M, Nagar S, Fleisher LN, McGahan MC (2009c) Distribution of ferritin chains in canine lenses with and without age-related nuclear cataracts. *Mol Vis* 15:2404–2410
- Gordeuk VR, Delanghe JR, Langlois MR, Boelaert JR (2001) Iron status and the outcome of HIV infection: an overview. *J Clin Virol* 20(3):111–115
- Gordeuk VR, Onojobi G, Schneider MF, Dawkins FW, Delapenha R, Voloshin Y et al (2006) The association of serum ferritin and transferrin receptor concentrations with mortality in women with human immunodeficiency virus infection. *Haematologica* 91(6):739–743
- Goris RJ (1990) Mediators of multiple organ failure. *Intensive Care Med* 16(Suppl 3):S192–196
- Gorria M, Tekpli X, Rissel M, Sergent O, Huc L, Landvik N et al (2008) A new lactoferrin- and iron-dependent lysosomal death pathway is induced by benzo[a]pyrene in hepatic epithelial cells. *Toxicol Appl Pharmacol* 228(2):212–224
- Götz M, Huttner WB (2005) The cell biology of neurogenesis. *Nat Rev Mol Cell Biol* 6(10):777–788
- Gotz ME, Double K, Gerlach M, Youdim MBH, Riederer P (2004) The relevance of iron in the pathogenesis of Parkinson's disease. *Redox-Active Metals Neurol Disord* 1012:193–208
- Gould E (2007) How widespread is adult neurogenesis in mammals? *Nat Rev Neurosci* 8(6):481–488
- Governa M, Amati M, Fontana S, Visona I, Botta GC, Mollo F et al (1999) Role of iron in asbestos-body-induced oxidant radical generation. *J Toxicol Environ Health A* 58(5):279–287
- Gozzelino R, Jeney V, Soares MP (2010) Mechanisms of cell protection by heme oxygenase-1. *Annu Rev Pharmacol Toxicol* 50:323–354

- Graf E, Eaton JW (1990) Antioxidant functions of phytic acid. *Free Radic Biol Med* 8(1):61–69
- Graf E, Eaton JW (1993) Suppression of colonic cancer by dietary phytic acid. *Nutr Cancer* 19(1):11–19
- Graf E, Empson KL, Eaton JW (1987) Phytic acid. A natural antioxidant. *J Biol Chem* 262(24):11647–11650
- Graf E, Mahoney JR, Bryant RG, Eaton JW (1984) Iron-catalyzed hydroxyl radical formation. Stringent requirement for free iron coordination site. *J Biol Chem* 259(6):3620–3624
- Grainger DJ (2007) TGF-beta and atherosclerosis in man. *Cardiovasc Res* 74(2):213–222
- Granic I, Dolga AM, Nijholt IM, van Dijk G, Eisel ULM (2009) Inflammation and NF-kappaB in Alzheimer's disease and diabetes. *J Alzheimers Dis* 16(4):809–821
- Grases F, Simonet RM, Prieto RM, March JG (2001) Dietary phytate and mineral bioavailability. *J Trace Elem Med Biol* 15(4):221–228
- Grasshoff C, Drexler B, Rudolph U, Antkowiak B (2006) Anaesthetic drugs: linking molecular actions to clinical effects. *Curr Pharm Des* 12(28):3665–3679
- Greenberg PL (2006) Myelodysplastic syndromes: iron overload consequences and current chelating therapies. *J Natl Compr Canc Netw* 4(1):91–96
- Greenberg PL, Attar E, Battiwalla M, Bennett JM, Bloomfield CD, DeCastro CM et al (2008) Myelodysplastic syndromes. *J Natl Compr Canc Netw* 6(9):902–926
- Greene BT, Thorburn J, Willingham MC, Thorburn A, Planalp RP, Brechbiel MW et al (2002) Activation of caspase pathways during iron chelator-mediated apoptosis. *J Biol Chem* 277(28):25568–25575
- Greene LS (1995) Asthma and oxidant stress—nutritional, environmental, and genetic risk factors. *J Am Coll Nutr* 14(4):317–324
- Grimaud J, Millar J, Thorpe JW, Moseley IF, McDonald WI, Miller DH (1995) Signal intensity on MRI of basal ganglia in multiple sclerosis. *J Neurol Neurosurg Psychiatry* 59(3):306–308
- Grootveld M, Halliwell B (1986) Aromatic hydroxylation as a potential measure of hydroxyl-radical formation in vivo. Identification of hydroxylated derivatives of salicylate in human body fluids. *Biochem J* 237(2):499–504
- Grossi C, Francese S, Casini A, Rosi MC, Luccarini I, Fiorentini A et al (2009) Clioquinol decreases amyloid-beta burden and reduces working memory impairment in a transgenic mouse model of Alzheimer's disease. *J Alzheimers Dis* 96(3):423–440
- Gu Y, Hua Y, Keep RF, Morgenstern LB, Xi G (2009) Deferoxamine reduces intracerebral hematoma-induced iron accumulation and neuronal death in piglets. *Stroke* 40(6):2241–2243
- Gudmundsdóttir KB, Sigurdarson S, Kristinsson J, Eiríksson T, Jóhannesson T (2006) Iron and iron/manganese ratio in forage from Icelandic sheep farms: relation to scrapie. *Acta Vet Scand* 48:16
- Gulumian M, van Wyk JA (1987) Hydroxyl radical production in the presence of fibres by a Fenton-type reaction. *Chem Biol Interact* 62(1):89–97
- Guo L, Morris DG, Liu XY, Vaslet C, Hurt RH, Kane AB (2007) Iron bioavailability and redox activity in diverse carbon nanotube samples. *Chem Mater* 19(14):3472–3478
- Gupta RC, Milatovic D (2009) Toxicants and neurodegenerative diseases special issue. *Toxicol Appl Pharmacol* 240(2):123
- Gürkan E, Ergun Y, Zorludemir S, Başlamışlı F, Koçak R (2005) Liver involvement in sickle cell disease. *Turk J Gastroenterol* 16(4):194–198
- Gusarov I, Shatalin K, Starodubtseva M, Nudler E (2009) Endogenous nitric oxide protects bacteria against a wide spectrum of antibiotics. *Science* 325(5946):1380–1384
- Gusella JF, MacDonald ME (2000) Unmasking polyglutamine triggers in neurodegenerative disease. *Nat Rev Neurosci* 1(2):109–115
- Gutierrez JM, Rucavado A, Escalante T, Lomonte B, Angulo Y, Fox JW (2010) Tissue pathology induced by snake venoms: how to understand a complex pattern of alterations from a systems biology perspective. *Toxicol* 55(1):166–170
- Gutteridge JMC, Mitchell J (1999) Redox imbalance in the critically ill. *Br Med Bull* 55(1):49–75
- Guyader D, Thirouard AS, Erdtmann L, Rakba N, Jacquelinet S, Danielou H et al (2007) Liver iron is a surrogate marker of severe fibrosis in chronic hepatitis C. *J Hepatol* 46(4):587–595
- Ha YM, Chung SW, Kim JM, Kim DH, Kim JY, Lee EK et al (2010) Molecular activation of NF-kappaB, pro-inflammatory mediators, and signal pathways in gamma-irradiated mice. *Biotechnol Lett* 32(3):373–378
- Haacke EM, Cheng NY, House MJ, Liu Q, Neelavalli J, Ogg RJ et al (2005) Imaging iron stores in the brain using magnetic resonance imaging. *Magn Reson Imaging* 23(1):1–25
- Haacke EM, Garbern J, Miao Y, Habib C, Liu M (2010) Iron stores and cerebral veins in MS studied by susceptibility weighted imaging. *Int Angiol* 29(2):149–157
- Haacke EM, Makki M, Ge Y, Maheshwari M, Sehgal V, Hu J et al (2009) Characterizing iron deposition in multiple sclerosis lesions using susceptibility weighted imaging. *J Magn Reson Imaging* 29(3):537–544
- Haas H, Eisendle M, Turgeon BG (2008) Siderophores in fungal physiology and virulence. *Annu Rev Phytopathol* 46:149–187
- Hadacek F, Bachmann G, Engelmeier D, Chobot V (2010) Hormesis and a chemical raison d'être for secondary plant metabolites. Dose-response (in press)
- Haddad JJ, Olver RE, Land SC (2000) Antioxidant/pro-oxidant equilibrium regulates HIF-1alpha and NF-kappa B redox sensitivity. Evidence for inhibition by glutathione oxidation in alveolar epithelial cells. *J Biol Chem* 275(28):21130–21139
- Hadley D, Orlin A, Brown G, Brucker AJ, Ho AC, Regillo CD et al (2010) Analysis of six genetic risk factors highly associated with AMD in the region surrounding ARMS2 and HTRA1 on chromosome 10, region q26. *Invest Ophthalmol Vis Sci* 51(4):2191–2196
- Hagen J (2006) *Industrial Catalysis: A Practical Approach*. Wiley VCH Chemie, Weinheim
- Haggarty SJ, Clemons PA, Schreiber SL (2003) Chemical genomic profiling of biological networks using graph theory and combinations of small molecule perturbations. *J Am Chem Soc* 125(35):10543–10545
- Hague T, Andrews PLR, Barker J, Naughton DP (2006) Dietary chelators as antioxidant enzyme mimetics: implications for dietary intervention in neurodegenerative diseases. *Behav Pharmacol* 17(5–6):425–430
- Hahn P, Milam AH, Dunaief JL (2003) Maculas affected by age-related macular degeneration contain increased chelatable iron in the retinal pigment epithelium and Bruch's membrane. *Arch Ophthalmol* 121(8):1099–1105
- Hahn P, Qian Y, Dentchev T, Chen L, Beard J, Harris ZL et al (2004) Disruption of ceruloplasmin and hephaestin in mice causes retinal iron overload and retinal degeneration with features of age-related macular degeneration. *Proc Natl Acad Sci U S A* 101(38):13850–13855
- Hallberg L, Brune M, Rossander L (1989) Iron absorption in man: ascorbic acid and dose-dependent inhibition by phytate. *Am J Clin Nutr* 49(1):140–144
- Halliwell B (2007) Dietary polyphenols: good, bad, or indifferent for your health? *Cardiovasc Res* 73(2):341–347
- Halliwell B (2009) The wanderings of a free radical. *Free Radic Biol Med* 46(5):531–542
- Halliwell B, Gutteridge JMC (2006) *Free radicals in biology and medicine*, 4th edn. Oxford University Press, Oxford
- Halliwell B, Rafter J, Jenner A (2005) Health promotion by flavonoids, tocopherols, tocotrienols, and other phenols: direct

- or indirect effects? Antioxidant or not? *Am J Clin Nutr* 81(1 Suppl):268S–276S
- Hammond KE, Metcalf M, Carvajal L, Okuda DT, Srinivasan R, Vigneron D et al (2008) Quantitative in vivo magnetic resonance imaging of multiple sclerosis at 7 Tesla with sensitivity to iron. *Ann Neurol* 64(6):707–713
- Han JY, Takeshita K, Utsumi H (2001) Noninvasive detection of hydroxyl radical generation in lung by diesel exhaust particles. *Free Radic Biol Med* 30(5):516–525
- Han O, Failla ML, Hill AD, Morris ER, Smith J C Jr (1994) Inositol phosphates inhibit uptake and transport of iron and zinc by a human intestinal cell line. *J Nutr* 124(4):580–587
- Handl J, Kell DB, Knowles J (2007) Multiobjective optimization in bioinformatics and computational biology. *IEEE Trans Comput Biol Bioinform* 4(2):279–292
- Hanson L, Roeytenberg A, Martinez PM, Coppes VG, Sweet DC, Rao RJ et al (2009) Intranasal deferoxamine provides increased brain exposure and significant protection in rat ischemic stroke. *J Pharmacol Exp Ther* 330:679–686
- Hanson LN, Engelman HM, Alekel DL, Schalinske KL, Kohut ML, Reddy MB (2006) Effects of soy isoflavones and phytate on homocysteine, C-reactive protein, and iron status in postmenopausal women. *Am J Clin Nutr* 84(4):774–780
- Hansson GK (2001) Immune mechanisms in atherosclerosis. *Arterioscler Thromb Vasc Biol* 21(12):1876–1890
- Hansson GK (2005) Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 352(16):1685–1695
- Haq RU, Wereley JP, Chitambar CR (1995) Induction of apoptosis by iron deprivation in human leukemic CCRF-CEM cells. *Exp Hematol* 23(5):428–432
- Hardy JA, Aust AE (1995a) The effect of iron binding on the ability of crocidolite asbestos to catalyze DNA single-strand breaks. *Carcinogenesis* 16(2):319–325
- Hardy JA, Aust AE (1995b) Iron in asbestos chemistry and carcinogenicity. *Chem Rev* 95(1):97–118
- Harvey PT (2003) Common eye diseases of elderly people: identifying and treating causes of vision loss. *Gerontology* 49(1):1–11
- Hasinoff BB, Rahimtula AD, Omar RF (1990) NADPH-cytochrome-P-450 reductase promoted hydroxyl radical production by the iron(III)-ochratoxin A complex. *Biochim Biophys Acta* 1036(1):78–81
- Hassoun EA, Wang X (1999) Time- and concentration-dependent production of superoxide anion, nitric oxide, DNA damage and cellular death by ricin in the J774A.1 macrophage cells. *J Biochem Mol Toxicol* 13(3–4):179–185
- Hastie T, Tibshirani R, Friedman J (2001) The elements of statistical learning: data mining, inference and prediction. Springer, Berlin
- Hattori N, Oda S, Sadahiro T, Nakamura M, Abe R, Shinozaki K et al (2009) YKL-40 identified by proteomic analysis as a biomarker of sepsis. *Shock* 32(4):393–400
- Haugen AC, Di Prospero NA, Parker JS, Fannin RD, Chou J, Meyer JN et al (2010) Altered gene expression and DNA damage in peripheral blood cells from Friedreich's ataxia patients: cellular model of pathology. *PLoS Genet* 6(1):e1000812
- Hawkins PT, Poyner DR, Jackson TR, Letcher AJ, Lander DA, Irvine RF (1993) Inhibition of iron-catalysed hydroxyl radical formation by inositol polyphosphates: a possible physiological function for myo-inositol hexakisphosphate. *Biochem J* 294(Pt 3):929–934
- Hayashi H, Wakusawa S, Yano M (2006) Iron removal by phlebotomy for the prophylaxis of fulminant hepatitis in a Wilson disease model of Long-Evans Cinnamon Rats. *Hepatol Res* 35(4):276–280
- Hayashi H, Yano M (2002) Iron cytotoxicity in chronic hepatitis C. *J Health Sci* 48(3):227–231
- Hayashi T, Ueno Y, Okamoto T (1993) Oxidoreductive regulation of nuclear factor kappa B. Involvement of a cellular reducing catalyst thioredoxin. *J Biol Chem* 268(15):11380–11388
- Hayes DP (2009) Nutritional hormesis and aging. *Dose Response* 8(1):10–15
- He X, Hahn P, Iacovelli J, Wong R, King C, Bhisitkul R et al (2007) Iron homeostasis and toxicity in retinal degeneration. *Prog Retin Eye Res* 26(6):649–673
- He X, Yablonskiy DA (2009) Biophysical mechanisms of phase contrast in gradient echo MRI. *Proc Natl Acad Sci USA* 106(32):13558–13563
- Heathcote EJ (2004) Prevention of hepatitis C virus-related hepatocellular carcinoma. *Gastroenterology* 127(5 Suppl 1):S294–S302
- Heckl S (2007) Future contrast agents for molecular imaging in stroke. *Curr Med Chem* 14(16):1713–1728
- Hegde ML, Bharathi P, Suram A, Venugopal C, Jagannathan R, Poddar P et al (2009) Challenges associated with metal chelation therapy in Alzheimer's disease. *J Alzheimers Dis* 17(3):457–468
- Hegreness M, Shores N, Damian D, Hartl D, Kishony R (2008) Accelerated evolution of resistance in multidrug environments. *Proc Natl Acad Sci USA* 105(37):13977–13981
- Heinrich R, Schuster S (1996) The regulation of cellular systems. Chapman & Hall, New York
- Heintz NH, Janssen-Heininger YM, Mossman BT (2010) Asbestos, lung cancers, and mesotheliomas: from molecular approaches to targeting tumor survival pathways. *Am J Respir Cell Mol Biol* 42(2):133–139
- Helal GK (2008) Systemic administration of Zn²⁺ during the reperfusion phase of transient cerebral ischaemia protects rat hippocampus against iron-catalysed postischaemic injury. *Clin Exp Pharmacol Physiol* 35(7):775–781
- Henshall TL, Tucker B, Lumsden AL, Nornes S, Lardelli MT, Richards RI (2009) Selective neuronal requirement for huntingtin in the developing zebrafish. *Hum Mol Genet* 18(24):4830–4842
- Herce-Pagliai C, Kotecha S, Shuker DEG (1998) Analytical methods for 3-nitrotyrosine as a marker of exposure to reactive nitrogen species: a review. *Nitric Oxide-Biol Chem* 2(5):324–336
- Herrgård MJ, Swainston N, Dobson P, Dunn WB, Arga KY, Arvas M et al (2008) A consensus yeast metabolic network obtained from a community approach to systems biology. *Nat Biotechnol* 26(10):1155–1160
- Hershko C, Weatherall DJ (1988) Iron-chelating therapy. *Crit Rev Clin Lab Sci* 26(4):303–345
- Hider RC, Liu ZD, Khodr HH (2001) Metal chelation of polyphenols. *Flavonoids Other Polyphenols* 335:190–203
- Hider RC, Ma Y, Molina-Holgado F, Gaeta A, Roy S (2008) Iron chelation as a potential therapy for neurodegenerative disease. *Biochem Soc Trans* 36(Pt 6):1304–1308
- Hilet D, Panayiotidis P, Hoffbrand AV (1995) Iron chelators induce apoptosis in proliferating cells. *Br J Haematol* 89(1):181–187
- Hill RH Jr, Todd GD, Kilbourne EM, Cline RE, McCraw J, Orti DL et al (1987) Gas chromatographic/mass spectrometric determination of aniline in food oils associated with the Spanish toxic oil syndrome. *Bull Environ Contam Toxicol* 39(3):511–515
- Hillmer AS, Putcha P, Levin J, Hogen T, Hyman BT, Kretzschmar H et al (2009) Converse modulation of toxic alpha-synuclein oligomers in living cells by N'-benzylidene-benzohydrazide derivatives and ferric iron. *Biochem Biophys Res Commun*
- Himmelfarb J, Stenvinkel P, Ikizler TA, Hakim RM (2002) The elephant in uremia: oxidant stress as a unifying concept of cardiovascular disease in uremia. *Kidney Int* 62(5):1524–1538
- Hininger I, Waters R, Osman M, Garrel C, Fernholz K, Roussel AM et al (2005) Acute prooxidant effects of vitamin C in EDTA chelation therapy and long-term antioxidant benefits of therapy. *Free Rad Biol Med* 38(12):1565–1570

- Hinson JA, Reid AB, McCullough SS, James LP (2004) Acetaminophen-induced hepatotoxicity: role of metabolic activation, reactive oxygen/nitrogen species, and mitochondrial permeability transition. *Drug Metab Rev* 36(3–4):805–822
- Hintze KJ, Theil EC (2006) Cellular regulation and molecular interactions of the ferritins. *Cell Mol Life Sci* 63(5):591–600
- Hippeli S, Elstner EF (1999) Transition metal ion-catalyzed oxygen activation during pathogenic processes. *FEBS Lett* 443(1):1–7
- Hirsch EC (2009) Iron transport in Parkinson's disease. *Parkinsonism Rel Disord* 15:S209–S211
- Hirsch EC, Faucheux BA (1998) Iron metabolism and Parkinson's disease. *Mov Disord* 13(Suppl 1):39–45
- Hoefnagel MH, Starrenburg MJ, Martens DE, Hugenholtz J, Kleerebezem M, Van S II et al (2002) Metabolic engineering of lactic acid bacteria, the combined approach: kinetic modeling, metabolic control and experimental analysis. *Microbiology* 148(Pt 4):1003–1013
- Hoffmann GR (2009) A perspective on the scientific, philosophical, and policy dimensions of hormesis. *Dose Response* 7(1):1–51
- Honda K, Casadesus G, Petersen RB, Perry G, Smith MA (2004) Oxidative stress and redox-active iron in Alzheimer's disease. *Ann NY Acad Sci* 1012:179–182
- Hoops S, Sahle S, Gauges R, Lee C, Pahle J, Simus N et al (2006) COPASI: a COMplex PATHway SIMulator. *Bioinformatics* 22(24):3067–3074
- Hopkins AL, Mason JS, Overington JP (2006) Can we rationally design promiscuous drugs? *Curr Opin Struct Biol* 16(1):127–136
- Hori A, Mizoue T, Kasai H, Kawai K, Matsushita Y, Nanri A et al (2010) Body iron store as a predictor of oxidative DNA damage in healthy men and women. *Cancer Sci* 101(2):517–522
- Horn KD (1998) Evolving strategies in the treatment of sepsis and systemic inflammatory response syndrome (SIRS). *QJM* 91(4):265–277
- Horváth I, Donnelly LE, Kiss A, Paredi P, Kharitonov SA, Barnes PJ (1998) Raised levels of exhaled carbon monoxide are associated with an increased expression of heme oxygenase-1 in airway macrophages in asthma: a new marker of oxidative stress. *Thorax* 53(8):668–672
- Horwitz LD, Rosenthal EA (1999) Iron-mediated cardiovascular injury. *Vasc Med* 4(2):93–99
- Horwitz LD, Sherman NA, Kong YN, Pike AW, Gobin J, Fennessey PV et al (1998) Lipophilic siderophores *Mycobacterium tuberculosis* prevent cardiac reperfusion injury. *Proc Natl Acad Sci* 95(9):5263–5268
- Hotchkiss RS, Karl IE (2003) The pathophysiology and treatment of sepsis. *N Engl J Med* 348(2):138–150
- Howes PS, Zacharski LR, Sullivan J, Chow B (2000) Role of stored iron in atherosclerosis. *J Vasc Nurs* 18(4):109–114 (quiz 115–106)
- Hsieh SR, Tsai DC, Chen JY, Tsai SW, Liou YM (2009) Green tea extract protects rats against myocardial infarction associated with left anterior descending coronary artery ligation. *PLugers Arch* 458(4):631–642
- Hua NW, Stoohs RA, Facchini FS (2001) Low iron status and enhanced insulin sensitivity in lacto-ovo vegetarians. *Br J Nutr* 86(4):515–519
- Hua Y, Keep RF, Hoff JT, Xi G (2008) Deferoxamine therapy for intracerebral hemorrhage. *Acta Neurochir Suppl* 105:3–6
- Huang J, Dizin E, Cowan JA (2008a) Mapping iron binding sites on human frataxin: implications for cluster assembly on the ISU Fe-S cluster scaffold protein. *J Biol Inorg Chem* 13(5):825–836
- Huang XJ, Li CT, Zhang WP, Lu YB, Fang SH, Wei EQ (2008b) Dihydroartemisinin potentiates the cytotoxic effect of temozolomide in rat C6 glioma cells. *Pharmacology* 82(1):1–9
- Huang KJ, Su IJ, Theron M, Wu YC, Lai SK, Liu CC et al (2005) An interferon-gamma-related cytokine storm in SARS patients. *J Med Virol* 75(2):185–194
- Huang ML, Becker EM, Whitnall M, Rahmanto YS, Ponka P, Richardson DR (2009) Elucidation of the mechanism of mitochondrial iron loading in Friedreich's ataxia by analysis of a mouse mutant. *Proc Natl Acad Sci USA* 106(38):16381–16386
- Hubel CA, Bodnar LM, Many A, Harger G, Ness RB, Roberts JM (2004) Nonglycosylated ferritin predominates in the circulation of women with preeclampsia but not intrauterine growth restriction. *Clin Chem* 50(5):948–951
- Hucka M, Finney A, Sauro HM, Bolouri H, Doyle JC, Kitano H et al (2003) The systems biology markup language (SBML): a medium for representation and exchange of biochemical network models. *Bioinformatics* 19(4):524–531
- Hull D, Pettifer SR, Kell DB (2008) Defrosting the digital library: bibliographic tools for the next generation web. *PLoS Comput Biol* 4(10):e1000204. doi:1000210.1001371/journal.pcbi.1000204
- Hur K, Kim JI, Choi SI, Choi EK, Carp RI, Kim YS (2002) The pathogenic mechanisms of prion diseases. *Mech Ageing Dev* 123(12):1637–1647
- Hurn PD, Koehler RC, Blizzard KK, Traystman RJ (1995) Deferoxamine reduces early metabolic failure associated with severe cerebral ischemic acidosis in dogs. *Stroke* 26(4):688–694 (discussion 694–685)
- Hwang D, Lee IY, Yoo H, Gehlenborg N, Cho JH, Petritis B et al (2009) A systems approach to prion disease. *Mol Syst Biol* 5:252
- Iannetti A, Pacifico F, Acquaviva R, Lavorgna A, Crescenzi E, Vascotto C et al. (2008) The neutrophil gelatinase-associated lipocalin (NGAL), a NF- κ B-regulated gene, is a survival factor for thyroid neoplastic cells. *Proc Natl Acad Sci USA* 105:14058–14063
- Ilbäck NG, Lindh U, Wesslén L, Fohlman J, Friman G (2000) Trace element distribution in heart tissue sections studied by nuclear microscopy is changed in Cocksackie virus B3 myocarditis in methyl mercury-exposed mice. *Biol Trace Elem Res* 78(1–3):131–147
- Imarisio S, Carmichael J, Korolchuk V, Chen CW, Saiki S, Rose C et al (2008) Huntington's disease: from pathology and genetics to potential therapies. *Biochem J* 412(2):191–209
- Imayoshi I, Sakamoto M, Ohtsuka T, Kageyama R (2009) Continuous neurogenesis in the adult brain. *Dev Growth Differ* 51(3):379–386
- Imlay JA (2008) Cellular defenses against superoxide and hydrogen peroxide. *Annu Rev Biochem* 77:755–776
- Imrie CE, Korre A, Munoz-Melendez G (2009) Spatial correlation between the prevalence of transmissible spongiform diseases and British soil geochemistry. *Environ Geochem Health* 31(1):133–145
- Inati A, Musallam KM, Wood JC, Taher AT (2010) Iron overload indices rise linearly with transfusion rate in patients with sickle cell disease. *Blood* 115(14): 2980–2981; author reply 2981–2982
- Ioannidis JP (2005) Why most published research findings are false. *PLoS Med* 2(8):e124
- Ioannidis JP (2008) Why most discovered true associations are inflated. *Epidemiology* 19(5):640–648
- Iqbal M, Okazaki Y, Okada S (2009) Curcumin attenuates oxidative damage in animals treated with a renal carcinogen, ferric nitrilotriacetate (Fe-NTA): implications for cancer prevention. *Mol Cell Biochem* 324(1–2):157–164
- Irazusta V, Cabisco E, Reverter-Branchat G, Ros J, Tamarit J (2006) Manganese is the link between frataxin and iron-sulfur deficiency in the yeast model of Friedreich ataxia. *J Biol Chem* 281(18):12227–12232
- Irazusta V, Moreno-Cermeno A, Cabisco E, Ros J, Tamarit J (2008) Major targets of iron-induced protein oxidative damage in frataxin-deficient yeasts are magnesium-binding proteins. *Free Radic Biol Med*

- Ishizaka N, Saito K, Mori I, Matsuzaki G, Ohno M, Nagai R (2005) Iron chelation suppresses ferritin upregulation and attenuates vascular dysfunction in the aorta of angiotensin II-infused rats. *Arterioscler Thromb Vasc Biol* 25(11):2282–2288
- Iwamaru Y, Shimizu Y, Imamura M, Murayama Y, Endo R, Tagawa Y et al (2008) Lactoferrin induces cell surface retention of prion protein and inhibits prion accumulation. *J Neurochem* 107(3):636–646
- Jabbour E, Garcia-Manero G, Taher A, Kantarjian HM (2009) Managing iron overload in patients with myelodysplastic syndromes with oral deferasirox therapy. *Oncologist* 14(5):489–496
- Jacob AK, Hotchkiss RS, DeMeester SL, Hiramatsu M, Karl IE, Swanson PE et al. (1997) Endothelial cell apoptosis is accelerated by inorganic iron and heat via an oxygen radical dependent mechanism. *Surgery* 122(2):243–253 (discussion 254)
- Jacob AK, Hotchkiss RS, Swanson PE, Tinsley KW, Karl IE, Buchman TG (2000) Injection of iron compounds followed by induction of the stress response causes tissue injury and apoptosis. *Shock* 14(4):460–464
- Jädersten M, Hellström-Lindberg E (2010) New clues to the molecular pathogenesis of myelodysplastic syndromes. *Exp Cell Res*
- Jaeschke H, Bajt ML (2006) Intracellular signaling mechanisms of acetaminophen-induced liver cell death. *Toxicol Sci* 89(1):31–41
- James LP, Mayeux PR, Hinson JA (2003) Acetaminophen-induced hepatotoxicity. *Drug Metab Dispos* 31(12):1499–1506
- Janson J, Laedtke T, Parisi JE, O'Brien P, Petersen RC, Butler PC (2004) Increased risk of type 2 diabetes in Alzheimer disease. *Diabetes* 53(2):474–481
- Janssen-Heininger YM, Mossman BT, Heintz NH, Forman HJ, Kalyanaraman B, Finkel T et al (2008) Redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radic Biol Med* 45(1):1–17
- Jarvis J, Seed MJ, Elton R, Sawyer L, Agius R (2005) Relationship between chemical structure and the occupational asthma hazard of low molecular weight organic compounds. *Occup Environ Med* 62(4):243–250
- Jaurand MC (1997) Mechanisms of fiber-induced genotoxicity. *Environ Health Perspect* 105(Suppl 5):1073–1084
- Javadi P, Buchman TG, Stromberg PE, Husain KD, Dunne WM, Woolsey CA et al (2004) High-dose exogenous iron following cecal ligation and puncture increases mortality rate in mice and is associated with an increase in gut epithelial and splenic apoptosis. *Crit Care Med* 32(5):1178–1185
- Javadi P, Buchman TG, Stromberg PE, Turnbull IR, Vyas D, Hotchkiss RS et al (2005) Iron dysregulation combined with aging prevents sepsis-induced apoptosis. *J Surg Res* 128(1):37–44
- Jayawardhana B, Kell DB, Rattray M (2008) Bayesian inference of the sites of perturbations in metabolic pathways via Markov Chain Monte Carlo. *Bioinformatics* 24:1191–1197. doi:10.1093/bioinformatics/btn103
- Jehn M, Clark JM, Guallar E (2004) Serum ferritin and risk of the metabolic syndrome in U.S. adults. *Diabetes Care* 27(10):2422–2428
- Jehn ML, Guallar E, Clark JM, Couper D, Duncan BB, Ballantyne CM et al (2007) A prospective study of plasma ferritin level and incident diabetes: the atherosclerosis risk in communities (ARIC) study. *Am J Epidemiol* 165(9):1047–1054
- Jellinger K, Paulus W, Grundke-Iqbal I, Riederer P, Youdim MB (1990) Brain iron and ferritin in Parkinson's and Alzheimer's diseases. *J Neural Transm Park Dis Dement Sect* 2(4):327–340
- Jellinger KA (1999) The role of iron in neurodegeneration—prospects for pharmacotherapy of Parkinson's disease. *Drugs Aging* 14(2):115–140
- Jellinger KA (2002) Recent developments in the pathology of Parkinson's disease. *J Neural Transm Suppl* (62):347–376
- Jellinger KA (2009) Recent advances in our understanding of neurodegeneration. *J Neural Transm* 116(9):1111–1162
- Jenner P (2003) Oxidative stress in Parkinson's disease. *Ann Neurol* 53(Suppl 3):S26–S36 (discussion S28–S36)
- Jenner P, Olanow CW (1996) Oxidative stress and the pathogenesis of Parkinson's disease. *Neurology* 47(6 Suppl 3):S161–S170
- Jeong SY, Rathore KI, Schulz K, Ponka P, Arosio P, David S (2009) Dysregulation of iron homeostasis in the CNS contributes to disease progression in a mouse model of amyotrophic lateral sclerosis. *J Neurosci* 29(3):610–619
- Jia J, Zhu F, Ma X, Cao ZW, Li YX, Chen YZ (2009) Mechanisms of drug combinations: interaction and network perspectives. *Nat Rev Drug Discov* 8(2):111–128
- Jiang D, Li X, Williams R, Patel S, Men L, Wang Y et al (2009) Ternary complexes of iron, amyloid- and nitrotriacetic acid: binding affinities, redox properties, and relevance to iron-induced oxidative stress in Alzheimer's Disease. *Biochemistry*
- Jiang XP, Wang F, Yang DC, Elliott RL, Head JF (2002) Induction of apoptosis by iron depletion in the human breast cancer MCF-7 cell line and the 13762NF rat mammary adenocarcinoma in vivo. *Anticancer Res* 22(5):2685–2692
- Jiao Y, Wilkinson JT, Di X, Wang W, Hatcher H, Kock ND et al (2009) Curcumin, a cancer chemopreventive and chemotherapeutic agent, is a biologically active iron chelator. *Blood* 113(2):462–469
- Jimenez-Del-Rio M, Guzman-Martinez C, Velez-Pardo C (2010) The effects of polyphenols on survival and locomotor activity in *Drosophila melanogaster* exposed to iron and paraquat. *Neurochem Res* 35(2):227–238
- Jimenez Del Río M, Vélez-Pardo C (2004) Transition metal-induced apoptosis in lymphocytes via hydroxyl radical generation, mitochondria dysfunction, and caspase-3 activation: an in vitro model for neurodegeneration. *Arch Med Res* 35(3):185–193
- Jin H, Terai S, Sakaida I (2007) The iron chelator deferoxamine causes activated hepatic stellate cells to become quiescent and to undergo apoptosis. *J Gastroenterol* 42(6):475–484
- Johansson C, Castoldi AF, Onishchenko N, Manzo L, Vahter M, Ceccatelli S (2007) Neurobehavioural and molecular changes induced by methylmercury exposure during development. *Neurotox Res* 11(3–4):241–260
- Johnson D, Mayers I (2001) Multiple organ dysfunction syndrome: a narrative review. *Can J Anaesth* 48(5):502–509
- Johnson JJ, Bailey HH, Mukhtar H (2010) Green tea polyphenols for prostate cancer chemoprevention: a translational perspective. *Phytomedicine* 17(1):3–13
- Johnson L (2008) Iron and siderophores in fungal-host interactions. *Mycol Res* 112(Pt 2):170–183
- Johnson S (2000) Iron catalyzed oxidative damage, in spite of normal ferritin and transferrin saturation levels and its possible role in Werner's syndrome, Parkinson's disease, cancer, gout, rheumatoid arthritis, etc. *Med Hypotheses* 55(3):242–244
- Jolkkonen J, Jokivarsi K, Laitinen T, Gröhn O (2007) Subacute hemorrhage and resolution of edema in Rose Bengal stroke model in rats coincides with improved sensorimotor functions. *Neurosci Lett* 428(2–3):99–102
- Jones BC, Beard JL, Gibson JN, Unger EL, Allen RP, McCarthy KA et al (2007) Systems genetic analysis of peripheral iron parameters in the mouse. *Am J Physiol Regul Integr Comp Physiol* 293(1):R116–124
- Jovanovic SV, Simic MG (2000) Antioxidants in nutrition. *Ann NY Acad Sci* 899:326–334
- Julius C, Hutter G, Wagner U, Seeger H, Kana V, Kranich J et al (2008) Transcriptional stability of cultured cells upon prion infection. *J Mol Biol* 375(5):1222–1233
- Jung T, Bader N, Grune T (2007) Lipofuscin: formation, distribution, and metabolic consequences. *Ann NY Acad Sci* 1119:97–111

- Jurenka JS (2009) Anti-inflammatory properties of curcumin, a major constituent of *Curcuma longa*: a review of preclinical and clinical research. *Altern Med Rev* 14(2):141–153
- Jurkschat K, Ji X, Crossley A, Compton RG, Banks CE (2007) Super-washing does not leave single walled carbon nanotubes iron-free. *Analyst* 132(1):21–23
- Justicia C, Ramos-Cabrer P, Hoehn M (2008) MRI detection of secondary damage after stroke: chronic iron accumulation in the thalamus of the rat brain. *Stroke* 39(5):1541–1547
- Kaelin WG Jr (2005) The concept of synthetic lethality in the context of anticancer therapy. *Nat Rev Cancer* 5(9):689–698
- Kagu MB, Khalil MI, Ahmed SG (2007) Bone marrow macrophage iron stores in patients with HIV infection and AIDS-associated Kaposi's sarcoma. *Afr J Med Med Sci* 36(2):125–128
- Kaiserová H, Šimůnek T, van der Vijgh WJ, Bast A, Kvasničková E (2007) Flavonoids as protectors against doxorubicin cardiotoxicity: role of iron chelation, antioxidant activity and inhibition of carbonyl reductase. *Biochim Biophys Acta* 1772(9):1065–1074
- Kaito M (2007) Molecular mechanism of iron metabolism and overload in chronic hepatitis C. *J Gastroenterol* 42(1):96–99
- Kaito M, Iwasa M, Kobayashi Y, Fujita N, Tanaka H, Gabazza EC et al (2006) Iron reduction therapy by phlebotomy reduces lipid peroxidation and oxidative stress in patients with chronic hepatitis C. *J Gastroenterol* 41(9):921–922
- Kakhlon O, Breuer W, Munnich A, Cabantchik ZI (2010) Iron redistribution as a therapeutic strategy for treating diseases of localized iron accumulation. *Can J Physiol Pharmacol* 88(3):187–196
- Kala SV, Hasinoff BB, Richardson JS (1996) Brain samples from Alzheimer's patients have elevated levels of loosely bound iron. *Int J Neurosci* 86(3–4):263–269
- Kallianpur AR (2005) Iron and oxidative injury—a commentary on “Fatty acid-mediated iron translocation: a synergistic mechanism of oxidative injury” by D. Yao et al. *Free Radic Biol Med* 39(10):1305–1309
- Kalpravidh RW, Siritanaratkul N, Insain P, Charoensakdi R, Panichkul N, Hatairaktham S et al (2010) Improvement in oxidative stress and antioxidant parameters in beta-thalassemia/Hb E patients treated with curcuminoids. *Clin Biochem* 43(4–5):424–429
- Kamp DW (2009) Asbestos-induced lung diseases: an update. *Transl Res* 153(4):143–152
- Kamp DW, Graceffa P, Pryor WA, Weitzman SA (1992) The role of free radicals in asbestos-induced diseases. *Free Radic Biol Med* 12(4):293–315
- Kamp DW, Israbian VA, Preusen SE, Zhang CX, Weitzman SA (1995a) Asbestos causes DNA strand breaks in cultured pulmonary epithelial cells: role of iron-catalyzed free radicals. *Am J Physiol* 268(3 Pt 1):L471–L480
- Kamp DW, Israbian VA, Yeldandi AV, Panos RJ, Graceffa P, Weitzman SA (1995b) Phytic acid, an iron chelator, attenuates pulmonary inflammation and fibrosis in rats after intratracheal instillation of asbestos. *Toxicol Pathol* 23(6):689–695
- Kamp DW, Panduri V, Weitzman SA, Chandel N (2002) Asbestos-induced alveolar epithelial cell apoptosis: role of mitochondrial dysfunction caused by iron-derived free radicals. *Mol Cell Biochem* 234–235(1–2):153–160
- Kamp DW, Weitzman SA (1997) Asbestosis: clinical spectrum and pathogenic mechanisms. *Proc Soc Exp Biol Med* 214(1):12–26
- Kamp DW, Weitzman SA (1999) The molecular basis of asbestos induced lung injury. *Thorax* 54(7):638–652
- Kanai Y, Endou H (2003) Functional properties of multispecific amino acid transporters and their implications to transporter-mediated toxicity. *J Toxicol Sci* 28(1):1–17
- Kandaswami C, Middleton E Jr (1994) Free radical scavenging and antioxidant activity of plant flavonoids. *Adv Exp Med Biol* 366:351–376
- Kang JO, Jones C, Brothwell B (1998) Toxicity associated with iron overload found in hemochromatosis: possible mechanism in a rat model. *Clin Lab Sci* 11(6):350–354
- Kaprelyants AS, Gottschal JC, Kell DB (1993) Dormancy in non-sporulating bacteria. *FEMS Microbiol Rev* 10:271–286
- Karbownik M, Reiter RJ, Garcia JJ, Tan D (2000) Melatonin reduces phenylhydrazine-induced oxidative damage to cellular membranes: evidence for the involvement of iron. *Int J Biochem Cell Biol* 32(10):1045–1054
- Karin M (2008) The IkappaB kinase—a bridge between inflammation and cancer. *Cell Res* 18(3):334–342
- Karin M, Greten FR (2005) NF-kappaB: linking inflammation and immunity to cancer development and progression. *Nat Rev Immunol* 5(10):749–759
- Karlberg T, Schagerlöf U, Gakh O, Park S, Ryde U, Lindahl M et al (2006) The structures of frataxin oligomers reveal the mechanism for the delivery and detoxification of iron. *Structure* 14(10):1535–1546
- Karlsen JR, Borregaard N, Cowland JB (2010) Induction of neutrophil gelatinase-associated lipocalin expression by co-stimulation with IL-17 and TNF- α is controlled by I κ B- ζ but neither by C/EBP- β nor by C/EBP- δ . *J Biol Chem*
- Kartikasari AE, Visseren FL, Marx JJ, van Mullekom S, Kats-Renaud JH, Asbeck BS et al (2009) Intracellular labile iron promotes firm adhesion of human monocytes to endothelium under flow and transendothelial migration: iron and monocyte-endothelial cell interactions. *Atherosclerosis* 205(2):369–375
- Kasarskis EJ, Tandon L, Lovell MA, Ehmann WD (1995) Aluminum, calcium, and iron in the spinal cord of patients with sporadic amyotrophic lateral sclerosis using laser microprobe mass spectroscopy: a preliminary study. *J Neurol Sci* 130(2):203–208
- Kato J, Kobune M, Nakamura T, Kuroiwa G, Takada K, Takimoto R et al (2001) Normalization of elevated hepatic 8-hydroxy-2'-deoxyguanosine levels in chronic hepatitis C patients by phlebotomy and low iron diet. *Cancer Res* 61(24):8697–8702
- Kato J, Miyanishi K, Kobune M, Nakamura T, Takada K, Takimoto R et al (2007) Long-term phlebotomy with low-iron diet therapy lowers risk of development of hepatocellular carcinoma from chronic hepatitis C. *J Gastroenterol* 42(10):830–836
- Katta S, Kaur I, Chakrabarti S (2009) The molecular genetic basis of age-related macular degeneration: an overview. *J Genet* 88(4):425–449
- Kaur D, Andersen J (2004) Does cellular iron dysregulation play a causative role in Parkinson's disease? *Ageing Res Rev* 3(3):327–343
- Kaur D, Rajagopalan S, Andersen JK (2009) Chronic expression of H-ferritin in dopaminergic midbrain neurons results in an age-related expansion of the labile iron pool and subsequent neurodegeneration: implications for Parkinson's disease. *Brain Res*
- Kaur D, Rajagopalan S, Chinta S, Kumar J, Di Monte D, Cherny RA et al (2007) Chronic ferritin expression within murine dopaminergic midbrain neurons results in a progressive age-related neurodegeneration. *Brain Res* 1140:188–194
- Kaur D, Yantiri F, Rajagopalan S, Kumar J, Mo JO, Boonplueang R et al (2003) Genetic or pharmacological iron chelation prevents MPTP-induced neurotoxicity in vivo: a novel therapy for Parkinson's disease. *Neuron* 37(6):899–909
- Kaur P, Aschner M, Syversen T (2006) Glutathione modulation influences methyl mercury induced neurotoxicity in primary cell cultures of neurons and astrocytes. *Neurotoxicology* 27(4):492–500
- Kaushal V, Schlichter LC (2008) Mechanisms of microglia-mediated neurotoxicity in a new model of the stroke penumbra. *J Neurosci* 28(9):2221–2230
- Kauwe JSK, Bertelsen S, Mayo K, Cruchaga C, Abraham R, Hollingworth P et al (2010) Suggestive synergy between genetic

- variants in TF and HFE as risk factors for Alzheimer's disease. *Am J Med Genet B Neuropsychiatr Genet* 153B(4):955–959
- Kawabata T, Ma Y, Yamadori I, Okada S (1997) Iron-induced apoptosis in mouse renal proximal tubules after an injection of a renal carcinogen, iron-nitritotriacetate. *Carcinogenesis* 18(7):1389–1394
- Kaysen GA (2009) Biochemistry and biomarkers of inflamed patients: why look, what to assess. *Clin J Am Soc Nephrol* 4(Suppl 1):S56–S63
- Kazi TG, Afridi HI, Kazi N, Jamali MK, Arain MB, Sarfraz RA et al (2008) Distribution of zinc, copper and iron in biological samples of Pakistani myocardial infarction (1st, 2nd and 3rd heart attack) patients and controls. *Clin Chim Acta* 389(1–2):114–119
- Ke Y, Qian ZM (2003) Iron misregulation in the brain: a primary cause of neurodegenerative disorders. *Lancet Neurol* 2(4):246–253
- Keel M, Trentz O (2005) Pathophysiology of polytrauma. *Injury* 36(6):691–709
- Kehrer JP (2000) The Haber-Weiss reaction and mechanisms of toxicity. *Toxicology* 149(1):43–50
- Keith CT, Borisy AA, Stockwell BR (2005) Multicomponent therapeutics for networked systems. *Nat Rev Drug Discov* 4(1):71–78
- Kell DB (2004) Metabolomics and systems biology: making sense of the soup. *Curr Op Microbiol* 7:296–307
- Kell DB (2006a) Metabolomics, modelling, machine learning in systems biology: towards an understanding of the languages of cells. The 2005 Theodor Bücher lecture. *FEBS J* 273:873–894
- Kell DB (2006b) Systems biology, metabolic modelling and metabolomics in drug discovery and development. *Drug Disc Today* 11(23/24):1085–1092
- Kell DB (2007) The virtual human: towards a global systems biology of multiscale, distributed biochemical network models. *IUBMB Life* 59(11):689–695
- Kell DB (2008) Progress being made on standards for use in data sharing. *Nature* 456(7218):29
- Kell DB (2009a) Iron behaving badly: inappropriate iron chelation as a major contributor to the aetiology of vascular and other progressive inflammatory and degenerative diseases. *BMC Med Genomics* 2:2. <http://www.biomedcentral.com/1755-8794/2/2/>
- Kell DB (2009b) A systems biologist ponders how disparate ideas can sometimes come together beautifully. *Nature* 460:669
- Kell DB, Darby RM, Draper J (2001) Genomic computing: explanatory analysis of plant expression profiling data using machine learning. *Plant Physiol.* 126:943–951
- Kell DB, Dobson PD (2009) The cellular uptake of pharmaceutical drugs is mainly carrier-mediated and is thus an issue not so much of biophysics but of systems biology. *Proceedings of the international Beilstein symposium on systems chemistry*, 149–168. <http://www.beilstein-institut.de/Bozen2008/Proceedings/Kell/Kell.pdf>. Logos Verlag, Berlin
- Kell DB, Kaprelyants AS, Grafen A (1995) On pheromones, social behaviour and the functions of secondary metabolism in bacteria. *Trends Ecol Evol* 10:126–129
- Kell DB, Kaprelyants AS, Weichart DH, Harwood CL, Barer MR (1998) Viability and activity in readily culturable bacteria: a review and discussion of the practical issues. *Antonie van Leeuwenhoek* 73:169–187
- Kell DB, King RD (2000) On the optimization of classes for the assignment of unidentified reading frames in functional genomics programmes: the need for machine learning. *Trends Biotechnol* 18(3):93–98
- Kell DB, Knowles JD (2006) The role of modeling in systems biology. In: Szallasi Z, Stelling J, Periwal V (eds) *System modeling in cellular biology: from concepts to nuts and bolts*. MIT Press, Cambridge, pp 3–18
- Kell DB, Mendes P (2008) The markup is the model: reasoning about systems biology models in the Semantic Web era. *J Theoret Biol* 252:538–543. doi: http://www.sciencedirect.com/science?_ob=MIimg&_imagekey=B6WMD-4R0KTJW-4-1&_cdi=6932&_user=121749&_orig=browse&_coverDate=10%2F27%2F2007&_sk=999999999&view=c&wchp=dGLbVtb-zSkWz&md5=464b923e674b26776e8ad407dd17f4ce&ie=/sarticle.pdf
- Kell DB, Oliver SG (2004) Here is the evidence, now what is the hypothesis? The complementary roles of inductive and hypothesis-driven science in the post-genomic era. *Bioessays* 26:99–105
- Kell DB, Westerhoff HV (1986a) Metabolic control theory: its role in microbiology and biotechnology. *FEMS Microbiol Rev* 39:305–320
- Kell DB, Westerhoff HV (1986b) Towards a rational approach to the optimization of flux in microbial biotransformations. *Trends Biotechnol* 4:137–142
- Kelleher P, Pacheco K, Newman LS (2000) Inorganic dust pneumonias: the metal-related parenchymal disorders. *Environ Health Perspect* 108(Suppl 4):685–696
- Kenyon CJ (2010) The genetics of ageing. *Nature* 464(7288):504–512
- Kermer P, Liman J, Weishaupt JH, Bahr M (2004) Neuronal apoptosis in neurodegenerative diseases: from basic research to clinical application. *Neurodegener Dis* 1(1):9–19
- Ketonen L, Kiebertz K, Kazee AM, Tuite M (1996) Putaminal iron deposition in HIV infection. *J NeuroAIDS* 1(2):33–40
- Kew MC, Asare GA (2007) Dietary iron overload in the African and hepatocellular carcinoma. *Liver Int* 27(6):735–741
- Khalil M, Enzinger C, Langkammer C, Tscherner M, Wallner-Blazek M, Jehna M et al (2009) Quantitative assessment of brain iron by R2* relaxometry in patients with clinically isolated syndrome and relapsing-remitting multiple sclerosis. *Mult Scler* 15(9):1048–1054
- Khan MF, Kaphalia BS, Ansari GA (1995) Erythrocyte-aniline interaction leads to their accumulation and iron deposition in rat spleen. *J Toxicol Environ Health* 44(4):415–421
- Khan MF, Wu X, Alcock NW, Boor PJ, Ansari GA (1999) Iron exacerbates aniline-associated splenic toxicity. *J Toxicol Environ Health A* 57(3):173–184
- Kibel A, Belovari T, Drenjančević-Perić I (2008) The role of transferrin in atherosclerosis. *Med Hypotheses* 70(4):793–797
- Kidd PM (2000) Parkinson's disease as multifactorial oxidative neurodegeneration: implications for integrative management. *Altern Med Rev* 5(6):502–529
- Kiechl S, Willeit J, Egger G, Poewe W, Oberhollenzer F (1997) Body iron stores and the risk of carotid atherosclerosis: prospective results from the Bruneck study. *Circulation* 96(10):3300–3307
- Kim BH, Jun YC, Jin JK, Kim JI, Kim NH, Leibold EA et al (2007) Alteration of iron regulatory proteins (IRP1 and IRP2) and ferritin in the brains of scrapie-infected mice. *Neurosci Lett* 422(3):158–163
- Kim D, Chun BG, Kim YK, Lee YH, Park CS, Jeon I et al (2008a) In vivo tracking of human mesenchymal stem cells in experimental stroke. *Cell Transpl* 16(10):1007–1012
- Kim EY, Ham SK, Shigenaga MK, Han O (2008b) Bioactive dietary polyphenolic compounds reduce nonheme iron transport across human intestinal cell monolayers. *J Nutr* 138(9):1647–1651
- Kim JI, Choi SI, Kim NH, Jin JK, Choi EK, Carp RI et al (2001) Oxidative stress and neurodegeneration in prion diseases. *Ann NY Acad Sci* 928:182–186
- Kim NH, Park SJ, Jin JK, Kwon MS, Choi EK, Carp RI et al (2000) Increased ferric iron content and iron-induced oxidative stress in the brains of scrapie-infected mice. *Brain Res* 884(1–2):98–103
- Kim YW, Lee SM, Shin SM, Hwang SJ, Brooks JS, Kang HE et al (2009) Efficacy of sauchinone as a novel AMPK-activating lignan for preventing iron-induced oxidative stress and liver injury. *Free Radic Biol Med* 47:1082–1092

- King RD, Rowland J, Oliver SG, Young M, Aubrey W, Byrne E et al (2009) The automation of science. *Science* 324(5923):85–89
- King RD, Whelan KE, Jones FM, Reiser PGK, Bryant CH, Muggleton SH et al (2004) Functional genomic hypothesis generation and experimentation by a robot scientist. *Nature* 427:247–252
- Kiokias S, Varzakas T, Oreopoulou V (2008) In vitro activity of vitamins, flavonoids, and natural phenolic antioxidants against the oxidative deterioration of oil-based systems. *Crit Rev Food Sci Nutr* 48(1):78–93
- Kirkwood TB (2008) A systematic look at an old problem. *Nature* 451(7179):644–647
- Kirsch R, Sijtsma HP, Tlali M, Marais AD, Hall P, de la M (2006) Effects of iron overload in a rat nutritional model of non-alcoholic fatty liver disease. *Liver Int* 26(10):1258–1267
- Kitagawa S, Tanaka Y, Tanaka M, Endo K, Yoshii A (2009) Enhanced skin delivery of quercetin by microemulsion. *J Pharm Pharmacol* 61(7):855–860
- Kitano H (2007) A robustness-based approach to systems-oriented drug design. *Nat Rev Drug Discov* 6(3):202–210
- Klipp E, Herwig R, Kowald A, Wierling C, Lehrach H (2005) Systems biology in practice: concepts, implementation and clinical application. Wiley/VCH, Berlin
- Klipp E, Liebermeister W, Helbig A, Kowald A, Schaber J (2007) Systems biology standards—the community speaks. *Nat Biotechnol* 25(4):390–391
- Knaapen AM, Borm PJ, Albrecht C, Schins RP (2004) Inhaled particles and lung cancer. Part A: mechanisms. *Int J Cancer* 109(6):799–809
- Knight CG, Platt M, Rowe W, Wedge DC, Khan F, Day P et al (2009) Array-based evolution of DNA aptamers allows modelling of an explicit sequence-fitness landscape. *Nucleic Acids Res* 37(1):e6
- Knight SA, Kim R, Pain D, Dancis A (1999) The yeast connection to Friedreich ataxia. *Am J Hum Genet* 64(2):365–371
- Knovich MA, Storey JA, Coffman LG, Torti SV, Torti FM (2009) Ferritin for the clinician. *Blood Rev* 23(3):95–104
- Ko C, Siddaiah N, Berger J, Gish R, Brandhagen D, Sterling RK et al (2007) Prevalence of hepatic iron overload and association with hepatocellular cancer in end-stage liver disease: results from the national hemochromatosis transplant registry. *Liver Int* 27(10):1394–1401
- Kobayashi K, Yoshino F, Takahashi SS, Todoki K, Maehata Y, Komatsu T et al (2008) Direct assessments of the antioxidant effects of propofol medium chain triglyceride/long chain triglyceride on the brain of stroke-prone spontaneously hypertensive rats using electron spin resonance spectroscopy. *Anesthesiology* 109(3):426–435
- Koc M, Nad'ová Z, Kovář J (2006) Sensitivity of cells to apoptosis induced by iron deprivation can be reversibly changed by iron availability. *Cell Prolif* 39(6):551–561
- Kocyigit A, Armutcu F, Gurel A, Ermis B (2004) Alterations in plasma essential trace elements selenium, manganese, zinc, copper, and iron concentrations and the possible role of these elements on oxidative status in patients with childhood asthma. *Biol Trace Elem Res* 97(1):31–41
- Kohanski MA, DePristo MA, Collins JJ (2010) Sublethal antibiotic treatment leads to multidrug resistance via radical-induced mutagenesis. *Mol Cell* 37(3):311–320
- Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ (2007) A common mechanism of cellular death induced by bactericidal antibiotics. *Cell* 130(5):797–810
- Kohanski MA, Dwyer DJ, Wierzbowski J, Cottarel G, Collins JJ (2008) Mistranslation of membrane proteins and two-component system activation trigger antibiotic-mediated cell death. *Cell* 135(4):679–690
- Kola I, Landis J (2004) Can the pharmaceutical industry reduce attrition rates? *Nat Rev Drug Discov* 3(8):711–715
- Kolberg JA, Jorgensen T, Gerwien RW, Hamren S, McKenna MP, Moler E et al (2009) Development of a type 2 diabetes risk model from a panel of serum biomarkers from the Inter99 cohort. *Diabetes Care* 32(7):1207–1212
- Kom GD, Schwedhelm E, Nielsen P, Böger RH (2006) Increased urinary excretion of 8-iso-prostaglandin F₂alpha in patients with HFE-related hemochromatosis: a case-control study. *Free Radic Biol Med* 40(7):1194–1200
- Kompala SD, Babbs CF, Blaho KE (1986) Effect of deferoxamine on late deaths following CPR in rats. *Ann Emerg Med* 15(4):405–407
- Kon K, Kim JS, Jaeschke H, Lemasters JJ (2004) Mitochondrial permeability transition in acetaminophen-induced necrosis and apoptosis of cultured mouse hepatocytes. *Hepatology* 40(5):1170–1179
- Končić MZ, Kremer D, Gruz J, Strnad M, Biševac G, Kosalec I et al (2010) Antioxidant and antimicrobial properties of *Moltkia petraea* (Tratt.) Griseb. flower, leaf and stem infusions. *Food Chem Toxicol* 48(6):1537–1542
- Kondapalli KC, Kok NM, Dancis A, Stemmler TL (2008) Drosophila frataxin: an iron chaperone during cellular Fe-S cluster bioassembly. *Biochemistry* 47(26):6917–6927
- Kontoghiorghes GJ, Kolnagou A, Peng CT, Shah SV, Aessopos A (2010) Safety issues of iron chelation therapy in patients with normal range iron stores including thalassaemia, neurodegenerative, renal and infectious diseases. *Expert Opin Drug Saf* 9(2):201–206
- Kooncumchoo P, Sharma S, Porter J, Govitrapong P, Ebadi M (2006) Coenzyme Q(10) provides neuroprotection in iron-induced apoptosis in dopaminergic neurons. *J Mol Neurosci* 28(2):125–141
- Koppenol WH, Moreno JJ, Pryor WA, Ischiropoulos H, Beckman JS (1992) Peroxynitrite, a cloaked oxidant formed by nitric oxide and superoxide. *Chem Res Toxicol* 5(6):834–842
- Kopterides P, Falagas ME (2009) Statins for sepsis: a critical and updated review. *Clin Microbiol Infect* 15(4):325–334
- Korkina LG, Afanas'ev IB (1997) Antioxidant and chelating properties of flavonoids. *Adv Pharmacol* 38:151–163
- Korkmaz A, Kunak ZI, Paredes SD, Yaren H, Tan DX, Reiter RJ (2008) The use of melatonin to combat mustard toxicity. REVIEW. *Neuro Endocrinol Lett* 29(5):614–619
- Korkmaz A, Reiter RJ, Topal T, Manchester LC, Oter S, Tan DX (2009) Melatonin: an established antioxidant worthy of use in clinical trials. *Mol Med* 15(1–2):43–50
- Korkmaz A, Yaren H, Topal T, Oter S (2006) Molecular targets against mustard toxicity: implication of cell surface receptors, peroxynitrite production, and PARP activation. *Arch Toxicol* 80(10):662–670
- Kostoff RN (2002) Overcoming specialization. *Bioscience* 52(10):937–941
- Kostoff RN, Delafuente JC (2006) The unknown impacts of combinations of large numbers of drugs. *Drug Saf* 29(3):183–185
- Kotamraju S, Chitambar CR, Kalivendi SV, Joseph J, Kalyanaraman B (2002) Transferrin receptor-dependent iron uptake is responsible for doxorubicin-mediated apoptosis in endothelial cells—role of oxidant-induced iron signaling in apoptosis. *J Biol Chem* 277(19):17179–17187
- Kotamraju S, Kalivendi SV, Konorev E, Chitambar CR, Joseph J, Kalyanaraman B (2004a) Oxidant-induced iron signaling in doxorubicin-mediated apoptosis. *Methods Enzymol* 378:362–382
- Kotamraju S, Tampo Y, Kalivendi SV, Joseph J, Chitambar CR, Kalyanaraman B (2004b) Nitric oxide mitigates peroxide-induced iron-signaling, oxidative damage, and apoptosis in endothelial cells: role of proteasomal function? *Arch Biochem Biophys* 423(1):74–80

- Koufaki M, Detsi A, Kiziridi C (2009) Multifunctional lipoic acid conjugates. *Curr Med Chem* 16(35):4728–4742
- Kourie JI (2002) Prion channel proteins and their role in vacuolation and neurodegenerative diseases. *Eur Biophys J* 31(5):409–416
- Kovář J, Stunz LL, Stewart BC, Kriegerbecková K, Ashman RF, Kemp JD (1997) Direct evidence that iron deprivation induces apoptosis in murine lymphoma 38C13. *Pathobiology* 65(2):61–68
- Kovář J, Valenta T, Štýbrová H (2001) Differing sensitivity of tumor cells to apoptosis induced by iron deprivation in vitro. *In Vitro Cell Dev Biol Anim* 37(7):450–458
- Kowdley KV (2010) The role of iron in nonalcoholic fatty liver disease: the story continues. *Gastroenterology* 138(3):817–819
- Krukoski DW, Comar SR, Claro LM, Leonart MS, do Nascimento AJ (2009) Effect of vitamin C, deferoxamine, quercetin and rutin against tert-butyl hydroperoxide oxidative damage in human erythrocytes. *Hematology* 14(3):168–172
- Kruman I, Bruce-Keller AJ, Bredesen D, Waeg G, Mattson MP (1997) Evidence that 4-hydroxynonenal mediates oxidative stress-induced neuronal apoptosis. *J Neurosci* 17(13):5089–5100
- Kruszewski M (2003) Labile iron pool: the main determinant of cellular response to oxidative stress. *Mutat Res* 531(1–2):81–92
- Kuhn M, Campillos M, Letunic I, Jensen LJ, Bork P (2010) A side effect resource to capture phenotypic effects of drugs. *Mol Syst Biol* 6:343
- Kujoth GC, Hiona A, Pugh TD, Someya S, Panzer K, Wohlgemuth SE et al (2005) Mitochondrial DNA mutations, oxidative stress, and apoptosis in mammalian aging. *Science* 309(5733):481–484
- Kumar O, Sugendran K, Vijayaraghavan R (2003) Oxidative stress associated hepatic and renal toxicity induced by ricin in mice. *Toxicol* 41(3):333–338
- Kumar V, Sinha AK, Makkar HPS, Becker K (2010) Dietary roles of phytate and phytase in human nutrition: a review. *Food Chem* 120(4):945–959
- Kummer S, Hayes GR, Gilbert RO, Beach DH, Lucas JJ, Singh BN (2008) Induction of human host cell apoptosis by *Trichomonas vaginalis* cysteine proteases is modulated by parasite exposure to iron. *Microb Pathog* 44(3):197–203
- Kunsch C, Medford RM (1999) Oxidative stress as a regulator of gene expression in the vasculature. *Circ Res* 85(8):753–766
- Kuo KL, Hung SC, Wei YH, Tarng DC (2008) Intravenous iron exacerbates oxidative DNA damage in peripheral blood lymphocytes in chronic hemodialysis patients. *J Am Soc Nephrol* 19(9):1817–1826
- Kurz T, Leake A, von Zglinicki T, Brunk UT (2004) Relocalized redox-active lysosomal iron is an important mediator of oxidative-stress-induced DNA damage. *Biochem J* 378:1039–1045
- Kurz T, Terman A, Brunk UT (2007) Autophagy, ageing and apoptosis: the role of oxidative stress and lysosomal iron. *Arch Biochem Biophys* 462(2):220–230
- Kurz T, Terman A, Gustafsson B, Brunk UT (2008a) Lysosomes and oxidative stress in aging and apoptosis. *Biochim Biophys Acta* 1780(11):1291–1303
- Kurz T, Terman A, Gustafsson B, Brunk UT (2008b) Lysosomes in iron metabolism, ageing and apoptosis. *Histochem Cell Biol* 129(4):389–406
- Kwiatkowski JL (2010) Oral iron chelators. *Hematol Oncol Clin North Am* 24(1):229–248
- Kwiatkowski TJ Jr, Bosco DA, Leclerc AL, Tamrazian E, Vandenberg CR, Russ C et al (2009) Mutations in the FUS/TLS gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science* 323(5918):1205–1208
- Lachili B, Hininger I, Faure H, Arnaud J, Richard MJ, Favier A et al (2001) Increased lipid peroxidation in pregnant women after iron and vitamin C supplementation. *Biol Trace Elem Res* 83(2):103–110
- Lagan AL, Melley DD, Evans TW, Quinlan GJ (2008) Pathogenesis of the systemic inflammatory syndrome and acute lung injury: role of iron mobilization and decompartmentalization. *Am J Physiol Lung Cell Mol Physiol* 294(2):L161–174
- Lakshmi Devi S, Anuradha CV (2009) Mitochondrial damage, cytotoxicity and apoptosis in iron-potentiated alcoholic liver fibrosis: amelioration by taurine. *Amino Acids*
- Lam WW, Au WY, Chu WC, Tam S, Ha SY, Pennell DJ (2008) One-stop measurement of iron deposition in the anterior pituitary, liver, and heart in thalassemia patients. *J Magn Reson Imaging* 28(1):29–33
- Lamey PJ, Biagioni PA (1995) Relationship between iron status and recrudescence of herpes labialis. *Eur J Clin Microbiol Infect Dis* 14(7):604–605
- Lancaster JR Jr (2008) Protein cysteine thiol nitrosation: marker or reactive nitrogen species-induced nonerythroid cellular signaling? *Nitric Oxide* 19(2):68–72
- Landino LM (2008) Protein thiol modification by peroxynitrite anion and nitric oxide donors. *Methods Enzymol* 440:95–109
- Lapenna D, Pierdomenico SD, Ciofani G, Ucchino S, Neri M, Giamberardino MA et al (2007) Association of body iron stores with low molecular weight iron and oxidant damage of human atherosclerotic plaques. *Free Rad Biol Med* 42(4):492–498
- Latunde-Dada GO, Vulpe CD, Anderson GJ, Simpson RJ, McKie AT (2004) Tissue-specific changes in iron metabolism genes in mice following phenylhydrazine-induced haemolysis. *Biochim Biophys Acta* 1690(2):169–176
- Lavados M, Guillón M, Mujica MC, Rojo LE, Fuentes P, Maccioni RB (2008) Mild cognitive impairment and Alzheimer patients display different levels of redox-active CSF iron. *J Alzheimers Dis* 13(2):225–232
- Le Novère N, Finney A, Hucka M, Bhalla US, Campagne F, Collado-Vides J et al (2005) Minimum information requested in the annotation of biochemical models (MIRIAM). *Nat Biotechnol* 23(12):1509–1515
- Le Novère N, Hucka M, Mi H, Moodie S, Schreiber F, Sorokin A et al (2009) The systems biology graphical notation. *Nat Biotechnol* 27(8):735–741
- Learidi A, Caraglia M, Selleri C, Pepe S, Pizzi C, Notaro R et al (1998) Desferioxamine increases iron depletion and apoptosis induced by ara-C of human myeloid leukaemic cells. *Br J Haematol* 102(3):746–752
- LeBel CP, Ali SF, Bondy SC (1992) Deferoxamine inhibits methyl mercury-induced increases in reactive oxygen species formation in rat brain. *Toxicol Appl Pharmacol* 112(1):161–165
- Lecube A, Hernandez C, Pelegri D, Simo R (2008) Factors accounting for high ferritin levels in obesity. *Int J Obes (Lond)*, doi:10.1038/ijo.2008.1154
- Lecube A, Hernández C, Simó R (2009) Glucose abnormalities in non-alcoholic fatty liver disease and chronic hepatitis C virus infection: the role of iron overload. *Diabetes Metab Res Rev* 25(5):403–410
- Lee RE, Balcerzak SP, Westerman MP (1967) Gaucher's disease. A morphologic study and measurements of iron metabolism. *Am J Med* 42(6):891–898
- Lee RE, Peters SP, Glew RH (1977) Gaucher's disease: clinical, morphologic, and pathogenetic considerations. *Pathol Annu* (12 Pt 2):309–339
- Lee FY, Lee TS, Pan CC, Huang AL, Chau LY (1998) Colocalization of iron and ceroid in human atherosclerotic lesions. *Atherosclerosis* 138(2):281–288
- Lee TS, Shiao MS, Pan CC, Chau LY (1999) Iron-deficient diet reduces atherosclerotic lesions in ApoE-deficient mice. *Circulation* 99(9):1222–1229
- Lee DW, Andersen JK (2010) Iron elevations in the aging Parkinsonian brain: a consequence of impaired iron homeostasis? *J Neurochem* 112(2):332–339

- Lee DH, Zacharski LR, Jacobs DR (2006a) Comparison of the serum ferritin and percentage of transferrin saturation as exposure markers of iron-driven oxidative stress-related disease outcomes. *Am Heart J* 151(6). doi:10.1016/j.ahj.2006.1003.1009
- Lee DW, Andersen JK, Kaur D (2006b) Iron dysregulation and neurodegeneration—the molecular connection. *Mol Interv* 6(2): 89–97
- Lee JC, Son YO, Choi KC, Jang YS (2006c) Hydrogen peroxide induces apoptosis of BJAB cells due to formation of hydroxyl radicals via intracellular iron-mediated Fenton chemistry in glucose oxidase-mediated oxidative stress. *Mol Cells* 22(1):21–29
- Lee MT, Piomelli S, Granger S, Miller ST, Harkness S, Brambilla DJ et al (2006d) Stroke prevention trial in sickle cell anemia (STOP): extended follow-up and final results. *Blood* 108(3):847–852
- Lee SK, Jang HJ, Lee HJ, Lee J, Jeon BH, Jun CD et al (2006e) p38 and ERK MAP kinase mediates iron chelator-induced apoptosis and -suppressed differentiation of immortalized and malignant human oral keratinocytes. *Life Sci* 79(15):1419–1427
- Lee SK, Lee JJ, Lee HJ, Lee J, Jeon BH, Jun CD et al (2006f) Iron chelator-induced growth arrest and cytochrome c-dependent apoptosis in immortalized and malignant oral keratinocytes. *J Oral Pathol Med* 35(4):218–226
- Lee KS, Raymond LD, Schoen B, Raymond GJ, Kett L, Moore RA et al (2007a) Hemin interactions and alterations of the subcellular localization of prion protein. *J Biol Chem* 282(50): 36525–36533
- Lee MS, Johansen L, Zhang Y, Wilson A, Keegan M, Avery W et al (2007b) The novel combination of chlorpromazine and pentamidine exerts synergistic antiproliferative effects through dual mitotic action. *Cancer Res* 67(23):11359–11367
- Leeson PD, Springthorpe B (2007) The influence of drug-like concepts on decision-making in medicinal chemistry. *Nat Rev Drug Discov* 6(11):881–890
- Lehár J, Krueger A, Zimmermann G, Borisy A (2008a) High-order combination effects and biological robustness. *Mol Syst Biol* 4:215
- Lehár J, Stockwell BR, Giaever G, Nislow C (2008b) Combination chemical genetics. *Nat Chem Biol* 4(11):674–681
- Lehár J, Zimmermann GR, Krueger AS, Molnar RA, Ledell JT, Heilbut AM et al (2007) Chemical combination effects predict connectivity in biological systems. *Mol Syst Biol* 3:80
- Lehmann DJ, Worwood M, Ellis R, Wimhurst VL, Merryweather-Clarke AT, Warden DR et al (2006) Iron genes, iron load and risk of Alzheimer's disease. *J Med Genet* 43(10):e52
- Lehmann S (2002) Metal ions and prion diseases. *Curr Opin Chem Biol* 6(2):187–192
- Leitch HA (2007) Improving clinical outcome in patients with myelodysplastic syndrome and iron overload using iron chelation therapy. *Leuk Res* 31(Suppl 3):S7–S9
- Leithead JA, Ferguson JW, Bates CM, Davidson JS, Lee A, Bathgate AJ et al (2009) The systemic inflammatory response syndrome is predictive of renal dysfunction in patients with non-paracetamol-induced acute liver failure. *Gut* 58(3):443–449
- Lemineur T, Deby-Dupont G, Preiser JC (2006) Biomarkers of oxidative stress in critically ill patients: what should be measured, when and how? *Curr Op Clin Nutr Metab Care* 9(6):704–710
- Leuner B, Kozorovitskiy Y, Gross CG, Gould E (2007) Diminished adult neurogenesis in the marmoset brain precedes old age. *Proc Natl Acad Sci USA* 104(43):17169–17173
- Levenson CW (2003) Iron and Parkinson's disease: chelators to the rescue? *Nutr Rev* 61(9):311–313
- Levenson CW (2005) Trace metal regulation of neuronal apoptosis: from genes to behavior. *Physiol Behav* 86(3):399–406
- Leveugle B, Spik G, Perl DP, Bouras C, Fillit HM, Hof PR (1997) Distribution of the iron-binding protein lactoferrin in the pathological lesions of neurodegenerative diseases. *Lactoferrin Interact Biol Funct* 28:119–124, 408
- LeVine SM (1997) Iron deposits in multiple sclerosis and Alzheimer's disease brains. *Brain Res* 760(1–2):298–303
- LeVine SM, Chakrabarty A (2004) The role of iron in the pathogenesis of experimental allergic encephalomyelitis and multiple sclerosis. *Ann NY Acad Sci* 1012:252–266
- Levesse V, Renier A, Levy F, Broaddus VC, Jaurand M (2000) DNA breakage in asbestos-treated normal and transformed (TSV40) rat pleural mesothelial cells. *Mutagenesis* 15(3):239–244
- Li D, Shimamura T, Ji H, Chen L, Haringsma HJ, McNamara K et al (2007) Bronchial and peripheral murine lung carcinomas induced by T790M–L858R mutant EGFR respond to HKI-272 and rapamycin combination therapy. *Cancer Cell* 12(1):81–93
- Li H, Rybicki AC, Suzuka SM, von Bonsdorff L, Breuer W, Hall CB et al (2010) Transferrin therapy ameliorates disease in beta-thalassemic mice. *Nat Med* 16(2):177–182
- Li H, Zhou Q, Wu Y, Fu J, Wang T, Jiang G (2009) Effects of waterborne nano-iron on medaka (*Oryzias latipes*): antioxidant enzymatic activity, lipid peroxidation and histopathology. *Ecotoxicol Environ Saf* 72(3):684–692
- Li L, Hölscher C (2007) Common pathological processes in Alzheimer disease and type 2 diabetes: a review. *Brain Res Rev* 56(2):384–402
- Li Q, Withoff S, Verma IM (2005) Inflammation-associated cancer: NF-kappaB is the lynchpin. *Trends Immunol* 26(6):318–325
- Li T-S, Marbán E (2010) Physiological levels of reactive oxygen species are required to maintain genomic stability in stem cells. *Stem Cells*
- Li XJ, Brazhnik O, Kamal A, Guo D, Lee C, Hoops S (2003) Databases and visualization for metabolomics. In: Harrigan GG, Goodacre R et al (eds) *Metabolic profiling: its role in biomarker discovery and gene function analysis*. Kluwer Academic Publishers, Boston, pp 293–309
- Libby P (2002) Inflammation in atherosclerosis. *Nature* 420(6917): 868–874
- Libby P, Ridker PM, Maseri A (2002) Inflammation and atherosclerosis. *Circulation* 105(9):1135–1143
- Liby KT, Yore MM, Sporn MB (2007) Triterpenoids and rexinoids as multifunctional agents for the prevention and treatment of cancer. *Nat Rev Cancer* 7(5):357–369
- Liebermeister W, Klipp E (2006) Bringing metabolic networks to life: integration of kinetic, metabolic, and proteomic data. *Theor Biol Med Model* 3:42
- Lill R (2009) Function and biogenesis of iron-sulphur proteins. *Nature* 460(7257):831–838
- Lill R, Muhlenhoff U (2008) Maturation of iron-sulfur proteins in eukaryotes: mechanisms, connected processes, and diseases. *Annu Rev Biochem* 77:669–700
- Lim CK, Kalinowski DS, Richardson DR (2008) Protection against hydrogen peroxide-mediated cytotoxicity in Friedreich's ataxia fibroblasts using novel iron chelators of the 2-pyridylcarboxaldehyde isonicotinoyl hydrazone class. *Mol Pharmacol* 74(1): 225–235
- Lim MK, Lee CK, Ju YS, Cho YS, Lee MS, Yoo B et al (2001) Serum ferritin as a serologic marker of activity in systemic lupus erythematosus. *Rheumatol Int* 20(3):89–93
- Lim ZY, Fiaccadori V, Gandhi S, Hayden J, Kenyon M, Ireland R et al (2010) Impact of pre-transplant serum ferritin on outcomes of patients with myelodysplastic syndromes or secondary acute myeloid leukaemia receiving reduced intensity conditioning allogeneic haematopoietic stem cell transplantation. *Leuk Res* 34(6):723–727

- Limdi JK, Crampton JR (2004) Hereditary haemochromatosis. *QJM* 97(6):315–324
- Lin MT, Beal MF (2006) Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443(7113):787–795
- Linden R, Martins VR, Prado MA, Cammarota M, Izquierdo I, Brentani RR (2008) Physiology of the prion protein. *Physiol Rev* 88(2):673–728
- Linseisen J, Rohrmann S (2008) Biomarkers of dietary intake of flavonoids and phenolic acids for studying diet-cancer relationship in humans. *Eur J Nutr* 47(Suppl 2):60–68
- Liu IF, Annamalai T, Sutherland JH, Tse-Dinh YC (2009a) Hydroxyl radicals are involved in cell killing by the bacterial topoisomerase I cleavage complex. *J Bacteriol* 191(16):5315–5319
- Liu G, Men P, Kudo W, Perry G, Smith MA (2009b) Nanoparticle-chelator conjugates as inhibitors of amyloid-beta aggregation and neurotoxicity: a novel therapeutic approach for Alzheimer disease. *Neurosci Lett* 455(3):187–190
- Liu G, Men P, Perry G, Smith MA (2009c) Metal chelators coupled with nanoparticles as potential therapeutic agents for Alzheimer's disease. *J Nanoneurosci* 1(1):42–55
- Liu G, Beri R, Mueller A, Kamp DW (2010a) Molecular mechanisms of asbestos-induced lung epithelial cell apoptosis. *Chem Biol Interact* (in press)
- Liu G, Men P, Perry G, Smith MA (2010b) Nanoparticle and iron chelators as a potential novel Alzheimer therapy. *Methods Mol Biol* 610:123–144
- Liu PK, Wang FS (2008) Inverse problems of biological systems using multi-objective optimization. *J Chin Inst Chem Eng* 39(5):399–406
- Ljung L (1987) System identification: theory for the user. Prentice Hall, Englewood Cliffs, NJ
- Lledo PM, Alonso M, Grubb MS (2006) Adult neurogenesis and functional plasticity in neuronal circuits. *Nat Rev Neurosci* 7(3):179–193
- Lleó A, Blesa R, Angelopoulos C, Pastor-Rubio P, Villa M, Oliva R et al (2002) Transferrin C2 allele, haemochromatosis gene mutations, and risk for Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 72(6):820–821
- Lloyd DR, Carmichael PL, Phillips DH (1998) Comparison of the formation of 8-hydroxy-2'-deoxyguanosine and single- and double-strand breaks in DNA mediated by Fenton reactions. *Chem Res Toxicol* 11(5):420–427
- Loft S, Poulsen HE (1996) Cancer risk and oxidative DNA damage in man. *J Mol Med* 74(6):297–312
- Loh A, Hadzihametovic M, Dunaief JL (2009) Iron homeostasis and eye disease. *Biochim Biophys Acta* 1790(7):637–649
- Loh KP, Huang SH, De Silva R, Tan BK, Zhu YZ (2006) Oxidative stress: apoptosis in neuronal injury. *Curr Alzheimer Res* 3(4):327–337
- Loke WM, Jenner AM, Proudfoot JM, McKinley AJ, Hodgson JM, Halliwell B et al (2009) A metabolite profiling approach to identify biomarkers of flavonoid intake in humans. *J Nutr* 139(12):2309–2314
- Lolis E, Bucala R (2003) Therapeutic approaches to innate immunity: severe sepsis and septic shock. *Nat Rev Drug Discov* 2(8):635–645
- Long LH, Clement MV, Halliwell B (2000) Artifacts in cell culture: Rapid generation of hydrogen peroxide on addition of (–)-epigallocatechin, (–)-epigallocatechin gallate, (+)-catechin, and quercetin to commonly used cell culture media. *Biochem Biophys Res Comm* 273(1):50–53
- Loomis LD, Raymond KN (1991) Solution equilibria of enterobactin and metal enterobactin complexes. *Inorg Chem* 30(5):906–911
- Lopes GKB, Schulman HM, Hermes-Lima M (1999) Polyphenol tannic acid inhibits hydroxyl radical formation from Fenton reaction by complexing ferrous ions. *Biochimica Et Biophysica Acta-General Subjects* 1472(1–2):142–152
- Lorber M (1960) The occurrence of intracellular iron in Gaucher's disease. *Ann Intern Med* 53:293–305
- Lorber M (1970) Adult-type Gaucher's disease: a secondary disorder of iron metabolism. *Mt Sinai J Med* 37(4):404–417
- Lorenz DR, Cantor CR, Collins JJ (2009) A network biology approach to aging in yeast. *Proc Natl Acad Sci USA* 106(4):1145–1150
- Lorrain B, Dangles O, Genot C, Dufour C (2010) Chemical modeling of heme-induced lipid oxidation in gastric conditions and inhibition by dietary polyphenols. *J Agric Food Chem* 58(1):676–683
- Lotery A, Trump D (2007) Progress in defining the molecular biology of age related macular degeneration. *Hum Genet* 122(3–4):219–236
- Lou M, Lieb K, Selim M (2009) The relationship between hematoma iron content and perihematoma edema: an MRI study. *Cerebrovasc Dis* 27(3):266–271
- Lovell MA, Robertson JD, Teesdale WJ, Campbell JL, Markesbery WR (1998) Copper, iron and zinc in Alzheimer's disease senile plaques. *J Neurol Sci* 158(1):47–52
- Lu C, Cortopassi G (2007) Frataxin knockdown causes loss of cytoplasmic iron-sulfur cluster functions, redox alterations and induction of heme transcripts. *Arch Biochem Biophys* 457(1):111–122
- Lu JJ, Meng LH, Cai YJ, Chen Q, Tong LJ, Lin LP et al (2008) Dihydroartemisinin induces apoptosis in HL-60 leukemia cells dependent of iron and p38 mitogen-activated protein kinase activation but independent of reactive oxygen species. *Cancer Biol Ther* 7(7):1017–1023
- Lüdtke N, Panzeri S, Brown M, Broomhead DS, Knowles J, Montemurro MA et al (2008) Information-theoretic Sensitivity Analysis: a general method for credit assignment in complex networks. *J Roy Soc Interface* 5(19):223–235. doi:10.1098/rsif.2007.1079
- Luft FC (2006) Doxorubicin toxicity in the Iron Age. *J Mol Med* 84(7):529–531
- Lukosz M, Jakob S, Buchner N, Zschauer TC, Altschmied J, Haendeler J (2010) Nuclear redox signaling. *Antioxid Redox Signal* 12(6):713–742
- Lumsden AL, Henshall TL, Dayan S, Lardelli MT, Richards RI (2007) Huntingtin-deficient zebrafish exhibit defects in iron utilization and development. *Hum Mol Genet* 16(16):1905–1920
- Lund LG, Aust AE (1991) Iron-catalyzed reactions may be responsible for the biochemical and biological effects of asbestos. *Biofactors* 3(2):83–89
- Lund LG, Aust AE (1992) Iron mobilization from crocidolite asbestos greatly enhances crocidolite-dependent formation of DNA single-strand breaks in phi X174 RFI DNA. *Carcinogenesis* 13(4):637–642
- Lusis AJ (2000) Atherosclerosis. *Nature* 407(6801):233–241
- Lwin A, Orvisky E, Goker-Alpan O, LaMarca ME, Sidransky E (2004) Glucocerebrosidase mutations in subjects with parkinsonism. *Mol Genet Metab* 81(1):70–73
- Ma'ayan A (2009) Insights into the organization of biochemical regulatory networks using graph theory analyses. *J Biol Chem* 284(9):5451–5455
- Ma H, Sorokin A, Mazein A, Selkov A, Selkov E, Demin O et al (2007) The Edinburgh human metabolic network reconstruction and its functional analysis. *Mol Syst Biol* 3:135
- Ma H, Wang J, Abdel-Rahman SZ, Boor PJ, Khan MF (2008) Oxidative DNA damage and its repair in rat spleen following subchronic exposure to aniline. *Toxicol Appl Pharmacol* 233(2):247–253

- Ma HW, Zeng AP (2003) The connectivity structure, giant strong component and centrality of metabolic networks. *Bioinformatics* 19(11):1423–1430
- Ma HW, Zhao XM, Yuan YJ, Zeng AP (2004) Decomposition of metabolic network into functional modules based on the global connectivity structure of reaction graph. *Bioinformatics* 20(12):1870–1876
- Ma YS, Wu SB, Lee WY, Cheng JS, Wei YH (2009) Response to the increase of oxidative stress and mutation of mitochondrial DNA in aging. *Biochim Biophys Acta* 1790(10):1021–1029
- MacKenzie EL, Iwasaki K, Tsuji Y (2008) Intracellular iron transport and storage: from molecular mechanisms to health implications. *Antioxid Redox Signal* 10(6):997–1030
- Madamanchi NR, Hakim ZS, Runge MS (2005a) Oxidative stress in atherogenesis and arterial thrombosis: the disconnect between cellular studies and clinical outcomes. *J Thromb Haemost* 3(2):254–267
- Madamanchi NR, Vendrov A, Runge MS (2005b) Oxidative stress and vascular disease. *Arterioscler Thromb Vasc Biol* 25(1):29–38
- Maeda S, Omata M (2008) Inflammation and cancer: role of nuclear factor-kappaB activation. *Cancer Sci* 99(5):836–842
- Maguire JL, deVeber G, Parkin PC (2007) Association between iron-deficiency anemia and stroke in young children. *Pediatrics* 120(5):1053–1057
- Maher B (2008) The case of the missing heritability. *Nature* 456(7218):18–21
- Mahesh S, Ginzburg Y, Verma A (2008) Iron overload in myelodysplastic syndromes. *Leuk Lymphoma* 49(3):427–438
- Mahmoud A, Musallam KM, Taher AT (2008) Iron and hepatitis C: what can we learn from thalassemia major? *Liver Int* 28(6):892–893; author reply 893
- Maioli MA, Alves LC, Campanini AL, Lima MC, Dorta DJ, Groppo M et al (2010) Iron chelating-mediated antioxidant activity of *Plectranthus barbatus* extract on mitochondria. *Food Chem*
- Mair SM, Weiss G (2009) New pharmacological concepts for the treatment of iron overload disorders. *Curr Med Chem* 16(5):576–590
- Majumdar AP, Banerjee S, Nautiyal J, Patel BB, Patel V, Du J et al (2009) Curcumin synergizes with resveratrol to inhibit colon cancer. *Nutr Cancer* 61(4):544–553
- Malcovati L (2009) Red blood cell transfusion therapy and iron chelation in patients with myelodysplastic syndromes. *Clin Lymphoma Myeloma* 9(Suppl 3):S305–S311
- Malecki EA, Connor JR (2002) The case for iron chelation and/or antioxidant therapy in Alzheimer's disease. *Drug Dev Res* 56(3):526–530
- Mallucci G, Dickinson A, Linehan J, Klohn PC, Brandner S, Collinge J (2003) Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science* 302(5646):871–874
- Mallucci GR, White MD, Farmer M, Dickinson A, Khatun H, Powell AD et al (2007) Targeting cellular prion protein reverses early cognitive deficits and neurophysiological dysfunction in prion-infected mice. *Neuron* 53(3):325–335
- Manach C, Scalbert A, Morand C, Remesy C, Jimenez L (2004) Polyphenols: food sources and bioavailability. *Am J Clin Nutr* 79(5):727–747
- Manach C, Mazur A, Scalbert A (2005a) Polyphenols and prevention of cardiovascular diseases. *Curr Opin Lipidol* 16(1):77–84
- Manach C, Williamson G, Morand C, Scalbert A, Remesy C (2005b) Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* 81(1 Suppl):230S–242S
- Managbanag JR, Witten TM, Bonchev D, Fox LA, Tsuchiya M, Kennedy BK et al (2008) Shortest-path network analysis is a useful approach toward identifying genetic determinants of longevity. *PLoS One* 3(11):e3802
- Mancino AM, Hindo SS, Kochi A, Lim MH (2009) Effects of clioquinol on metal-triggered amyloid-beta aggregation revisited. *Inorg Chem* 48(20):9596–9598
- Mandel S, Grunblatt E, Riederer P, Youdim MB (2003) Genes and oxidative stress in Parkinsonism: cDNA microarray studies. *Adv Neurol* 91:123–132
- Mandel S, Amit T, Bar-Am O, Youdim MB (2007) Iron dysregulation in Alzheimer's disease: multimodal brain permeable iron chelating drugs, possessing neuroprotective-neurorescue and amyloid precursor protein-processing regulatory activities as therapeutic agents. *Prog Neurobiol* 82(6):348–360
- Mandel S, Weinreb O, Amit T, Youdim MB (2005) Mechanism of neuroprotective action of the anti-Parkinson drug rasagiline and its derivatives. *Brain Res Brain Res Rev* 48(2):379–387
- Mandel S, Weinreb O, Amit T, Youdim MB (2004) Cell signaling pathways in the neuroprotective actions of the green tea polyphenol (-)-epigallocatechin-3-gallate: implications for neurodegenerative diseases. *J Neurochem* 88(6):1555–1569
- Mandel S, Youdim MBH (2004) Catechin polyphenols: neurodegeneration and neuroprotection in neurodegenerative diseases. *Free Radic Biol Med* 37(3):304–317
- Mandel SA, Amit T, Weinreb O, Reznichenko L, Youdim MB (2008) Simultaneous manipulation of multiple brain targets by green tea catechins: a potential neuroprotective strategy for Alzheimer and Parkinson diseases. *CNS Neurosci Ther* 14(4):352–365
- Mandemakers W, Morais VA, De Strooper B (2007) A cell biological perspective on mitochondrial dysfunction in Parkinson disease and other neurodegenerative diseases. *J Cell Sci* 120(Pt 10):1707–1716
- Mansour MA (2000) Protective effects of thymoquinone and desferrioxamine against hepatotoxicity of carbon tetrachloride in mice. *Life Sci* 66(26):2583–2591
- Markesbery WR (1997) Oxidative stress hypothesis in Alzheimer's disease. *Free Radic Biol Med* 23(1):134–147
- Markesbery WR, Lovell MA (1998) 4-hydroxynonenal, a product of lipid peroxidation, is increased in the brain in Alzheimer's disease. *Neurobiol Aging* 19(1):33–36
- Marmolino D, Acquaviva F (2009) Friedreich's Ataxia: from the (GAA)_n repeat mediated silencing to new promising molecules for therapy. *Cerebellum* 8(3):245–259
- Marniemi J, Alanen E, Impivaara O, Seppanen R, Hakala P, Rajala T et al (2005) Dietary and serum vitamins and minerals as predictors of myocardial infarction and stroke in elderly subjects. *Nutr Metab Cardiovasc Dis* 15(3):188–197
- Martin-Murphy BV, Holt MP, Ju C (2010) The role of damage associated molecular pattern molecules in acetaminophen-induced liver injury in mice. *Toxicol Lett* 192(3):387–394
- Martin GS, Mannino DM, Eaton S, Moss M (2003) The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 348(16):1546–1554
- Maruyama H, Morino H, Ito H, Izumi Y, Kato H, Watanabe Y et al (2010) Mutations of optineurin in amyotrophic lateral sclerosis. *Nature* 465(7295):223–226
- Maruyama Y, Nakayama M, Yoshimura K, Nakano H, Yamamoto H, Yokoyama K et al (2007) Effect of repeated intravenous iron administration in haemodialysis patients on serum 8-hydroxy-2'-deoxyguanosine levels. *Nephrol Dial Transpl* 22(5):1407–1412
- Marx JJM (2002) Prevention of organ failure in hereditary haemochromatosis. *Neth J Med* 60(11):419–422
- Marx JJM, Kartikasari AER, Georgiou NA (2008) Can iron chelators influence the progression of atherosclerosis? *Hemoglobin* 32(1–2):123–134
- Mascitelli L, Pezzetta F (2006) High iron stores and risk of ischemic stroke in persons with metabolic syndrome. *Intern Med* 45(5):347
- Mascitelli L, Pezzetta F, Goldstein MR (2009) Iron, type 2 diabetes mellitus, and Alzheimer's disease. *Cell Mol Life Sci* 66:2943

- Massie HR, Aiello VR, Banziger V (1983) Iron accumulation and lipid peroxidation in aging C57bl/6j mice. *Exp Gerontol* 18(4):277–285
- Masubuchi Y, Suda C, Horie T (2005) Involvement of mitochondrial permeability transition in acetaminophen-induced liver injury in mice. *J Hepatol* 42(1):110–116
- Mateo-Gallego R, Calmarza P, Jarauta E, Burillo E, Cenarro A, Civeira F (2010) Serum ferritin is a major determinant of lipid phenotype in familial combined hyperlipidemia and familial hypertriglyceridemia. *Metabolism* 59(2):154–158
- Matthews AJ, Vercellotti GM, Menchaca HJ, Bloch PH, Michalek VN, Marker PH et al (1997) Iron and atherosclerosis: inhibition by the iron chelator deferiprone (L1). *J Surg Res* 73(1):35–40
- Mattson MP (2004) Metal-catalyzed disruption of membrane protein and lipid signaling in the pathogenesis of neurodegenerative disorders. *Ann N Y Acad Sci* 1012:37–50
- Mattson MP (2006) Neuronal life-and-death signaling, apoptosis, and neurodegenerative disorders. *Antioxid Redox Signal* 8(11–12):1997–2006
- Mattson MP (2008a) Awareness of hormesis will enhance future research in basic and applied neuroscience. *Crit Rev Toxicol* 38(7):633–639
- Mattson MP (2008b) Hormesis defined. *Ageing Res Rev* 7(1):1–7
- Matusch A, Depboylu C, Palm C, Wu B, Hoglinger GU, Schafer MK et al (2010) Cerebral bioimaging of Cu, Fe, Zn, and Mn in the MPTP mouse model of Parkinson's disease using laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS). *J Am Soc Mass Spectrom* 21(1):161–171
- Mazumdar M, Heeney MM, Sox CM, Lieu TA (2007) Preventing stroke among children with sickle cell anemia: an analysis of strategies that involve transcranial Doppler testing and chronic transfusion. *Pediatrics* 120(4):e1107–1116
- McDermid JM, Jaye A, Schim van der Loeff MF, Todd J, Bates C, Austin S et al (2007) Elevated iron status strongly predicts mortality in West African adults with HIV infection. *J Acquir Immune Defic Syndr* 46(4):498–507
- McDermid JM, Prentice AM (2006) Iron and infection: effects of host iron status and the iron-regulatory genes haptoglobin and NRAMPI (SLC11A1) on host-pathogen interactions in tuberculosis and HIV. *Clin Sci (Lond)* 110(5):503–524
- McDermid JM, van der Loeff MF, Jaye A, Hennig BJ, Bates C, Todd J et al (2009) Mortality in HIV infection is independently predicted by host iron status and SLC11A1 and HP genotypes, with new evidence of a gene-nutrient interaction. *Am J Clin Nutr* 90(1):225–233
- McLaren GD, Gordeuk VR (2009) Hereditary hemochromatosis: insights from the hemochromatosis and iron overload screening (HEIRS) study. *Hematol Am Soc Hematol Educ Program* 195–206
- McLean CA (2008) Review. The neuropathology of kuru and variant Creutzfeldt-Jakob disease. *Philos Trans R Soc Lond B Biol Sci* 363(1510):3685–3687
- McNeill A, Birchall D, Hayflick SJ, Gregory A, Schenk JF, Zimmerman EA et al (2008) T2* and FSE MRI distinguishes four subtypes of neurodegeneration with brain iron accumulation. *Neurology* 70(18):1614–1619
- McRae R, Bagchi P, Sumalekshmy S, Fahrni CJ (2009) In situ imaging of metals in cells and tissues. *Chem Rev* 109(10):4780–4827
- Mehdiratta M, Kumar S, Hackney D, Schlaug G, Selim M (2008) Association between serum ferritin level and perihematoma edema volume in patients with spontaneous intracerebral hemorrhage. *Stroke* 39(4):1165–1170
- Mehrotra B, Li XJ, Kamal A, Hoops S, Swainston N, Mendes P (2010) B-Net: a relational data model for biochemical networks. *BMC Bioinform*
- Mehta SH, Webb RC, Ergul A, Tawak A, Dorrance AM (2004) Neuroprotection by tempol in a model of iron-induced oxidative stress in acute ischemic stroke. *Am J Physiol* 286(2):R283–R288
- Mellert W, Deckardt K, Gemhardt C, Zwierner-Baier I, Jackh R, van Ravenzwaay B (2004) Aniline: early indicators of toxicity in male rats and their relevance to spleen carcinogenicity. *Hum Exp Toxicol* 23(8):379–389
- Mendes P, Kell DB (1996) On the analysis of the inverse problem of metabolic pathways using artificial neural networks. *Biosystems* 38:15–28
- Mendes P, Kell DB (1998) Non-linear optimization of biochemical pathways: applications to metabolic engineering and parameter estimation. *Bioinformatics* 14:869–883
- Menke A, Fernández-Real JM, Muntner P, Guallar E (2009) The association of biomarkers of iron status with peripheral arterial disease in US adults. *BMC Cardiovasc Disord* 9(1):34
- Messer JG, Kilbarger AK, Erikson KM, Kipp DE (2009) Iron overload alters iron-regulatory genes and proteins, down-regulates osteoblastic phenotype, and is associated with apoptosis in fetal rat calvaria cultures. *Bone*
- Messner DJ, Sivam G, Kowdley KV (2009) Curcumin reduces the toxic effects of iron loading in rat liver epithelial cells. *Liver Int* 29(1):63–72
- Méthy D, Bertrand N, Prigent-Tessier A, Mossiat C, Stanimirovic D, Beley A et al (2008) Beneficial effect of dipyrindyl, a liposoluble iron chelator against focal cerebral ischemia: in vivo and in vitro evidence of protection of cerebral endothelial cells. *Brain Res* 1193:136–142
- Meyer D (2006) Iron chelation as therapy for HIV and *Mycobacterium tuberculosis* co-infection under conditions of iron overload. *Curr Pharm Des* 12(16):1943–1947
- Michalik A, Van Broeckhoven C (2003) Pathogenesis of polyglutamine disorders: aggregation revisited. *Hum Mol Genet* 12(2):R173–R186
- Miele G, Manson J, Clinton M (2001) A novel erythroid-specific marker of transmissible spongiform encephalopathies. *Nature Med* 7(3):361–364
- Miethke M, Marahiel MA (2007) Siderophore-based iron acquisition and pathogen control. *Microbiol Mol Biol Rev* 71(3):413–451
- Migliore L, Fontana I, Colognato R, Coppede F, Siciliano G, Murri L (2005) Searching for the role and the most suitable biomarkers of oxidative stress in Alzheimer's disease and in other neurodegenerative diseases. *Neurobiol Aging* 26(5):587–595
- Milhavet O, Lehmann S (2002) Oxidative stress and the prion protein in transmissible spongiform encephalopathies. *Brain Res Rev* 38(3):328–339
- Millán M, Sobrino T, Arenillas JF, Rodríguez-Yáñez M, Garcia M, Nombela F et al (2008) Biological signatures of brain damage associated with high serum ferritin levels in patients with acute ischemic stroke and thrombolytic treatment. *Dis Markers* 25(3):181–188
- Millan M, Sobrino T, Castellanos M, Nombela F, Arenillas JF, Riva E et al (2007) Increased body iron stores are associated with poor outcome after thrombolytic treatment in acute stroke. *Stroke* 38(1):90–95
- Millan MJ (2006) Multi-target strategies for the improved treatment of depressive states: Conceptual foundations and neuronal substrates, drug discovery and therapeutic application. *Pharmacol Ther* 110(2):135–370
- Miller DM, Buettner GR, Aust SD (1990) Transition metals as catalysts of "autoxidation" reactions. *Free Radic Biol Med* 8(1):95–108
- Miller ER III, Pastor-Barriuso R, Dalal D, Riemersma RA, Appel LJ, Guallar E (2005) Meta-analysis: high-dosage vitamin E supplementation may increase all-cause mortality. *Ann Intern Med* 142(1):37–46

- Miller RL, James-Kracke M, Sun GY, Sun AY (2009) Oxidative and inflammatory pathways in Parkinson's disease. *Neurochem Res* 34(1):55–65
- Millerot-Serruot E, Bertrand N, Mossiat C, Faure P, Prigent-Tessier A, Garnier P et al (2008) Temporal changes in free iron levels after brain ischemia Relevance to the timing of iron chelation therapy in stroke. *Neurochem Int* 52(8):1442–1448
- Milo R, Jorgensen P, Moran U, Weber G, Springer M (2010) BioNumbers—the database of key numbers in molecular and cell biology. *Nucleic Acids Res* 38(Database issue):D750–D753
- Milton NGN (2004) Role of hydrogen peroxide in the aetiology of Alzheimer's disease—implications for treatment. *Drugs Aging* 21(2):81–100
- Minihane AM, Rimbach G (2002) Iron absorption and the iron binding and anti-oxidant properties of phytic acid. *Int J Food Sci Technol* 37(7):741–748
- Minqin R, Watt F, Huat BTK, Halliwell B (2003) Correlation of iron and zinc levels with lesion depth in newly formed atherosclerotic lesions. *Free Radic Biol Med* 34(6):746–752
- Miranda S, Opazo C, Larrondo LF, Munoz FJ, Ruiz F, Leighton F et al (2000) The role of oxidative stress in the toxicity induced by amyloid beta-peptide in Alzheimer's disease. *Progr Neurobiol* 62(6):633–648
- Mirre E, Brousse V, Berteloot L, Lambot-Juhan K, Verlhac S, Boulat C et al (2010) Feasibility and efficacy of chronic transfusion for stroke prevention in children with sickle cell disease. *Eur J Haematol* 84(3):259–265
- Mirshafiey A, Mohsenzadegan M (2009) Antioxidant therapy in multiple sclerosis. *Immunopharmacol Immunotoxicol* 31(1):13–29
- Mishra V (2007) Oxidative stress and role of antioxidant supplementation in critical illness. *Clin Lab* 53(3–4):199–209
- Mitchell KM, Dotson AL, Cool KM, Chakrabarty A, Benedict SH, LeVine SM (2007) Deferiprone, an orally deliverable iron chelator, ameliorates experimental autoimmune encephalomyelitis. *Mult Scler* 13(9):1118–1126
- Miyamoto S, Kuwata G, Imai M, Nagao A, Terao J (2000) Protective effect of phytic acid hydrolysis products on iron-induced lipid peroxidation of liposomal membranes. *Lipids* 35(12):1411–1413
- Mladěnka P, Kalinowski DS, Hašková P, Bobrovová Z, Hrdina R, Šimůnek T et al (2009) The novel iron chelator, 2-pyridylcarboxaldehyde 2-thiophenecarboxyl hydrazone, reduces catecholamine-mediated myocardial toxicity. *Chem Res Toxicol* 22(1):208–217
- Moalem S, Percy ME, Andrews DF, Kruck TP, Wong S, Dalton AJ et al (2000) Are hereditary hemochromatosis mutations involved in Alzheimer disease? *Am J Med Genet* 93(1):58–66
- Mogilner A, Wollman R, Marshall WF (2006) Quantitative modeling in cell biology: what is it good for? *Dev Cell* 11(3):279–287
- Mohkam M, Shamsian BS, Gharib A, Nariman S, Arzanian MT (2008) Early markers of renal dysfunction in patients with beta-thalassemia major. *Pediatr Nephrol* 23(6):971–976
- Moles CG, Mendes P, Banga JR (2003) Parameter estimation in biochemical pathways: a comparison of global optimization methods. *Genome Res* 13(11):2467–2474
- Molfino A, Kushta I, Tommasi V, Fanelli FR, Muscaritoli M (2009) Amyotrophic lateral sclerosis, enteral nutrition and the risk of iron overload. *J Neurol* 256(6):1015–1016
- Moore JH, Asselbergs FW, Williams SM (2010) Bioinformatics challenges for genome-wide association studies. *Bioinformatics* 26(4):445–455
- Moreira PI, Duarte AI, Santos MS, Rego AC, Oliveira CR (2009) An integrative view of the role of oxidative stress, mitochondria and insulin in Alzheimer's disease. *J Alzheimers Dis* 16(4):741–761
- Morel I, Lescoat G, Cillard P, Cillard J (1994) Role of flavonoids and iron chelation in antioxidant action. *Methods Enzymol* 234:437–443
- Morgan C, Colombres M, Nunez MT, Inestrosa NC (2004) Structure and function of amyloid in Alzheimer's disease. *Prog Neurobiol* 74(6):323–349
- Morgan MA, Hoffbrand AV, Laulich M, Luck W, Knowles S (1983) Serum ferritin concentration in Gaucher's disease. *Br Med J (Clin Res Ed)* 286(6381):1864
- Morohashi M, Winn AE, Borisuk MT, Bolouri H, Doyle J, Kitano H (2002) Robustness as a measure of plausibility in models of biochemical networks. *J Theor Biol* 216(1):19–30
- Morphy R, Rankovic Z (2007) Fragments, network biology and designing multiple ligands. *Drug Discov Today* 12(3–4):156–160
- Mossman BT, Churg A (1998) Mechanisms in the pathogenesis of asbestosis and silicosis. *Am J Respir Crit Care Med* 157(5 Pt 1):1666–1680
- Mossman BT, Marsh JP, Shatos MA, Doherty J, Gilbert R, Hill S (1987) Implication of active oxygen species as second messengers of asbestos toxicity. *Drug Chem Toxicol* 10(1–2):157–180
- Mu D, Chang YS, Vexler ZS, Ferriero DM (2005) Hypoxia-inducible factor 1alpha and erythropoietin upregulation with deferoxamine salvage after neonatal stroke. *Exp Neurol* 195(2):407–415
- Mueller S, Afdhal NH, Schuppan D (2006) Iron, HCV, and liver cancer: hard metal setting the pace? *Gastroenterology* 130(7):2229–2234
- Muldoon DF, Hassoun EA, Stohs SJ (1996) Role of iron in ricin-induced lipid peroxidation and superoxide production. *Res Commun Mol Pathol Pharmacol* 92(1):107–118
- Mullenix PS, Andersen CA, Starnes BW (2005) Atherosclerosis as inflammation. *Ann Vasc Surg* 19(1):130–138
- Muller FL, Lustgarten MS, Jang Y, Richardson A, Van Remmen H (2007) Trends in oxidative aging theories. *Free Radic Biol Med* 43(4):477–503
- Muñoz X, Cruz MJ, Freixa A, Guardino X, Morell F (2009) Occupational asthma caused by metal arc welding of iron. *Respiration* 78(4):455–459
- Murakami K, Ishida K, Watakabe K, Tsubouchi R, Naruse M, Yoshino M (2006) Maltol/iron-mediated apoptosis in HL60 cells: participation of reactive oxygen species. *Toxicol Lett* 161(2):102–107
- Murphy MP, Packer MA, Scarlett JL, Martin SW (1998) Peroxynitrite: a biologically significant oxidant. *Gen Pharmacol* 31(2):179–186
- Murray AR, Kisin E, Leonard SS, Young SH, Kommineni C, Kagan VE et al (2009) Oxidative stress and inflammatory response in dermal toxicity of single-walled carbon nanotubes. *Toxicology* 257(3):161–171
- Nagababu E, Rifkind JM, Boindala S, Nakka L (2010) Assessment of antioxidant activity of eugenol *in vitro* and *in vivo*. *Methods Mol Biol* 610:165–180
- Nagai Y, Fujikake N, Ohno K, Higashiyama H, Popiel HA, Rahadian J et al (2003) Prevention of polyglutamine oligomerization and neurodegeneration by the peptide inhibitor QBP1 in *Drosophila*. *Hum Mol Genet* 12(11):1253–1259
- Naghii MR (2002) Sulfur mustard intoxication, oxidative stress, and antioxidants. *Mil Med* 167(7):573–575
- Nahon P, Sutton A, Rufat P, Zioli M, Thabut G, Schischmanoff PO et al (2008) Liver iron, HFE gene mutations, and hepatocellular carcinoma occurrence in patients with cirrhosis. *Gastroenterology* 134(1):102–110
- Nakamura T, Keep RF, Hua Y, Nagao S, Hoff JT, Xi G (2006) Iron-induced oxidative brain injury after experimental intracerebral hemorrhage. *Acta Neurochir Suppl* 96:194–198
- Nakano M, Kawanishi Y, Kamohara S, Uchida Y, Shiota M, Inatomi Y et al (2003) Oxidative DNA damage (8-hydroxydeoxyguanosine) and body iron status: a study on 2507 healthy people. *Free Radic Biol Med* 35(7):826–832

- Napier I, Ponka P, Richardson DR (2005) Iron trafficking in the mitochondrion: novel pathways revealed by disease. *Blood* 105(5):1867–1874
- Napoli E, Taroni F, Cortopassi GA (2006) Frataxin, iron-sulfur clusters, heme, ROS, and aging. *Antioxid Redox Signal* 8(3–4):506–516
- Narula MK, Ahuja GK, Whig J, Narang AP, Soni RK (2007) Status of lipid peroxidation and plasma iron level in bronchial asthmatic patients. *Indian J Physiol Pharmacol* 51(3):289–292
- Nathans AB, Neff MJ, Jurkovich GJ, Klotz P, Farver K, Ruzinski JT et al (2002) Randomized, prospective trial of antioxidant supplementation in critically ill surgical patients. *Ann Surg* 236(6):814–822
- Natoli M, Felsani A, Ferruzza S, Sambuy Y, Canali R, Scarino ML (2009) Mechanisms of defence from Fe(II) toxicity in human intestinal Caco-2 cells. *Toxicol In Vitro* 23(8):1510–1515
- Neale BM, Fagerness J, Reynolds R, Sobrin L, Parker M, Raychaudhuri S et al (2010) Genome-wide association study of advanced age-related macular degeneration identifies a role of the hepatic lipase gene (LIPC). *Proc Natl Acad Sci USA*
- Neema M, Arora A, Healy BC, Guss ZD, Brass SD, Duan Y et al (2009a) Deep gray matter involvement on brain MRI scans is associated with clinical progression in multiple sclerosis. *J Neuroimaging* 19(1):3–8
- Neema M, Goldberg-Zimring D, Guss ZD, Healy BC, Guttmann CR, Houtchens MK et al (2009b) 3 T MRI relaxometry detects T2 prolongation in the cerebral normal-appearing white matter in multiple sclerosis. *Neuroimage* 46(3):633–641
- Neema M, Stankiewicz J, Arora A, Dandamudi VS, Batt CE, Guss ZD et al (2007a) T1- and T2-based MRI measures of diffuse gray matter and white matter damage in patients with multiple sclerosis. *J Neuroimaging* 17(Suppl 1):16S–21S
- Neema M, Stankiewicz J, Arora A, Guss ZD, Bakshi R (2007b) MRI in multiple sclerosis: what's inside the toolbox? *Neurotherapeutics* 4(4):602–617
- Nelson DE, Ihekweba AEC, Elliott M, Gibney CA, Foreman BE, Nelson G et al (2004) Oscillations in NF- κ B signalling control the dynamics of target gene expression. *Science* 306:704–708
- Nemery B (1990) Metal toxicity and the respiratory tract. *Eur Respir J* 3(2):202–219
- Neumann H, Wang K, Davis L, Garcia-Alai M, Chin JW (2010) Encoding multiple unnatural amino acids via evolution of a quadruplet-decoding ribosome. *Nature* 464(7287):441–444
- Neumann KF, Rojo L, Navarrete LP, Farias G, Reyes P, Maccioni RB (2008) Insulin resistance and Alzheimer's disease: molecular links & clinical implications. *Curr Alzheimer Res* 5(5):438–447
- Neupane GP, Kim DM (2010) In vitro time-kill activities of ciprofloxacin alone and in combination with the iron chelator deferasirox against *Vibrio vulnificus*. *Eur J Clin Microbiol Infect Dis* 29(4):407–410
- Nguyen HB, Rivers EP, Abrahamian FM, Moran GJ, Abraham E, Trzeciak S et al (2006) Severe sepsis and septic shock: Review of the literature and emergency department management guidelines. *Ann Emerg Med* 48(1):28–54
- Nguyen T, Hamby A, Massa SM (2005) Clonidine down-regulates mutant huntingtin expression in vitro and mitigates pathology in a Huntington's disease mouse model. *Proc Natl Acad Sci USA* 102(33):11840–11845
- Nidhi G, Glick M, Davies JW, Jenkins JL (2006) Prediction of biological targets for compounds using multiple-category Bayesian models trained on chemogenomics databases. *J Chem Inf Model* 46(3):1124–1133
- Nighoghossian N, Wiart M, Berthezene Y (2008) Novel applications of magnetic resonance imaging to image tissue inflammation after stroke. *J Neuroimaging* 18(4):349–352
- Nigro J, Osman N, Dart AM, Little PJ (2006) Insulin resistance and atherosclerosis. *Endocr Rev* 27(3):242–259
- Nijveldt RJ, van Nood E, van Hoorn DEC, Boelens PG, van Norren K, van Leeuwen PAM (2001) Flavonoids: a review of probable mechanisms of action and potential applications. *Am J Clin Nutr* 74(4):418–425
- Nishi T, Shimizu N, Hiramoto M, Sato I, Yamaguchi Y, Hasegawa M et al (2002) Spatial redox regulation of a critical cysteine residue of NF- κ B in vivo. *J Biol Chem* 277(46):44548–44556
- Nishina S, Hino K, Korenaga M, Vecchi C, Pietrangelo A, Mizukami Y et al (2008) Hepatitis C virus-induced reactive oxygen species raise hepatic iron level in mice by reducing hepcidin transcription. *Gastroenterology* 134(1):226–238
- Noble D (2002) The rise of computational biology. *Nat Rev Mol Cell Biol* 3(6):460–463
- Noble D (2006) *The music of life: biology beyond genes*. Oxford University Press, Oxford
- Noble D (2010) *Biophysics and systems biology*. *Philos Transact A Math Phys Eng Sci* 368(1914):1125–1139
- Northrop-Clewes CA (2008) Interpreting indicators of iron status during an acute phase response—lessons from malaria and human immunodeficiency virus. *Ann Clin Biochem* 45(Pt 1):18–32
- Nowicki M, Kosacka J, Spaniel-Borowski K, Borlak J (2009) Deferoxamine-induced neurite outgrowth and synapse formation in postnatal rat dorsal root ganglion (DRG) cell cultures. *Eur J Cell Biol* 88(10):551–562
- Nunomura A, Castellani RJ, Zhu X, Moreira PI, Perry G, Smith MA (2006) Involvement of oxidative stress in Alzheimer disease. *J Neuropathol Exp Neurol* 65(7):631–641
- Nunomura A, Perry G, Aliev G, Hirai K, Takeda A, Balraj EK et al (2001) Oxidative damage is the earliest event in Alzheimer disease. *J Neuropathol Exp Neurol* 60(8):759–767
- Nymark P, Wikman H, Hienonen-Kempas T, Anttila S (2008) Molecular and genetic changes in asbestos-related lung cancer. *Cancer Lett* 265(1):1–15
- O'Donnell VB, Freeman BA (2001) Interactions between nitric oxide and lipid oxidation pathways—implications for vascular disease. *Circ Res* 88(1):12–21
- O'Hagan A (2006) Bayesian analysis of computer code outputs: a tutorial. *Reliab Eng Syst Saf* 91(10–11):1290–1300
- O'Hagan S, Dunn WB, Broadhurst D, Williams R, Ashworth JA, Cameron M et al (2007) Closed-loop, multi-objective optimisation of two-dimensional gas chromatography (GCxGC-tof-MS) for serum metabolomics. *Anal Chem* 79(2):464–476
- O'Hagan S, Dunn WB, Brown M, Knowles JD, Kell DB (2005) Closed-loop, multiobjective optimisation of analytical instrumentation: gas-chromatography-time-of-flight mass spectrometry of the metabolomes of human serum and of yeast fermentations. *Anal Chem* 77:290–303
- O'Neill HA, Gakh O, Park S, Cui J, Mooney SM, Sampson M et al (2005) Assembly of human frataxin is a mechanism for detoxifying redox-active iron. *Biochemistry* 44(2):537–545
- O'Rourke D, McCarthy R, Lynch B (2008) Iron deficiency anaemia and childhood stroke. *Ir Med J* 101(6):165
- Oakley AE, Collingwood JF, Dobson J, Love G, Perrott HR, Edwardson JA et al (2007) Individual dopaminergic neurons show raised iron levels in Parkinson disease. *Neurology* 68(21):1820–1825
- Oates AC, Gorfinkiel N, Gonzalez-Gaitan M, Heisenberg CP (2009) Quantitative approaches in developmental biology. *Nat Rev Genet* 10(8):517–530
- Oda K, Kitano H (2006) A comprehensive map of the toll-like receptor signaling network. *Mol Syst Biol* 2:2006 0015
- Ogeturk M, Kus I, Kavakli A, Zararsiz I, Ilhan N, Sarsilmaz M (2004) Effects of melatonin on carbon tetrachloride-induced changes in rat serum. *J Physiol Biochem* 60(3):205–210
- Ohnuma S, Harris WA (2003) Neurogenesis and the cell cycle. *Neuron* 40(2):199–208

- Okauchi M, Hua Y, Keep RF, Morgenstern LB, Schallert T, Xi G (2010) Deferoxamine treatment for intracerebral hemorrhage in aged rats: therapeutic time window and optimal duration. *Stroke* 41(2):375–382
- Okauchi M, Hua Y, Keep RF, Morgenstern LB, Xi G (2009) Effects of deferoxamine on intracerebral hemorrhage-induced brain injury in aged rats. *Stroke* 40(5):1858–1863
- Okouchi M, Ekshyyan O, Maracine M, Aw TY (2007) Neuronal apoptosis in neurodegeneration. *Antioxid Redox Signal* 9(8):1059–1096
- Oksenberg JR, Baranzini SE, Sawcer S, Hauser SL (2008) The genetics of multiple sclerosis: SNPs to pathways to pathogenesis. *Nat Rev Genet* 9(7):516–526
- Olaleye MT, Rocha BT (2008) Acetaminophen-induced liver damage in mice: effects of some medicinal plants on the oxidative defense system. *Exp Toxicol Pathol* 59(5):319–327
- Olanow CW, Arendash GW (1994) Metals and free radicals in neurodegeneration. *Curr Opin Neurol* 7(6):548–558
- Oldroyd D (1986) The arch of knowledge: an introduction to the history of the philosophy and methodology of science. Methuen, New York
- Olivares D, Huang X, Branden L, Greig NH, Rogers JT (2009) Physiological and pathological role of alpha-synuclein in Parkinson's disease through iron mediated oxidative stress; the role of a putative iron-responsive element. *Int J Mol Sci* 10(3):1226–1260
- Omar RF, Hasinoff BB, Mejilla F, Rahimtula AD (1990) Mechanism of ochratoxin A stimulated lipid peroxidation. *Biochem Pharmacol* 40(6):1183–1191
- Ong WY, Farooqui AA (2005) Iron, neuroinflammation, and Alzheimer's disease. *J Alzheimers Dis* 8(2):183–200
- Orhan H, van Holland B, Krab B, Moeken J, Vermeulen NP, Hollander P et al (2004) Evaluation of a multi-parameter biomarker set for oxidative damage in man: increased urinary excretion of lipid, protein and DNA oxidation products after one hour of exercise. *Free Radic Res* 38(12):1269–1279
- Orrenius S, Gogvadze A, Zhivotovsky B (2007) Mitochondrial oxidative stress: implications for cell death. *Ann Rev Pharmacol Toxicol* 47:143–183
- Ossowska K, Śmiałowska M, Kuter K, Wierońska J, Zięba B, Wardas J et al (2006) Degeneration of dopaminergic mesocortical neurons and activation of compensatory processes induced by a long-term paraquat administration in rats: implications for Parkinson's disease. *Neuroscience* 141(4):2155–2165
- Ostrova-Golts N, Petrucelli L, Hardy J, Lee JM, Farer M, Wolozin B (2000) The A53T alpha-synuclein mutation increases iron-dependent aggregation and toxicity. *J Neurosci* 20(16):6048–6054
- Othman AI, El-Missiry MA, Amer MA, Arafa M (2008) Melatonin controls oxidative stress and modulates iron, ferritin, and transferrin levels in adriamycin treated rats. *Life Sci* 83(15–16):563–568
- Otogawa K, Ogawa T, Shiga R, Nakatani K, Ikeda K, Nakajima Y et al (2008) Attenuation of acute and chronic liver injury in rats by iron-deficient diet. *Am J Physiol Regul Integr Comp Physiol* 294(2):R311–320
- Pachauri V, Saxena G, Mehta A, Mishra D, Flora SJ (2009) Combinational chelation therapy abrogates lead-induced neurodegeneration in rats. *Toxicol Appl Pharmacol* 240(2):255–264
- Pacher P, Beckman JS, Liaudet L (2007) Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87(1):315–424
- Packard RR, Libby P (2008) Inflammation in atherosclerosis: from vascular biology to biomarker discovery and risk prediction. *Clin Chem* 54(1):24–38
- Pae HO, Lee YC, Chung HT (2008) Heme oxygenase-1 and carbon monoxide: emerging therapeutic targets in inflammation and allergy. *Recent Pat Inflamm Allergy Drug Discov* 2(3):159–165
- Palau F (2001) Friedreich's ataxia and frataxin: molecular genetics, evolution and pathogenesis (Review). *Int J Mol Med* 7(6):581–589
- Palsson BØ (2006) Systems biology: properties of reconstructed networks. Cambridge University Press, Cambridge
- Pamplona R, Naudi A, Gavin R, Pastrana MA, Sajani G, Ilieva EV et al (2008) Increased oxidation, glycooxidation, and lipoxidation of brain proteins in prion disease. *Free Radic Biol Med* 45(8):1159–1166
- Pan YJ, Hopkins RG, Loo G (2004) Increased GADD153 gene expression during iron chelation-induced apoptosis in Jurkat T-lymphocytes. *Biochim Biophys Acta* 1691(1):41–50
- Pandey M, Gupta S (2009) Green tea and prostate cancer: from bench to clinic. *Front Biosci (Elite Ed)* 1:13–25
- Pandolfo M (2009) Friedreich ataxia: the clinical picture. *J Neurol* 256(Suppl 1):3–8.
- Pandolfo M, Pastore A (2009) The pathogenesis of Friedreich ataxia and the structure and function of frataxin. *J Neurol* 256(Suppl 1):9–17
- Panduri V, Surapureddi S, Soberanes S, Weitzman SA, Chandel N, Kamp DW (2006) P53 mediates amosite asbestos-induced alveolar epithelial cell mitochondria-regulated apoptosis. *Am J Respir Cell Mol Biol* 34(4):443–452
- Pantopoulos K (2008) Function of the hemochromatosis protein HFE: Lessons from animal models. *World J Gastroenterol* 14(45):6893–6901
- Paoletti R, Gotto AM Jr, Hajjar DP (2004) Inflammation in atherosclerosis and implications for therapy. *Circulation* 109(23 Suppl 1):III20–III26
- Papanikolaou G, Pantopoulos K (2005) Iron metabolism and toxicity. *Toxicol Appl Pharmacol* 202(2):199–211
- Paris I, Martinez-Alvarado P, Cárdenas S, Perez-Pastene C, Graumann R, Fuentes P et al (2005) Dopamine-dependent iron toxicity in cells derived from rat hypothalamus. *Chem Res Toxicol* 18(3):415–419
- Park SH, Aust AE (1998) Participation of iron and nitric oxide in the mutagenicity of asbestos in hgp⁺, gpt⁺ Chinese hamster V79 cells. *Cancer Res* 58(6):1144–1148
- Park S, Gakh O, Mooney SM, Isaya G (2002) The ferroxidase activity of yeast frataxin. *J Biol Chem* 277(41):38589–38595
- Park JH, Lee KH, Kim TY, Lee SY (2007) Metabolic engineering of *Escherichia coli* for the production of L-valine based on transcriptome analysis and in silico gene knockout simulation. *Proc Natl Acad Sci USA* 104(19):7797–7802
- Park JH, Lee SY, Kim TY, Kim HU (2008a) Application of systems biology for bioprocess development. *Trends Biotechnol* 26(8):404–412
- Park SJ, Kim NH, Jeong BH, Jin JK, Choi JK, Park YJ et al (2008b) The effect of Fenton reaction on protease-resistant prion protein (PrP^{Sc}) degradation and scrapie infectivity. *Brain Res* 1238:172–180
- Park SJ, Chung HY, Lee JH (2010) Rapid in vivo screening system for anti-oxidant activity using bacterial redox sensor strains. *J Appl Microbiol* 108:1217–1225
- Pasinelli P, Brown RH (2006) Molecular biology of amyotrophic lateral sclerosis: insights from genetics. *Nat Rev Neurosci* 7(9):710–723
- Pastore C, Franzese M, Sica F, Temussi P, Pastore A (2007) Understanding the binding properties of an unusual metal-binding protein—a study of bacterial frataxin. *FEBS J* 274(16):4199–4210
- Patel N, Adewoyin T, Chong NV (2008) Age-related macular degeneration: a perspective on genetic studies. *Eye (Lond)* 22(6):768–776
- Patel PI, Isaya G (2001) Friedreich ataxia: from GAA triplet-repeat expansion to frataxin deficiency. *Am J Hum Genet* 69(1):15–24

- Paterniti I, Genovese T, Crisafulli C, Mazzon E, Di Paola R, Galuppo M et al (2009) Treatment with green tea extract attenuates secondary inflammatory response in an experimental model of spinal cord trauma. *Naunyn Schmiedebergs Arch Pharmacol* 380(2):179–192
- Patil BS, Jayaprakasha GK, Chidambara Murthy KN, Vikram A (2009) Bioactive compounds: historical perspectives, opportunities, and challenges. *J Agric Food Chem* 57(18):8142–8160
- Patt A, Horesh IR, Berger EM, Harken AH, Repine JE (1990) Iron depletion or chelation reduces ischemia/reperfusion-induced edema in gerbil brains. *J Pediatr Surg* 25(2):224–227 (discussion 227–228)
- Pavlica S, Gebhardt R (2010) Protective effects of flavonoids and two metabolites against oxidative stress in neuronal PC12 cells. *Life Sci* 86(3–4):79–86
- Pavlovic R, Santaniello E (2007) Peroxynitrite and nitrosoperoxy-carbonate, a tightly connected oxidizing-nitrating couple in the reactive nitrogen-oxygen species family: new perspectives for protection from radical-promoted injury by flavonoids. *J Pharm Pharmacol* 59(12):1687–1695
- Pedchenko TV, LeVine SM (1998) Desferrioxamine suppresses experimental allergic encephalomyelitis induced by MBP in SJL mice. *J Neuroimmunol* 84(2):188–197
- Peden AH, Ironside JW (2004) Review: pathology of variant Creutzfeldt-Jakob disease. *Folia Neuropathol* 42(Suppl A):85–91
- Peng CT, Chang JS, Wu KH, Tsai CH, Lin HS (2008) Mechanisms of and obstacles to iron cardiomyopathy in thalassemia. *Front Biosci* 13:5975–5987
- Peng J, Peng L, Stevenson FF, Doctrow SR, Andersen JK (2007) Iron and paraquat as synergistic environmental risk factors in sporadic Parkinson's disease accelerate age-related neurodegeneration. *J Neurosci* 27(26):6914–6922
- Peng J, Stevenson FF, Oo ML, Andersen JK (2009) Iron-enhanced paraquat-mediated dopaminergic cell death due to increased oxidative stress as a consequence of microglial activation. *Free Radic Biol Med* 46(2):312–320
- Peng Y, Wang C, Xu HH, Liu YN, Zhou F (2010) Binding of alpha-synuclein with Fe(III) and with Fe(II) and biological implications of the resultant complexes. *J Inorg Biochem* 104(4):365–370
- Pennell DJ, Porter JB, Cappellini MD, El-Beshlawy A, Chan LL, Aydinok Y et al (2010) Efficacy of deferasirox in reducing and preventing cardiac iron overload in {beta}-thalassemia. *Blood* 115:2364–2371
- Pérez-Jiménez J, Neveu V, Vos F, Scalbert A (2010) Systematic analysis of the content of 502 polyphenols in 452 foods and beverages: an application of the phenol-explorer database. *J Agric Food Chem* 58:4959–4969
- Perez CA, Tong Y, Guo M (2008) Iron chelators as potential therapeutic agents for Parkinson's disease. *Curr Bioactive Comp* 4(3):150–158
- Perez LR, Franz KJ (2010) Minding metals: tailoring multifunctional chelating agents for neurodegenerative disease. *Dalton Trans* 39(9):2177–2187
- Pérez VI, Bokov A, Van Remmen H, Mele J, Ran Q, Ikeno Y et al (2009a) Is the oxidative stress theory of aging dead? *Biochim Biophys Acta* 1790(10):1005–1014
- Pérez VI, Van Remmen H, Bokov A, Epstein CJ, Vijg J, Richardson A (2009b) The overexpression of major antioxidant enzymes does not extend the lifespan of mice. *Aging Cell* 8(1):73–75
- Perez VP, de Lima MN, da Silva RS, Dornelles AS, Vedana G, Bogo MR et al (2010) Iron leads to memory impairment that is associated with a decrease in acetylcholinesterase pathways. *Curr Neurovasc Res* 7(1):15–22
- Perlstein EO, Ruderfer DM, Ramachandran G, Haggarty SJ, Kruglyak L, Schreiber SL (2006) Revealing complex traits with small molecules and naturally recombinant yeast strains. *Chem Biol* 13(3):319–327
- Perron NR, Brumaghim JL (2009) A review of the antioxidant mechanisms of polyphenol compounds related to iron binding. *Cell Biochem Biophys* 53(2):75–100
- Perron NR, Hodges JN, Jenkins M, Brumaghim JL (2008) Predicting how polyphenol antioxidants prevent DNA damage by binding to iron. *Inorg Chem* 47(14):6153–6161
- Persson HL (2005) Iron-dependent lysosomal destabilization initiates silica-induced apoptosis in murine macrophages. *Toxicol Lett* 159(2):124–133
- Persson HL, Yu Z, Tirosh O, Eaton JW, Brunk UT (2003) Prevention of oxidant-induced cell death by lysosomotropic iron chelators. *Free Radic Biol Med* 34(10):1295–1305
- Perutz MF, Windle AH (2001) Cause of neural death in neurodegenerative diseases attributable to expansion of glutamine repeats. *Nature* 412(6843):143–144
- Petersen RB, Siedlak SL, Lee HG, Kim YS, Nunomura A, Tagliavini F et al (2005) Redox metals and oxidative abnormalities in human prion diseases. *Acta Neuropathologica* 110(3):232–238
- Petri S, Calingasan NY, Alsaied OA, Wille E, Kiaei M, Friedman JE et al (2007) The lipophilic metal chelators DP-109 and DP-460 are neuroprotective in a transgenic mouse model of amyotrophic lateral sclerosis. *J Neurochem* 102(3):991–1000
- Petronilho F, Constantino L, de Souza B, Reinke A, Martins MR, Fraga CM et al (2009) Efficacy of the combination of N-acetylcysteine and desferrioxamine in the prevention and treatment of gentamicin-induced acute renal failure in male Wistar rats. *Nephrol Dial Transpl* 24(7):2077–2082
- Petti S, Scully C (2009) Polyphenols, oral health and disease: a review. *J Dent* 37(6):413–423
- Pettifer SR, Thorne D, McDermott P, Marsh J, Villéger A, Kell DB et al (2009) Visualising biological data: a semantic approach to tool and database integration. *BMC Bioinform* 10(Suppl 6):S19
- Pfefferbaum A, Adalsteinsson E, Rohlfing T, Sullivan EV (2009) MRI estimates of brain iron concentration in normal aging: comparison of field-dependent (FDMI) and phase (SWI) methods. *Neuroimage* 47(2):493–500
- Phillips DC, Dias HK, Kitas GD, Griffiths HR (2010) Aberrant reactive oxygen and nitrogen species generation in rheumatoid arthritis (RA): causes and consequences for immune function, cell survival, and therapeutic intervention. *Antioxid Redox Signal* 12(6):743–785
- Phillips R, Milo R (2009) A feeling for the numbers in biology. *Proc Natl Acad Sci USA* 106(51):21465–21471
- Pierron J, Malan C, Creus M, Gradinaru J, Hafner I, Ivanova A et al (2008) Artificial metalloenzymes for asymmetric allylic alkylation on the basis of the biotin-avidin technology. *Angew Chem Int Ed Engl* 47(4):701–705
- Pietrangelo A (2006) Hereditary hemochromatosis. *Biochim Biophys Acta* 1763(7):700–710
- Pietta PG (2000) Flavonoids as antioxidants. *J Nat Prod* 63(7):1035–1042
- Pinho RA, Silveira PC, Silva LA, Luiz Streck E, Dal-Pizzol F, JC FM (2005) N-acetylcysteine and deferoxamine reduce pulmonary oxidative stress and inflammation in rats after coal dust exposure. *Environ Res* 99(3):355–360
- Pirko I, Johnson AJ, Lohrey AK, Chen Y, Ying J (2009) Deep gray matter T2 hypointensity correlates with disability in a murine model of MS. *J Neurol Sci* 282(1–2):34–38
- Platt M, Rowe W, Wedge DC, Kell DB, Knowles J, Day PJR (2009) Aptamer evolution for array-based diagnostics. *Anal Biochem* 390:203–205
- Pócsi I, Jeney V, Kertai P, Pócsi I, Emri T, Gyémánt G et al (2008) Fungal siderophores function as protective agents of LDL

- oxidation and are promising anti-atherosclerotic metabolites in functional food. *Mol Nutr Food Res* 52(12):1434–1447
- Poland A, Knutson JC (1982) 2,3,7,8-tetrachlorodibenzo-*p*-dioxin and related halogenated aromatic hydrocarbons: examination of the mechanism of toxicity. *Annu Rev Pharmacol Toxicol* 22:517–554
- Polla AS, Polla LL, Polla BS (2003) Iron as the malignant spirit in successful ageing. *Ageing Res Rev* 2(1):25–37
- Pollera C, Lucchini B, Formentin E, Bareggi S, Poli G, Ponti W (2005) Evaluation of anti-prionic activity of clioquinol in an in vivo model (*Mesocricetus auratus*). *Vet Res Commun* 29(Suppl 2): 253–255
- Ponraj D, Makjanic J, Thong PSP, Tan BKH, Watt F (1999) The onset of atherosclerotic lesion formation in hypercholesterolemic rabbits is delayed by iron depletion. *FEBS Lett* 459(2):218–222
- Popa C, Netea MG, van Riel PL, van der Meer JW, Stalenhoef AF (2007) The role of TNF- α in chronic inflammatory conditions, intermediary metabolism, and cardiovascular risk. *J Lipid Res* 48(4):751–762
- Popescu BF, Pickering IJ, George GN, Nichol H (2007) The chemical form of mitochondrial iron in Friedreich's ataxia. *J Inorg Biochem* 101(6):957–966
- Pordea A, Ward TR (2008) Chemogenetic protein engineering: an efficient tool for the optimization of artificial metalloenzymes. *Chem Commun (Camb)* (36):4239–4249
- Porter J, Galanello R, Saglio G, Neufeld EJ, Vichinsky E, Cappellini MD et al (2008) Relative response of patients with myelodysplastic syndromes and other transfusion-dependent anaemias to deferasirox (ICL670): a 1-yr prospective study. *Eur J Haematol* 80(2):168–176
- Porter MJ, Roberts SM, Skidmore J (1999) Polyamino acids as catalysts in asymmetric synthesis. *Bioorg Med Chem* 7(10):2145–2156
- Poser I, Rahman Q, Lohani M, Yadav S, Becker HH, Weiss DG et al (2004) Modulation of genotoxic effects in asbestos-exposed primary human mesothelial cells by radical scavengers, metal chelators and a glutathione precursor. *Mutat Res* 559(1–2):19–27
- Postgate JR (1967) Viability measurements and the survival of microbes under minimum stress. *Adv Micr Physiol* 1:1–23
- Postgate JR (1976) Death in microbes and macrobes. In: Gray TRG, Postgate JR (eds) *In the survival of vegetative microbes*. Cambridge University Press, Cambridge, pp 1–19
- Prass K, Ruscher K, Karsch M, Isaev N, Megow D, Priller J et al (2002) Desferrioxamine induces delayed tolerance against cerebral ischemia in vivo and in vitro. *J Cereb Blood Flow Metab* 22(5):520–525
- Price L, Kowdley KV (2009) The role of iron in the pathophysiology and treatment of chronic hepatitis C. *Can J Gastroenterol* 23(12):822–828
- Proctor CJ, Gray DA (2010) GSK3 and p53—is there a link in Alzheimer's disease? *Mol Neurodegener* 5:7
- Proctor CJ, Tsigotou M, Gray DA (2007) An in silico model of the ubiquitin-proteasome system that incorporates normal homeostasis and age-related decline. *BMC Syst Biol* 1:17
- Protti A, Singer M (2006) Bench-to-bedside review: Potential strategies to protect or reverse mitochondrial dysfunction in sepsis-induced organ failure. *Crit Care* 10(5):227–233
- Prusiner SB (1998) Prions. *Proc Natl Acad Sci* 95(23):13363–13383
- Prusiner SB (2001) Shattuck lecture—neurodegenerative diseases and prions. *N Engl J Med* 344(20):1516–1526
- Pryor WA, Squadrito GL (1995) The chemistry of peroxynitrite: a product from the reaction of nitric oxide with superoxide. *Am J Physiol* 268(5 Pt 1): L699–L722
- Puccio H (2009) Multicellular models of Friedreich ataxia. *J Neurol* 256(Suppl 1):18–24
- Pulliam JF, Jennings CD, Kryscio RJ, Davis DG, Wilson D, Montine TJ et al (2003) Association of HFE mutations with neurodegeneration and oxidative stress in Alzheimer's disease and correlation with APOE. *Am J Med Genet B Neuropsychiatr Genet* 119B(1):48–53
- Purdey M (2000) Ecosystems supporting clusters of sporadic TSEs demonstrate excesses of the radical-generating divalent cation manganese and deficiencies of antioxidant co factors Cu, Se, Fe, Zn. Does a foreign cation substitution at prion protein's Cu domain initiate TSE? *Med Hypotheses* 54(2):278–306
- Que L Jr, Tolman WB (2008) Biologically inspired oxidation catalysis. *Nature* 455(7211):333–340
- Quinlan GJ, Chen Y, Evans TW, Gutteridge JMC (2001) Iron signalling regulated directly and through oxygen: implications for sepsis and the acute respiratory distress syndrome. *Clinical Science* 100(2):169–182
- Quinlan GJ, Evans TW, Gutteridge JM (2002) Iron and the redox status of the lungs. *Free Radic Biol Med* 33(10):1306–1313
- Quinlan TR, Marsh JP, Janssen YM, Borm PA, Mossman BT (1994) Oxygen radicals and asbestos-mediated disease. *Environ Health Perspect* 102(Suppl 10):107–110
- Quintana C (2007) Contribution of analytical microscopies to human neurodegenerative diseases research (PSP and AD). *Mini Rev Med Chem* 7(9):961–975
- Quintana C, Bellefqih S, Laval JY, Guerquin-Kern JL, Wu TD, Avila J et al (2006) Study of the localization of iron, ferritin, and hemosiderin in Alzheimer's disease hippocampus by analytical microscopy at the subcellular level. *J Struct Biol* 153(1):42–54
- Quintanilla RA, Johnson GVW (2009) Role of mitochondrial dysfunction in the pathogenesis of Huntington's disease. *Brain Res Bull* 80(4–5):242–247
- Qureshi AI, Mendelow AD, Hanley DF (2009) Intracerebral haemorrhage. *Lancet* 373(9675):1632–1644
- Qureshi M, Brown RH Jr, Rogers JT, Cudkovic ME (2008) Serum ferritin and metal levels as risk factors for amyotrophic lateral sclerosis. *Open Neurol J* 2:51–54
- Radak Z, Chung HY, Koltai E, Taylor AW, Goto S (2008) Exercise, oxidative stress and hormesis. *Ageing Res Rev* 7(1):34–42
- Rader DJ, Daugherty A (2008) Translating molecular discoveries into new therapies for atherosclerosis. *Nature* 451(7181):904–913
- Radi R, Cassina A, Hodara R, Quijano C, Castro L (2002) Peroxynitrite reactions and formation in mitochondria. *Free Radic Biol Med* 33(11):1451–1464
- Radi R, Peluffo G, Alvarez MN, Naviliat M, Cayota A (2001) Unraveling peroxynitrite formation in biological systems. *Free Radic Biol Med* 30(5):463–488
- Rajendran R, Ren M, Ning P, Tan Kwong Huat B, Halliwell B, Watt F (2007) Promotion of atherogenesis by copper or iron—which is more likely? *Biochem Biophys Res Commun* 353(1):6–10
- Rajpathak SN, Crandall JP, Wylie-Rosett J, Kabat GC, Rohan TE, Hu FB (2009) The role of iron in type 2 diabetes in humans. *Biochim Biophys Acta* 1790:671–681
- Ramagopalan SV, Cukjati M, Cernilec M, DeLuca GC, Dymont DA, Degenhardt A et al (2008) Mutations in the hemochromatosis gene and the clinical outcome of multiple sclerosis. *J Neuroimmunol* 203(1):104–107
- Ramakrishna G, Rooke TW, Cooper LT (2003) Iron and peripheral arterial disease: revisiting the iron hypothesis in a different light. *Vasc Med* 8(3):203–210
- Ramesh E, Jayakumar T, Elanchezian R, Sakthivel M, Geraldine P, Thomas PA (2009) Green tea catechins, alleviate hepatic lipidemic-oxidative injury in Wistar rats fed an atherogenic diet. *Chem Biol Interact* 180(1):10–19
- Rana A, Gnaneswari D, Bansal S, Kundu B (2009) Prion metal interaction: is prion pathogenesis a cause or a consequence of metal imbalance? *Chem Biol Interact* 181(3):282–291
- Rao PS, Liu XK, Das DK, Weinstein GS, Tyras DH (1991) Protection of ischemic heart from reperfusion injury by myo-inositol

- hexaphosphate, a natural antioxidant. *Ann Thorac Surg* 52(4):908–912
- Ratan RR, Siddiq A, Aminova L, Langley B, McConoughey S, Karpisheva K et al (2008) Small molecule activation of adaptive gene expression: tilorone or its analogs are novel potent activators of hypoxia inducible factor-1 that provide prophylaxis against stroke and spinal cord injury. *Ann NY Acad Sci* 1147:383–394
- Rattan SI (2008) Hormesis in aging. *Ageing Res Rev* 7(1):63–78
- Rattan SI, Demirovic D (2009) Hormesis can and does work in humans. *Dose Response* 8(1):58–63
- Rattan SIS (2004) Aging, anti-aging, and hormesis. *Mech Ageing Dev* 125(4):285–289
- Rattner A, Nathans J (2006) Macular degeneration: recent advances and therapeutic opportunities. *Nat Rev Neurosci* 7(11):860–872
- Ravindran J, Prasad S, Aggarwal BB (2009) Curcumin and cancer cells: how many ways can curry kill tumor cells selectively? *AAPS J* 11(3):495–510
- Rawat R, Humphrey J, Ntozini R, Mutasa K, Iliff P, Stoltzfus R (2008) Elevated iron stores are associated with HIV disease severity and mortality among postpartum women in Zimbabwe. *Public Health Nutr* 12:1–9
- Rayman MP, Barlis J, Evans RW, Redman CW, King LJ (2002) Abnormal iron parameters in the pregnancy syndrome pre-eclampsia. *Am J Obstet Gynecol* 187(2):412–418
- Raymond KN, Dertz EA, Kim SS (2003) Enterobactin: an archetype for microbial iron transport. *Proc Natl Acad Sci USA* 100(7):3584–3588
- Reddy PH (2008) Mitochondrial medicine for aging and neurodegenerative diseases. *Neuromol Med* 10(4):291–315
- Reddy PH, Beal MF (2008) Amyloid beta, mitochondrial dysfunction and synaptic damage: implications for cognitive decline in aging and Alzheimer's disease. *Trends Mol Med* 14(2):45–53
- Reddy VP, Zhu X, Perry G, Smith MA (2009) Oxidative stress in diabetes and Alzheimer's disease. *J Alzheimers Dis* 16(4):763–774
- Redl H, Gasser H, Schlag G, Marzi I (1993) Involvement of oxygen radicals in shock related cell injury. *Br Med Bull* 49(3):556–565
- Reid AB, Kurten RC, McCullough SS, Brock RW, Hinson JA (2005) Mechanisms of acetaminophen-induced hepatotoxicity: role of oxidative stress and mitochondrial permeability transition in freshly isolated mouse hepatocytes. *J Pharmacol Exp Ther* 312(2):509–516
- Reif DW (1992) Ferritin as a source of iron for oxidative damage. *Free Radic Biol Med* 12(5):417–427
- Reiter RJ, Paredes SD, Manchester LC, Tan DX (2009) Reducing oxidative/nitrosative stress: a newly-discovered genre for melatonin. *Crit Rev Biochem Mol Biol* 44(4):175–200
- Ren MQ, Rajendran R, Pan N, Tan BKH, Ong WY, Watt F et al (2005) The iron chelator desferrioxamine inhibits atherosclerotic lesion development and decreases lesion iron concentrations in the cholesterol-fed rabbit. *Free Radic Biol Med* 38(9):1206–1211
- Ren MQ, Watt F, Huat BTK, Halliwell B (2003) Trace elemental distributions in induced atherosclerotic lesions using nuclear microscopy. *Nuclear Instr Meth Phys Res B* 210:336–342
- Reynolds A, Laurie C, Mosley RL, Gendelman HE (2007) Oxidative stress and the pathogenesis of neurodegenerative disorders. *Int Rev Neurobiol* 82:297–325
- Reynolds FD, Dauchy R, Blask D, Dietz PA, Lynch D, Zuckerman R (2003) The pineal gland hormone melatonin improves survival in a rat model of sepsis/shock induced by zymosan A. *Surgery* 134(3):474–479
- Reznichenko L, Amit T, Zheng H, Avramovich-Tirosh Y, Youdim MB, Weinreb O et al (2006) Reduction of iron-regulated amyloid precursor protein and beta-amyloid peptide by (-)-epigallocatechin-3-gallate in cell cultures: implications for iron chelation in Alzheimer's disease. *J Neurochem* 97(2):527–536
- Reznichenko L, Kalfon L, Amit T, Youdim MB, Mandel SA (2010) Low dosage of rasagiline and epigallocatechin gallate synergistically restored the nigrostriatal axis in MPTP-induced Parkinsonism. *Neurodegener Dis*
- Rice-Evans CA, Miller J, Paganga G (1997) Antioxidant properties of phenolic compounds. *Trends Plant Sci* 2(4):152–159
- Rice-Evans CA, Packer L (eds) (2003) *Flavonoids in health and disease*, 2nd edn. Marcel Dekker, New York
- Richardson DR, Huang ML, Whitnall M, Becker EM, Ponka P, Suryo Rahmanto Y (2009a) The ins and outs of mitochondrial iron-loading: the metabolic defect in Friedreich's ataxia. *J Mol Med*
- Richardson DR, Kalinowski DS, Lau S, Jansson PJ, Lovejoy DB (2009b) Cancer cell iron metabolism and the development of potent iron chelators as anti-tumour agents. *Biochim Biophys Acta* 1790(7):702–717
- Richer S, Rudy D, Statkute L, Karofy K, Frankowski J (2002) Serum iron, transferrin saturation, ferritin, and dietary data in age-related macular degeneration. *Am J Ther* 9(1):25–28
- Ridker PM, Brown NJ, Vaughan DE, Harrison DG, Mehta JL (2004) Established and emerging plasma biomarkers in the prediction of first atherothrombotic events. *Circulation* 109(25 Suppl 1):6–19
- Riedl MA, Nel AE (2008) Importance of oxidative stress in the pathogenesis and treatment of asthma. *Curr Opin Allergy Clin Immunol* 8(1):49–56
- Rigamonti C, Andorno S, Maduli E, Capelli F, Boldorini R, Sartori M (2005) Gender and liver fibrosis in chronic hepatitis: the role of iron status. *Aliment Pharmacol Ther* 21(12):1445–1451
- Rimbach G, Pallauf J, Moehring J, Kraemer K, Minihane AM (2008) Effect of dietary phytate and microbial phytase on mineral and trace element bioavailability: a literature review. *Curr Topics Nutraceut Res* 6:131–144
- Ristow M, Zarse K (2010) How increased oxidative stress promotes longevity and metabolic health: The concept of mitochondrial hormesis (mitohormesis). *Exp Gerontol*
- Ritter C, Andrades ME, Reinke A, Menna-Barreto S, Moreira JC, Dal-Pizzol F (2004) Treatment with N-acetylcysteine plus deferoxamine protects rats against oxidative stress and improves survival in sepsis. *Crit Care Med* 32(2):342–349
- Ritter C, da Cunha AA, Echer IC, Andrades M, Reinke A, Lucchiari N et al (2006) Effects of N-acetylcysteine plus deferoxamine in lipopolysaccharide-induced acute lung injury in the rat. *Crit Care Med* 34(2):471–477
- Rittirsch D, Flierl MA, Ward PA (2008) Harmful molecular mechanisms in sepsis. *Nat Rev Immunol* 8(10):776–787
- Rival T, Page RM, Chandraratna DS, Sendall TJ, Ryder E, Liu B et al (2009) Fenton chemistry and oxidative stress mediate the toxicity of the beta-amyloid peptide in a Drosophila model of Alzheimer's disease. *Eur J Neurosci* 29(7):1335–1347
- Rivera-Espinoza Y, Muriel P (2009) Pharmacological actions of curcumin in liver diseases or damage. *Liver Int* 29(10):1457–1466
- Robinson SR, Dang TN, Dringen R, Bishop GM (2009) Hemin toxicity: a preventable source of brain damage following hemorrhagic stroke. *Redox Rep* 14(6):228–235
- Robson KJ, Lehmann DJ, Wilmhurst VL, Livesey KJ, Combrinck M, Merryweather-Clarke AT et al (2004) Synergy between the C2 allele of transferrin and the C282Y allele of the haemochromatosis gene (HFE) as risk factors for developing Alzheimer's disease. *J Med Genet* 41(4):261–265
- Rodríguez-Caso C, Corominas-Murtra B, Solé RV (2009) On the basic computational structure of gene regulatory networks. *Mol Biosyst* 5(12):1617–1629
- Rodríguez-Fernandez M, Mendes P, Banga JR (2006) A hybrid approach for efficient and robust parameter estimation in biochemical pathways. *Biosystems* 83(2–3):248–265

- Roijers RB, Dutta RK, Mutsaers PHA, Gijbels MJJ, de Winther MPJ, de Goeij JJM et al (2005) Spatial correlation of trace elements with morphological features of atherosclerotic plaques. *Nuclear Instrum Methods Phys Res Sect B-Beam Interact Mater Atoms* 231:239–244
- Rokushima M, Omi K, Araki A, Kyokawa Y, Furukawa N, Itoh F et al (2007a) A toxicogenomic approach revealed hepatic gene expression changes mechanistically linked to drug-induced hemolytic anemia. *Toxicol Sci* 95(2):474–484
- Rokushima M, Omi K, Imura K, Araki A, Furukawa N, Itoh F et al (2007b) Toxicogenomics of drug-induced hemolytic anemia by analyzing gene expression profiles in the spleen. *Toxicol Sci* 100(1):290–302
- Romeo AM, Christen L, Niles EG, Kosman DJ (2001) Intracellular chelation of iron by bipyridyl inhibits DNA virus replication: ribonucleotide reductase maturation as a probe of intracellular iron pools. *J Biol Chem* 276(26):24301–24308
- Roostae AR, Roostae MH, Roucou X (2010) An update on prion biology and proteomics. *Curr Proteom* 7(1):36–48
- Rooyackers TM, Stroes ES, Kooistra MP, van Faassen EE, Hider RC, Rabelink TJ et al (2002) Ferric saccharate induces oxygen radical stress and endothelial dysfunction in vivo. *Eur J Clin Invest* 32(Suppl 1):9–16
- Roriz-Filho JS, Sa-Roriz TM, Rosset I, Camozzato AL, Santos AC, Chaves ML et al (2009) (Pre)diabetes, brain aging, and cognition. *Biochim Biophys Acta* 1792(5):432–443
- Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A et al (1993) Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 362(6415):59–62
- Ross CA (1995) When more is less: pathogenesis of glutamine repeat neurodegenerative diseases. *Neuron* 15(3):493–496
- Ross OA, Meschia JF (2009) Genetics of ischemic stroke: inheritance of a sporadic disorder. *Curr Neurol Neurosci Rep* 9(1):19–27
- Ross R (1999) Atherosclerosis: an inflammatory disease. *N Engl J Med* 340(2):115–126
- Rötig A, Sidi D, Munnich A, Rustin P (2002) Molecular insights into Friedreich's ataxia and antioxidant-based therapies. *Trends Mol Med* 8(5):221–224
- Rottkamp CA, Nunomura A, Raina AK, Sayre LM, Perry G, Smith MA (2000) Oxidative stress, antioxidants, and Alzheimer disease. *Alzheimer Dis Assoc Disord* 14(Suppl 1):S62–S66
- Rouault TA (2006) The role of iron regulatory proteins in mammalian iron homeostasis and disease. *Nat Chem Biol* 2(8):406–414
- Rowe W, Platt M, Wedge D, Day PJ, Kell DB, Knowles J (2010) Analysis of a complete DNA-protein affinity landscape. *J R Soc Interface* 7(44):397–408
- Rubbo H, O'Donnell V (2005) Nitric oxide, peroxynitrite and lipoxigenase in atherogenesis: mechanistic insights. *Toxicology* 208(2):305–317
- Rubbo H, Trostchansky A, O'Donnell VB (2009) Peroxynitrite-mediated lipid oxidation and nitration: mechanisms and consequences. *Arch Biochem Biophys* 484(2):167–172
- Sadrzadeh SM, Saffari Y (2004) Iron and brain disorders. *Am J Clin Pathol* 121(Suppl):S64–S70
- Sahle S, Mendes P, Hoops S, Kummer U (2008) A new strategy for assessing sensitivities in biochemical models. *Phil Trans Royal Soc A* 366:3619–3631. doi:10.1098/rsta.2008.0108
- Saito K, Ishizaka N, Aizawa T, Sata M, Iso-o N, Noiri E et al (2005) Iron chelation and a free radical scavenger suppress angiotensin II-induced upregulation of TGF-beta1 in the heart. *Am J Physiol Heart Circ Physiol* 288(4):H1836–1843
- Sakaïda I, Kayano K, Wasaki S, Nagatomi A, Matsumura Y, Okita K (1995) Protection against acetaminophen-induced liver injury in vivo by an iron chelator, deferoxamine. *Scand J Gastroenterol* 30(1):61–67
- Sakudo A, Ikuta K (2009a) Fundamentals of prion diseases and their involvement in the loss of function of cellular prion protein. *Protein Pept Lett* 16(3):217–229
- Sakudo A, Ikuta K (2009b) Prion protein functions and dysfunction in prion diseases. *Curr Med Chem* 16(3):380–389
- Saleh A, Schroeter M, Ringelstein A, Hartung HP, Siebler M, Modder U et al (2007) Iron oxide particle-enhanced MRI suggests variability of brain inflammation at early stages after ischemic stroke. *Stroke* 38(10):2733–2737
- Salonen JT, Nyssonen K, Korpela H, Tuomilehto J, Seppanen R, Salonen R (1992) High stored iron levels are associated with excess risk of myocardial infarction in eastern Finnish men. *Circulation* 86(3):803–811
- Saltelli A, Ratto M, Andres T, Campolongo F, Cariboni J, Gatelli D et al (2008) Global sensitivity analysis: the primer. WileyBlackwell, New York
- Saltelli A, Tarantola S, Campolongo F, Ratto M (2004) Sensitivity analysis in practice: a guide to assessing scientific models. Wiley, New York
- Salvador GA (2010) Iron in neuronal function and dysfunction. *Biofactors* 36:103–110
- Sampietro M, Caputo L, Casatta A, Merigalli M, Pellagatti A, Tagliabue J et al (2001) The hemochromatosis gene affects the age of onset of sporadic Alzheimer's disease. *Neurobiol Aging* 22(4):563–568
- Sandberg AS, Brune M, Carlsson NG, Hallberg L, Skoglund E, Rossander-Hulthen L (1999) Inositol phosphates with different numbers of phosphate groups influence iron absorption in humans. *Am J Clin Nutr* 70(2):240–246
- Sandy M, Butler A (2009) Microbial iron acquisition: marine and terrestrial siderophores. *Chem Rev* 109(10):4580–4595
- Sanfeliu C, Sebastià J, Cristòfol R, Rodríguez-Farré E (2003) Neurotoxicity of organomercurial compounds. *Neurotox Res* 5(4):283–305
- Santhosh K, Kesavadas C, Thomas B, Gupta AK, Thamburaj K, Kapilamoorthy TR (2009) Susceptibility weighted imaging: a new tool in magnetic resonance imaging of stroke. *Clin Radiol* 64(1):74–83
- Santos R, Lefevre S, Sliwa D, Seguin A, Camadro JM, Lesuisse E (2010) Friedreich's Ataxia: molecular mechanisms, redox considerations and therapeutic opportunities. *Antioxid Redox Signal*
- Sarafian T, Verity MA (1991) Oxidative mechanisms underlying methyl mercury neurotoxicity. *Int J Dev Neurosci* 9(2):147–153
- Sarafian TA, Vartavarian L, Kane DJ, Bredesen DE, Verity MA (1994) bcl-2 expression decreases methyl mercury-induced free-radical generation and cell killing in a neural cell line. *Toxicol Lett* 74(2):149–155
- Sarkar FH, Li Y (2009) Harnessing the fruits of nature for the development of multi-targeted cancer therapeutics. *Cancer Treat Rev* 35(7):597–607
- Sarkar FH, Li Y, Wang Z, Kong D (2009) Cellular signaling perturbation by natural products. *Cell Signal* 21(11):1541–1547
- Savarino A, Pescarmona GP, Boelaert JR (1999) Iron metabolism and HIV infection: reciprocal interactions with potentially harmful consequences? *Cell Biochem Funct* 17(4):279–287
- Sawyers CL (2007) Cancer: mixing cocktails. *Nature* 449(7165):993–996
- Saxena K, Ranalli M, Khan N, Blanchong C, Kahwash SB (2005) Fatal stroke in a child with severe iron deficiency anemia and multiple hereditary risk factors for thrombosis. *Clin Pediatr (Phila)* 44(2):175–180
- Sayre LM, Moreira PI, Smith MA, Perry G (2005) Metal ions and oxidative protein modification in neurological disease. *Ann Ist Super Sanita* 41(2):143–164

- Sayre LM, Perry G, Atwood CS, Smith MA (2000a) The role of metals in neurodegenerative diseases. *Cell Mol Biol (Noisy-le-grand)* 46(4):731–741
- Sayre LM, Perry G, Harris PLR, Liu YH, Schubert KA, Smith MA (2000b) *In situ* oxidative catalysis by neurofibrillary tangles and senile plaques in Alzheimer's disease: A central role for bound transition metals. *J Neurochem* 74(1):270–279
- Scalbert A, Manach C, Morand C, Remesy C, Jimenez L (2005) Dietary polyphenols and the prevention of diseases. *Crit Rev Food Sci Nutr* 45(4):287–306
- Scalbert A, Williamson G (2000) Dietary intake and bioavailability of polyphenols. *J Nutr* 130(8S Suppl):2073S–2085S
- Schagerlöf U, Elmlund H, Gakh O, Nordlund G, Hebert H, Lindahl M et al (2008) Structural basis of the iron storage function of frataxin from single-particle reconstruction of the iron-loaded oligomer. *Biochemistry* 47(17):4948–4954
- Schenck JF, Zimmerman EA (2004) High-field magnetic resonance imaging of brain iron: birth of a biomarker? *NMR Biomed* 17(7):433–445
- Schiano TD, Grinberg M, Nawabi I, Grabowski G (1993) Blue nasal secretions: a presentation of Gaucher's disease and concurrent hemosiderosis. *Am J Hematol* 44(3):219–220
- Schlawe D, Majdalani A, Velcicky J, Hessler E, Wieder T, Prokop A et al (2004) Iron-containing nucleoside analogues with pronounced apoptosis-inducing activity. *Angew Chem Int Ed Engl* 43(13):1731–1734
- Schleicher E, Friess U (2007) Oxidative stress, AGE, and atherosclerosis. *Kidney Int Suppl*(106):S17–S26
- Schlemmer U, Frolich W, Prieto RM, Grases F (2009) Phytate in foods and significance for humans: food sources, intake, processing, bioavailability, protective role and analysis. *Mol Nutr Food Res* 53(Suppl 2):S330–S375
- Schmitt-Ulms G, Ehsani S, Watts JC, Westaway D, Wille H (2009) Evolutionary descent of prion genes from the ZIP family of metal ion transporters. *PLoS One* 4(9):e7208
- Schnaider Beeri M, Goldbourt U, Silverman JM, Noy S, Schmeidler J, Ravona-Springer R et al (2004) Diabetes mellitus in midlife and the risk of dementia three decades later. *Neurology* 63(10):1902–1907
- Schnellmann JG, Pumford NR, Kusewitt DF, Bucci TJ, Hinson JA (1999) Deferoxamine delays the development of the hepatotoxicity of acetaminophen in mice. *Toxicol Lett* 106(1):79–88
- Schrag M, McAuley G, Pomakian J, Jiffry A, Tung S, Mueller C et al (2009) Correlation of hypointensities in susceptibility-weighted images to tissue histology in dementia patients with cerebral amyloid angiopathy: a postmortem MRI study. *Acta Neuropathol*
- Schrattenholz A, Šoškić V (2008) What does systems biology mean for drug development? *Curr Med Chem* 15(15):1520–1528
- Schreck R, Albermann K, Baeuerle PA (1992a) Nuclear factor kappa-B—an oxidative stress-responsive transcription factor of eukaryotic cells (a Review). *Free Rad Res Comm* 17(4):221–237
- Schreck R, Meier B, Mannel DN, Droge W, Baeuerle PA (1992b) Dithiocarbamates as potent inhibitors of nuclear factor kappa B activation in intact cells. *J Exp Med* 175(5):1181–1194
- Schreck R, Baeuerle PA (1994) Assessing oxygen radicals as mediators in activation of inducible eukaryotic transcription factor NF-kappa B. *Methods Enzymol* 234:151–163
- Schreck R, Rieber P, Baeuerle PA (1991) Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-Kappa-B transcription Factor and HIV-1. *EMBO J* 10(8):2247–2258
- Schulz JB, Boesch S, Burk K, Durr A, Giunti P, Mariotti C et al (2009) Diagnosis and treatment of Friedreich ataxia: a European perspective. *Nat Rev Neurol* 5(4):222–234
- Schulz M, Uhlenendorf J, Klipp E, Liebermeister W (2006) SBML-merge, a system for combining biochemical network models. *Genome Inform* 17(1):62–71
- Scott LE, Orvig C (2009) Medicinal inorganic chemistry approaches to passivation and removal of aberrant metal ions in disease. *Chem Rev* 109(10):4885–4910
- Sebai H, Ben-Attia M, Sani M, Aouani E, Ghanem-Boughanmi N (2009) Protective effect of resveratrol in endotoxemia-induced acute phase response in rats. *Arch Toxicol* 83(4):335–340
- Sebastiani G, Walker AP (2007) HFE gene in primary and secondary hepatic iron overload. *World J Gastroenterol* 13(35):4673–4689
- Seet RC, Lee CY, Lim EC, Tan JJ, Quek AM, Chong WL et al (2010) Oxidative damage in Parkinson disease: measurement using accurate biomarkers. *Free Radic Biol Med* 48(4):560–566
- Seguin A, Sutak R, Bulteau AL, Garcia-Serres R, Oddou JL, Lefevre S et al (2010) Evidence that yeast frataxin is not an iron storage protein *in vivo*. *Biochim Biophys Acta* 1802(6):531–538
- Seifried HE, Anderson DE, Fisher EI, Milner JA (2007) A review of the interaction among dietary antioxidants and reactive oxygen species. *J Nutr Biochem* 18(9):567–579
- Selim M (2009) Deferoxamine mesylate: a new hope for intracerebral hemorrhage: from bench to clinical trials. *Stroke* 40(3 Suppl):S90–91
- Selim MH, Ratan RR (2004) The role of iron neurotoxicity in ischemic stroke. *Ageing Res Rev* 3(3):345–353
- Sen CK, Packer L (1996) Antioxidant and redox regulation of gene transcription. *FASEB J* 10(7):709–720
- Sener A, Ozsavci D, Bingol-Ozakpinar O, Cevik O, Yanikkaya-Demirel G, Yardimci T (2009) Oxidized-LDL and Fe³⁺/Ascorbic Acid-Induced Oxidative Modifications and Phosphatidylserine Exposure in Human Platelets are Reduced by Melatonin. *Folia Biol (Praha)* 55(2):45–52
- Serra JA, Domínguez RO, Marschoff ER, Guareschi EM, Famulari AL, Boveris A (2009) Systemic oxidative stress associated with the neurological diseases of aging. *Neurochem Res*
- Shah SV, Alam MG (2003) Role of iron in atherosclerosis. *Am J Kidney Dis* 41(3 Suppl 1):S80–S83
- Shamoto-Nagai M, Maruyama W, Yi H, Akao Y, Tribl F, Gerlach M et al (2006) Neuromelanin induces oxidative stress in mitochondria through release of iron: mechanism behind the inhibition of 26S proteasome. *J Neural Transm* 113(5):633–644
- Shamsuddin AM (1995) Inositol phosphates have novel anticancer function. *J Nutr* 125(3 Suppl):725S–732S
- Shanker G, Aschner M (2003) Methylmercury-induced reactive oxygen species formation in neonatal cerebral astrocytic cultures is attenuated by antioxidants. *Brain Res Mol Brain Res* 110(1):85–91
- Shannahan JH, Schladweiler MC, Richards JH, Ledbetter AD, Ghio AJ, Kodavanti UP (2010) Pulmonary oxidative stress, inflammation, and dysregulated iron homeostasis in rat models of cardiovascular disease. *J Toxicol Environ Health A* 73(10):641–656
- Sharifi F, Nasab NM, Zadeh HJ (2008) Elevated serum ferritin concentrations in prediabetic subjects. *Diab Vasc Dis Res* 5(1):15–18
- Sharma S, Haldar C (2009) Comparative effect of melatonin and vitamin E on phenylhydrazine-induced toxicity in the spleen of *Funambulus pennanti*. *Environ Toxicol* 24(1):1–9
- Sharom JR, Bellows DS, Tyers M (2004) From large networks to small molecules. *Curr Opin Chem Biol* 8(1):81–90
- Shatos MA, Doherty JM, Marsh JP, Mossman BT (1987) Prevention of asbestos-induced cell death in rat lung fibroblasts and alveolar macrophages by scavengers of active oxygen species. *Environ Res* 44(1):103–116
- Shcherbatykh I, Carpenter DO (2007) The role of metals in the etiology of Alzheimer's disease. *J Alzheimers Dis* 11(2):191–205

- She H, Xiong S, Lin M, Zandi E, Giulivi C, Tsukamoto H (2002) Iron activates NF-kappaB in Kupffer cells. *Am J Physiol Gastrointest Liver Physiol* 283(3):G719–726
- Sheftel A, Stehling O, Lill R (2010) Iron-sulfur proteins in health and disease. *Trends Endocrinol Metab* 21(5):302–314
- Shen Z, Parker VD, Aust AE (1995) Mediated, thin-layer cell, coulometric determination of redox-active iron on the surface of asbestos fibers. *Anal Chem* 67(2):307–311
- Sheth S, Brittenham GM (2000) Genetic disorders affecting proteins of iron metabolism: clinical implications. *Annu Rev Med* 51:443–464
- Shi T, Knaapen AM, Begerow J, Birmili W, Borm PJ, Schins RP (2003) Temporal variation of hydroxyl radical generation and 8-hydroxy-2'-deoxyguanosine formation by coarse and fine particulate matter. *Occup Environ Med* 60(5):315–321
- Shi Z, Nie G, Duan XL, Rouault T, Wu WS, Ning B et al (2010) Neuroprotective mechanism of mitochondrial ferritin on 6-hydroxydopamine induced dopaminergic cell damage: implication for neuroprotection in Parkinson's disease. *Antioxid Redox Signal*
- Shin SM, Kim SG (2009) Inhibition of arachidonic acid and iron-induced mitochondrial dysfunction and apoptosis by oltipraz and novel 1, 2-dithiole-3-thione congeners. *Mol Pharmacol* 75(1):242–253
- Shiomi M, Ito T, Tsukada T, Tsujita Y, Horikoshi H (1999) Combination treatment with troglitazone, an insulin action enhancer, and pravastatin, an inhibitor of HMG-CoA reductase, shows a synergistic effect on atherosclerosis of WHHL rabbits. *Atherosclerosis* 142(2):345–353
- Shoelson SE, Goldfine AB (2009) Getting away from glucose: fanning the flames of obesity-induced inflammation. *Nat Med* 15(4):373–374
- Shoelson SE, Herrero L, Naaz A (2007) Obesity, inflammation, and insulin resistance. *Gastroenterology* 132(6):2169–2180
- Shore SA (2008) Obesity and asthma: possible mechanisms. *J Allergy Clin Immunol* 121(5):1087–1093 (quiz 1094–1085)
- Shukla A, Gulumian M, Hei TK, Kamp D, Rahman Q, Mossman BT (2003a) Multiple roles of oxidants in the pathogenesis of asbestos-induced diseases. *Free Radic Biol Med* 34(9):1117–1129
- Shukla A, Jung M, Stern M, Fukagawa NK, Taatjes DJ, Sawyer D et al (2003b) Asbestos induces mitochondrial DNA damage and dysfunction linked to the development of apoptosis. *Am J Physiol Lung Cell Mol Physiol* 285(5):L1018–L1025
- Sidransky E, Nalls MA, Aasly JO, Aharon-Peretz J, Annesi G, Barbosa ER et al (2009) Multicenter analysis of glucocerebrosidase mutations in Parkinson's disease. *N Engl J Med* 361(17):1651–1661
- Signorini C, Ciccoli L, Leoncini S, Carloni S, Perrone S, Comporti M et al (2009) Free iron, total F-isoprostanes and total F-neuroprostanes in a model of neonatal hypoxic-ischemic encephalopathy: neuroprotective effect of melatonin. *J Pineal Res* 46(2):148–154
- Sikorska K, Stalke P, Izycka-Swieszewska E, Romanowski T, Bielawski KP (2010) The role of iron overload and HFE gene mutations in the era of pegylated interferon and ribavirin treatment of chronic hepatitis C. *Med Sci Monit* 16(3):CR137–CR143
- Silver H, Chertkow Y, Weinreb O, Danovich L, Youdim M (2009) Multifunctional pharmacotherapy: what can we learn from study of selective serotonin reuptake inhibitor augmentation of antipsychotics in negative-symptom schizophrenia? *Neurotherapeutics* 6(1):86–93
- Silver LL (2007) Multi-targeting by monotherapeutic antibacterials. *Nat Rev Drug Discov* 6(1):41–55
- Silvestri L, Camaschella C (2008) A potential pathogenetic role of iron in Alzheimer's Disease. *J Cell Mol Med* 12(5a):1548–1550
- Simka M, Rybak Z (2008) Hypothetical molecular mechanisms by which local iron overload facilitates the development of venous leg ulcers and multiple sclerosis lesions. *Med Hypotheses* 71(2):293–297
- Simmonds RE, Foxwell BM (2008) NF- κ B and its relevance to arthritis and inflammation. *Rheumatology (Oxford)*
- Simmons-Willis TA, Koh AS, Clarkson TW, Ballatori N (2002) Transport of a neurotoxicant by molecular mimicry: the methylmercury-L-cysteine complex is a substrate for human L-type large neutral amino acid transporter (LAT) 1 and LAT2. *Biochem J* 367(Pt 1):239–246
- Simonart T, Degraef C, Andrei G, Mosselmans R, Hermans P, Van Vooren JP et al (2000) Iron chelators inhibit the growth and induce the apoptosis of Kaposi's sarcoma cells and of their putative endothelial precursors. *J Invest Dermatol* 115(5):893–900
- Šimůnek T, Štěrbá M, Popelova O, Adamcova M, Hrdina R, Geršl V (2009) Anthracycline-induced cardiotoxicity: overview of studies examining the roles of oxidative stress and free cellular iron. *Pharmacol Rep* 61(1):154–171
- Singh AV, Zamboni P (2009) Anomalous venous blood flow and iron deposition in multiple sclerosis. *J Cereb Blood Flow Metab* 29(12):1867–1878
- Singh N, Pillay V, Choonara YE (2007) Advances in the treatment of Parkinson's disease. *Prog Neurobiol* 81(1):29–44
- Singh A, Isaac AO, Luo X, Mohan ML, Cohen ML, Chen F et al (2009a) Abnormal brain iron homeostasis in human and animal prion disorders. *PLoS Pathog* 5(3):e1000336
- Singh A, Kong Q, Luo X, Petersen RB, Meyerson H, Singh N (2009b) Prion protein (PrP) knock-out mice show altered iron metabolism: a functional role for PrP in iron uptake and transport. *PLoS One* 4(7):e6115
- Singh A, Mohan ML, Isaac AO, Luo X, Petrak J, Vyoral D et al (2009c) Prion protein modulates cellular iron uptake: a novel function with implications for prion disease pathogenesis. *PLoS ONE* 4(2):e4468
- Singh N, Das D, Singh A, Mohan ML (2010a) Prion protein and metal interaction: physiological and pathological implications. *Curr Issues Mol Biol* 12(2):99–108
- Singh N, Singh A, Das D, Mohan ML (2010b) Redox control of prion and disease pathogenesis. *Antioxid Redox Signal* 12:1271–1294
- Skinner AC, Steiner MJ, Henderson FW, Perrin EM (2010) Multiple markers of inflammation and weight status: cross-sectional analyses throughout childhood. *Pediatrics* 125(4):e801–809
- Slemmer JE, Shacka JJ, Sweeney MI, Weber JT (2008) Antioxidants and free radical scavengers for the treatment of stroke, traumatic brain injury and aging. *Curr Med Chem* 15(4):404–414
- Smallbone K, Simeonidis E, Broomhead DS, Kell DB (2007) Something from nothing: bridging the gap between constraint-based and kinetic modelling. *FEBS J* 274:5576–5585
- Smirnova GV, Samoylova ZY, Muzyka NG, Oktyabrsky ON (2009) Influence of polyphenols on *Escherichia coli* resistance to oxidative stress. *Free Radic Biol Med* 46(6):759–768
- Smith AG, Chernova T (2009) Disruption of heme synthesis by polyhalogenated aromatics. *Adv Mol Toxicol* 3:169–218
- Smith AG, Clothier B, Robinson S, Scullion MJ, Carthew P, Edwards R et al (1998) Interaction between iron metabolism and 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin in mice with variants of the Ahr gene: a hepatic oxidative mechanism. *Mol Pharmacol* 53(1):52–61
- Smith AG, Elder GH (2010) Complex gene-chemical interactions: hepatic uroporphyrin as a paradigm. *Chem Res Toxicol* 23(4):712–723

- Smith C, Mitchinson MJ, Aruoma OI, Halliwell B (1992) Stimulation of lipid peroxidation and hydroxyl radical generation by the contents of human atherosclerotic lesions. *Biochem J* 286: 901–905
- Smith DG, Cappai R, Barnham KJ (2007) The redox chemistry of the Alzheimer's disease amyloid beta peptide. *Biochim Biophys Acta* 1768(8):1976–1990
- Smith MA, Perry G, Richey PL, Sayre LM, Anderson VE, Beal MF et al (1996) Oxidative damage in Alzheimer's. *Nature* 382(6587):120–121
- Smith MA, Harris PLR, Sayre LM, Perry G (1997a) Iron accumulation in Alzheimer disease is a source of redox-generated free radicals. *Proc Natl Acad Sci* 94(18):9866–9868
- Smith MA, Richey Harris PL, Sayre LM, Beckman JS, Perry G (1997b) Widespread peroxynitrite-mediated damage in Alzheimer's disease. *J Neurosci* 17(8):2653–2657
- Smith MA, Rottkamp CA, Nunomura A, Raina AK, Perry G (2000) Oxidative stress in Alzheimer's disease. *Biochim Biophys Acta* 1502(1):139–144
- Smith MA, Zhu X, Tabaton M, Liu G, McKeel DW Jr, Cohen ML et al (2010) Increased iron and free radical generation in preclinical Alzheimer disease and mild cognitive impairment. *J Alzheimers Dis* 19(1):363–372
- Sobrin L, Maller JB, Neale BM, Reynolds RC, Fagerness JA, Daly MJ et al (2010) Genetic profile for five common variants associated with age-related macular degeneration in densely affected families: a novel analytic approach. *Eur J Hum Genet* 18(4):496–501
- Sofic E, Paulus W, Jellinger K, Riederer P, Youdim MB (1991) Selective increase of iron in substantia nigra zona compacta of parkinsonian brains. *J Neurochem* 56(3):978–982
- Sohn YS, Breuer W, Munnich A, Cabantchik ZI (2008) Redistribution of accumulated cell iron: a modality of chelation with therapeutic implications. *Blood* 111(3):1690–1699
- Soloniuk DS, Perkins E, Wilson JR (1992) Use of allopurinol and deferoxamine in cellular protection during ischemia. *Surg Neurol* 38(2):110–113
- Song MK, Chung JS, Seol YM, Shin HJ, Choi YJ, Cho GJ (2009) Elevation of serum ferritin is associated with the outcome of patients with newly diagnosed multiple myeloma. *Korean J Intern Med* 24(4):368–373
- Sørensen AR, Thulstrup AM, Hansen J, Ramlau-Hansen CH, Meersohn A, Skytthe A et al (2007) Risk of lung cancer according to mild steel and stainless steel welding. *Scand J Work Environ Health* 33(5):379–386
- Sørensen JG, Holmstrup M, Sarup P, Loeschcke V (2009) Evolutionary theory and studies of model organisms predict a cautiously positive perspective on the therapeutic use of hormesis for healthy aging in humans. *Dose Response* 8(1):53–57
- Sow FB, Alvarez GR, Gross RP, Satoskar AR, Schlesinger LS, Zwilling BS et al (2009) Role of STAT1, NF-kappaB, and C/EBPbeta in the macrophage transcriptional regulation of hepcidin by mycobacterial infection and IFN-gamma. *J Leukoc Biol* 86(5):1247–1258
- Sparaco M, Gaeta LM, Santorelli FM, Passarelli C, Tozzi G, Bertini E et al (2009) Friedreich's ataxia: oxidative stress and cytoskeletal abnormalities. *J Neurol Sci* 287(1–2):111–118
- Spasojević I, Stević Z, Nikolić-Kokić A, Jones DR, Blagojević D, Spasić MB (2010) Different roles of radical scavengers—ascorbate and urate in the cerebrospinal fluid of amyotrophic lateral sclerosis patients. *Redox Rep* 15(2):81–86
- Spencer JP, Abd El Mohsen MM, Minihane AM, Mathers JC (2008) Biomarkers of the intake of dietary polyphenols: strengths, limitations and application in nutrition research. *Br J Nutr* 99(1):12–22
- Sporn MB, Liby K, Yore MM, Suh N, Albini A, Honda T et al (2007) Platforms and networks in triterpenoid pharmacology. *Drug Dev Res* 68(4):174–182
- Squadrito GL, Pryor WA (1998) Oxidative chemistry of nitric oxide: the roles of superoxide, peroxynitrite, and carbon dioxide. *Free Radic Biol Med* 25(4–5):392–403
- Sriram K, Lin GX, Jefferson AM, Roberts JR, Chapman RS, Chen BT et al (2010) Dopaminergic neurotoxicity following pulmonary exposure to manganese-containing welding fumes. *Arch Toxicol*
- Sriskandan S, Altmann DM (2008) The immunology of sepsis. *J Pathol* 214(2):211–223
- Stacey AR, Norris PJ, Qin L, Haygreen EA, Taylor E, Heitman J et al (2009) Induction of a striking systemic cytokine cascade prior to peak viremia in acute human immunodeficiency virus type 1 infection, in contrast to more modest and delayed responses in acute hepatitis B and C virus infections. *J Virol* 83(8):3719–3733
- Stadler N, Lindner RA, Davies MJ (2004) Direct detection and quantification of transition metal ions in human atherosclerotic plaques: evidence for the presence of elevated levels of iron and copper. *Arterioscler Thromb Vasc Biol* 24(5):949–954
- Stadler N, Stanley N, Heeneman S, Vacata V, Daemen MJ, Bannon PG et al (2008) Accumulation of zinc in human atherosclerotic lesions correlates with calcium levels but does not protect against protein oxidation. *Arterioscler Thromb Vasc Biol* 28(5):1024–1030
- Stankiewicz J, Panter SS, Neema M, Arora A, Batt CE, Bakshi R (2007) Iron in chronic brain disorders: imaging and neurotherapeutic implications. *Neurotherapeutics* 4(3):371–386
- Stanley N, Stadler N, Woods AA, Bannon PG, Davies MJ (2006) Concentrations of iron correlate with the extent of protein, but not lipid, oxidation in advanced human atherosclerotic lesions. *Free Radic Biol Med* 40(9):1636–1643
- Starley BQ, Calcagno CJ, Harrison SA (2010) Nonalcoholic fatty liver disease and hepatocellular carcinoma: a weighty connection. *Hepatology* 51(5):1820–1832
- Stegmann J, Grohmann G (2003) Hypothesis generation guided by co-word clustering. *Scientometrics* 56(1):111–135
- Stein SE, Heller SR, Tchekhovski D (2003) An open standard for chemical structure representation—the IUPAC chemical identifier. In: *Proceedings of 2003 Nimes international chemical information conference*, pp 131–143
- Stelling J, Sauer U, Szallasi Z, Doyle F J 3rd, Doyle J (2004) Robustness of cellular functions. *Cell* 118(6):675–685
- Stevens KB, Del Rio Vilas VJ, Guitian J (2009) Classical sheep scrapie in Great Britain: spatial analysis and identification of environmental and farm-related risk factors. *BMC Vet Res* 5:33
- Stintzi A, Barnes C, Xu J, Raymond KN (2000) Microbial iron transport via a siderophore shuttle: a membrane ion transport paradigm. *Proc Natl Acad Sci USA* 97(20):10691–10696
- Stocker R, Keaney JF (2004) Role of oxidative modifications in atherosclerosis. *Physiol Rev* 84(4):1381–1478
- Stocker R, Keaney JF (2005) New insights on oxidative stress in the artery wall. *J Thromb Haemostasis* 3(8):1825–1834
- Stohs SJ, Bagchi D (1995) Oxidative mechanisms in the toxicity of metal ions. *Free Radic Biol Med* 18(2):321–336
- Stolovitzky G, Califano A (2006) Reverse engineering biological networks: opportunities and challenges in computational methods for pathway inference, vol 1158. *New York Academy of Sciences*, New York
- Stolovitzky G, Kahlem P, Califano A (2009) The challenges of systems biology: community efforts to harness biological complexity, vol 1158. *New York Academy of Sciences*, New York
- Stommel JM, Kimmelman AC, Ying H, Nabioullin R, Ponugoti AH, Wiedemeyer R et al (2007) Coactivation of receptor tyrosine kinases affects the response of tumor cells to targeted therapies. *Science* 318(5848):287–290

- Storey JA, Connor RF, Lewis ZT, Hurd D, Pomper G, Keung YK et al (2009) The transplant iron score as a predictor of stem cell transplant survival. *J Hematol Oncol* 2:44
- Størmer FC, Hoiby EA (1996) Citrinin, ochratoxin A and iron. Possible implications for their biological function and induction of nephropathy. *Mycopathologia* 134(2):103–107
- Streck EL, Comim CM, Barichello T, Quevedo J (2008) The septic brain. *Neurochem Res* 33(11):2171–2177
- Sturm B, Bistrich U, Schranzhofer M, Sarsero JP, Rauhen U, Scheiber-Mojdehkar B et al (2005) Friedreich's ataxia, no changes in mitochondrial labile iron in human lymphoblasts and fibroblasts: a decrease in antioxidative capacity? *J Biol Chem* 280(8):6701–6708
- Subramanian V, Ferrante AW Jr (2009) Obesity, inflammation, and macrophages. *Nestle Nutr Workshop Ser Pediatr Program* 63:151–159 (discussion 159–162, 259–168)
- Suchy J, Chan A, Shea TB (2009) Dietary supplementation with a combination of alpha-lipoic acid, acetyl-L-carnitine, glycerophosphocoline, docosahexaenoic acid, and phosphatidylserine reduces oxidative damage to murine brain and improves cognitive performance. *Nutr Res* 29(1):70–74
- Sugihara N, Arakawa T, Ohnishi M, Furuno K (1999) Anti- and pro-oxidative effects of flavonoids on metal-induced lipid hydroperoxide-dependent lipid peroxidation in cultured hepatocytes loaded with alpha-linolenic acid. *Free Radic Biol Med* 27(11–12):1313–1323
- Sugimoto R, Fujita N, Tomosugi N, Hara N, Miyachi H, Tanaka H et al (2009) Impaired regulation of serum hepcidin during phlebotomy in patients with chronic hepatitis C. *Hepatol Res* 39(6):619–624
- Suh KS, Chon S, Oh S, Kim SW, Kim JW, Kim YS et al (2009) Prooxidative effects of green tea polyphenol (–)-epigallocatechin-3-gallate on the HIT-T15 pancreatic beta cell line. *Cell Biol Toxicol*
- Sullivan JL (1981) Iron and the sex difference in heart-disease risk. *Lancet* 1(8233):1293–1294
- Sullivan JL (2001) Misconceptions in the debate on the iron hypothesis. *J Nutr Biochem* 12(1):33–37
- Sullivan JL (2003) Are menstruating women protected from heart disease because of, or in spite of, estrogen? Relevance to the iron hypothesis. *Amer Heart J* 145(2):190–194
- Sullivan JL (2004) Is stored iron safe? *J Lab Clin Med* 144(6):280–284
- Sullivan JL (2007) Macrophage iron, hepcidin, and atherosclerotic plaque stability. *Exp Biol Med (Maywood)* 232(8):1014–1020
- Sullivan JL (2009) Iron in arterial plaque: a modifiable risk factor for atherosclerosis. *Biochim Biophys Acta* 1790(7):718–723
- Sumida Y, Yoshikawa T, Okanou T (2009) Role of hepatic iron in non-alcoholic steatohepatitis. *Hepatol Res* 39(3):213–222
- Sun L, Franco OH, Hu FB, Cai L, Yu Z, Li H et al (2008) Ferritin concentrations, metabolic syndrome, and type 2 diabetes in middle-aged and elderly chinese. *J Clin Endocrinol Metab* 93(12):4690–4696
- Sun X, Ge R, Cai Z, Sun H, He QY (2009) Iron depletion decreases proliferation and induces apoptosis in a human colonic adenocarcinoma cell line, CaCO₂. *J Inorg Biochem* 103(7):1074–1081
- Suntharalingam G, Perry MR, Ward S, Brett SJ, Castello-Cortes A, Brunner MD et al (2006) Cytokine storm in a phase 1 trial of the anti-CD28 monoclonal antibody TGN1412. *N Engl J Med* 355(10):1018–1028
- Surh YJ, Kundu JK, Na HK, Lee JS (2005) Redox-sensitive transcription factors as prime targets for chemoprevention with anti-inflammatory and antioxidative phytochemicals. *J Nutr* 135(12 Suppl):2993S–3001S
- Surh YJ, Na HK (2008) NF-kappaB and Nrf2 as prime molecular targets for chemoprevention and cytoprotection with anti-inflammatory and antioxidant phytochemicals. *Genes Nutr* 2(4):313–317
- Sutedja NA, Sinke RJ, Van Vught PW, Van der Linden MW, Wokke JH, Van Duijn CM et al (2007) The association between H63D mutations in HFE and amyotrophic lateral sclerosis in a Dutch population. *Arch Neurol* 64(1):63–67
- Sweeny GD, Jones KG, Cole FM, Basford D, Krestynski F (1979) Iron deficiency prevents liver toxicity of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin. *Science* 204(4390):332–335
- Switzer JA, Hess DC, Nichols FT, Adams RJ (2006) Pathophysiology and treatment of stroke in sickle-cell disease: present and future. *Lancet Neurol* 5(6):501–512
- Szabo C (1996) The pathophysiological role of peroxynitrite in shock, inflammation, and ischemia-reperfusion injury. *Shock* 6(2):79–88
- Szabó C, Ischiropoulos H, Radi R (2007) Peroxynitrite: biochemistry, pathophysiology and development of therapeutics. *Nat Rev Drug Discov* 6(8):662–680
- Szajerka T, Jablecki J (2007) Kaposi's sarcoma revisited. *AIDS Rev* 9(4):230–236
- Tabner BJ, El-Agnaf OM, German MJ, Fullwood NJ, Allsop D (2005) Protein aggregation, metals and oxidative stress in neurodegenerative diseases. *Biochem Soc Trans* 33(Pt 5):1082–1086
- Tacchini L, Gammella E, De Ponti C, Recalcati S, Cairo G (2008) Role of HIF-1 and NF-kappaB transcription factors in the modulation of transferrin receptor by inflammatory and anti-inflammatory signals. *J Biol Chem* 283(30):20674–20686
- Taher AT, Musallam KM, Karimi M, El-Beshlawy A, Belhoul K, Daar S et al (2010) Overview on practices in thalassemia intermedia management aiming for lowering complication rates across a region of endemicity: the OPTIMAL CARE study. *Blood* 115(10):1886–1892
- Tamgüney G, Giles K, Glidden DV, Lessard P, Wille H, Tremblay P et al (2008) Genes contributing to prion pathogenesis. *J Gen Virol* 89(Pt 7):1777–1788
- Tan JS, Wang JJ, Flood V, Rohtchina E, Smith W, Mitchell P (2008) Dietary antioxidants and the long-term incidence of age-related macular degeneration: the Blue Mountains eye study. *Ophthalmology* 115(2):334–341
- Tan KT, Lip GY (2008) Imaging of the unstable plaque. *Int J Cardiol* 127(2):157–165
- Tanaka N, Horiuchi A, Yamaura T, Komatsu M, Tanaka E, Kiyosawa K (2007) Efficacy and safety of 6-month iron reduction therapy in patients with hepatitis C virus-related cirrhosis: a pilot study. *J Gastroenterol* 42(1):49–55
- Tanaka N, Horiuchi A, Yamaura T, Komatsu M, Yokoyama T, Okaniwa S et al (2009) Efficacy and safety of addition of minor bloodletting (petit phlebotomy) in hepatitis C virus-infected patients receiving regular glycyrrhizin injections. *J Gastroenterol* 44(6):577–582
- Tanaka N, Kiyosawa K (2004) Phlebotomy: a promising treatment for chronic hepatitis C. *J Gastroenterol* 39(6):601–603
- Tang TY, Muller KH, Graves MJ, Li ZY, Walsh SR, Young V et al (2009) Iron oxide particles for atherosclerosis imaging. *Arterioscler Thromb Vasc Biol* 29(7):1001–1008
- Taqueti VR, Mitchell RN, Lichtman AH (2006) Protecting the pump: controlling myocardial inflammatory responses. *Annu Rev Physiol* 68:67–95
- Taupin P (2007) Protocols for studying adult neurogenesis: insights and recent developments. *Regen Med* 2(1):51–62
- Tedgui A, Mallat Z (2006) Cytokines in atherosclerosis: pathogenic and regulatory pathways. *Physiol Rev* 86(2):515–581
- Teixeira KC, Soares FS, Rocha LG, Silveira PC, Silva LA, Valença SS et al (2008) Attenuation of bleomycin-induced lung injury

- and oxidative stress by N-acetylcysteine plus deferoxamine. *Pulm Pharmacol Ther* 21(2):309–316
- Tenopoulou M, Doulias PT, Barbouti A, Brunk U, Galaris D (2005) Role of compartmentalized redox-active iron in hydrogen peroxide-induced DNA damage and apoptosis. *Biochem J* 387(Pt 3):703–710
- Tenopoulou M, Kurz T, Doulias PT, Galaris D, Brunk UT (2007) Does the calcein-AM method assay the total cellular 'labile iron pool' or only a fraction of it? *Biochem J* 403(2):261–266
- Terao J (2009) Dietary flavonoids as antioxidants. *Forum Nutr* 61:87–94
- Terman A, Brunk UT (2004) Lipofuscin. *Int J Biochem Cell Biol* 36(8):1400–1404
- Terman A, Brunk UT (2006) Oxidative stress, accumulation of biological 'garbage', and aging. *Antioxid Redox Signal* 8(1–2):197–204
- Terman A, Gustafsson B, Brunk UT (2006a) The lysosomal-mitochondrial axis theory of postmitotic aging and cell death. *Chem Biol Interact* 163(1–2):29–37
- Terman A, Kurz T, Gustafsson B, Brunk UT (2006b) Lysosomal labilization. *IUBMB Life* 58(9):531–539
- Terman A, Kurz T, Navratil M, Arriaga EA, Brunk UT (2010) Mitochondrial turnover and aging of long-lived postmitotic cells: the mitochondrial-lysosomal axis theory of aging. *Antioxid Redox Signal* 12(4):503–535
- Thannickal VJ, Fanburg BL (2000) Reactive oxygen species in cell signaling. *Am J Physiol Lung Cell Mol Physiol* 279(6):L1005–1028
- Theil EC (2007) Coordinating responses to iron and oxygen stress with DNA and mRNA promoters: the ferritin story. *Biometals* 20(3–4):513–521
- Thephinlap C, Phisalaphong C, Fucharoen S, Porter JB, Srichairatanakool S (2009) Efficacy of curcuminoids in alleviation of iron overload and lipid peroxidation in thalassemic mice. *Med Chem* 5:475–482
- Thiele I, Palsson BØ (2010) A protocol for generating a high-quality genome-scale metabolic reconstruction. *Nat Protoc* 5(1):93–121
- Thomas M, Jankovic J (2004) Neurodegenerative disease and iron storage in the brain. *Curr Opin Neurol* 17(4):437–442
- Thomas C, Mackey MM, Diaz AA, Cox DP (2009a) Hydroxyl radical is produced via the Fenton reaction in submitochondrial particles under oxidative stress: implications for diseases associated with iron accumulation. *Redox Rep* 14(3):102–108
- Thomas P, Wang YJ, Zhong JH, Kosaraju S, O'Callaghan NJ, Zhou XF et al (2009b) Grape seed polyphenols and curcumin reduce genomic instability events in a transgenic mouse model for Alzheimer's disease. *Mutat Res* 661(1–2):25–34
- Thomas SR, Schulz E, Keane JF (2006) Hydrogen peroxide restrains endothelium-derived nitric oxide bioactivity—role for iron-dependent oxidative stress. *Free Radic Biol Med* 41(4):681–688
- Thompson KJ, Shoham S, Connor JR (2001) Iron and neurodegenerative disorders. *Brain Res Bull* 55(2):155–164
- Thougaard AV, Langer SW, Hainau B, Grauslund M, Juhl BR, Jensen PB et al (2010) A murine experimental anthracycline extravasation model: pathology and study of the involvement of topoisomerase II alpha and iron in the mechanism of tissue damage. *Toxicology* 269(1):67–72
- Tian H, Maddox IS, Ferguson LR, Shu Q (2010) Evaluation of the cytoprotective effects of bovine lactoferrin against intestinal toxins using cellular model systems. *Biometals* 23:589–592
- Tielsch JM, Javitt JC, Coleman A, Katz J, Sommer A (1995) The prevalence of blindness and visual impairment among nursing home residents in Baltimore. *N Engl J Med* 332(18):1205–1209
- Tjoa CW, Benedict RH, Weinstock-Guttman B, Fabiano AJ, Bakshi R (2005) MRI T2 hypointensity of the dentate nucleus is related to ambulatory impairment in multiple sclerosis. *J Neurol Sci* 234(1–2):17–24
- Tomlin CJ, Axelrod JD (2007) Biology by numbers: mathematical modelling in developmental biology. *Nat Rev Genet* 8(5):331–340
- Torreilles F, Salman-Tabcheh S, Guérin M, Torreilles J (1999) Neurodegenerative disorders: the role of peroxynitrite. *Brain Res Brain Res Rev* 30(2):153–163
- Toyokuni S (2002) Iron and carcinogenesis: from Fenton reaction to target genes. *Redox Rep* 7(4):189–197
- Toyokuni S (2009a) Mechanisms of asbestos-induced carcinogenesis. *Nagoya J Med Sci* 71(1–2):1–10
- Toyokuni S (2009b) Role of iron in carcinogenesis: cancer as a ferrotoxic disease. *Cancer Sci* 100(1):9–16
- Toyokuni S, Sagripanti JL (1996) Association between 8-hydroxy-2'-deoxyguanosine formation and DNA strand breaks mediated by copper and iron. *Free Radic Biol Med* 20(6):859–864
- Trachootham D, Alexandre J, Huang P (2009) Targeting cancer cells by ROS-mediated mechanisms: a radical therapeutic approach? *Nat Rev Drug Discov* 8(7):579–591
- Trachootham D, Lu W, Ogasawara MA, Nilsa RD, Huang P (2008) Redox regulation of cell survival. *Antioxid Redox Signal* 10(8):1343–1374
- Traoré, H. N., & Meyer, D. (2004). The effect of iron overload on in vitro HIV-1 infection. *J Clin Virol* 31(Suppl 1):S92-S98.
- Traoré HN, Meyer D (2007) Necrosis of host cells and survival of pathogens following iron overload in an in vitro model of co-infection with human immunodeficiency virus (HIV) and *Mycobacterium tuberculosis*. *Int J Antimicrob Agents* 29(4):465–470
- Trinder D, Ayonrinde OT, Olynyk JK (2008) HCV, iron, and oxidative stress: the new choreography of hepcidin. *Gastroenterology* 134(1):348–351
- Truksa J, Kovář J, Valenta T, Ehrlichová M, Polák J, Naumann PW (2003) Iron deprivation induces apoptosis independently of p53 in human and murine tumour cells. *Cell Prolif* 36(4):199–213
- Tsimikas S, Willeit J, Knoflach M, Mayr M, Egger G, Notdurfter M et al (2009) Lipoprotein-associated phospholipase A2 activity, ferritin levels, metabolic syndrome, and 10-year cardiovascular and non-cardiovascular mortality: results from the Bruneck study. *Eur Heart J* 30(1):107–115
- Tung BY, Emond MJ, Bronner MP, Raaka SD, Cotler SJ, Kowdley KV (2003) Hepatitis C, iron status, and disease severity: relationship with HFE mutations. *Gastroenterology* 124(2):318–326
- Tuntipipat S, Zeder C, Siriprapa P, Charoenkiatkul S (2009) Inhibitory effects of spices and herbs on iron availability. *Int J Food Sci Nutr* 60(Suppl 1):43–55
- Tuomainen TP, Loft S, Nyysönen K, Punnonen K, Salonen JT, Poulsen HE (2007) Body iron is a contributor to oxidative damage of DNA. *Free Radic Res* 41(3):324–328
- Tuomainen TP, Punnonen K, Nyysönen K, Salonen JT (1998) Association between body iron stores and the risk of acute myocardial infarction in men. *Circulation* 97(15):1461–1466
- Turner NJ (2009) Directed evolution drives the next generation of biocatalysts. *Nat Chem Biol* 5(8):567–573
- Ulloa L, Brunner M, Ramos L, Deitch EA (2009) Scientific and clinical challenges in sepsis. *Curr Pharm Des* 15(16):1918–1935
- Ulrich-Merzenich G, Zeitler H, Vetter H, Kraft K (2009) Synergy research: vitamins and secondary plant components in the maintenance of the redox-homeostasis and in cell signaling. *Phytomedicine* 16(1):2–16
- Upadhyay D, Kamp DW (2003) Asbestos-induced pulmonary toxicity: role of DNA damage and apoptosis. *Exp Biol Med* (Maywood) 228(6):650–659

- Usenko CY, Harper SL, Tanguay RL (2008) Fullerene C60 exposure elicits an oxidative stress response in embryonic zebrafish. *Toxicol Appl Pharmacol* 229(1):44–55
- Usuki F, Yasutake A, Umehara F, Tokunaga H, Matsumoto M, Eto K et al (2001) In vivo protection of a water-soluble derivative of vitamin E, Trolox, against methylmercury-intoxication in the rat. *Neurosci Lett* 304(3):199–203
- Utzschneider KM, Kowdley KV (2010) Hereditary hemochromatosis and diabetes mellitus: implications for clinical practice. *Nat Rev Endocrinol* 6(1):26–33
- Valavanidis A, Vlachogianni T, Fiotakis C (2009) 8-hydroxy-2'-deoxyguanosine (8-OHdG): a critical biomarker of oxidative stress and carcinogenesis. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev* 27(2):120–139
- Valavanidis A, Vlahoyianni T, Fiotakis K (2005) Comparative study of the formation of oxidative damage marker 8-hydroxy-2'-deoxyguanosine (8-OHdG) adduct from the nucleoside 2'-deoxyguanosine by transition metals and suspensions of particulate matter in relation to metal content and redox reactivity. *Free Radic Res* 39(10):1071–1081
- Valdmanis PN, Daoud H, Dion PA, Rouleau GA (2009) Recent advances in the genetics of amyotrophic lateral sclerosis. *Curr Neurol Neurosci Rep* 9(3):198–205
- Valenti L, Dongiovanni P, Fargion S, Fracanzani AL (2008) Iron genes, dysmetabolism and fibrosis in chronic hepatitis C. *J Hepatol* 48(3):513–514 (author reply 514–515)
- Valenti L, Swinkels DW, Burdick L, Dongiovanni P, Tjalsma H, Motta BM et al (2010) Serum ferritin levels are associated with vascular damage in patients with nonalcoholic fatty liver disease. *Nutr Metab Cardiovasc Dis* (in press)
- Valenti L, Valenti G, Como G, Burdick L, Santorelli G, Dongiovanni P et al (2007) HFE gene mutations and oxidative stress influence serum ferritin, associated with vascular damage, in hemodialysis patients. *Am J Nephrol* 27(1):101–107
- Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J (2007) Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 39(1):44–84
- Valko M, Morris H, Cronin MTD (2005) Metals, toxicity and oxidative stress. *Curr Med Chem* 12(10):1161–1208
- Valvassori SS, Petronilho FC, Reus GZ, Steckert AV, Oliveira VB, Boeck CR et al (2008) Effect of N-acetylcysteine and/or deferoxamine on oxidative stress and hyperactivity in an animal model of mania. *Prog Neuropsychopharmacol Biol Psychiatry* 32(4):1064–1068
- van Asbeck BS, Georgiou NA, van der Bruggen T, Oudshoorn M, Nottet HS, Marx JJ (2001) Anti-HIV effect of iron chelators: different mechanisms involved. *J Clin Virol* 20(3):141–147
- van Laar VS, Berman SB (2009) Mitochondrial dynamics in Parkinson's disease. *Exp Neurol* 218(2):247–256
- van Leuven SI, Franssen R, Kastelein JJ, Levi M, Stroes ES, Tak PP (2008) Systemic inflammation as a risk factor for atherothrombosis. *Rheumatology (Oxford)* 47(1):3–7
- van Oostrom AJ, van Wijk J, Cabezas MC (2004) Lipaemia, inflammation and atherosclerosis: novel opportunities in the understanding and treatment of atherosclerosis. *Drugs* 64(Suppl 2):19–41
- Vance C, Rogelj B, Hortobagyi T, De Vos KJ, Nishimura AL, Sreedharan J et al (2009) Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323(5918):1208–1211
- Vanden Berghe T, Vanlangenakker N, Parthoens E, Deckers W, Devos M, Festjens N et al (2009) Necroptosis, necrosis and secondary necrosis converge on similar cellular disintegration features. *Cell Death Differ* 17:922–930
- Vanlangenakker N, Vanden Berghe T, Krysko DV, Festjens N, Vandenabeele P (2008) Molecular mechanisms and pathophysiology of necrotic cell death. *Curr Mol Med* 8(3):207–220
- Vaupel JW (2010) Biodemography of human ageing. *Nature* 464(7288):536–542
- Vaz SM, Augusto O (2008) Inhibition of myeloperoxidase-mediated protein nitration by tempol: kinetics, mechanism, and implications. *Proc Natl Acad Sci USA* 105(24):8191–8196
- Vázquez-Manrique RP, González-Cabo P, Ros S, Aziz H, Baylis HA, Palau F (2006) Reduction of *Caenorhabditis elegans* frataxin increases sensitivity to oxidative stress, reduces lifespan, and causes lethality in a mitochondrial complex II mutant. *Faseb J* 20(1):172–174
- Vázquez Roncero A, Janer del Valle C, Maestro Durán R, Graciani Constanse E (1983) New aniline derivatives in cooking oils associated with toxic oil syndrome. *Lancet* 2(8357):1024–1025
- Vélez-Pardo C, Del Río MJ, Verschuere H, Ebinger G, Vauquelin G (1997) Dopamine and iron induce apoptosis in PC12 cells. *Pharmacol Toxicol* 80(2):76–84
- Venturini D, Simao AN, Barbosa DS, Lavado EL, Narciso VE, Dichi I et al (2010) Increased oxidative stress, decreased total antioxidant capacity, and iron overload in untreated patients with chronic hepatitis C. *Dig Dis Sci* 55:1120–1127
- Verduzco LA, Nathan DG (2009) Sick cell disease and stroke. *Blood* 114(25):5117–5125
- Vichinsky E, Butensky E, Fung E, Hudes M, Theil E, Ferrell L et al (2005) Comparison of organ dysfunction in transfused patients with SCD or beta thalassemia. *Am J Hematol* 80(1):70–74
- Victor VM, Rocha M, De la Fuente M (2004) Immune cells: free radicals and antioxidants in sepsis. *Int Immunopharmacol* 4(3):327–347
- Victor VM, Rocha M, Esplugues JV, De la Fuente M (2005) Role of free radicals in sepsis: antioxidant therapy. *Curr Pharm Des* 11(24):3141–3158
- Villéger AC, Pettifer SR, Kell DB (2010) Arcadia: a visualization tool for metabolic pathways. *Bioinformatics* 20(11):1470–1471
- Visseren FLJ, Verkerk MSA, van der Bruggen T, Marx JJM, van Asbeck BS, Diepersloot RJA (2002) Iron chelation and hydroxyl radical scavenging reduce the inflammatory response of endothelial cells after infection with *Chlamydia pneumoniae* or influenza A. *Eur J Clin Invest* 32(Suppl 1):84–90
- Vlachodimitropoulou E, Naftalin RJ, Sharp PA (2010) Quercetin is a substrate for the transmembrane oxidoreductase, Dcytb. *Free Radic Biol Med* 48:1366–1369
- Vlašić Ž, Dodig S, Čepelak I, Zrinski Topić RZ, Živčić J, Nogalo B et al (2009) Iron and ferritin concentrations in exhaled breath condensate of children with asthma. *J Asthma* 46(1):81–85
- Vllesis AA, Goldman RK, Trunkey DD (1995) New concepts in the pathophysiology of oxygen metabolism during sepsis. *Br J Surg* 82(7):870–876
- Voloboueva LA, Killilea DW, Atamna H, Ames BN (2007) N-tert-butyl hydroxylamine, a mitochondrial antioxidant, protects human retinal pigment epithelial cells from iron overload: relevance to macular degeneration. *FASEB J* 21(14):4077–4086
- Vucenic I, Shamsuddin AM (2006) Protection against cancer by dietary IP6 and inositol. *Nutr Cancer* 55(2):109–125
- Vucic S, Kiernan MC (2009) Pathophysiology of neurodegeneration in familial amyotrophic lateral sclerosis. *Curr Mol Med* 9(3):255–272
- Vysheirsky V, Girolami MA (2008) Bayesian ranking of biochemical system models. *Bioinformatics* 24(6):833–839
- Wagner A (2005) Robustness, evolvability, and neutrality. *FEBS Lett* 579(8):1772–1778
- Wagner KR, Sharp FR, Ardizzone TD, Lu A, Clark JF (2003) Heme and iron metabolism: role in cerebral hemorrhage. *J Cereb Blood Flow Metab* 23(6):629–652

- Wahba ZZ, Murray WJ, Stohs SJ (1990) Desferrioxamine-induced alterations in hepatic iron distribution, DNA damage and lipid peroxidation in control and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin-treated rats. *J Appl Toxicol* 10(2):119–124
- Waldron KJ, Rutherford JC, Ford D, Robinson NJ (2009) Metalloproteins and metal sensing. *Nature* 460(7257):823–830
- Walker NM, Stuart KA, Ryan RJ, Desai S, Saab S, Nicol JA et al (2010) Serum ferritin concentration predicts mortality in patients awaiting liver transplantation. *Hepatology* 51(5):1683–1691
- Walter U, Wagner S, Horowski S, Benecke R, Zettl UK (2009) Transcranial brain sonography findings predict disease progression in multiple sclerosis. *Neurology* 73(13):1010–1017
- Walters AS, Rye DB (2009) Review of the relationship of restless legs syndrome and periodic limb movements in sleep to hypertension, heart disease, and stroke. *Sleep* 32(5):589–597
- Walters RH, Murphy RM (2009) Examining polyglutamine peptide length: a connection between collapsed conformations and increased aggregation. *J Mol Biol* 393(4):978–992
- Wandersman C, Delepelaire P (2004) Bacterial iron sources: from siderophores to hemophores. *Annu Rev Microbiol* 58:611–647
- Wang H, Ma S (2008) The cytokine storm and factors determining the sequence and severity of organ dysfunction in multiple organ dysfunction syndrome. *Am J Emerg Med* 26(6):711–715
- Wang XS, Lee S, Simmons Z, Boyer P, Scott K, Liu W et al (2004) Increased incidence of the Hfe mutation in amyotrophic lateral sclerosis and related cellular consequences. *J Neurol Sci* 227(1):27–33
- Wang J, Gines S, MacDonald ME, Gusella JF (2005) Reversal of a full-length mutant huntingtin neuronal cell phenotype by chemical inhibitors of polyglutamine-mediated aggregation. *BMC Neurosci* 6:1
- Wang J, Wang G, Ansari GA, Khan MF (2008) Activation of oxidative stress-responsive signaling pathways in early splenotoxic response of aniline. *Toxicol Appl Pharmacol* 230(2):227–234
- Wang T, Craig EA (2008) Binding of yeast frataxin to the scaffold for Fe-S cluster biogenesis. *Isu. J Biol Chem* 283(18):12674–12679
- Wang X, Zhao X (2009) Contribution of oxidative damage to antimicrobial lethality. *Antimicrob Agents Chemother* 53(4):1395–1402
- Wang HQ, Xu YX, Zhao XY, Zhao H, Yan J, Sun XB et al (2009a) Overexpression of F₀F₁-ATP synthase alpha suppresses mutant huntingtin aggregation and toxicity in vitro. *Biochem Biophys Res Commun* 390(4):1294–1298
- Wang J, Du XX, Jiang H, Xie JX (2009b) Curcumin attenuates 6-hydroxydopamine-induced cytotoxicity by anti-oxidation and nuclear factor-kappaB modulation in MES23.5 cells. *Biochem Pharmacol* 78(2):178–183
- Wang ZY, Zhao KK, Song ZM, Shen LJ, Qu J (2009c) Erythropoietin as a novel therapeutic agent for atrophic age-related macular degeneration. *Med Hypotheses* 72(4):448–450
- Wang YC, Chen BS (2010) Integrated cellular network of transcription regulations and protein-protein interactions. *BMC Syst Biol* 4(1):20
- Wang J, Ma H, Boor PJ, Ramanujam VM, Ansari GA, Khan MF (2010a) Up-regulation of heme oxygenase-1 in rat spleen after aniline exposure. *Free Radic Biol Med* 48(4):513–518
- Wang W, Knovich MA, Coffman LG, Torti FM, Torti SV (2010b) Serum ferritin: past, present and future. *Biochim Biophys Acta*
- Wang X, Moualla D, Wright JA, Brown DR (2010c) Copper binding regulates intracellular alpha-synuclein localisation, aggregation and toxicity. *J Neurochem* 113(3):704–714
- Wardman P, Candeias LP (1996) Fenton chemistry: an introduction. *Rad Res* 145(5):523–531
- Warkentin LM, Auriat AM, Wowk S, Colbourne F (2010) Failure of deferoxamine, an iron chelator, to improve outcome after collagenase-induced intracerebral hemorrhage in rats. *Brain Res* 1309:95–103
- Watanabe J, Nakamachi T, Ogawa T, Naganuma A, Nakamura M, Shioda S et al (2009) Characterization of antioxidant protection of cultured neural progenitor cells (NPC) against methylmercury (MeHg) toxicity. *J Toxicol Sci* 34(3):315–325
- Watt F, Rajendran R, Ren MQ, Tan BKH, Halliwell B (2006) A nuclear microscopy study of trace elements Ca, Fe, Zn and Cu in atherosclerosis. *Nucl Instr Meth Phys Res B* 249:646–652
- Watterson S, Marshall S, Ghazal P (2008) Logic models of pathway biology. *Drug Discov Today* 13(9–10):447–456
- Watts JC, Westaway D (2007) The prion protein family: diversity, rivalry, and dysfunction. *Biochim Biophys Acta* 1772(6):654–672
- Wayne Martin WR (2009) Quantitative estimation of regional brain iron with magnetic resonance imaging. *Parkinsonism Rel Disord* 15S3:S215–S218
- Wedge D, Kell DB (2008) Rapid prediction of optimum population size in genetic programming using a novel genotype-fitness correlation. In: Keizer M et al (eds) GECCO 2008, pp 1315–1322
- Wedge D, Rowe W, Kell DB, Knowles J (2009) In silico modelling of directed evolution: implications for experimental design and stepwise evolution. *J Theor Biol* 257:131–141
- Weinberg ED (1989) Iron, asbestos, and carcinogenicity. *Lancet* 1(8651):1399–1400
- Weinberg ED (2004) Exposing the hidden dangers of iron: what every medical professional should know about the impact of iron on the disease process. Cumberland House
- Weinberg F, Chandel NS (2009) Reactive oxygen species-dependent signaling regulates cancer. *Cell Mol Life Sci* 66(23):3663–3673
- Weinberg GA (2006) Iron status and mortality in HIV infection. *Haematologica* 91(6):721
- Weininger D (1988) SMILES, a chemical language and information system. 1. Introduction to methodology and encoding rules. *J Chem Inf Comput Sci* 28(1):31–36
- Weinreb O, Amit T, Mandel S, Youdim MB (2009) Neuroprotective molecular mechanisms of (–)-epigallocatechin-3-gallate: a reflective outcome of its antioxidant, iron chelating and neurotogenic properties. *Genes Nutr*
- Weinreb O, Amit T, Youdim MB (2007) A novel approach of proteomics and transcriptomics to study the mechanism of action of the antioxidant-iron chelator green tea polyphenol (–)-epigallocatechin-3-gallate. *Free Radic Biol Med* 43(4):546–556
- Weinreb O, Mandel S, Bar-Am O, Yogev-Falach M, Avramovich-Tirosh Y, Amit T et al (2009b) Multifunctional neuroprotective derivatives of rasagiline as anti-Alzheimer's disease drugs. *Neurotherapeutics* 6(1):163–174
- Weisberger J, Emmons F, Gorczyca W (2004) Cytochemical diagnosis of Gaucher's disease by iron stain. *Br J Haematol* 124(6):696
- Weiss G (2010) Genetic mechanisms and modifying factors in hereditary hemochromatosis. *Nat Rev Gastroenterol Hepatol* 7(1):50–58
- Weitzman SA, Graceffa P (1984) Asbestos catalyzes hydroxyl and superoxide radical generation from hydrogen peroxide. *Arch Biochem Biophys* 228(1):373–376
- Weng LC, Yeh WT, Bai CH, Chen HJ, Chuang SY, Chang HY et al (2008) Is ischemic stroke risk related to folate status or other nutrients correlated with folate intake? *Stroke* 39(12):3152–3158
- Wermuth CG (2004) Selective optimization of side activities: another way for drug discovery. *J Med Chem* 47(6):1303–1314
- West AK, Hidalgo J, Eddins D, Levin ED, Aschner M (2008) Metallothionein in the central nervous system: roles in

- protection, regeneration and cognition. *Neurotoxicology* 29(3):489–503
- Westaway D, DeArmond SJ, Cayetano-Canlas J, Groth D, Foster D, Yang SL et al (1994) Degeneration of skeletal muscle, peripheral nerves, and the central nervous system in transgenic mice overexpressing wild-type prion proteins. *Cell* 76(1):117–129
- Westergaard L, Christensen HM, Harris DA (2007) The cellular prion protein (PrP(C)): its physiological function and role in disease. *Biochim Biophys Acta* 1772(6):629–644
- Westerhoff HV, Kell DB (2007) The methodologies of systems biology. In: Boogerd FC, Bruggeman FJ, Hofmeyr J-HS, Westerhoff HV (eds) *Systems biology: philosophical foundations*. Elsevier, Amsterdam, pp 23–70
- White BC, Nayini NR, Krause GS, Aust SD, March GG, Bicknell JS et al (1988) Effect on biochemical markers of brain injury of therapy with deferoxamine or superoxide dismutase following cardiac arrest. *Am J Emerg Med* 6(6):569–576
- White CR, Brock TA, Chang LY, Crapo J, Briscoe P, Ku D et al (1994) Superoxide and peroxynitrite in atherosclerosis. *Proc Natl Acad Sci USA* 91(3):1044–1048
- Whitehurst AW, Bodemann BO, Cardenas J, Ferguson D, Girard L, Peyton M et al (2007) Synthetic lethal screen identification of chemosensitizer loci in cancer cells. *Nature* 446(7137):815–819
- Wijesekera LC, Leigh PN (2009) Amyotrophic lateral sclerosis. *Orphanet J Rare Dis* 4:3
- Wilkinson DJ (2007) Bayesian methods in bioinformatics and computational systems biology. *Brief Bioinform* 8(2):109–116
- Wilkinson SJ, Benson N, Kell DB (2008) Proximate parameter tuning for biochemical networks with uncertain kinetic parameters. *Mol Biosyst* 4:74–97. doi:10.1039/b707506e
- Willerson JT, Ridker PM (2004) Inflammation as a cardiovascular risk factor. *Circulation* 109(21 Suppl 1):II2–II10
- Williams A, Lucassen PJ, Ritchie D, Bruce M (1997) PrP deposition, microglial activation, and neuronal apoptosis in murine scrapie. *Exp Neurol* 144(2):433–438
- Wilson JG, Lindquist JH, Grambow SC, Crook ED, Maher JF (2003) Potential role of increased iron stores in diabetes. *Am J Med Sci* 325(6):332–339
- Wilson RB (2006) Iron dysregulation in Friedreich ataxia. *Semin Pediatr Neurol* 13(3):166–175
- Wimazal F, Nosslinger T, Baumgartner C, Sperr WR, Pfeilstocker M, Valent P (2009) Deferasirox induces regression of iron overload in patients with myelodysplastic syndromes. *Eur J Clin Invest* 39(5):406–411
- Winkelmann G (2002) Microbial siderophore-mediated transport. *Biochem Soc Trans* 30(4):691–696
- Winterbourn CC (1995) Toxicity of iron and hydrogen peroxide: the Fenton reaction. *Toxicol Lett* 82–83:969–974
- Wisniewski T, Sigurdsson EM (2007) Therapeutic approaches for prion and Alzheimer's diseases. *FEBS J* 274(15):3784–3798
- Wolff B, Volzke H, Ludemann J, Robinson D, Vogelgesang D, Staudt A et al (2004) Association between high serum ferritin levels and carotid atherosclerosis in the study of health in Pomerania (SHIP). *Stroke* 35(2):453–457
- Wolozin B, Golts N (2002) Iron and Parkinson's disease. *Neuroscientist* 8(1):22–32
- Won JE, Jeong SH, Chung JI, Lee JH, Hwang SH, Kim JW et al (2009) Hepatic iron, serum ferritin, HFE mutation, and hepatic fibrosis in chronic hepatitis C. *Intervirology* 52(5):239–246
- Wondrak GT (2009) Redox-directed cancer therapeutics: molecular mechanisms and opportunities. *Antioxid Redox Signal* 11(12):3013–3069
- Wong BS, Brown DR, Pan T, Whiteman M, Liu T, Bu X et al (2001a) Oxidative impairment in scrapie-infected mice is associated with brain metals perturbations and altered antioxidant activities. *J Neurochem* 79(3):689–698
- Wong BS, Brown DR, Sy MS (2001b) A Yin-Yang role for metals in prion disease. *Panminerva Medica* 43(4):283–287
- Wong PK, Yu F, Shahangian A, Cheng G, Sun R, Ho CM (2008) Closed-loop control of cellular functions using combinatorial drugs guided by a stochastic search algorithm. *Proc Natl Acad Sci USA* 105(13):5105–5110
- Wong RW, Richa DC, Hahn P, Green WR, Dunaief JL (2007) Iron toxicity as a potential factor in AMD. *Retina* 27(8):997–1003
- Woo PC, Tung ET, Chan KH, Lau CC, Lau SK, Yuen KY (2010) Cytokine profiles induced by the novel swine-origin influenza A/H1N1 virus: implications for treatment strategies. *J Infect Dis* 201(3):346–353
- Wu J, Hua Y, Keep RF, Nakamura T, Hoff JT, Xi G (2003) Iron and iron-handling proteins in the brain after intracerebral hemorrhage. *Stroke* 34(12):2964–2969
- Wu X, Kannan S, Ramanujam VM, Khan MF (2005) Iron release and oxidative DNA damage in splenic toxicity of aniline. *J Toxicol Environ Health A* 68(8):657–666
- Wu X, Patel D, Hasinoff BB (2004) The iron chelating cardioprotective prodrug dexrazoxane does not affect the cell growth inhibitory effects of bleomycin. *J Inorg Biochem* 98(11):1818–1823
- Wu XD, Kumar V, Quinlan JR, Ghosh J, Yang Q, Motoda H et al (2008) Top 10 algorithms in data mining. *Knowl Inf Syst* 14(1):1–37
- Wypijewska A, Galazka-Friedman J, Bauminger ER, Wszolek ZK, Schweitzer KJ, Dickson DW et al (2010) Iron and reactive oxygen species activity in parkinsonian substantia nigra. *Parkinsonism Relat Disord*
- Xiong SG, She HY, Takeuchi H, Han B, Engelhardt JF, Barton CH et al (2003) Signaling role of intracellular iron in NF-kappa B activation. *J Biol Chem* 278(20):17646–17654
- Xu A, Wu LJ, Santella RM, Hei TK (1999) Role of oxyradicals in mutagenicity and DNA damage induced by crocidolite asbestos in mammalian cells. *Cancer Res* 59(23):5922–5926
- Xu Q, Kanthasamy AG, Reddy MB (2008a) Neuroprotective effect of the natural iron chelator, phytic acid in a cell culture model of Parkinson's disease. *Toxicology* 245(1–2):101–108
- Xu X, Satak R, Richardson DR (2008b) Iron chelation by clinically relevant anthracyclines: alteration in expression of iron-regulated genes and atypical changes in intracellular iron distribution and trafficking. *Mol Pharmacol* 73(3):833–844
- Xu WL, von Strauss E, Qiu CX, Winblad B, Fratiglioni L (2009) Uncontrolled diabetes increases the risk of Alzheimer's disease: a population-based cohort study. *Diabetologia* 52(6):1031–1039
- Xu J, Marzetti E, Seo AY, Kim JS, Prolla TA, Leeuwenburgh C (2010) The emerging role of iron dyshomeostasis in the mitochondrial decay of aging. *Mech Ageing Dev*
- Yadav AS, Bhatnagar D (2007a) Free radical scavenging activity, metal chelation and antioxidant power of some of the Indian spices. *Biofactors* 31(3–4):219–227
- Yadav AS, Bhatnagar D (2007b) Modulatory effect of spice extracts on iron-induced lipid peroxidation in rat liver. *Biofactors* 29(2–3):147–157
- Yadav AS, Bhatnagar D (2010) Inhibition of iron induced lipid peroxidation and antioxidant activity of Indian spices and *Acacia* *in vitro*. *Plant Foods Hum Nutr* 65:18–24
- Yamamoto K, Kawanishi S (1992) Site-specific DNA damage by phenylhydrazine and phenelzine in the presence of Cu(II) ion or Fe(III) complexes: roles of active oxygen species and carbon radicals. *Chem Res Toxicol* 5(3):440–446
- Yamashita T, Ando Y, Nakamura M, Obayashi K, Terazaki H, Haraoka K et al (2004) Inhibitory effect of α -tocopherol on methylmercury-induced oxidative stress. *Environ Health Prev Med* 9(3):111–117

- Yang F, Stonehuerner JG, Richards JH, Nguyen NB, Callaghan KD, Haile DJ et al (2010) Deficiency in the divalent metal transporter 1 increases bleomycin-induced lung injury. *Biometals*
- Yasuda T, Mochizuki H (2010) The regulatory role of alpha-synuclein and parkin in neuronal cell apoptosis; possible implications for the pathogenesis of Parkinson's disease. *Apoptosis*
- Yeager MP, Coleman RA (2010) In silico evidence for glutathione- and iron-related pathogenesis in Parkinson's disease. *J Neurosci Methods*
- Yee S, Choi BH (1994) Methylmercury poisoning induces oxidative stress in the mouse brain. *Exp Mol Pathol* 60(3):188–196
- Yee S, Choi BH (1996) Oxidative stress in neurotoxic effects of methylmercury poisoning. *Neurotoxicology* 17(1):17–26
- Yeh P, Tschumi AI, Kishony R (2006) Functional classification of drugs by properties of their pairwise interactions. *Nat Genet* 38(4):489–494
- Yeh PJ, Hegreness MJ, Aiden AP, Kishony R (2009) Drug interactions and the evolution of antibiotic resistance. *Nat Rev Microbiol* 7(6):460–466
- Yin H, Cheng L, Holt M, Hail N Jr, Maclaren R, Ju C (2010) Lactoferrin protects against acetaminophen-induced liver injury in mice. *Hepatology* 51(3):1007–1016
- Yin Z, Milatovic D, Aschner JL, Syversen T, Rocha JB, Souza DO et al (2007) Methylmercury induces oxidative injury, alterations in permeability and glutamine transport in cultured astrocytes. *Brain Res* 1131(1):1–10
- Yin ZB, Jiang HY, Syversen T, Rocha JBT, Farina M, Aschner M (2008) The methylmercury-L-cysteine conjugate is a substrate for the L-type large neutral amino acid transporter. *J Neurochem* 107(4):1083–1090
- Ying QL, Wray J, Nichols J, Batlle-Morera L, Doble B, Woodgett J et al (2008) The ground state of embryonic stem cell self-renewal. *Nature* 453(7194):519–523
- Yokoyama H, Kuroiwa H, Yano R, Araki T (2008) Targeting reactive oxygen species, reactive nitrogen species and inflammation in MPTP neurotoxicity and Parkinson's disease. *Neurol Sci* 29(5):293–301
- Yoneda M, Nozaki Y, Endo H, Mawatari H, Iida H, Fujita K et al (2010) Serum ferritin is a clinical biomarker in Japanese patients with nonalcoholic steatohepatitis (NASH) independent of HFE gene mutation. *Dig Dis Sci* 55:808–814
- Yoshino M, Murakami K (1998) Interaction of iron with polyphenolic compounds: application to antioxidant characterization. *Anal Biochem* 257(1):40–44
- You SA, Archacki SR, Angheloiu G, Moravec CS, Rao S, Kinter M et al (2003) Proteomic approach to coronary atherosclerosis shows ferritin light chain as a significant marker: evidence consistent with iron hypothesis in atherosclerosis. *Physiol Genomics* 13(1):25–30
- You SA, Wang Q (2005) Ferritin in atherosclerosis. *Clin Chim Acta* 357(1):1–16
- Youdim MB (2003) What have we learnt from CDNA microarray gene expression studies about the role of iron in MPTP induced neurodegeneration and Parkinson's disease? *J Neural Transm Suppl* (65):73–88
- Youdim MBH, Fridkin M, Zheng H (2004a) Novel bifunctional drugs targeting monoamine oxidase inhibition and iron chelation as an approach to neuroprotection in Parkinson's disease and other neurodegenerative diseases. *J Neural Transm* 111(10–11):1455–1471
- Youdim MBH, Stephenson G, Ben Shachar D (2004b) Ironing iron out in Parkinson's disease and other neurodegenerative diseases with iron chelators—a lesson from 6-hydroxydopamine and iron chelators, desferal and VK-28. *Redox-Active Metals Neurol Disord* 1012:306–325
- Young JL, Libby P, Schonbeck U (2002) Cytokines in the pathogenesis of atherosclerosis. *Thromb Haemost* 88(4):554–567
- Yu Z, Persson HL, Eaton JW, Brunk UT (2003) Intralysosomal iron: a major determinant of oxidant-induced cell death. *Free Radic Biol Med* 34(10):1243–1252
- Yuan XM, Li W (2008) Iron involvement in multiple signaling pathways of atherosclerosis: a revisited hypothesis. *Curr Med Chem* 15(21):2157–2172
- Yue H, Brown M, Knowles J, Wang H, Broomhead DS, Kell DB (2006) Insights into the behaviour of systems biology models from dynamic sensitivity and identifiability analysis: a case study of an NF-kappaB signalling pathway. *Mol Biosyst* 2:640–649
- Yue H, Wang Y, Broomhead DS, Brown M, Kell DB (2007) Sensitivity analysis of an oscillatory signal transduction pathway. *FOSBE* 2007:99–104
- Yue H, Brown M, He F, Jia J, Kell DB (2008) Sensitivity analysis and robust experimental design of a signal transduction pathway system. *Int J Chem Kinet* 40(11):730–741
- Yun SW, Gerlach M, Riederer P, Klein MA (2006) Oxidative stress in the brain at early preclinical stages of mouse scrapie. *Exp Neurol* 201(1):90–98
- Zacharski LR, Chow B, Lavori PW, Howes PS, Bell MR, DiTommaso MA et al (2000) The iron (Fe) and atherosclerosis study (FeAST): a pilot study of reduction of body iron stores in atherosclerotic peripheral vascular disease. *Am Heart J* 139(2):337–345
- Zacharski LR, Chow BK, Howes PS, Shamayeva G, Baron JA, Dalman RL et al (2008) Decreased cancer risk after iron reduction in patients with peripheral arterial disease: results from a randomized trial. *J Natl Cancer Inst* 100(14):996–1002
- Zacharski LR, Gerhard GS (2003) Atherosclerosis: a manifestation of chronic iron toxicity? *Vasc Med* 8(3):153–155
- Zamboni P (2006) The big idea: iron-dependent inflammation in venous disease and proposed parallels in multiple sclerosis. *J R Soc Med* 99(11):589–593
- Zamin LL, Filippi-Chiela EC, Dillenburg-Pilla P, Horn F, Salbego C, Lenz G (2009) Resveratrol and quercetin cooperate to induce senescence-like growth arrest in C6 rat glioma cells. *Cancer Sci* 100(9):1655–1662
- Zandman-Goddard G, Shoenfeld Y (2008) Hyperferritinemia in autoimmunity. *Isr Med Assoc J* 10(1):83–84
- Zanella I, Derosas M, Corrado M, Cocco E, Cavadini P, Biasiotto G et al (2008) The effects of frataxin silencing in HeLa cells are rescued by the expression of human mitochondrial ferritin. *Biochim Biophys Acta* 1782(2):90–98
- Zapelini PH, Rezin GT, Cardoso MR, Ritter C, Klamt F, Moreira JC et al (2008) Antioxidant treatment reverses mitochondrial dysfunction in a sepsis animal model. *Mitochondrion* 8(3): 211–218
- Zarembek KA, Cruz AR, Huang CY, Gallin JI (2009) Antifungal activities of natural and synthetic iron chelators alone and in combination with azole and polyene antibiotics against *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 53(6):2654–2656
- Zatta P, Drago D, Bolognin S, Sensi SL (2009) Alzheimer's disease, metal ions and metal homeostatic therapy. *Trends Pharmacol Sci* 30(7):346–355
- Zecca L, Youdim MBH, Riederer P, Connor JR, Crichton RR (2004) Iron, brain ageing and neurodegenerative disorders. *Nat Rev Neurosci* 5(11):863–873
- Zhang J, Perry G, Smith MA, Robertson D, Olson SJ, Graham DG et al (1999) Parkinson's disease is associated with oxidative damage to cytoplasmic DNA and RNA in substantia nigra neurons. *Am J Pathol* 154(5):1423–1429
- Zhang HY (2006) Same causes, same cures. *Biochem Biophys Res Commun* 351(3):578–581

- Zhang SB, Wang Y, Liu CL (2009) Metal-based therapeutic strategies of neurodegenerative diseases. *Progr Chem* 21(5):903–910
- Zhang J, Adrian FJ, Jahnke W, Cowan-Jacob SW, Li AG, Iacob RE et al (2010a) Targeting Bcr-Abl by combining allosteric with ATP-binding-site inhibitors. *Nature* 463(7280):501–506
- Zhang J, Zhang Y, Wang J, Cai P, Luo C, Qian Z et al (2010b) Characterizing iron deposition in Parkinson's disease using susceptibility-weighted imaging: an in vivo MR study. *Brain Res*
- Zhang X, Smith DL, Meriin AB, Engemann S, Russel DE, Roark M et al (2005a) A potent small molecule inhibits polyglutamine aggregation in Huntington's disease neurons and suppresses neurodegeneration in vivo. *Proc Natl Acad Sci USA* 102(3):892–897
- Zhang X, Xie W, Qu S, Pan T, Wang X, Le W (2005b) Neuroprotection by iron chelator against proteasome inhibitor-induced nigral degeneration. *Biochem Biophys Res Commun* 333(2):544–549
- Zhang Z, Rigas B (2006) NF-kappaB, inflammation and pancreatic carcinogenesis: NF-kappaB as a chemoprevention target (review). *Int J Oncol* 29(1):185–192
- Zhang Z, Wei T, Hou J, Li G, Yu S, Xin W (2003) Iron-induced oxidative damage and apoptosis in cerebellar granule cells: attenuation by tetramethylpyrazine and ferulic acid. *Eur J Pharmacol* 467(1–3):41–47
- Zhao C, Deng W, Gage FH (2008) Mechanisms and functional implications of adult neurogenesis. *Cell* 132(4):645–660
- Zhao J, Geng C, Tao L, Zhang D, Jiang Y, Tang K et al (2010) Reconstruction and analysis of human liver-specific metabolic network based on CNHLPP data. *J Proteome Res*
- Zhao J, Yu H, Luo JH, Cao ZW, Li YX (2006) Hierarchical modularity of nested bow-ties in metabolic networks. *BMC Bioinform* 7:386
- Zhao M, Laissie JA, Zimmermann A (1997) Hepatocyte apoptosis in hepatic iron overload diseases. *Histol Histopathol* 12(2):367–374
- Zhao R, Planalp RP, Ma R, Greene BT, Jones BT, Brechbiel MW et al (2004) Role of zinc and iron chelation in apoptosis mediated by tachpyridine, an anti-cancer iron chelator. *Biochem Pharmacol* 67(9):1677–1688
- Zheng H, Weiner LM, Bar-Am O, Epsztejn S, Cabantchik ZI, Warshawsky A et al (2005) Design, synthesis, and evaluation of novel bifunctional iron-chelators as potential agents for neuroprotection in Alzheimer's, Parkinson's, and other neurodegenerative diseases. *Bioorg Med Chem* 13(3):773–783
- Zhu X, Su B, Wang X, Smith MA, Perry G (2007) Causes of oxidative stress in Alzheimer disease. *Cell Mol Life Sci* 64(17):2202–2210
- Ziegler JL, Simonart T, Snoeck R (2001) Kaposi's sarcoma, oncogenic viruses, and iron. *J Clin Virol* 20(3):127–130
- Zimmermann GR, Lehar J, Keith CT (2007) Multi-target therapeutics: when the whole is greater than the sum of the parts. *Drug Discov Today* 12(1–2):34–42
- Zimmet JM, Hare JM (2006) Nitroso-redox interactions in the cardiovascular system. *Circulation* 114(14):1531–1544
- Zivadinov R, Bakshi R (2004) Role of MRI in multiple sclerosis II: brain and spinal cord atrophy. *Front Biosci* 9:647–664
- Zivadinov R, Minagar A (2009) Evidence for gray matter pathology in multiple sclerosis: a neuroimaging approach. *J Neurosci* 28(1–2):1–4
- Zuliani G, Cherubini A, Ranzini M, Ruggiero C, Atti AR, Fellin R (2006) Risk factors for short-term mortality in older subjects with acute ischemic stroke. *Gerontology* 52(4):231–236
- Zurich MG, Monnet-Tschudi F (2009) Contribution of in vitro neurotoxicology studies to the elucidation of neurodegenerative processes. *Brain Res Bull* 80(4–5):211–216