

Gata2 haploinsufficiency promotes proliferation and functional decline of hematopoietic stem cells with myeloid bias during aging

Ali Abdelfattah,^{1,2} Antonia Hughes-Davies,¹ Liam Clayfield,¹ Juan Bautista Menendez-Gonzalez,^{1,3,4} Alhomidi Almotiri,^{1,5} Badi Alotaibi,¹ Alex Tonks,⁶ and Neil P. Rodrigues¹

¹European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Cardiff, United Kingdom; ²Department of Medical Laboratory Sciences, Faculty of Applied Medical Sciences, The Hashemite University, Zarqa, Jordan; ³Department of Stem Cell and Regenerative Biology, Harvard Stem Cell Institute, Harvard University, Cambridge, MA; ⁴Center for Regenerative Medicine, Massachusetts General Hospital, Boston, MA; ⁵College of Applied Medical Sciences-Dawadmi, Shaqra University, Dawadmi, Saudi Arabia; and ⁶Department of Hematology, Division of Cancer and Genetics, School of Medicine, Cardiff University, Cardiff, United Kingdom

Key Points

- *Gata2* haploinsufficiency promotes HSC proliferation, monocytosis, and decreases CLP generation during aging.
- Elderly *Gata2* haploinsufficient HSCs are functionally impaired with evidence of myeloid bias.

During aging, hematopoietic stem cell (HSC) function wanes with important biological and clinical implications for benign and malignant hematology, and other comorbidities, such as cardiovascular disease. However, the molecular mechanisms regulating HSC aging remain incompletely defined. *GATA2* haploinsufficiency driven clinical syndromes initially result in primary immunodeficiencies and routinely evolve into hematologic malignancies on acquisition of further epigenetic mutations in both young and older patients. Using a conditional mouse model of *Gata2* haploinsufficiency, we discover that during aging *Gata2* promotes HSC proliferation, monocytosis, and loss of the common lymphoid progenitor. Aging of *Gata2* haploinsufficient mice also offsets enhanced HSC apoptosis and decreased granulocyte-macrophage progenitor number normally observed in young *Gata2* haploinsufficient mice. Transplantation of elderly *Gata2* haploinsufficient HSCs impairs HSC function with evidence of myeloid bias. Our data demonstrate that *Gata2* regulates HSC aging and suggest the mechanisms by which *Gata2* mediated HSC aging has an impact on the evolution of malignancies in *GATA2* haploinsufficiency syndromes.

Introduction

Aging of the hematopoietic system is marked by an irrevocable decrease in blood cell function accompanied by fundamental alterations including an increased abundance of hematopoietic stem cells (HSCs), which have diminished function, and imbalanced differentiation potential toward myeloid cell production at the expense of lymphoid cells.¹⁻³ Increasing HSC numbers manifest during aging with a rise in proliferation and a decrease in apoptosis,^{1,2} yet reduction in their function appears to be associated with upregulation of stress and inflammatory responses and downregulation of genome integrity, including DNA repair, and chromatin modifications.⁴ Notably, aged HSCs highly express genes associated with myeloid malignancy.³ In the elderly, a reduced number of HSC clones contribute to hematopoiesis, a phenomenon known as clonal hematopoiesis of indeterminate potential (CHIP), and this occurs together with frequent mutations in epigenetic regulators that pose an increased risk for both myeloid malignancy and atherosclerosis.⁵ Thus, overall, aging of HSCs has important biological and clinical implications for hematologic malignancy, anemia, clotting disorders, immunodeficiency, and other comorbidities, such as cardiovascular disease. However, the molecular mechanisms regulating aging of HSCs remain incompletely defined.

Submitted 12 March 2021; accepted 10 June 2021; prepublished online on *Blood Advances* First Edition 8 September 2021; final version published online 26 October 2021. DOI 10.1182/bloodadvances.2021004726.

Requests for data sharing should be made to the corresponding author: rodriguesn@cardiff.ac.uk.

The full-text version of this article contains a data supplement.

© 2021 by The American Society of Hematology. Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

Gata2, a zinc-finger transcription factor, has been identified as critical for HSC function in both the embryo and adult.⁶⁻⁹ Deregulation of GATA2 expression leads to hematopoietic aplasia, dysplasia, and neoplasia, underscoring the functional importance of GATA2 in maintaining normal hematopoiesis.^{10,11} For example, overexpression of *GATA2* is observed in myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML), and gain-of-function mutations in chronic myeloid leukemia correlate with poor prognosis.^{10,12,13} In contrast, sporadic or germline *GATA2* haploinsufficiency mutations in coding or enhancer regions give rise to MDS or AML that is typically preceded by a phase of immunodeficiency.^{10,11,13,14} Given that up to 20% of *GATA2* haploinsufficient patients develop MDS or AML after middle age^{10,13} and little is known about the impact of *Gata2* in this setting, we sought to evaluate *Gata2* haploinsufficiency during hematopoietic aging.

Methods

Animals

All mice were housed at the Heath Park Unit, Cardiff University. All animal experiments were executed under PPL 30/3380 in accordance with the 1986 Animals (Scientific Procedures) Act.

Flow cytometry

Stained cells were analyzed on an LSR Fortessa (Becton Dickinson Biosciences) flow cytometer, and fluorescence-activated cell sorter plots were generated using FlowJo 10.6.1 software (Tree Star, Inc).

Transplantation

Transplantation was performed as described previously.¹⁵

Statistical analysis

Statistical analysis was performed with GraphPad Prism 8.3.0 software (GraphPad Software, Inc.).

Further details for these methods are provided in supplemental Methods.

Results and discussion

We first assessed *Gata2* expression in prospectively isolated hematopoietic cell populations from aged wild-type bone marrow (BM). Consistent with their young BM counterparts,^{8,9} *Gata2* expression was highest in HSCs, attenuated during lympho-myeloid differentiation (hematopoietic progenitor cell 1 [HPC1], hematopoietic progenitor cell 2 [HPC2], lymphoid-primed multipotent progenitor [LMPP], common myeloid progenitor [CMP], granulocyte-macrophage progenitor [GMP], and megakaryocyte-erythroid progenitor [MEP]; and mature myeloid cells), and was notably absent in the common lymphoid progenitor (CLP) and mature lymphoid (T-cell and B-cell) compartments (Figure 1A).

To explore the impact of *Gata2* haploinsufficiency in hematopoiesis during aging, one allele of the floxed *Gata2* gene was deleted in hematopoietic cells using the pan-hematopoietic *Vav-iCre* promoter.¹⁶ *Gata2*^{+/+}; *Vav-iCre*⁺ with *Gata2*^{fl/fl}; *Vav-iCre*⁻ were bred to produce *Gata2*^{+/fl}; *Vav-iCre*⁺ (heterozygote) and *Gata2*^{+/fl}; *Vav-iCre*⁻ (control) littermates. Genomic DNA polymerase chain reaction genotyping of ear notch biopsies and BM cells confirmed *Gata2* haploinsufficiency and control status (supplemental Figure 1A). *Gata2*^{+/fl}; *Vav-iCre*⁺ and control (*Gata2*^{+/fl}; *Vav-iCre*⁻) mice were subsequently

aged for 18 to 20 months old, which correlates with humans aged 56 to 69 years (Figure 1B).¹⁷ In agreement with previous data,¹⁻³ skewed differentiation capacity toward myeloid lineages and away from lymphoid maturation was observed over time in the peripheral blood (PB) of both aged *Gata2*^{+/fl}; *Vav-iCre*⁺ and control mice (data not shown). At 18 to 20 months, the proportion of myeloid cells (Mac1⁺Gr1⁺ and Mac1⁺Gr1⁻), B cells (B220⁺), and T cells (CD4⁺ and CD8⁺) in PB was also similar between genotypes (supplemental Figure 1B). Although most *GATA2* haploinsufficiency patients present with immunodeficiency, a substantial proportion of patients display normal hematologic profiles before acquiring infections that characterize disease progression.^{10,13} Thus, unchanged hematologic parameters observed in our mouse model of *Gata2* haploinsufficiency may reflect what is seen in this subset of patients. Yet, when comparing the proportion of myeloid and lymphoid cells in PB from aged vs young *Gata2*^{+/fl}; *Vav-iCre*⁺ mice, a significant increase in Mac1⁺Gr1⁻ myeloid cells was noted in aged *Gata2*^{+/fl}; *Vav-iCre*⁺ mice (supplemental Figure 1C), demonstrating that *Gata2* haploinsufficiency causes steady-state accumulation of monocytes during aging.

Age-related changes in cellular composition of BM and extramedullary hematopoietic organs occur during aging.¹⁸ However, cellularity and mature myeloid, erythroid, and lymphoid cell proportions were found to be similar in BM, spleen, and thymus of aged *Gata2*^{+/fl}; *Vav-iCre*⁺ and control mice (supplemental Figure 1D-H).

Evaluating the impact of *Gata2* haploinsufficiency on the frequencies of hematopoietic stem/progenitor cells (HSPCs) (HSCs, MPPs, HPC1, HPC2) and committed progenitors (CMPs, GMPs, MEPs, CLPs) in BM,^{19,20} we noted a decrease only in HSC, MPP, LMPP, and CLP populations of aged *Gata2*^{+/fl}; *Vav-iCre*⁺ (Figure 1C-D). Although reduction of HSCs in young *Gata2* haploinsufficient mice is well established,⁶⁻⁹ the reduction in HSCs was more pronounced in older *Gata2*^{+/fl}; *Vav-iCre*⁺ vs younger *Gata2*^{+/fl}; *Vav-iCre*⁺ mice (supplemental Figure 1I). Of further note, a decrease in CLP abundance in aged *Gata2*^{+/fl}; *Vav-iCre*⁺ mice contrasted with the lack of impact of *Gata2* haploinsufficiency on CLPs from young mice (data not shown) and an overall twofold reduction in CLPs compared with their young counterparts (supplemental Figure 1I). GMP abundance was reduced in the context of young *Gata2* haploinsufficient mice²¹ (and data not shown). Unexpectedly, GMP abundance was comparable between aged *Gata2*^{+/fl}; *Vav-iCre*⁺ and control mice, demonstrating that aging of *Gata2* haploinsufficient mice normalizes numbers of GMPs (Figure 1C-D). Therefore, during aging, haploinsufficiency of *Gata2* selectively drives attrition of both HSC and CLP compartments and rescues the number of GMPs.

To discern the cellular mechanisms leading to reduced HSCs during aging of *Gata2*^{+/fl}; *Vav-iCre*⁺ mice, we evaluated their apoptosis and cell cycle status. Differing from young *Gata2* haploinsufficient mice, where HSCs display increased apoptosis (supplemental Figure 1K),⁷ no significant difference was observed in early- and late-stage apoptosis in HSCs from aged *Gata2*^{+/fl}; *Vav-iCre*⁺ mice compared with their control counterparts, as judged by the Annexin V assay (supplemental Figure 1J-K). Similarly, a significant increase in late-stage apoptosis in MPPs of young *Gata2* haploinsufficient mice was lost during aging (supplemental Figure 1J-K). In stark contrast to enhanced quiescence observed in young *Gata2* haploinsufficient mice (Figure 1F),⁷ Ki-67 analysis revealed a significant decrease in the proportion of G₀ HSCs together with an increase in the frequency of HSCs in S/G₂/M phases in aged *Gata2*^{+/fl}; *Vav-iCre*⁺ mice




Figure 1. Long-term *Gata2* haploinsufficiency enhances HSC proliferation, decreases HSC abundance, and causes a reduction in common lymphoid progenitors. (A) Relative expression of *Gata2* mRNA in purified hematopoietic compartments, including HSCs (LSK_{CD150}⁺CD48⁻), HPC1 (LSK_{CD150}⁻CD48⁺), HPC2 (LSK_{CD150}⁺CD48⁺), LMPP (LSK_{CD34}⁺CD135^{hi}), CMP (LK_{CD34}⁺CD16/32⁻), GMP (LK_{CD34}⁺CD16/32⁺), MEP (LK_{CD34}⁻CD16/32⁻), CLP (Lin⁻c-kit^{lo}Sca1^{lo}CD127⁺CD135⁺), myeloid cells (Mac1⁺Gr1⁺), T cells (CD3_β), and B cells (B220). n = 2 replicates per population. (B) Experimental design for analysis of long-term and short-term *Gata2* haploinsufficiency. *Gata2*^{+/fl}; *Vav-iCre*⁻ (control) and *Gata2*^{+/fl}; *Vav-iCre*⁺ mice were analyzed at 18 to 20 months and at 8 to 12 weeks old. (C-D) Representative immunophenotypic HSPC analysis and gating scheme for *Gata2*^{+/fl}; *Vav-iCre*⁻ (control) and *Gata2*^{+/fl}; *Vav-iCre*⁺ mice at 18 to 20 months (C) and absolute cell count of primitive and committed hematopoietic populations (D) from control (n = 9) and *Gata2*^{+/fl}; *Vav-iCre*⁺ (n = 7) mice. MPPs are LSK_{CD150}⁻CD48⁻. Percentages represent a frequency of live nucleated BM cells. Data from 3 independent experiments. (E) Representative flow cytometry plots for cell cycle analysis of BM HSCs from *Gata2*^{+/fl}; *Vav-iCre*⁻ (control) and *Gata2*^{+/fl}; *Vav-iCre*⁺ mice at 18 to 20 months using Ki-67/4',6-diamidino-2-phenylindole (DAPI). Bivariate plots showing the frequency of HSCs in G0 (Ki-67⁻DAPI⁺), G1 (Ki-67⁺DAPI⁻), and S/G2/M (Ki-67⁺DAPI⁺) phases from control (n = 9) and *Gata2*^{+/fl}; *Vav-iCre*⁺ (n = 7) mice from 3 independent experiments. (F) The percentage of BM HSCs in G0, G1, and S/G2/M cell cycle phases from aged mice (n = 9 control and 7 *Gata2*^{+/fl}; *Vav-iCre*⁺) and young mice (n = 6 for each genotype) from 3 independent experiments for each condition. Data presented as mean ± standard error of the mean (SEM). Statistical analysis is performed using Mann-Whitney U test. Significant data: *P < .05; **P < .01; ***P < .001.




Figure 2. Elderly *Gata2* haploinsufficient HSCs have a functional defect in reconstitution of multilineage hematopoietic compartments with a myeloid bias. (A) Schematic representation of competitive HSC transplantation experiment. Three hundred HSCs from aged control or *Gata2*^{+*fl*}; *Vav-iCre*⁺ mice (CD45.2⁺) together with 2 × 10⁵ unfractionated BM competitor cells (CD45.1⁺) were transplanted into lethally irradiated (9.5 Gy) recipient mice (CD45.1⁺). Four independent biological replicates were used for each genotype. Donor chimerism in PB was tested every 4 weeks until week 16 after transplant. (B) Proportion of CD45.2 donor-derived cells in PB after transplantation of donor cells from control (n = 8 recipients) or *Gata2*^{+*fl*}; *Vav-iCre*⁺ mice (n = 8 recipients). (C-E) Percentages of CD45.2 donor-derived cells contribution to mature cells in PB (C) BM HSPCs (D) and BM committed myeloid/lymphoid progenitors (E) at week 16 after transplantation of control or *Gata2*^{+*fl*}; *Vav-iCre*⁺ donor cells. n = 8 recipients for each genotype from 2 independent experiments. (F) Fold change ratios of CD45.2 donor-derived cell contribution to mature PB cells in aged mice (n = 8 control and 8 *Gata2*^{+*fl*}; *Vav-iCre*⁺) in relation to young mice (n = 8 control and 8 *Gata2*^{+*fl*}; *Vav-iCre*⁺) donor cells after normalizing to their control counterparts. 2 to 3 independent experiments were performed for each condition. Data is presented as mean ± SEM. Statistical analysis is performed using the Mann-Whitney U test. Significant data: **P* < .05; ***P* < .01; ****P* < .001.

(Figure 1E-F), indicating *Gata2* haploinsufficiency promotes HSC proliferation during aging.

To assess the impact of enhanced HSC proliferation on the function of aged *Gata2*^{+*fl*}; *Vav-iCre*⁺ HSCs, we performed competitive transplantation experiments. To do this, 300 CD45.2 purified aged HSCs from control or *Gata2*^{+*fl*}; *Vav-iCre*⁺ mice were cotransplanted with 2×10^5 CD45.1 competitor BM cells into lethally irradiated CD45.1 recipients (Figure 2A). Engraftment analysis was conducted every 4 weeks and revealed that the frequency of PB donor contribution was significantly lower in recipients of aged *Gata2*^{+*fl*}; *Vav-iCre*⁺ derived cells compared with control derived cells during the entire 16-week posttransplant period (Figure 2B), which was reflected in lower donor contribution to all mature myeloid and lymphoid lineages in BM, PB, and spleen (Figure 2C; supplemental Figure 2) and lower HSPC and committed progenitor engraftment in BM at week 16 (Figure 2D-E). Notably, engraftment of the T-cell compartment was more affected than myeloid or B-cell compartments (Figure 2C), suggesting myeloid bias after transplantation of elderly *Gata2* haploinsufficient HSCs. This contention is supported by data showing a twofold expansion in myeloid cells and an approximately twofold decrease in CD4⁺ T cells in the PB of recipients of aged *Gata2*^{+*fl*}; *Vav-iCre*⁺ HSCs compared with recipients of HSCs from young *Gata2*^{+*fl*}; *Vav-iCre*⁺ mice (Figure 2F). Thus, aged *Gata2* haploinsufficient HSCs fail to reconstitute multilineage hematopoiesis and functionally impart myeloid bias on aged HSCs in vivo.

In this report, we identify *Gata2* as a critical determinant of HSC function during aging. Although HSCs continue to decline in prevalence and function in *Gata2* haploinsufficient mice during aging, we demonstrate significant alterations in HSC and progenitor cell behavior that selectively occur in aged *Gata2* haploinsufficient mice rather than in their young counterparts, including (1) augmented HSC proliferation (contrasting with enhanced quiescence in young *Gata2* haploinsufficient HSCs), (2) normalization of HSC apoptosis (whereas HSC apoptosis was enhanced in young *Gata2* haploinsufficient HSCs), (3) normalization of GMP abundance (compared with the decrease in GMPs observed in young *Gata2* haploinsufficient mice), (4) monocytosis, and (5) decreased lymphoid progenitor capacity exemplified by loss of CLPs. These data offer insights into the specific HSC fates that are mediated by *Gata2* during aging, setting the stage for studies that further decipher the transcriptional underpinnings of *Gata2* regulation in the aging hematopoietic system.

Our data also have implications for *GATA2* haploinsufficiency-driven clinical syndromes. Aged *Gata2*^{+*fl*}; *Vav-iCre*⁺ mice fail to fully recapitulate the immunodeficiency and progression to MDS/AML observed clinically, which may be ascribed to a number of factors. First, our hematopoietic-specific conditional *Gata2* mouse model does not assess niche-driven effects of *Gata2* haploinsufficiency on the hematopoietic system. Second, in striking contrast to mice, *GATA2* haploinsufficient patients are routinely exposed to infectious agents that push the hematopoietic system to exhaustion where they either succumb or

clonally adapt to this inflammatory environment, acquire secondary mutations, and develop MDS/AML. Further work will be needed to assess whether infectious or inflammatory stimuli will induce disease progression in *Gata2*^{+*fl*}; *Vav-iCre*⁺ mice. These caveats notwithstanding, atypical sequela includes monocytosis and B-cell loss in *GATA2*-related chronic myelomonocytic leukemia and MDS,^{11,22} which mirrors monocytosis observed here in aged *Gata2* haploinsufficient mice. Where monocytosis occurs in young patients, this may be caused by premature aging of HSCs, diminishing their function and promoting myeloid bias. Such myeloid bias may explain the presence of lymphoid progenitor deficiency in *GATA2* haploinsufficiency despite the absence of *Gata2* expression in lymphoid progenitors and mature lymphoid cells.²²⁻²⁴ During aging, *Gata2* haploinsufficient HSCs were driven out of quiescence and into proliferation, and they relinquish their apoptotic advantage; if such a mechanism operates in any *GATA2* haploinsufficient patient, it may cause a gradual accrual of DNA damage and a proliferative environment conducive for acquiring secondary epigenetic mutations with clonal dominance in developing MDS/AML. Finally, paralleling recent findings that inflammatory processes driven by atherosclerosis enhance both HSC proliferation and CHIP,²⁵ it will be of interest to investigate whether monocytosis and increased HSC proliferation observed in *Gata2* haploinsufficient mice during aging are directly linked and clinically relevant to malignant progression of *GATA2* haploinsufficiency syndromes.

Acknowledgments

A. Abdelfattah was supported by The Hashemite University, Zarqa, Jordan. N.P.R. laboratory is supported by the Saudi Arabian Cultural Bureau, Leukemia Cancer Society and Leukemia Research Appeal for Wales.

Authorship

Contribution: A. Abdelfattah designed and performed experiments, analyzed and interpreted data, prepared figures, and contributed to writing the manuscript; A.H.-D., L.C., A. Almotiri, and B.A. performed experiments and contributed to data analysis; J.B.M.-G. performed experiments and contributed to data analysis and interpretation; A.T. contributed to experimental design, data analysis, and interpretation; N.P.R. conceived and supervised the project, designed experiments, analyzed and interpreted the data, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: A.A., 0000-0003-0294-0568; A.A., 0000-0002-5906-6544; N.P.R., 0000-0002-1925-7733.

Correspondence: Neil P. Rodrigues, European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, United Kingdom; e-mail: rodriguesn@cardiff.ac.uk.

References

1. Pearce DJ, Anjos-Afonso F, Ridler CM, Eddaoudi A, Bonnet D. Age-dependent increase in side population distribution within hematopoiesis: implications for our understanding of the mechanism of aging. *Stem Cells*. 2007;25(4):828-835.
2. Morrison SJ, Wandycz AM, Akashi K, Globerson A, Weissman IL. The aging of hematopoietic stem cells. *Nat Med*. 1996;2(9):1011-1016.

3. Rossi DJ, Bryder D, Zahn JM, et al. Cell intrinsic alterations underlie hematopoietic stem cell aging. *Proc Natl Acad Sci USA*. 2005;102(26):9194-9199.
4. Rossi DJ, Bryder D, Seita J, Nussenzweig A, Hoeijmakers J, Weissman IL. Deficiencies in DNA damage repair limit the function of haematopoietic stem cells with age. *Nature*. 2007;447(7145):725-729.
5. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371(26):2488-2498.
6. Ling KW, Ottersbach K, van Hamburg JP, et al. GATA-2 plays two functionally distinct roles during the ontogeny of hematopoietic stem cells. *J Exp Med*. 2004;200(7):871-882.
7. Rodrigues NP, Janzen V, Forkert R, et al. Haploinsufficiency of GATA-2 perturbs adult hematopoietic stem-cell homeostasis. *Blood*. 2005;106(2):477-484.
8. Menendez-Gonzalez JB, Vukovic M, Abdelfattah A, et al. Gata2 as a crucial regulator of stem cells in adult hematopoiesis and acute myeloid leukemia. *Stem Cell Reports*. 2019;13(2):291-306.
9. Guo G, Luc S, Marco E, et al. Mapping cellular hierarchy by single-cell analysis of the cell surface repertoire. *Cell Stem Cell*. 2013;13(4):492-505.
10. Wlodarski MW, Collin M, Horwitz MS. GATA2 deficiency and related myeloid neoplasms. *Semin Hematol*. 2017;54(2):81-86.
11. Spinner MA, Sanchez LA, Hsu AP, et al. GATA2 deficiency: a protean disorder of hematopoiesis, lymphatics, and immunity. *Blood*. 2014;123(6):809-821.
12. Zhang SJ, Ma LY, Huang QH, et al. Gain-of-function mutation of GATA-2 in acute myeloid transformation of chronic myeloid leukemia. *Proc Natl Acad Sci USA*. 2008;105(6):2076-2081.
13. Collin M, Dickinson R, Bigley V. Haematopoietic and immune defects associated with GATA2 mutation. *Br J Haematol*. 2015;169(2):173-187.
14. Johnson KD, Hsu AP, Ryu M-J, et al. Cis-element mutated in GATA2-dependent immunodeficiency governs hematopoiesis and vascular integrity. *J Clin Invest*. 2012;122(10):3692-3704.
15. Duran-Struuck R, Dysko RC. Principles of bone marrow transplantation (BMT): providing optimal veterinary and husbandry care to irradiated mice in BMT studies. *J Am Assoc Lab Anim Sci*. 2009;48(1):11-22.
16. Stadtfeld M, Graf T. Assessing the role of hematopoietic plasticity for endothelial and hepatocyte development by non-invasive lineage tracing. *Development*. 2005;132(1):203-213.
17. Jackson SJ, Andrews N, Ball D, et al. Does age matter? The impact of rodent age on study outcomes. *Lab Anim*. 2017;51(2):160-169.
18. Young K, Borikar S, Bell R, Kuffler L, Philip V, Trowbridge JJ. Progressive alterations in multipotent hematopoietic progenitors underlie lymphoid cell loss in aging. *J Exp Med*. 2016;213(11):2259-2267.
19. Akashi K, Traver D, Miyamoto T, Weissman IL. A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature*. 2000;404(6774):193-197.
20. Oguro H, Ding L, Morrison SJ. SLAM family markers resolve functionally distinct subpopulations of hematopoietic stem cells and multipotent progenitors. *Cell Stem Cell*. 2013;13(1):102-116.
21. Rodrigues NP, Boyd AS, Fugazza C, et al. GATA-2 regulates granulocyte-macrophage progenitor cell function. *Blood*. 2008;112(13):4862-4873.
22. Nováková M, Žaliová M, Suková M, et al. Loss of B cells and their precursors is the most constant feature of GATA-2 deficiency in childhood myelodysplastic syndrome. *Haematologica*. 2016;101(6):707-716.
23. Harada N, Hasegawa A, Hirano I, Yamamoto M, Shimizu R. GATA2 hypomorphism induces chronic myelomonocytic leukemia in mice. *Cancer Sci*. 2019;110(4):1183-1193.
24. Laurenti E, Doulatov S, Zandi S, et al. The transcriptional architecture of early human hematopoiesis identifies multilevel control of lymphoid commitment. *Nat Immunol*. 2013;14(7):756-763.
25. Heyde A, Rohde D, McAlpine CS, et al. Increased stem cell proliferation in atherosclerosis accelerates clonal hematopoiesis. *Cell*. 2021;184(5):1348-1361.