



Original article

Nosema ceranae infection in honeybee samples from Tuscanian Archipelago (Central Italy) investigated by two qPCR methods

Giovanni Cilia^{a,b,1}, Simona Sagona^{b,1}, Matteo Giusti^b, Pedro Emanuel Jarmela dos Santos^a, Antonio Nanetti^a, Antonio Felicioli^{b,*}

^aCREA Research Centre for Agriculture and Environment, Via di Saliceto 80, 40128 Bologna, Italy

^bDepartment of Veterinary Sciences, University of Pisa, Viale delle Piagge 2, 20159 Pisa, Italy



ARTICLE INFO

Article history:

Received 15 June 2018

Revised 7 November 2018

Accepted 25 November 2018

Available online 26 November 2018

Keywords:

Nosema ceranae

Nosemosis

qPCR

Hsp70 and *16S rRNA* gene

Apis mellifera

Tuscanian Archipelago

ABSTRACT

Nosema apis and *Nosema ceranae* are microsporidian parasite worldwide spread causing an emerging infectious disease of European honeybee *Apis mellifera*. The *Nosema* presence was deeply investigated in several countries but low information are presents about islands. In this investigation was evaluated the presence *N. ceranae* and *N. apis* in apiaries located in Tuscanian Archipelago islands (Central Italy). For *N. ceranae* detection, two different Real-Time PCR (qPCR) methods, the *16S rRNA* and *Hsp70* gene amplification qPCR, were performed on honey bee samples; while, for *N. apis* only the *16S rRNA* qPCR amplification was performed. On all islands, only *N. ceranae* was present, while *N. apis* was not found in the samples. The two qPCR showed significant difference ($p < 0.040$) in *N. ceranae* spores quantification. The single-copy *Hsp70* gene method qPCR assay systematically detected a lower amount of *N. ceranae* copies compared to the multi-copy *16S rRNA* gene method.

© 2018 Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Microsporidia are unicellular eukaryotes infecting vertebrate and invertebrate animal hosts. So far, three microsporidia species have been described associated to the honey bees, all belonging to the *Nosema* genus. The infections are known as nosemosis irrespective they are caused by *Nosema apis* Zander (Fries, 1993), *Nosema ceranae* (Fries et al., 1996) or *Nosema neumannii* (Chemurot et al., 2017). The first two species are spread worldwide and responsible for known symptoms (Fries et al., 2006; Higes et al., 2006; Klee et al., 2007; Martin-Hernandez et al., 2007). However, *N. neumannii* was recently detected only in Ugandan bees and its consequences for the host bees have not been described yet.

N. apis and *N. ceranae* are generally considered original parasites of European and Asian honey bees, respectively (Botías et al., 2012). However, the second showed capable to effectively infect

Apis mellifera individuals also, which allowed its spread far beyond the Asian continent (Fries et al., 2006).

It was suggested that *N. ceranae* may have replaced *N. apis* in Italy (Ferroglio et al., 2013) and in other countries (Ansari et al., 2017; Calderón et al., 2008; Fries et al., 2006; Giersch et al., 2009; Higes et al., 2009b; Invernizzi et al., 2009; Klee et al., 2007; Pacini et al., 2016; Tapasztó et al., 2009), although the possibility of a direct competition between the two species is still debated.

In Italy, *N. ceranae* was found in honey bee colonies reared in all regions (Ferroglio et al., 2013; Klee et al., 2007; Maiolino et al., 2014; Mutinelli et al., 2010; Porrini et al., 2016), but small islands have never been investigated in this respect. Thus, no data are available for the islands of Tuscanian Archipelago: Giannutri (area: 26 ha), Montecristo (103 ha), Pianosa (104 ha), Capraia (193 ha), Giglio (204 ha), Gorgona (223 ha) and Elba (22.350 ha).

Both *Nosema* species do infect the epithelial cells of the honey bee ventriculum (Fries et al., 1996; Higes et al., 2007) with effects recognizable both at individual and colony level, like reduced lifespan, lethargic behaviour and poor honey and pollen harvest (Eiri et al., 2015; Higes et al., 2009c, 2008).

Diarrhoeic faeces, high mortality at the hive entrance and swollen bee abdomens are commonly seen in *N. apis* infected colonies (Bourgeois et al., 2010), but *N. ceranae* infections tend to produce subtle symptoms that are difficult to spot in the field. Nevertheless,



Production and hosting by Elsevier

<https://doi.org/10.1016/j.sjbs.2018.11.017>

1319-562X/© 2018 Production and hosting by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

damage may be severe and result into adult population decline until colony collapse (Fries et al., 1996; Giersch et al., 2009; Higes et al., 2006).

Nosema spp. spores can be detected by light microscopy in the honey bee ventriculum, but reliable species determination requires advanced techniques. Although individual spore morphology may be examined by transmission electron microscope with this purpose (Chen et al., 2009; Forsgren and Fries, 2010; Fries et al., 1996), biomolecular bioassays are generally preferred for practical reasons and because they allow accurate quantification by the Real-Time PCR (qPCR) approach.

The *16S rRNA* gene is commonly used in qPCR assays where species discrimination and quantification are required (Chen et al., 2008; Forsgren and Fries, 2010; Higes et al., 2006; Klee et al., 2007; Martin-Hernandez et al., 2007). However, this is a multi-copy gene characterized by a variable number of sequences in the *N. ceranae* genome. This suggested to develop a new qPCR method, which is based on the single-copy *N. ceranae* gene *Hsp70* (Cilia et al., 2018).

The aim of this investigation was to seek the presence of *Nosema* spp. in managed colonies from the Tuscanian Archipelago islands and, concurrently, compare the results of the qPCR methods based on the *16S rRNA* and *Hsp70* genes.

2. Materials and methods

In July and August of 2017, honey bee workers were sampled on the islands of Tuscanian Archipelago in order to determine the *Nosema* infection.

Samples were taken on asymptomatic apiaries from the island of Pianosa (42°35'05.2"N 10°05'22.3"E), Capraia (43°01'54.4"N 9°29'59.8"E), Gorgona (43°26'00.1"N 9°54'18.0"E), and Elba (42°51'43.3"N 10°25'03.1"E) islands (Fig. 1). On Giannutri and Montecristo, no managed apiaries are known. On Giglio, the contacted beekeepers refused to take samples for this research.

On each island, two colonies from the same apiary were sampled. Each sample consisted on 50 forager bees, which were refrigerated until analysis.

The samples were divided in two subsamples of 25 bees, which were extracted separately. For extraction, the gastrum (later referred to as abdomen as of common use) was carefully dissected with tweezers. The abdomens of each subsample were pooled and homogenized (in 1 ml of DNA free water) with a Tissue Lyser II (Qiagen, Hilden, Germany) operated for 3 min at 30 Hz.

Total DNA was extracted with High Pure PCR Template Preparation Kit (Roche, Mannheim, Germany) following the manufacturer's instructions.

Two aliquots from each DNA extract were taken and analysed by different qPCR methods. In one case, primers and probes for *N. apis* and *N. ceranae* designed on a *16S rRNA* gene sequence were used as described by Forsgren and Fries (2010). The optimized amplification and quantitation protocol was: 15 min at 98 °C followed by 40 cycles of denaturation for 5 s at 98 °C, annealing/extension for 10 s at 63 °C and melt curve analysis from 65 to 95 °C (in 0.5 °C increments) 10 sec/step. The other aliquot served to quantify *N. ceranae* copies in each sample using the *Hsp70* gene-based primers and probes described by Cilia et al. (2018). In this case, amplification and quantitation were performed according to the following cycling conditions: initial activation step 95 °C for 10 min, PCR cycling (40 cycles of 95 °C for 15 s, 56 °C for 60 s).

All the analyses were conducted in duplicate and the final infection data reported as average of *Nosema* number copies per honey bee.

Data obtained were statistically analysed through a three-way ANOVA. Number of *N. ceranae* copies were tested against colony, qPCR method and replicate. Statistical significance threshold was set at $P = 0.05$. Where applicable, multiple comparison of means was made by using Newman-Keuls test.

Regression was calculated on the averages of the replicate analyses from each colony to compare the two qPCR methods.

All the statistics were calculated with the package STATISTICA (Sistema software di analisi dei dati), ver. 7.1, StatSoft Italia srl (2005).




Fig. 1. Map of Tuscanian Archipelago (Central Italy) with the distances in kilometres (km) between each island and the nearest land.

3. Results

All of the samples resulted in DNA amplification below the detection limit for *N. apis*. Table 1 reports the number of *N. ceranae* copies resulting from each replicate analysis performed with the two methods. No detectable *N. ceranae* infections were found in the samples from Elba Island, whereas the samples from Gorgona, Capraia and Pianosa samples were positive.

Samples positive for *N. ceranae* contained an average of 13.99 (± 10.34 s.e.; s.d. = 25.34) * 10⁶ copies per bee with the 16S rRNA-based method, but only 1.33 (± 0.71 s.e.; s.d. = 1.74) * 10⁶ by using the *Hsp70* sequences.

A principal effects ANOVA conducted on the data of Table 1 resulted in a significant effect of colony ($F(5, 16) = 4.020$, $p = 0.015$) and analytical method ($F(1, 16) = 5.380$, $p = 0.034$) but not of replicate ($F(1, 16) = 0.004$, $p = 0.950$).

On average, the 16S rRNA-based technique detected 8.59 (± 2.91 s.e.) times the *N. ceranae* copies found with the *Hsp70* method, and the ratio between the copies detected with the two methods (16S rRNA/*Hsp70*) was in the range 1.35–16.53.

Results show both high between-sample variability of this ratio (s.d. = 7.12) and insufficient correlation between the paired replicate averages to make direct conversion between the two methods possible (Pearson $r = 0.832$ ($F(1, 4) = 9.020$, $p = 0.040$)).

In a regression analysis of data ($y = b_0 + b_1 * x$), the number of copies detected with 16S rRNA and *Hsp70* methods were respectively considered dependent and independent variables, with good fit to the linear model ($F(1,4) = 9.02p < 0.040$), $b_0 = -2,061,789$ ($p = 0.817$) and $b_1 = 12,09$ ($p = 0.040$). However, low determination index (Adjusted $R^2 = 0.616$) confirms that major factors other than *Hsp70* estimate should be considered to account for the variability in the number of *N. ceranae* copies detected with the 16S rRNA-based method.

4. Discussion

The results of this investigation allow to include Tuscanian Archipelago among the Italian areas where *N. ceranae* can be found in honey bee colonies (Ferroglio et al., 2013; Klee et al., 2007; Maiolino et al., 2014; Mutinelli et al., 2010; Porrini et al., 2016). This confirms that, despite the likely infrequent exchanges with the mainland populations, this pathogen may find suitable conditions to thrive in small Mediterranean islands. In this respect, samples collected from the comparatively larger Elba island Elba should be considered more occasional finding than a realistic representation of the overall situation.

N. apis was not detected in any of the investigated colonies. Although, the results of this investigation do not clarify this aspect, seasonality (Martin-Hernandez et al., 2007; Ondrus, 2017), different plasticity to temperature (Fries et al., 2006; Higes et al., 2009a,b; Klee et al., 2007; Paxton et al., 2007) and/or competition between the two investigated species (Chen et al., 2012; Forsgren and Fries, 2010; Martín-Hernández et al., 2012) may account for the difference in prevalence between the two *Nosema* species.

In the three islands where positive samples could be found, the number of *N. ceranae* copies detected with both qPCR methods was in the order of magnitude of 10⁴ or higher, which is compatible with fully developed infections and suggestive of consolidated pathogen populations in those islands.

As resulted by the three-way ANOVA test, colony was a significant source of variability, suggesting heterogeneous levels of infection in the considered hives. Conversely, the replicate did not result into significant effect. This is compatible with high repeatability of both qPCR methods (Cilia et al., 2018; Martin-Hernandez et al., 2007). However, *Hsp70* qPCR assay systematically detected less *N. ceranae* copies than the alternate method, whose results were one or two orders of magnitude higher. This and the unsteady 16S rRNA/*Hsp70* ratio calculated at sample level agrees with the variable number of sequences of the 16S rRNA gene of *N. ceranae* (Sagastume et al., 2016). Molecular diagnostics may take advantage of multi-copy genes in that they allow designing high sensitivity tests (more possibility to detect the target gene), but the fluctuations in the number of 16S rRNA sequences in the *N. ceranae* genome may ultimately affect the quantification reliability (Sagastume et al., 2011). Conversely, the *Hsp70* method bases on a highly-conserved region of the *N. ceranae* genome (Gomez-Moracho et al., 2014; Wang et al., 2017), making it suitable to accurate quantification.

The results of the 16S rRNA amplification may be correlated to the highly variable number of copies of multi-copy genes. The PCR primers may affect their capability to anneal to the template influencing the assay sensitivity. On the other hand, the results of the *Hsp70* gene amplification, detecting a lower number of copies, may be correlated to its highly conserved sequence. The designed PCR primers anneal to the template once, with high sensitivity and specificity.

In conclusion, the results of this investigation provide evidence that, as well known, *N. ceranae* may successfully proliferate in asymptomatic honey bee colonies living in small Mediterranean islands and indicate that *N. apis* might not find similarly suitable conditions to thrive. In this respect, further research is required to elucidate the long-term host-parasite relationship in limited

Table 1
Replicates of *Nosema ceranae* gene copies detected with 16S rRNA and *Hsp70* qPCR assay in honey bee samples from Tuscanian Archipelago.

Island	Colony	Replicate	16S gene copies/bee	<i>Hsp70</i> gene copies/bee
Capraia	C1	1	3,16E+05	2,28E+04
		2	3,28E+05	2,44E+04
	C2	1	1,16E+07	3,12E+06
		2	1,28E+07	3,24E+06
Gorgona	G1	1	4,52E+04	3,36E+04
		2	4,96E+04	3,64E+04
	G2	1	5,88E+06	3,96E+05
		2	6,12E+06	4,16E+05
Pianosa	P1	1	5,52E+05	3,80E+05
		2	5,80E+05	4,28E+05
	P2	1	6,36E+07	3,88E+06
		2	6,60E+07	3,96E+06
Elba	E1	1	<LOD	<LOD
		2	<LOD	<LOD
	E2	1	<LOD	<LOD
		2	<LOD	<LOD

environments characterized by genetic isolation. A comparison between different q-PCR assays showed profound differences in the results when the primers are designed on single-copy or multi-copy sequences. The choice of analytical method may then be critical when reliable quantification is required.

References

- Ansari, M.J., Al-Ghamdi, A., Nuru, A., Khan, K.A., Alattal, Y., 2017. Geographical distribution and molecular detection of *Nosema ceranae* from indigenous honey bees of Saudi Arabia. *Saudi J. Biol. Sci.* 24, 983–991. <https://doi.org/10.1016/j.sjbs.2017.01.054>.
- Botías, C., Anderson, D.L., Meana, A., Garrido-Bailón, E., Martín-Hernández, R., Higes, M., 2012. Further evidence of an oriental origin for *Nosema ceranae* (Microsporidia: Nosematidae). *J. Invertebr. Pathol.* 110, 108–113. <https://doi.org/10.1016/j.jip.2012.02.014>.
- Bourgeois, A.L., Rinderer, T.E., Beaman, L.D., Danka, R.G., 2010. Genetic detection and quantification of *Nosema apis* and *N. ceranae* in the honey bee. *J. Invertebr. Pathol.* 103, 53–58. <https://doi.org/10.1016/j.jip.2009.10.009>.
- Calderón, R.A., Sanchez, L.A., Yañez, O., Fallas, N., 2008. Presence of *Nosema ceranae* in Africanized honey bee colonies in Costa Rica. *J. Apic. Res.* 47, 328–329. <https://doi.org/10.1080/00218839.2008.11101485>.
- Chemurot, M., De Smet, L., Brunain, M., De Ryck, R., de Graaf, D.C., 2017. *Nosema neumanni* n. sp. (Microsporidia, Nosematidae), a new microsporidian parasite of honeybees, *Apis mellifera* in Uganda. *Eur. J. Protistol.* 61, 13–19. <https://doi.org/10.1016/j.ejop.2017.07.002>.
- Chen, Y., Evans, J.D., Smith, I.B., Pettis, J.S., 2008. *Nosema ceranae* is a long-present and wide-spread microsporidian infection of the European honey bee (*Apis mellifera*) in the United States. *J. Invertebr. Pathol.* 97, 186–188. <https://doi.org/10.1016/j.jip.2007.07.010>.
- Chen, Y.P., Evans, J.D., Murphy, C., Gutell, R., Zuker, M., Gundensen-Rindal, D., Pettis, J.S., 2009. Morphological, molecular, and phylogenetic characterization of *Nosema ceranae*, a microsporidian parasite isolated from the European honey bee, *Apis mellifera*. *J. Eukaryot. Microbiol.* 56, 142–147. <https://doi.org/10.1111/j.1550-7408.2008.00374.x>.
- Chen, Y.W., Chung, W.P., Wang, C.H., Solter, L.F., Huang, W.F., 2012. *Nosema ceranae* infection intensity highly correlates with temperature. *J. Invertebr. Pathol.* 111, 264–267. <https://doi.org/10.1016/j.jip.2012.08.014>.
- Cilia, G., Cabbri, R., Maiorana, G., Cardaio, I., Dall'Olio, R., Nanetti, A., 2018. A novel TaqMan[®] assay for *Nosema ceranae* quantification in honey bee, based on the protein coding gene Hsp70. *Eur. J. Protistol.* 63, 44–50. <https://doi.org/10.1016/j.ejop.2018.01.007>.
- Eiri, D.M., Suwannapong, G., Endler, M., Nieh, J.C., 2015. *Nosema ceranae* can infect honey bee larvae and reduces subsequent adult longevity. *PLoS One* 10. <https://doi.org/10.1371/journal.pone.0126330> e0126330.
- Ferroglio, E., Zanet, S., Peraldo, N., Tachis, E., Trisciuglio, A., Laurino, D., Porporato, M., 2013. *Nosema ceranae* has been infecting honey bees *Apis mellifera* in Italy since at least 1993. *J. Apic. Res.* 52, 60–61. <https://doi.org/10.3896/IBRA.1.52.2.11>.
- Forsgren, E., Fries, I., 2010. Comparative virulence of *Nosema ceranae* and *Nosema apis* in individual European honey bees. *Vet. Parasitol.* 170, 212–217. <https://doi.org/10.1016/j.vetpar.2010.02.010>.
- Fries, I., 1993. *Nosema Apis* - a parasite in the honey bee colony. *Bee World* 74, 5–19. <https://doi.org/10.1080/0005772X.1993.11099149>.
- Fries, I., Feng, F., da Silva, A., Slemenda, S.B., Pieniazek, N.J., 1996. *Nosema ceranae* n. sp. (Microsporida, Nosematidae), morphological and molecular characterization of a microsporidian parasite of the Asian honey bee *Apis cerana* (Hymenoptera, Apidae). *Eur. J. Protistol.* 32, 356–365. [https://doi.org/10.1016/S0932-4739\(96\)80059-9](https://doi.org/10.1016/S0932-4739(96)80059-9).
- Fries, I., Martín, R., Meana, A., García-Palencia, P., Higes, M., 2006. Natural infections of *Nosema ceranae* in European honey bees. *J. Apic. Res.* 45, 230–233. <https://doi.org/10.1080/00218839.2006.11101355>.
- Giersch, T., Berg, T., Galea, F., Hornitzky, M., 2009. *Nosema ceranae* infects honey bees (*Apis mellifera*) and contaminates honey in Australia. *Apidologie* 40, 117–123. <https://doi.org/10.1051/apido/2008065>.
- Gomez-Moracho, T., Maside, X., Martín-Hernández, R., Higes, M., Bartolomé, C., 2014. High levels of genetic diversity in *Nosema ceranae* within *Apis mellifera* colonies. *Parasitology* 141, 475–481. <https://doi.org/10.1017/S0031182013001790>.
- Higes, M., García-Palencia, P., Martín-Hernández, R., Meana, A., 2007. Experimental infection of *Apis mellifera* honeybees with *Nosema ceranae* (Microsporida). *J. Invertebr. Pathol.* 94, 211–217. <https://doi.org/10.1016/j.jip.2006.11.001>.
- Higes, M., Martín-Hernández, R., Botías, C., Bailón, E.G., González-Porto, A.V., Barrios, L., Del Nozal, M.J., Bernal, J.L., Jiménez, J.J., Palencia, P.G., Meana, A., 2008. How natural infection by *Nosema ceranae* causes honeybee colony collapse. *Environ. Microbiol.* 10, 2659–2669. <https://doi.org/10.1111/j.1462-2920.2008.01687.x>.
- Higes, M., Martín-Hernández, R., Garci-a-Palencia, P., Mari-n, P., Meana, A., 2009a. Horizontal transmission of *Nosema ceranae* (Microsporida) from worker honeybees to queens (*Apis mellifera*). *Environ. Microbiol. Rep.* 1, 495–498. <https://doi.org/10.1111/j.1758-2229.2009.00052.x>.
- Higes, M., Martín-Hernández, R., Garrido-Bailón, E., Botías, C., Meana, A., 2009b. The presence of *Nosema ceranae* (Microsporida) in North African honey bees (*Apis mellifera intermissa*). *J. Apic. Res.* 48, 217–219. <https://doi.org/10.3896/IBRA.1.48.3.12>.
- Higes, M., Martín-Hernández, R., Garrido-Bailón, E., González-Porto, A.V., García-Palencia, P., Meana, A., del Nozal, M.J., Mayo, R., Bernal, J.L., Kamer, Y., Klinberg, E., Zioni, N., Inbar, S., Chejanovsky, N., Alaux, C., 2009c. Honeybee colony collapse due to *Nosema ceranae* in professional apiaries. *Environ. Microbiol. Rep.* 1, 110–113. <https://doi.org/10.1111/j.1758-2229.2009.00014.x>.
- Higes, M., Martín, R., Meana, A., 2006. *Nosema ceranae*, a new microsporidian parasite in honeybees in Europe. *J. Invertebr. Pathol.* 92, 93–95. <https://doi.org/10.1016/j.jip.2006.02.005>.
- Invernizzi, C., Abud, C., Tomasco, I.H., Harriet, J., Ramallo, G., Campá, J., Katz, H., Gardiol, G., Mendoza, Y., 2009. Presence of *Nosema ceranae* in honeybees (*Apis mellifera*) in Uruguay. *J. Invertebr. Pathol.* 101, 150–153. <https://doi.org/10.1016/j.jip.2009.03.006>.
- Klee, J., Besana, A.M., Genersch, E., Gisder, S., Nanetti, A., Tam, D.Q., Chinh, T.X., Puerta, F., Ruz, J.M., Kryger, P., Message, D., Hatjina, F., Korpela, S., Fries, I., Paxton, R.J., 2007. Widespread dispersal of the microsporidian *Nosema ceranae*, an emergent pathogen of the western honey bee, *Apis mellifera*. *J. Invertebr. Pathol.* 96, 1–10. <https://doi.org/10.1016/j.jip.2007.02.014>.
- Maiolino, P., Iafigliola, L., Rinaldi, L., De Leva, G., Restucci, B., Martano, M., 2014. Histopathological findings of the midgut in European honey bee (*Apis mellifera* L.) naturally infected by *Nosema* spp. *Vet. Med. Anim. Sci.* 2, 4. <https://doi.org/10.7243/2054-3425-2-4>.
- Martín-Hernández, R., Botías, C., Bailón, E.G., Martínez-Salvador, A., Prieto, L., Meana, A., Higes, M., 2012. Microsporidia infecting *Apis mellifera*: coexistence or competition. Is *Nosema ceranae* replacing *Nosema apis*? *Environ. Microbiol.* 14, 2127–2138. <https://doi.org/10.1111/j.1462-2920.2011.02645.x>.
- Martín-Hernández, R., Meana, A., Prieto, L., Salvador, A.M., Garrido-Bailón, E., Higes, M., 2007. Outcome of colonization of *Apis mellifera* by *Nosema ceranae*. *Appl. Environ. Microbiol.* 73, 6331–6338. <https://doi.org/10.1128/AEM.00270-07>.
- Mutinelli, F., Costa, C., Lodesani, M., Baggio, A., Medrzycki, P., Formato, G., Porrini, C., 2010. Honey bee colony losses in Italy. *J. Apic. Res.* 49, 119–120. <https://doi.org/10.3896/IBRA.1.49.1.24>.
- Ondrus, P., 2017. The seasonality, thermal dependence, and spread of *Nosema ceranae* in new locales. *Bee World* 94, 105–109. <https://doi.org/10.1080/0005772X.2017.1382041>.
- Pacini, A., Mira, A., Molineri, A., Giacobino, A., Bulacio Cagnolo, N., Aignasse, A., Zago, L., Izaguirre, M., Merke, J., Orellano, E., Bertozzi, E., Pietronave, H., Russo, R., Scannapieco, A., Lanzavecchia, S., Schnitter, L., Signorini, M., 2016. Distribution and prevalence of *Nosema apis* and *N. ceranae* in temperate and subtropical eco-regions of Argentina. *J. Invertebr. Pathol.* 141, 34–37. <https://doi.org/10.1016/j.jip.2016.11.002>.
- Paxton, R.J., Klee, J., Korpela, S., Fries, I., 2007. *Nosema ceranae* has infected *Apis mellifera* in Europe since at least 1998 and may be more virulent than *Nosema apis*. *Apidologie* 38, 558–565. <https://doi.org/10.1051/apido:2007037>.
- Porrini, C., Mutinelli, F., Bortolotti, L., Granato, A., Laurensen, L., Roberts, K., Gallina, A., Silvester, N., Medrzycki, P., Renzi, T., Sgolastra, F., Lodesani, M., 2016. The status of honey bee health in Italy: results from the nationwide bee monitoring network. *PLoS One* 11. <https://doi.org/10.1371/journal.pone.0155411> e0155411.
- Sagastume, S., del Águila, C., Martín-Hernández, R., Higes, M., Henriques-Gil, N., 2011. Polymorphism and recombination of rDNA in the putatively asexual microsporidian *Nosema ceranae*, a pathogen of honeybees. *Environ. Microbiol.* 13, 84–95. <https://doi.org/10.1111/j.1462-2920.2010.02311.x>.
- Sagastume, S., Martín-Hernández, R., Higes, M., Henriques-Gil, N., 2016. Genotype diversity in the honey bee parasite *Nosema ceranae*: multi-strain isolates, cryptic sex or both? *BMC Evol. Biol.* 16, 216. <https://doi.org/10.1186/s12862-016-0797-7>.
- Tapasztó, Z., Forgách, P., Kóvágo, C., Békési, L., Bakonyi, T., Rusvai, M., 2009. First detection and dominance of *Nosema ceranae* in Hungarian honeybee colonies. *Acta Vet. Hung.* 57, 383–388. <https://doi.org/10.1556/AVet.57.2009.3.4>.
- Wang, Y., Ru, Y., Liu, W., Wang, D., Zhou, J., Jiang, Y., Shi, S., Qin, L., 2017. Morphological characterization and HSP70-, IGS-based phylogenetic analysis of two microsporidian parasites isolated from *Antheraea pernyi*. *Parasitol. Res.* 116, 971–977. <https://doi.org/10.1007/s00436-017-5373-6>.