

# In Vivo MR Imaging of Intraarterially Delivered Magnetically Labeled Mesenchymal Stem Cells in a Canine Stroke Model

Shan-shan Lu<sup>1</sup>✉, Sheng Liu<sup>1</sup>✉, Qing-quan Zu<sup>1</sup>, Xiao-quan Xu<sup>1</sup>, Jing Yu<sup>1</sup>, Jian-wei Wang<sup>1</sup>, Yu Zhang<sup>2</sup>, Hai-bin Shi<sup>1\*</sup>

**1** Department of Radiology, The First Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu Province, China, **2** State Key Laboratory of Bioelectronics, Jiangsu Key Laboratory for Biomaterials and Devices, School of Biological Science & Medical Engineering, Southeast University, Nanjing, Jiangsu Province, China

## Abstract

**Background:** This study aimed to evaluate the feasibility of intraarterial (IA) delivery and in vivo MR imaging of superparamagnetic iron oxide (SPIO)-labeled mesenchymal stem cells (MSCs) in a canine stroke model.

**Methodology:** MSCs harvested from beagles' bone marrow were labeled with home-synthesized SPIO. Adult beagle dogs (n = 12) were subjected to left proximal middle cerebral artery (MCA) occlusion by autologous thrombus, followed by two-hour left internal carotid artery (ICA) occlusion with 5 French vertebral catheter. One week later, dogs were classified as three groups before transplantation: group A: complete MCA recanalization, group B: incomplete MCA recanalization, group C: no MCA recanalization.  $3 \times 10^6$  labeled-MSCs were delivered through left ICA. Series in vivo MRI images were obtained before cell grafting, one and 24 hours after transplantation and weekly thereafter until four weeks. MRI findings were compared with histological studies at the time point of 24 hours and four weeks.

**Principal Findings:** Home-synthesized SPIO was useful to label MSCs without cell viability compromise. MSCs scattered widely in the left cerebral hemisphere in group A, while fewer grafted cells were observed in group B and no cell was detected in group C at one hour after transplantation. A larger infarction on the day of cell transplantation was associated with more grafted cells in the brain. Grafted MSCs could be tracked effectively by MRI within four weeks and were found in peri-infarction area by Prussian blue staining.

**Conclusion:** It is feasible of IA MSCs transplantation in a canine stroke model. Both the ipsilateral MCA condition and infarction volume before transplantation may affect the amount of grafted cells in target brain. In vivo MR imaging is useful for tracking IA delivered MSCs after SPIO labeling.

**Citation:** Lu S-s, Liu S, Zu Q-q, Xu X-q, Yu J, et al. (2013) In Vivo MR Imaging of Intraarterially Delivered Magnetically Labeled Mesenchymal Stem Cells in a Canine Stroke Model. PLoS ONE 8(2): e54963. doi:10.1371/journal.pone.0054963

**Editor:** Marcel Daadi, Stanford University School of Medicine, United States of America

**Received:** August 15, 2012; **Accepted:** December 17, 2012; **Published:** February 7, 2013

**Copyright:** © 2013 Lu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This study was supported by National Natural Science Foundation of China (30870710 to Hai-bin Shi, 81000653 to Sheng Liu) and foundation of Research and Innovation Program for postgraduates in Jiangsu Province (CXZZ11.0718 to Shan-shan Lu). <http://www.nsf.gov.cn/Portal0/default152.htm>; [http://www.ecjss.edu.cn/art/2011/6/11/art\\_4266\\_26988.html](http://www.ecjss.edu.cn/art/2011/6/11/art_4266_26988.html). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: shihb@vip.sina.com


✉ These authors contributed equally to this work.

## Introduction

Stem cell based therapies have been shown to improve functional outcome in many experimental stroke. The likely mechanisms have been suggested as the neuroprotection [1–3], angiogenesis [4–6], axon-myelin remodeling [7], endogenous cell proliferation and possible replacement of damaged cells [8]. In the past years, because of relatively low immunogenicity and easy way of isolation, many studies used mesenchymal stem cells (MSCs) for stroke and suggested their effectiveness in cerebral ischemia. However, most of those studies were performed in rodents with intravenous or stereotactic injection of stem cells [4,9,10]. Unfortunately, the rodent brain is not gyrencephalic like the human brain. Besides, intravenously delivered cells pass through

the systemic and pulmonary circulation systems which significantly reduce cells homing to the injured brain. Intracerebral injection is highly invasive and requires craniotomy [11]. Intuitively, the intra-arterial (IA) route of delivery, given that the first pass of MSCs would be brain, seems better and enables widespread cerebral engraftment of cells. Furthermore, IA delivery of MSCs is clinically available because transfemoral catheterization of selected arteries has already been widely performed [12–14].

Magnetic resonance imaging (MRI) has been used to evaluate distribution and migration of stem cells after labeled with superparamagnetic iron oxide (SPIO) contrast agents [15–17]. To our knowledge, few studies have addressed in vivo MRI tracking of intra-arterially delivered MSCs in large animals like beagle dogs which are structurally similar to the human brain.



**Figure 1. The transmission electronic microscopy image of home-synthesized  $\text{Fe}_3\text{O}_4$  nanoparticles.** The average size of  $\text{Fe}_3\text{O}_4$  nanoparticles calculated from 100 particles at least was 12 nm. doi:10.1371/journal.pone.0054963.g001

The aims of the present study were as follows: 1. Is it feasible of MSCs transplantation by IA route in an experimental canine model of cerebral infarction? 2. Can 3.0 Tesla (T) MRI effectively track the IA transplanted SPIO-labeled MSCs?

## Materials and Methods

### Synthesis of $\text{Fe}_3\text{O}_4$ -DMSA-PLL (SPIO)

0.5 mol·L<sup>-1</sup> FeCl<sub>3</sub> and 0.25 mol·L<sup>-1</sup> FeSO<sub>4</sub>·7H<sub>2</sub>O were dissolved in 100 ml deionized water in a three-neck flask followed by adding 2 mol·L<sup>-1</sup> NH<sub>3</sub>·H<sub>2</sub>O solution with stirring at 70°C under a N<sub>2</sub> atmosphere. Then NH<sub>3</sub>·H<sub>2</sub>O was progressively added to adjust the aqueous dispersion pH to 11.0 - 12.0. The temperature was raised to 65°C and the system was then mixed for two hours at 65°C. 30 ml oleic acid (OA) was slowly added to the solution. The pH was adjusted with dilute hydrochloric acid (HCL) solution to 6.0 - 7.0, and the temperature was raised to 80°C for one hour, and then cooled down to room temperature with the pH controlled at 3.4 - 4.0. Black precipitates were collected by filtration and carefully washed with deionized water and ethanol several times. The obtained  $\text{Fe}_3\text{O}_4$ -OA nanoparticles were dissolved in 100 ml hexanol.

0.65 mmol  $\text{Fe}_3\text{O}_4$ -OA and 0.3 mmol meso-2,3-Dimercaptosuccinic acid (DMSA) were dissolved in 50 ml acetone with stirring at 60°C for four hours. Then  $\text{Fe}_3\text{O}_4$ -DMSA nanoparticles were collected by magnetic separation, washed with ethanol and deionized water several times and dissolved in 20 ml deionized water.

To 20 ml of 0.5 mg/ml poly-L-lysine (PLL) solution, 20 ml of 0.175 mg/ml  $\text{Fe}_3\text{O}_4$ -DMSA were added under stirring. The mixture was sonicated for 20 minutes and then further stirred for two hours. The  $\text{Fe}_3\text{O}_4$ -DMSA-PLL nanoparticles were obtained by removing free PLL via ultrafiltration.

### Cell Culture and Labeling

Bone marrow was obtained from the humerus of dog two weeks before embolization. Autologous MSCs were isolated and purified by density gradient centrifugation and adhering to the wall. Briefly

speaking, the bone marrow was diluted (1:1) in phosphate buffered saline (PBS), layered onto a density gradient solution (Ficoll-Paque, TBDscience, Tianjin, China) and centrifuged for 25 minutes at 300 grams. After that, enriched cells were removed from plasma-solution interphase, washed twice with PBS, and cultured in low-glucose Dulbecco's modified Eagle's medium (DMEM, Gibco, Karlsruhe, Germany) supplemented with 0.2 mmol/ml of L-glutamine, 100 U of penicillin, 100 g/ml of streptomycin and 10% fetal bovine serum (FBS, Gibco, Karlsruhe, Germany) at 37°C and 5% CO<sub>2</sub> [18]. Three days later, the culture solution was replaced for the first time, and nonadherent cells were removed at the same time. The medium was changed every three days. After the cultures reached 90% confluence, the cells were harvested with 0.25% trypsin (Gibco, Karlsruhe, Germany) and replated.


Passage three cells were co-cultured in fresh medium containing various concentrations of SPIO (5, 10, 20, 40, 80 µg/ml) for 24 hours according to the previous reports [19] and then were washed three times in PBS to eliminate extracellular SPIO.

After that, prussian blue (PB) staining was performed to show intracellular iron. The cells were incubated with 2% potassium ferrocyanide in 6% HCL for 15 minutes and then counterstained with nuclear fast red for one minute. The intracellular iron content was also quantified before and after cell labeling by atomic absorption spectrometry (Shengyang, China) as described previously [19]. The average iron content per cell was calculated according to the cell numbers.

### Cellular Viability and Apoptosis

For assessment of cell viability, cultures were incubated with 0.4% trypan blue (Sigma, St. Louis, MO, USA). Cell viability was determined according to the percentage of cells not internalizing the dye.

For assessment of cell apoptosis, labeled cells ( $1 \times 10^6$ ) and unlabeled cells ( $1 \times 10^6$ ) were collected, washed twice with cool PBS and resuspended in 50 µl annexin medium, followed by adding 5 µl annexin V-fluorescein isothiocyanate (AV-FITC). The suspension was then kept in dark for 15 minutes at room temperature. 10 µl propidium iodide (PI) was added and the



**Figure 2. Photomicrography of MSCs after incubated with  $\text{Fe}_3\text{O}_4$ -DMSA-PLL.** Prussian blue stained smear of labeled MSCs showed most of the cells were prussian blue positive (A). Intracytoplasmic blue particles were clearly visible under inverted phase contrast microscope (B).  
doi:10.1371/journal.pone.0054963.g002


suspension was again stored in dark for five minutes. After that, the cells were washed twice again with cool PBS and resuspended with 300  $\mu\text{l}$  annexin. Flow cytometer (BD FACSCalibur, CA, USA) was used to determine the percentage of cells undergoing early apoptosis (annexin<sup>+</sup>/PI<sup>-</sup>) and late apoptosis or death (annexin<sup>+</sup>/PI<sup>+</sup>).

#### Animal Ischemia Model Establishment and the Transplantation Procedure


All the experimental procedures were approved by the local Animal Experiment Ethical Committee. 14 beagle dogs (body weight,  $13.9 \pm 1.0$  kg) were enrolled and randomized to transplantation group (n = 12) and control group (n = 2). They were

anesthetized with 3 mg/kg pentobarbital intravenously during all the procedure. Rectal temperature was continuously monitored and maintained at 37°C to 39°C.

The left proximal MCA was embolized with autologous thrombus (1.7 mm in diameter and 5 mm in length) prepared approximately two hours before as described in our previous study [20]. Under angiography (Axiom Artis; Siemens AG, Muenchen, Germany), a 5 French vertebral catheter (Terumo Medical Corporation, Somerset, NJ, USA) was advanced 2 cm into the ascending part of left internal carotid artery (ICA). One thrombus was injected into the ICA with 3 ml of 0.9% saline. Complete left proximal MCA occlusion (MCAO) was verified by following angiography. Then the catheter was kept in the left ICA for two



**Figure 3. AV-FITC-PI double staining for MSCs apoptosis after labeling.** Early apoptotic percent (lower right quadrant) were 4.95% and 6.36% while late apoptotic and necrotic percent (upper right quadrant) were 1.89% and 2.93%, respectively in unlabeled cells (left) and labeled cells (right). The cross inside was gate.  
doi:10.1371/journal.pone.0054963.g003



**Figure 4. Three types of ischemic infarction and the amount of grafted MSCs in the target brain.** Moderate size of cerebral infarction involving the left temporal lobe (A), large infarction in left MCA territory (B) and small infarction located in subcortical white matter (C) were got after combining left proximal MCA embolization with two-hour left ICA blood flow occluding. All these three dogs (dog 5, 12 and 13) had complete left MCA recanalization before transplantation (not shown). After intra-carotid MSCs transplantation, dog with large infarction had more obvious MSCs in the brain than that with moderate size of infarction (D-E), while dog with very small infarction was shown fewer scattered cells (F). doi:10.1371/journal.pone.0054963.g004

hours for continuously occluding the blood flow. All the dogs were kept warm with heating blanket and observed continuously until they had fully recovered from anesthesia. Intramuscular ampicillin (20 mg/kg) was given to all of them daily for up to one week.

One week after ischemia, all the animals in experimental group underwent angiography before MSCs transplantation. They were then classified as three groups according to the angiography findings. Group A: dogs with complete left MCA recanalization and distal blood flow; group B: dogs with incomplete left MCA recanalization regardless whether there was distal blood flow or not; group C: dogs with persistent left proximal MCA occlusion without any distal blood flow. A bolus of cell suspension with about  $3 \times 10^6$  MSCs inside was slowly injected over a five-minute period into all the dogs using a 10 ml syringe through a 5 F vertebral catheter in the left ICA, followed by 5 ml 0.9% saline. The catheter was left in place for 10 minutes before withdrawal to minimize cell backflow.

#### In vivo MR Imaging

3.0 T MRI scanner (Magnetom Trio, Siemens Medical Systems, Germany) and an eight-channel knee coil were used in the study. Dogs were positioned supinely in the MR scanner with head in the center of the coil.

T2 weighted imaging (T2WI) and diffusion weighted imaging (DWI) were performed at six hours, 24 hours and one week after MCA embolization and temporal ICA occlusion. T2\* weighted

imaging (T2\*WI), T2\* map and susceptibility weighted imaging (SWI) were added to in vivo MRI imaging and were obtained just before cell grafting and at one hour, 24 hours and weekly thereafter until four weeks after IA transplantation. The parameters were as follows: T2WI: TR 5000 ms, TE 60 ms, matrix  $320 \times 320$ ; T2\*WI: TR 1460 ms, six TEs (20, 25.2, 30.4, 35.4, 40.8, 46.0 ms), matrix  $320 \times 320$ ; T2\* map: TR 80 ms, TE 20 ms, matrix  $256 \times 256$ ; SWI: TR 28 ms, TE 20 ms, matrix  $448 \times 448$ , and DWI: TR 5500 ms, TE 96 ms, matrix  $192 \times 192$ , b value 0 and  $1000 \text{ sec/mm}^2$ . Other parameters were the same: FOV 200 mm, 22 slices, and slice thickness 2.0 mm with no gap.

#### MRI Data Processing

One experienced radiologist (XQ Xu) measured visible ischemic lesions by drawing regions of interest around the lesions on T2WI images at 24 hours and one week after embolization. Image brightness and contrast were optimized between areas of abnormal tissue and normal appearing brain. Lesion volumes were obtained by multiplying the lesion areas by the section and gap thickness.

T2\* value of transplantation group was measured from T2\* maps by one radiologist (QQ Zu). The ROIs were selected to avoid susceptibility effects arising from air tissue interface and hemorrhage. Each ROI area was  $20 \text{ mm}^2$ . ROIs were placed on the peri-infarction (PI) area, ipsilateral normal parenchyma (INP) and infarction itself, according to the corresponding T2WI and SWI, respectively. T2\* ratios were got after normalizing with the

**Table 1.** Dog grouping and infarction volume on T2WI after embolization.

Grouping *	Dog ID	Infarct volume on T2WI after embolization (mm <sup>3</sup> )	
		24 hours	1 week
Group A	Dog 2	841.8	410.54
	Dog 5	819.94	319.74
	Dog 8	857.9	430.26
	Dog 12	3777.9	3185.16
	Dog 13	64.84	51.14
Group B	Dog 1	725.6	587.28
	Dog 9	468.36	319.36
	Dog 10	876.56	478.12
	Dog 3	230.26	304.1
Group C	Dog 14	251.36	423.64
	Dog 7	5902	3367.96
	Dog 11	318.52	312.52
Control group	Dog 4	3255.32	1960.9
	Dog 6	604.8	323.76

Note: \* Group A: complete left MCA recanalization with distal blood flow before transplantation; Group B: incomplete left MCA recanalization regardless whether there was distal blood flow or not before transplantation; Group C: persistent left MCA occlusion without any distal blood flow before transplantation; Control group: dogs without cell transplantation. doi:10.1371/journal.pone.0054963.t001

contralateral mirror region. Due to the heterogeneity of T2\* maps, we measured T2\* ratio of each area three times in each dog. The mean value of T2\* ratio was used for further analyses.

### Histopathological Analysis

Two dogs (dog 8, 10) were sacrificed at 24 hours after transplantation to localize the MSCs. Other dogs were sacrificed at the fourth week. The brains were removed and immersed in 4% paraformaldehyde in PBS at 4°C for two days and then cut into 2-mm thick coronal blocks, followed by embedded in paraffin. Four  $\mu$ m thick coronal sections were sliced from each block and stained for hematoxylin and eosin (HE staining).

To visualize the transplanted MSCs in the host brain, sections were processed for PB staining. The sections were incubated for 30

minutes with 2% potassium ferrocyanide in 6% HCL, counterstained with eosin and viewed under light microscope (model AH3, Olympus, Tokyo, Japan).

### Neurological Scoring

Neurological scoring was performed using a standardized categorical rating scale according to previous report [21]. It was consisted of evaluation for motor function (no deficit = 1, hemiparetic but able to walk = 2, stands only with assistance = 3, hemiplegic and unable to stand = 4, comatose or dead, not testable = 4), consciousness (normal = 1, mildly reduced = 2, severely reduced = 3, comatose or dead = 4), head turning (absent = 0, posturing and turns toward side of infarct = 1, unable to lift head, comatose, or dead = 1), circling (absent = 0, present = 1, does not walk or dead = 1) and hemianopsia (absent = 0, present = 1, unable to test because of reduced consciousness or death = 1). Dogs were evaluated at 24 hours, one week after ischemia and then weekly until sacrifice.

### Statistical Analysis

All data were reported as mean  $\pm$  standard deviation (SD). The student *t* test was used to determine whether there was difference between SPIO-labeled MSCs and control group. One-way analysis of variance followed by a post-hoc test was applied to assess the T2\* ratio differences among the three groups and between any two groups one hour after IA transplantation. The *P*-value was two-sided and value of less than 0.05 was considered statistically significant. All analyses were performed using SPSS (Version 13.0, Chicago, Illinois).


## Results

### Fe<sub>3</sub>O<sub>4</sub>-DMSA-PLL Nanoparticles


The prepared Fe<sub>3</sub>O<sub>4</sub>-DMSA-PLL nanoparticles were stable and fully water-soluble. The transmission electronic microscopy (TEM, JEOL JEM-2100) image showed the Fe<sub>3</sub>O<sub>4</sub> nanoparticles with the average size of 12.2 $\pm$ 4.3 nm (Figure 1). Hydrodynamic size and zeta potential of Fe<sub>3</sub>O<sub>4</sub>-DMSA-PLL nanoparticles were 152.8 $\pm$ 2.2 nm and +43.43 $\pm$ 6.1 mV.

### Cell Culture, Labeling, Viability and Apoptosis

After PB staining, blue particles were seen in almost every MSC (Figure 2A–B). As the concentration of SPIO increased from 5–80  $\mu$ g/ml, the blue cytoplasmic inclusions increased proportionally. Low concentration (20  $\mu$ g/ml) was favored for further use



**Figure 5. Comparison of three sequences for detecting IA transplanted MSCs.** Grafted MSCs were shown obvious hypointensity and diffuse distribution on both SWI (A) and T2WI images (B). T2WI image allowed a proper visualization of the ischemic brain tissue but it failed to detect the MSCs in the brain (C). doi:10.1371/journal.pone.0054963.g005



**Figure 6. The amount of engrafted MSCs in the brain in three groups.** (A-B) Large amounts of transplanted MSCs scattered in a wide range in the left cerebral hemisphere after IA delivery in group A with complete MCA recanalization (long arrow); (C-D) Fewer amounts of MSCs in the brain were observed in group B with incomplete MCA recanalization (empty arrow); (E-F) No definite hypointensity of MSC was observed in group C with continuous MCA occlusion before transplantation (short arrow).  
doi:10.1371/journal.pone.0054963.g006

considering of safety. The mean intracellular iron was  $33.14 \pm 4.35$  pg/cell in labeled cells in comparison with  $1.56 \pm 0.47$  pg/cell in the unlabeled cells.

Trypan blue staining showed SPIO loading had no effect on cell viability when compared with control group ( $t = 1.682$ ,  $P > 0.05$ ). The mean viability was  $94.71\% \pm 1.48\%$  and  $96.14\% \pm 1.46\%$ , respectively. Cell apoptosis test showed that total apoptosis and necrosis rate were  $8.86\% \pm 3.56\%$  and  $7.66\% \pm 1.31\%$  in the labeled and control group, respectively (Figure 3). There was no statistical differences between them ( $t = 0.71$ ,  $P > 0.05$ ).

**Table 2. T2\* ratios of infarction, PI and INP one hour after transplantation.**

Area	T2* ratio			P-value #
	Group A	Group B	Group C	
PI	$0.45 \pm 0.19$	$0.78 \pm 0.07$	$1.02 \pm 0.03$	0.001
INP	$0.46 \pm 0.23$	$0.82 \pm 0.15$	$1.02 \pm 0.04$	0.004
Infarction	$1.04 \pm 0.18$	$1.11 \pm 0.15$	$1.06 \pm 0.09$	0.777

Note: PI: peri-infarction area, INP: ipsilateral normal parenchyma. # One-way analysis of variance was performed for the three groups. Data was reported as mean  $\pm$  standard deviation.

doi:10.1371/journal.pone.0054963.t002


### Establishment of Cerebral Ischemia

After combining left proximal MCAO with two-hour left ICA blood flow occluding, moderate size of cerebral infarction, involving left basal ganglia and left temporal lobe including cortex and subcortical white matter, was observed in 10 dogs (Figure 4A). The mean volume on T2WI was  $599.45 \pm 262.50$  mm<sup>3</sup> at 24 hours and decreased to  $390.93 \pm 92.82$  mm<sup>3</sup> at one week (before transplantation). Three dogs (dog 4, 7, 12) had large MCA territory infarction before transplantation, respectively (Figure 4B). One dog (dog 13) only had small infarction involving the subcortical white matter of left temporal lobe, with the volume of  $61.14$  mm<sup>3</sup> (Figure 4C).

One week after ischemia, there were five, four and three dogs distributed to group A, B and C, respectively, according to the angiography findings before MSCs transplantation in experimental group. The infarction volume on T2WI at 24 hours and one week after embolization and the details of dogs grouping before transplantation were summarized in table 1.

### In vivo MRI Detection of Transplanted MSCs

Both T2\*WI and SWI could detect the MSCs in vivo clearly (Figure 5A–B). On T2\*WI, the grafted cells were shown more obvious signal intensity loss along with TE time increasing from 20 ms to 46 ms while the signal-to-noise ratio of images decreased at the same time. T2WI allowed a proper visualization of the



**Figure 7. Serial T2\*WI images from one hour to four weeks after cell transplantation.** Hypointensity of MSCs generally kept decreasing but was still visible within four weeks (A-F). Hypointensity of the PI area (black arrow in D-F) showed a trend of slower decline than that of the INP area. doi:10.1371/journal.pone.0054963.g007

ischemic brain tissue but it failed to detect the MSCs in all the dogs (Figure 5C).

In group A, large amounts of MSCs scattered discretely as small clusters in a wide range in the left cerebral hemisphere in four dogs one hour after IA administration (Figure 6A–B), except dog 13 which had very small infarction and was shown fewer scattered cell clusters (Figure 4F). Moreover, dog 12 with large infarction had more grafted MSCs than those with moderate size of infarction (Figure 4D–E). In group B, dogs were shown fewer amounts of MSCs in the brain by visual evaluation (Figure 6C–D). Grafted MSCs of two dogs (dog 3, 10) were mainly observed in the left basal ganglia while only small amount of cells were detected in the

INP area. In group C, no visible MSC was observed by in vivo MRI imaging (Figure 6E–F).


The corresponding T2\* ratios of infarction, the PI and INP area in transplantation groups at one hour after transplantation were shown in table 2. There was significant difference of PI T2\* ratios among the three groups and between any two groups ( $F=17.71$ ,  $P<0.01$ ). The significant differences of INP T2\* ratios were found between group A and B as well as group A and C ( $F=10.54$ ,  $P<0.01$ ).

During the series tracking in four weeks, the hypointensity of MSCs generally kept decreasing but was still visible, accompanied with T2\* ratio increasing. The hypointensity of the PI area showed a trend of slower decline than that of the INP area (Figure 7A–F).

**Table 3. Neurological evaluation before and after MSCs transplantation.**

Grouping	Before transplantation *		After transplantation			
	24 hours	1 week	1 week	2 weeks	3 weeks	4 weeks
Group A	6.20±2.17	4.60±1.52	4.00±1.83	3.50±1.29	3.00±1.15	2.50±0.58
Group B	5.75±0.96	4.50±1.29	3.33±0.58	2.67±0.58	2.33±0.58	2.33±0.58
Group C	6.67±2.89	5.33±2.52	4.00±2.00	3.67±2.08	3.33±1.53	3.00±1.00
Control group	7.50±2.12	6.50±2.12	5.50±2.12	4.50±2.12	4.50±2.12	3.50±0.71

Note: \* After completing the procedure of left MCA embolization and temporal ICA occlusion. Data was reported as mean ± standard deviation. doi:10.1371/journal.pone.0054963.t003




**Figure 8. HE staining after infarction.** (A) White matter involvement was observed after infarction ( $\times 100$ ). (B) Prominent neuronal cell loss, necrotic neurons and nuclear karyolysis were shown, accompanied with aggregated macrophages and small amount of lymphocytes (dotted box in A,  $\times 400$ ).

doi:10.1371/journal.pone.0054963.g008

Two dogs (dog 2, 12) developed new cerebral infarction on DWI within 24 hours after transplantation. The volume was  $549.68 \text{ mm}^3$  (dog 2) and  $358.08 \text{ mm}^3$  (dog 12) respectively. Both of the two dogs had high cell engraftments in the brain. The new infarction appeared around the area of cell clusters. Another dog (dog 10) was suffered from epilepsy attack after recovery from


anesthesia although the amount of grafted cells in the brain was low. It was sacrificed at 24 hours after transplantation.



**Figure 9. Histological findings at four weeks after MSCs transplantation.** (A) Microscopic view of specimen (white box in the gross image) showed clear boundary between infarction and peri-infarction area (HE staining  $\times 100$ ). (B) Neuronal cell necrosis, tissue loss and aggravated macrophages (black arrows) were observed inside of infarction (dotted box in A, HE staining  $\times 400$ ). (C) No obvious macrophage was observed in the peri-infarction area (solid line box in A, HE staining  $\times 400$ ). (D) PB staining showed congregate PB-positive cells nearby lesion boundary (PB staining  $\times 400$ ).

doi:10.1371/journal.pone.0054963.g009





**Figure 10. PB staining of the control group.** No PB-positive cell was observed in the peri-infarction area (PI). doi:10.1371/journal.pone.0054963.g010

### Neurological Scoring and Histological Findings

Twenty-four hours after the onset of ischemia, eight animals had mildly reduced consciousness and right side hemiparetic but able to walk, while three dogs (dog 4, 7, 12) had severely reduced consciousness and were unable to stand. Three dogs didn't show any motor function deficit but had mildly reduced responsiveness. The circling movements, head turning and hemianopsia were observed in 11, 8 and 8 dogs, respectively. One week later, these signs were improved. Continuous improvement was observed within four weeks after MSCs transplantation. The mean neurological scorings before and after transplantation were summarized in Table 3.

By HE staining, infarction involving the white matter was observed. Prominent neuronal cell loss and nuclear karyolysis were shown (Figure 8). PB stain-positive cells were observed mainly around the vessels at 24 hours after transplantation. Congregate PB-positive cells were observed nearby infarction boundary four weeks after transplantation (Figure 9A, C, D). Macrophages gathered inside of the infarction rather than the PI area (Figure 9B, C). Some scattered PB-positive cells were also found in brain parenchyma in dogs of group C although they were not detected by MRI in vivo. PB stain of PI area was negative in the control group (Figure 10).

### Discussion

Intra-arterially delivered MSCs was non-invasively and dynamically investigated by 3.0 T MRI after labeling with  $\text{Fe}_3\text{O}_4$ -DMSA-PLL in the present study. We showed that MRI could visualize the cells as soon as one hour after IA injection and was effective for tracking grafted MSCs within four weeks.

We choose beagle dog for IA transplantation because it is structurally similar to the human brain, has little variation in the size of the brain specimens and in the diameters of intracranial arteries despite the range of weight and technically easy for transfemoral catheterization of carotid artery [20,22]. In our previous study [20], an embolic stroke model resembling lacunar infarction was got by proximal MCA occlusion in beagle dogs. In order to enlarge the cerebral infarction, proximal MCA occlusion in combination with temporal occlusion of the ipsilateral ICA which had been reported previously was used in our current study [23,24]. Collateral supply was restricted by temporary ICA

occlusion. Cerebral blood flow in the MCA territory was further reduced into the ischemic range which gave rise to the increase of infarction volume compared with single-thrombus approach. However, due to the variability of collateral circulation in dogs, heterogeneous infarction was observed in our study.

Results of labeling MSCs in vitro with home-synthesized SPIO demonstrated its usefulness to label MSCs without cell viability compromise which was compatible with widely used ferumoxides-PLL complex [25,26]. Moreover, such labeled MSCs could be detected in vivo efficiently after IA transplantation by 3.0 T MRI. In agreement with previous studies [15,27,28], IA administration could produce a large number of MSCs in the target brain. The injected cells mainly distributed in the ipsilateral cerebral hemisphere of infarction while only few MSCs were found in the contralateral side.

However, no MSC was shown in dogs with occluded MCA at the time of transplantation. We considered two reasons for this. First, the main approach for MSCs entering brain may be the ipsilateral MCA. Thus, the flow status of that MCA before transplantation may play an important role in distribution and the amount of cells in the target tissue. Fewer MSCs detected in group B than group A could also confirm our speculation. Second, we considered that there should be a small amount of MSCs in the brain which was delivered to the host brain through other access, like anterior cerebral artery, as confirmed by PB staining in group C. However, our imaging methods may not be sensitive enough to detect small amount cells or tiny cell clusters.

In the present study, labeled MSCs were transplanted at one week after onset of ischemia. Serbian et al considered the blood-brain barrier (BBB) was continuously open for several weeks after focal cerebral ischemia [29]. Komatsu et al also found that BBB may largely break for at least two weeks after cerebral infarction and remain insufficient even four weeks [1]. Thus, we think that deficit BBB after cerebral ischemia may account for the mechanical trapping of MSCs from vessel to brain parenchyma as well as the possible cells immigration after transplantation.

The gradually fading of low signal intensity in the host brain was not only due to the cell division but also biodegradation and entry of iron into metabolic pathways [25,30]. Even so, SPIO-labeled MSCs will retain the iron nanoparticles to a sufficient degree to produce hypointensity within four weeks on SWI and  $T2^*WI$ .  $T2^*$  ratio could reflect the amount of grafted cells in the brain. But it is still difficult to directly extrapolate the exact number due to heterogeneity in the structure and composition of brain tissue and differences in MRI sequence and parameters [31].

The slower fading of hypointensity in PI area than in INP may support the concept that chemoattractive factors may be released in the site of brain lesion which gave rise to reservation and specific homing of transplanted cells to the PI area [32,33]. PB staining confirmed that there were MSCs in the PI area.

Our results also showed that a smaller infarction on the day of cell transplantation seemed to be associated with fewer cells in the brain after IA delivery while dog with large infarction had more engrafted cells. But due to the small number of dogs, we didn't perform statistical analyze. Li et al reported animal with smaller lesions (less than 10% of brain volume) at the time of transplantation have fewer grafted cells into the parenchyma [15]. Our finding was consistent with theirs. The possible reason may be due to the following two reasons: First, small infarction was associated with mild BBB deficit which led to fewer mechanically trapped MSCs. Second, recruitment forces such as chemoattractive factors released by small infarction may not enough to attract lots of MSCs whereas these chemoattractant forces and possible other

factors like brain edema caused excessive cell accumulation in the large infarction group [27].

Although IA administration showed early arrival and more transplanted MSCs in the target brain, there were still some risks. New infarction developed within 24 hours in two dogs. The likely cause was microembolism caused by transplanted MSCs which in turn led to local impeded cerebral blood flow [27,34]. Besides, exogenous labeled-MSCs may cause epilepsy after transplantation which should be paid more attention in the future research.

There were some limitations of our study. First, the numbers of dogs with large and small cerebral infarction in group A were small, limiting the investigation of the relationship between initial ischemic volume and the amount of grafted cells in the brain. Second, the amount of grafted cells in the brain was analyzed by visual evaluation and  $T2^*$  ratio.  $T2^*$  value was measured by manually drawn ROIs which may lead to some bias. Third, MRI sensitivity of detecting labeled MSCs was affected by many factors, such as MRI protocol and software. Our imaging methods may not be sensitive enough for small amount of engrafted cells or tiny cell clusters. Finally, more animals were needed to confirm the reproducibility of our study.

## References

- Komatsu K, Honmou O, Suzuki J, Houkin K, Hamada H, et al. (2010) Therapeutic time window of mesenchymal stem cells derived from bone marrow after cerebral ischemia. *Brain Res* 1334: 84–92.
- Guzman R, Choi R, Gera A, De Los Angeles A, Andres RH, et al. (2008) Intravascular cell replacement therapy for stroke. *Neurosurg Focus* 24: E15.
- Bliss T, Guzman R, Daadi M, Steinberg GK (2007) Cell transplantation therapy for stroke. *Stroke* 38: 817–826.
- Chen J, Zhang ZG, Li Y, Wang L, Xu YX, et al. (2003) Intravenous administration of human bone marrow stromal cells induces angiogenesis in the ischemic boundary zone after stroke in rats. *Circ Res* 92: 692–699.
- Li L, Jiang Q, Zhang L, Ding G, Wang L, et al. (2006) Ischemic cerebral tissue response to subventricular zone cell transplantation measured by iterative self-organizing data analysis technique algorithm. *J Cereb Blood Flow Metab* 26: 1366–1377.
- Toyama K, Honmou O, Harada K, Suzuki J, Houkin K, et al. (2009) Therapeutic benefits of angiogenic gene-modified human mesenchymal stem cells after cerebral ischemia. *Exp Neurol* 216: 47–55.
- Shen LH, Li Y, Chen J, Zhang J, Vanguri P, et al. (2006) Intracarotid transplantation of bone marrow stromal cells increases axon-myelin remodeling after stroke. *Neuroscience* 137: 393–399.
- Chopp M, Li Y (2002) Treatment of neural injury with marrow stromal cells. *Lancet Neurol* 1: 92–100.
- Chen J, Li Y, Wang L, Zhang Z, Lu D, et al. (2001) Therapeutic benefit of intravenous administration of bone marrow stromal cells after cerebral ischemia in rats. *Stroke* 32: 1005–1011.
- Rice HE, Hsu EW, Sheng H, Evenson DA, Freemerman AJ, et al. (2007) Superparamagnetic iron oxide labeling and transplantation of adipose-derived stem cells in middle cerebral artery occlusion-injured mice. *AJR Am J Roentgenol* 188: 1101–1108.
- Lu D, Li Y, Wang L, Chen J, Mahmood A, et al. (2001) Intraarterial administration of marrow stromal cells in a rat model of traumatic brain injury. *J Neurotrauma* 18: 813–819.
- Criado FJ, Lingelbach JM, Ledesma DF, Lucas PR (2002) Carotid artery stenting in a vascular surgery practice. *J Vasc Surg* 35: 430–434.
- Rink C, Christoforidis G, Abduljalil A, Kontzialis M, Bergdall V, et al. (2008) Minimally invasive neuroradiologic model of preclinical transient middle cerebral artery occlusion in canines. *Proc Natl Acad Sci U S A* 105: 14100–14105.
- Weon YC, Kang SG, Chung JW, Kim YI, Park JH, et al. (2006) Technical feasibility and biocompatibility of a newly designed separating stent-graft in the normal canine aorta. *AJR Am J Roentgenol* 186: 1148–1154.
- Li L, Jiang Q, Ding G, Zhang L, Zhang ZG, et al. (2010) Effects of administration route on migration and distribution of neural progenitor cells transplanted into rats with focal cerebral ischemia, an MRI study. *J Cereb Blood Flow Metab* 30: 653–662.
- Song M, Kim Y, Ryu S, Song I, Kim SU, et al. (2009) MRI tracking of intravenously transplanted human neural stem cells in rat focal ischemia model. *Neurosci Res* 64: 235–239.
- Zhou B, Shan H, Li D, Jiang ZB, Qian JS, et al. (2010) MR tracking of magnetically labeled mesenchymal stem cells in rats with liver fibrosis. *Magn Reson Imaging* 28: 394–399.
- Alhadlaq A, Mao JJ (2004) Mesenchymal stem cells: isolation and therapeutics. *Stem Cells Dev* 13: 436–448.
- Ju S, Teng G, Zhang Y, Ma M, Chen F, et al. (2006) In vitro labeling and MRI of mesenchymal stem cells from human umbilical cord blood. *Magn Reson Imaging* 24: 611–617.
- Liu S, Hu WX, Zu QQ, Lu SS, Xu XQ, et al. (2012) A novel embolic stroke model resembling lacunar infarction following proximal middle cerebral artery occlusion in beagle dogs. *J Neurosci Methods* 209: 90–96.
- Kang BT, Jang DP, Gu SH, Lee JH, Jung DI, et al. (2009) MRI features in a canine model of ischemic stroke: correlation between lesion volume and neurobehavioral status during the subacute stage. *Comp Med* 59: 459–464.
- Kang BT, Lee JH, Jung DI, Park C, Gu SH, et al. (2007) Canine model of ischemic stroke with permanent middle cerebral artery occlusion: clinical and histopathological findings. *J Vet Sci* 8: 369–376.
- Chen ST, Hsu CY, Hogan EL, Maricq H, Balentine JD (1986) A model of focal ischemic stroke in the rat: reproducible extensive cortical infarction. *Stroke* 17: 738–743.
- Traystman RJ (2003) Animal models of focal and global cerebral ischemia. *ILAR J* 44: 85–95.
- Arbab AS, Bashaw LA, Miller BR, Jordan EK, Lewis BK, et al. (2003) Characterization of biophysical and metabolic properties of cells labeled with superparamagnetic iron oxide nanoparticles and transfection agent for cellular MR imaging. *Radiology* 229: 838–846.
- Yocum GT, Wilson LB, Ashari P, Jordan EK, Frank JA, et al. (2005) Effect of human stem cells labeled with ferumoxides-poly-L-lysine on hematologic and biochemical measurements in rats. *Radiology* 235: 547–552.
- Walczak P, Zhang J, Gilad AA, Kedziorek DA, Ruiz-Cabello J, et al. (2008) Dual-modality monitoring of targeted intraarterial delivery of mesenchymal stem cells after transient ischemia. *Stroke* 39: 1569–1574.
- Li Y, Chen J, Wang L, Lu M, Chopp M (2001) Treatment of stroke in rat with intracarotid administration of marrow stromal cells. *Neurology* 56: 1666–1672.
- Strbian D, Durukan A, Pitkonen M, Marinkovic I, Tatlisumak E, et al. (2008) The blood-brain barrier is continuously open for several weeks following transient focal cerebral ischemia. *Neuroscience* 153: 175–181.
- Pawelczyk E, Arbab AS, Pandit S, Hu E, Frank JA (2006) Expression of transferrin receptor and ferritin following ferumoxides-protamine sulfate labeling of cells: implications for cellular magnetic resonance imaging. *NMR Biomed* 19: 581–592.
- Thu MS, Najbauer J, Kendall SE, Harutyunyan I, Sangalang N, et al. (2009) Iron labeling and pre-clinical MRI visualization of therapeutic human neural stem cells in a murine glioma model. *PLoS One* 4: e7218.
- Ben-Hur T, Einstein O, Mizrahi-Kol R, Ben-Menachem O, Reinhartz E, et al. (2003) Transplanted multipotential neural precursor cells migrate into the inflamed white matter in response to experimental autoimmune encephalomyelitis. *Glia* 41: 73–80.
- Imitola J, Raddassi K, Park KI, Mueller FJ, Nieto M, et al. (2004) Directed migration of neural stem cells to sites of CNS injury by the stromal cell-derived factor 1 $\alpha$ /CXCL12 chemokine receptor 4 pathway. *Proc Natl Acad Sci U S A* 101: 18117–18122.
- Parr AM, Tator CH, Keating A (2007) Bone marrow-derived mesenchymal stromal cells for the repair of central nervous system injury. *Bone Marrow Transplant* 40: 609–619.

In conclusion, it is feasible to transplant MSCs through IA route for cerebral infarction in a canine model. Successful IA administration showed diffuse distribution pattern, and large amounts of transplanted MSCs in the target brain. Both the flow status of ipsilateral MCA and infarction volume before transplantation may play an important role in the amount of grafted cells in the brain. In vivo MR imaging is useful to track SPIO-labeled MSCs for at least four weeks. However, more attention should be paid on the safety of IA approach considering the high ratio of adverse consequences.

## Acknowledgments

We thank Li-xin Jin at Siemens institute of Shanghai and De-feng Wang at the Chinese University of Hong Kong, China, for their technical supports.

## Author Contributions

Conceived and designed the experiments: S-sL SL H-bS. Performed the experiments: S-sL SL Q-qZ J-wW X-qX JY. Analyzed the data: S-sL Q-qZ X-qX. Contributed reagents/materials/analysis tools: YZ. Wrote the paper: S-sL SL.