

RESEARCH ARTICLE

Cardiovascular benefits of *Eruca sativa* mill. Defatted seed meal extract: Potential role of hydrogen sulfide

Lara Testai^{1,2,3}  | Eleonora Pagnotta⁴ | Eugenia Piragine¹ | Lorenzo Flori¹ |
Valentina Citi¹ | Alma Martelli^{1,2,3}  | Lorenzo Di Cesare Mannelli⁵  |
Carla Ghelardini⁵ | Roberto Matteo⁴ | Serafino Suriano⁶ | Antonio Troccoli⁶ |
Nicola Pecchioni⁶ | Vincenzo Calderone^{1,2,3} 

¹Department of Pharmacy, University of Pisa, Pisa, Italy

²Interdepartmental Research Center Nutrafood “Nutraceuticals and Food for Health”, University of Pisa, Pisa, Italy

³Interdepartmental Research Centre of Ageing Biology and Pathology, University of Pisa, Pisa, Italy

⁴CREA-Council for Agricultural Research and Economics, Research Centre for Cereal and Industrial Crops, Bologna, Italy

⁵Department of Neuroscience, Psychology, Drug Research and Child Health–Neurofarba– Pharmacology and Toxicology Section, University of Florence, Florence, Italy

⁶CREA-Council for Agricultural Research and Economics, Research Centre for Cereal and Industrial Crops, Foggia, Italy

Correspondence

Lara Testai, Department of Pharmacy,
University of Pisa, 56120 Pisa, Italy.
Email: lara.testai@unipi.it

Funding information

“Contratto per l’affidamento del servizio di consulenza farmacologica per la caratterizzazione cinetico-chimica del rilascio di acido solfidrico (H₂S) da farmaci e molecole di origine naturale in soluzione, in vitro e in vivo e per la valutazione degli effetti di una modulazione della concentrazione di H₂S in modelli in vitro e in vivo di patologie cardiovascolari e tumorali” funded by CREA Council for Agricultural Research and Economics, Italy, Grant/Award Number: Grant decree COMETA prot. n. 1741 of 05/07/2018; Hydrogen Sulfide in the Vascular Inflammation-Aging: role, therapeutic Opportunities and development of novel pharmacological tools for age-related cardiovascular diseases (SVAgO), Grant/Award Number: PRIN 2017_2017XP72RF; Botanicals in the Treatment of Ageing-Related Cardiovascular Diseases, Grant/Award Number: PRA2017_26; MIUR-DAAD Joint Mobility Program (Prog. n. 32678, entitled: “RBC and EC NOS3 in exercise dependent cardioprotection: focus on mitochondria”), Grant/Award Number: MIUR-

Abstract

Eruca sativa Mill. is an edible plant belonging to the *Brassicaceae* botanical family with a long story as a medicinal material, mainly linked to the presence of glucoerucin. One of the main products of this glucosinolate is erucin, a biologically active isothiocyanate recently recognized as a hydrogen sulfide (H₂S) donor. In this work, an *Eruca sativa* extract has been obtained from a defatted seed meal (DSM), achieving a powder rich in thiofunctionalized glucosinolates, glucoerucin, and glucoraphanin, accounting for 95% and 5% of the total glucosinolate content (17% on a dry weight basis), associated with 13 identified phenolic acids and flavonoids accounting for 2.5%. In a cell-free model, *Eruca sativa* DSM extract slowly released H₂S. Moreover, this extract promoted significant hypotensive effects in hypertensive rats, and evoked dose-dependent cardioprotection in in vivo model of acute myocardial infarct, obtained through a reversible coronary occlusion. This latter effect was sensitive to blockers of mitochondrial KATP and Kv7.4 potassium channels, suggesting a potential role of these mitochondrial channels in the protective effects of *Eruca sativa* DSM extract. Accordingly, *Eruca sativa* DSM extract reduced calcium uptake and apoptotic cell death in isolated cardiac mitochondria. Taken together, these results demonstrate that *Eruca sativa* DSM extract is endowed with an interesting nutraceutical profile on the cardiovascular system due to, at least in part, its H₂S releasing properties. These results pave the way for future investigations on active metabolites.

Lara Testai and Eleonora Pagnotta equally contributed to the manuscript.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Phytotherapy Research* published by John Wiley & Sons Ltd.

DAAD Joint Mobility Program -Prog. n. 32678;
PON "Ricerca e Innovazione"
2014-2020-Azione II, by the COMETA
research project "Colture autoctone
mediterranee e loro valorizzazione con
tecnologie avanzate di chimica verde" (Native
Mediterranean crops and their enhancement
with advanced green chemistry technologies),
Grant/Award Number: ARS01_00606

KEYWORDS

cardiovascular system, *Eruca sativa* mill., erucin, glucoerucin, hydrogen sulfide

1 | INTRODUCTION

Eruca sativa Mill, a synonym of *E. vesicaria* subsp. *sativa* (Miller) Thell, is the only taxon of *Eruca vesicaria* (L.) Cav. to be cultivated worldwide. It is a diploid annual or biennial herb belonging to *Brassicaceae* family and in Europe, it is known as salad rocket (Great Britain), salatrauke (Germany), oruga or eruca (Spain), roquette (France), rucola (Italy), and while in the USA, it is frequently reported as arugula. *E. sativa* sel. Nemat (Italy) is a selection that produces a bigger seed size than *E. vesicaria*, a difference that permits to classify Nemat as an industrial oleaginous crop ready to be cultivated by agricultural mechanization, so that it can be considered for industrial biorefinery strategies. It spontaneously grows in the Mediterranean basin with different accessions and is cultivated as leaf vegetable or for oilseeds in Northern America, Europe, and Asia (Mirzabe, Hajiahmad, & Asadollahzadeh, 2021).

In Western countries, *E. sativa*'s leaves are used as a salad or garnish and they are sold in both processed and fresh markets, where it has become very popular as "baby leaf" greens over the last two decades (Bell & Wagstaff, 2019; Di Gioia, Avato, Serio, & Argentieri, 2018). In India and Pakistan, special ecotypes of *E. sativa* are widely used as oilseed, forage, and fodder crops (Bhandari & Chandel, 1996).

As regards the beneficial effects in humans, *E. sativa* is known since Roman times for its aphrodisiac properties and other biological activities, including diuretic, stomachic, carminative, and to alleviate abdominal discomfort and improve digestion (Khan & Khan, 2014). Furthermore, this plant possesses anti-secretory, cytoprotective, and anti-ulcer activity against experimentally induced gastric lesions (Khan & Khan, 2014). Fuentes and colleagues highlighted anti-platelet and anti-thrombotic activity mediated through the inhibition of NF- κ B pathway and then through a reduced release of inflammatory mediators (Fuentes, Alarcón, Fuentes, Carrasco, & Palomo, 2014).

All the reported health benefits provided by *E. sativa* may be ascribed to the presence of different phytochemicals in its tissues. 4-(methylthio)-butyl-GSL, glucoerucin, is the main GSL present in *E. sativa* seeds, together with its oxidized analogue 4-methylsulfinylbutyl-GSL, glucoraphanin (Franco et al., 2016). *E. sativa* seeds also contain polyphenols (Sarwar, Kaur, Jabbar, Javed, & Athar, 2007), flavonoids, and carotenoids (Barillari et al., 2005).

Due to the presence of the high amount of phenolic compounds and GSLs, extract from *E. sativa* seeds is considered a rich source of antioxidants and anti-inflammatory agents (Villatoro-Pulido et al., 2012).

In particular, *E. sativa* extract inhibits COX2, TLR4, NLRP3 inflammasome pathways, and pro-inflammatory cytokines and the activation of apoptotic mechanisms in murine motor neurons exposed to the culture medium of lipopolysaccharide-stimulated murine macrophage cell lines (Gugliandolo et al., 2018).

Recently, *E. sativa* DSM has been demonstrated to be effective in the management of neuropathic pain, in an experimental model of diabetes. Glucoerucin has been identified as hydrogen sulfide (H_2S) donor moiety and responsible, at least in part, for these beneficial effects (Lucarini et al., 2019).

In humans, H_2S has been recognized as a gasotransmitter deeply involved in cardiovascular homeostasis and it is implicated also in regulating metabolism, cognitive, respiratory, and reproductive functions (Martelli et al., 2012); whereas in plants, it acts as an antioxidant, improves the performance of the organism under stress conditions and plays significant roles in several physiological processes, including seed germination, morphogenesis, photosynthesis, and flowering (Hancock & Whiteman, 2014). Notable, erucin, likewise glucoerucin, has been demonstrated to promote a slow release of H_2S in cell-free and cell-based models (Citi et al., 2014; Martelli et al., 2020).

Recently, *E. sativa* DSM showed hypoglycaemic and anti-obesity effects in a rodent model of metabolic disorder (Piragine et al., 2021). Moreover, aerial parts *E. sativa* extract produced endothelium-mediated vasorelaxation in normotensive conditions, probably due to activation of muscarinic receptors; on the other hand, in hypertensive conditions, an endothelium-independent vasorelaxation was observed (Salma, Khan, & Shah, 2018).

Anyway, the effects of *Brassicaceae* deriving compounds on the cardiovascular system still need to be fully elucidated; therefore, the main goal of this work was the production and characterization of an *E. sativa* DSM extract, and the evaluation of its implications on the cardiovascular system.

2 | MATERIALS AND METHODS

2.1 | Plant material and preparation of *Eruca sativa* DSM extract

Eruca sativa Mill. sel. NEMAT (Lazzeri, Errani, Leoni, & Venturi, 2004) was cultivated during the 2018 season, within a plot with a size of around 1,100 m². The cultivation was carried out at an experimental farm located at Budrio (Bologna) in the Po Valley area (Emilia Romagna region, 44°32'00" N; 11°29'33" E, altitude 28 m a.s.l.). The field

was fertilized by a tertiary fertilizer NPK (11–11–18) at a dose of 700 kg ha⁻¹. *E. sativa* seeds were harvested using a plot combine, cleaned with a fixed Small-Scale Threshing Equipment (Cicoria Srl, Italy), and finally air-dried at room temperature. *E. sativa* seeds were defatted by a small continuous seed crusher machine (Bracco Company model Elle.Gi type 0.90, Italy) as previously described (Lucarini et al., 2019) and defatted seed meal (DSM) was characterized for moisture; residual oil, using Soxhlet extraction method with hexane as a solvent; GSL, and phenolic acids and flavonoids according to section 2. The characterized *E. sativa* DSM was ground to 500 µm grain size by the Ultra Centrifugal Mill ZM200 (Retsch GMBH, Germany) and 500 g of homogeneous DSM was extracted in 6 L 70% ethanol for 10 min at 80°C with an Ultra-Turrax homogenizer (IKA-Werk, Staufen, Germany). The extract was cooled in the ice bath and then centrifuged at 25,900g for 30 min at 10°C. The supernatant was reduced about three-fold at 40°C, stored at -20°C for 48 hr, centrifuged, and finally lyophilized (DLAB 500, Italian Vacuum Technology, Italy).

2.2 | HPLC analysis of Desulfo-Glucosinolates in *Eruca sativa* DSM extract

For GSL analysis (EN ISO 9167:2019; Franco et al., 2016), 40–50 mg fine-powdered lyophilized *E. sativa* DSM extract was diluted in 4 ml pure water and 1 ml of the diluted extract and filtered extract (cellulose acetate 0.45 µm syringe filter) was loaded onto a mini-column filled with DEAE Sephadex A-25 anion-exchange resin (GE Healthcare, Germany), conditioned with 25 mM sodium acetate buffer (pH 5.6). 1 ml of the diluted extract was loaded onto a mini-column filled with DEAE Sephadex A-25 anion-exchange resin (GE Healthcare, Germany), conditioned with 25 mM sodium acetate buffer (pH 5.6). After washing with the same buffer (3 ml), sulfatase (200 µL, 0.35 U ml⁻¹), purified from *Helix pomatia* (Sigma Aldrich, USA) according to ISO 9167-1 (EN ISO 9167:2019), was loaded onto the mini-column, which was then left overnight at 30°C. The desulfated GSLs were finally eluted with four 1 ml portions of ultrapure water and water was allowed to drain after each addition. The desulfated GSLs were detected by HPLC-UV monitoring their absorbance at 229 nm and identified with respect to their retention times and UV spectra according to our library (Franco et al., 2016). Their amounts were estimated using isolated sinigrin as an internal standard. Sinigrin was isolated as previously reported at HPLC purity >99% and stored at -20°C until required; the response factor for desulfated glucoerucin and glucoraphanin was according to Wathelet et al. (Wathelet et al., 2004). Each dilution of the powdered extract and analysis was performed in triplicate. GSL storage stability of the *E. sativa* DSM extract was conducted for a sample of extract kept at -20°C for about 1 year storage. GSL profile was analyzed at about 1, 2, 3, and 12 months from the date of lyophilisation, as above described.

2.3 | HPLC analysis of phenolic acids and flavonoids in *E. sativa* DSM extract

Phenolic acids and flavonoids were analyzed by HPLC (1,290 UHPLC series Rapid Resolution; Agilent Technologies, USA), with a diode array detector. 500 mg of *E. sativa* DSM was extracted twice with 5 ml methanol/ 1 N HCl (85:15) solution by 20 min sonication, with nitrogen gas introduced into the tubes, and centrifugation (4,500 rpm, 10 min, 4°C). Supernatants were recovered and stored at -20°C to precipitate the large molecules. For soluble conjugated phenolic compounds, 5 ml of the methanolic extracts was digested with 6 ml 4 N NaOH in a sonication bath for 10 min. The solutions were then brought to pH 2 with 6 M HCl and extracted twice with 10 ml of diethyl ether/ethyl acetate (1:1 v/v). The extracts were clarified for centrifugation, evaporated, resuspended in 2 ml of mobile phase (phosphoric acid 10⁻³ M / acetonitrile 95:5 v/v), and filtered with a 0.22 µm PTFE membrane before HPLC analysis. The insoluble bound phenolics were obtained from the solid residue of methanolic extracts and were further digested in 12 ml 4 N NaOH in sonication bath for 30 min, vortexing every 5 min. After centrifugation, the supernatant was brought to pH 2 with 6 N HCl and extracted twice with 10 ml of diethyl ether/ethyl ether (1:1 v/v). After separation, the organic phase was clarified for centrifugation, evaporated, resuspended in 2 ml of mobile phase (phosphoric acid 10⁻³ M / acetonitrile 95:5 v/v), and analyzed by HPLC. The final soluble and insoluble phenolic acid extracts were also analyzed for flavonoids. 500 mg of *E. sativa* extract was completely dissolved in methanol/1 N HCl (85:15) solution for soluble phenolic acids and flavonoids extraction. Phenolic acids and flavonoids chromatographic separation was carried out with a 50 mm × 2.1 mm × 1.8 µm column (Eclipse Plus-C18 RRHD; Agilent). The temperature of the column oven was set at 40°C. The flow rate of the mobile phase was 0.3 ml min⁻¹, and the injection volume was 1 µl. The mobile phase was 10⁻³ M phosphoric acid (solution A), acetonitrile (solution B). The gradient, in terms of eluent B was: 0–2 min, 5% B (isocratic); from 2 to 4.5 min at 10% B; at 5.5 min, 15% B; at 9 min, 35% B; at 11 min, 55% B; from 12 min to 13 min at 70% B (isocratic); after each analysis, the initial mobile phase conditions were set, and the system was allowed to equilibrate for 3 min. Detection wavelengths were 280 nm and 320 nm. For peak quantification, calibration curves of the following standards (Sigma-Aldrich, Saint Louis, MO, USA) were constructed: gallic acid, protocatechuic acid, *p*-hydroxybenzoic acid, vanillic acid, syringic acid, and vanillin (between 5 and 40 mg L⁻¹); caffeic acid, *p*-coumaric acid, sinapic acid, *trans*-cinnamic acid, luteolin, vitexin, apigenin, naringenin, rutin, and quercetin-3 glucoside (between 2.5 and 20 mg L⁻¹). The correlation coefficient for each analyte was in the range (0.97–0.99) (see Figure S1 supplementary materials). Sample compounds were identified based on the retention times and confirmed by their spectra in comparison with standards and were quantified by comparing their peak areas with those of standard curves.

2.4 | Amperometric approach

The H₂S-generating property of *E. sativa* DSM extract has been evaluated by an amperometric approach, through an Apollo-4,000 Free Radical Analyzer (WPI) detector and H₂S-selective mini-electrodes at room temperature as reported previously (Testai et al., 2016). Briefly, a “PBS buffer 10×” was prepared (NaH₂PO₄·H₂O 1.28 g, Na₂HPO₄·12H₂O 5.97 g, and NaCl 43.88 g in 500 ml of H₂O) and stocked at 4°C. Immediately, before the experiments, the “PBS buffer 10×” was diluted in distilled water (1:10) to obtain the assay buffer (AB); pH was adjusted to 7.4. The H₂S-selective minielectrode was equilibrated in 10 ml of the AB, until the recovery of a stable baseline. Then, 100 µl of a dimethyl sulfoxide (DMSO) solution of *E. sativa* DSM extract was added at the final concentration of 1 mg ml⁻¹. The generation of H₂S was observed for 30 min. In order to explore the contribution of thiol groups, naturally present in the biological environment, in the release of H₂S, 4 mM L-cysteine was added, before the extract application.

2.4.1 | Data analysis

The correct relationship between the amperometric currents (recorded in pA) and the corresponding concentrations of H₂S was determined by opportune calibration curves with applying increasing concentrations of NaHS (1, 3, 5, and 10 µM) at pH 4.0. The lowest limit of reliable quantitative determination was 0.3 µM. The vertical bars represent the standard error ($n = 9$).

2.5 | In vivo systolic blood pressure values recordings

All the procedures involving animals were carried out following the guidelines of the European Community Council Directive and in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki, EU Directive 2010/63/EU for animal experiments). The experiments were authorized by the Ethical Committee of the University of Pisa and by the Italian Ministry of Health (authorization number 751/2018-PR).

The effects of *E. sativa* DSM extract and its vehicle (DMSO 1 ml kg⁻¹) on blood pressure were tested on 12-week-old male normotensive, spontaneously hypertensive rats (SHR) and rats in which the hypertension was induced by the administration of a single i.p. injection of phenylephrine (PE) 150 µg kg⁻¹ (300–350 g). Briefly, all the rats were anaesthetized with sodium thiopental 60 mg kg⁻¹ i.p. and after the administration of the anesthetic drug, they were kept on a heated platform (about 30°C) for 20 min to induce slight vasodilation of the tail artery, in order to allow an easier recording of the basal systolic blood pressure by the “tail-cuff” method by a BP recorder (BP-2000 Blood Pressure Analysis System, Series II, Visitech System, Apex NC, USA). The basal level of systolic blood pressure (Psys) was recorded for 20 min, at 5 min intervals. Then, for rats in which

hypertension was pharmacologically induced, and after the PE administration, the rise of systolic blood pressure values was monitored for 20 min every 5 min and, at the reaching of stable levels of hypertension, the *E. sativa* DSM extract was administered as following. Briefly, *E. sativa* DSM extracts 10 mg kg⁻¹, 100 mg kg⁻¹, or the corresponding vehicle, were administered i.p. to different groups (normotensive, SHR, PE-hypertensive rats). Starting from the administration of the tested compounds, the Psys values were recorded, for 2 hours after 10, 30, 60, 90, and 120 min. Rats were maintained under anesthesia for all experiments, in order to avoid interferences correlated to the stress of experimental procedures.

2.5.1 | Data analysis

Basal Psys was expressed as a mean of the four measurements carried out in each rat before the administration of tested compounds. Change in systolic blood pressure, recorded after ES extract or vehicle administration, was expressed as a percentage of the basal Psys and also calculated as the mean value of the five recordings (at 10, 30, 60, 90, and 120 min) carried out after the drug administration. Blood pressure measurements were carried out in 6 animals/group ($n = 6$ for each treatment both in normotensive and SHR groups).

All the experimental data were analyzed by a computer fitting procedure (GraphPad Prism 6.0, La Jolla, CA, USA) and expressed as mean ± SEM of at least six independent experiments. ANOVA and Bonferroni's post-test were selected as statistical analyses, and $p < .05$ was considered representative of significant statistical differences.

2.6 | In vivo acute myocardial infarct

12-week-old male Wistar rats (300–320 g) were used accordingly with DL26/2014 (protocol number 909/2016-PR, 04/10/2016). Two hours before the experimental procedures, animals received an i.p. injection (about 0.35 mL) of *E. sativa* DSM extract (10, 50 or 100 mg kg⁻¹) or vehicle (DMSO).

Then, rats were anesthetized with sodium thiopental (70 mg kg⁻¹, i.p.) and the experimental protocol for coronary occlusion–reperfusion was performed as previously described (Testai et al., 2016). The acute infarct protocol consisted of 30 min occlusion/120 min reperfusion; successful occlusion was confirmed by observing regional cyanosis downstream of the ligation, and by ST elevation in the ECG recording. In a group of animals, the thread was inserted without ligation (sham). In another experimental set, the potassium channel blockers, 5-decanoic acid (5-HD) 10 mg kg⁻¹ or XE991 2 mg kg⁻¹, selective inhibitors of mitoKATP and mitoKv7, respectively, were i.p. administered 20 min before the administration of *E. sativa* DSM extract (100 mg kg⁻¹). A group of vehicle-pretreated animals was submitted to an ischemic preconditioning (IPC) procedure, achieved by two cycles of 5 min occlusion/10 min reperfusion, followed by 30 min coronary occlusion and 120 min

reperfusion, and it was used, according to literature, as cardioprotective protocol.

At the end of reperfusion, rats were euthanized by an overdose of sodium thiopental, then hearts were quickly excised, mounted on a Langendorff apparatus (Radnoti, Monrovia, CA, USA) and perfused for 10 min with Krebs solution at 37°C to wash out the coronary blood vessels. Then, left ventricular tissue was dried, frozen at -20°C, and cut into 4–5 transverse slices from the apex to the base of equal thickness (about 2 mm). The slices were then incubated in a TTC solution in a phosphate buffer (pH 7.4) at 37°C for 20 min. TTC reacts with NADH in the presence of dehydrogenase enzymes, to form a formazan derivative, which stain the viable cells with intense red color. Then, the slices were fixed overnight in 10% formaldehyde and finally photographed. In the viable area, red-stained viable tissue was easily distinguished from the white-unstained necrotic tissue.

2.6.1 | Data analysis

The ischemic area (A_i) was planimetrically evaluated using an image analyzer program (The GIMP 2). The infarct size was calculated as a percentage of the whole left ventricle area (A_{LV}).

All the values were expressed as a mean \pm SEM for six different experiments. Data were statistically analyzed by ANOVA and Bonferroni's post-test. P -values lower than 0.05 were considered as indicative of significant differences.

2.7 | Calcium-uptake evaluation on isolated cardiac mitochondria

2.7.1 | Isolation procedure

Rat cardiac mitochondria were isolated by differential centrifugation in accordance with the authorization registered to the Italian Minister of Health (protocol number 45,972, September 21, 2016) and as previously described (Testai et al., 2016), with minor modifications. Briefly, 12 week-old male Wistar rats (300–320 g) were sacrificed by sodium thiopental overdose, hearts were immediately removed and placed in an ice-cold isolation buffer (IB1, composition mM: sucrose 250, Tris 5, EGTA 1, pH 7.4 adjusted with HCl). The atria were removed, and the ventricular tissue was finely minced with surgical scissors (about 2 mm³ pieces) and homogenized by using an Ultra-Turrax homogenizer (20 mL of isolation buffer per heart, IKA1-Werke GmbH & Co., Staufen, Germany).

Three homogenization cycles (each of 20 s) were performed on ice, and then the suspension was centrifuged at 1075 g for 3 min at 4°C (EuroClone, Speed Master 14 R centrifuge, Milano, Italy). The resulting supernatant was centrifuged at 11,950 g for 10 min at 4°C. The pellet containing the mitochondrial fraction was further re-suspended in the isolation buffer (without EGTA, IB2) and centrifuged at 11,950 g for 10 min at 4°C; this step was repeated once more. The final mitochondrial pellet was re-suspended in a minimal volume of

400 ml of the IB2 and stored on ice throughout the experiments, which were performed within 2 h. Mitochondrial protein concentrations were determined using the usual Bradford reaction.

2.7.2 | Calcium-uptake measurement

Mitochondrial calcium-uptake was measured by potentiometric technique, as previously described (Testai et al., 2016). In particular, the changes of the calcium concentration in the medium (i.e., extra-mitochondrial calcium) were continuously measured with a calcium-selective mini-electrode, coupled with a reference electrode (WPI, FL, USA), using a data acquisition software (Biopac Systems Inc., Goleta, CA, USA). In order to correlate the potentiometric measurements (in mV) with the corresponding concentrations of calcium ions in the solution, calibration curves were generated before each experiment, by using known concentrations of CaCl₂. Mitochondria (1 mg protein ml⁻¹) were added, under gentle stirring, to the IM plus CaCl₂ 100 μM in the presence of vehicle (DMSO 1%) or *E. sativa* DSM extract (10, 30 and 100 μmol L⁻¹ of total GSLs). After the addition of mitochondria, the maximal decrease of the calcium concentration in the medium, related to its accumulation in the mitochondrial matrix, was measured.

2.7.3 | Data analysis


Mitochondrial calcium uptake was evaluated by measuring the reduction of the extra-mitochondrial CaCl₂ concentration, following the addition of mitochondria into a calcium-rich buffer. All data are expressed as a mean \pm SEM. Data were statistically analyzed by ANOVA and Bonferroni's post-test. p -values lower than .05 were considered as indicative of significant differences. Each result was obtained with mitochondria isolated from the hearts of six different animals.

3 | RESULTS

3.1 | Characterization of the *E. sativa* DSM extract

Starting from an *E. sativa* DSM characterized by 8.6% relative moisture, 16.9 \pm 0.8% residual oil, and 125 \pm 3 μmol g⁻¹ total GSLs, with 92.8% glucoerucin and the remaining glucoraphanin, 95 g of brown powder containing 400 \pm 15 μmol g⁻¹ total GSLs were obtained. Total GSLs in *E. sativa* lyophilized extract corresponded to 20 \pm 1 μmol g⁻¹ glucoraphanin and 380 \pm 12 μmol g⁻¹ glucoerucin, accounting for a 0.9% and a 16% on a weight basis, respectively. The final recovery in GSLs, referring to initial GSL content in *E. sativa* DSM, after a single ethanol extraction attested to 60%. For about 12 months (403 days) under chilled storage conditions (-20 \pm 2°C), the *E. sativa* DSM lyophilized extract seemed homogeneous without sign of color changes and exhibited only minor changes in GSL content (Figure 1).

FIGURE 1 Glucoerucin (a), glucoraphanin (b) and total glucosinolate (c)(GSL), glucoraphanin (GRA) and glucoerucin (GER) content in *E. sativa* defatted seed meal (DSM) lyophilized extract for a period of storage of about 12 months at -20°C and means \pm standard deviation (SD) ($n = 3$) were presented



The polyphenol complete characterization was reported in Table 1 and Table 2, and the total soluble polyphenol fraction identified accounted for 2.5% on a weight basis in lyophilized *E. sativa* DSM extract. Flavonoids (57.6%) were the most representative of the soluble polyphenols fraction in *E. sativa* DSM, followed by phenyl carboxylic (24.5%) and phenyl cinnamic (17.9%) acids; the insoluble fraction was almost totally represented by the phenyl cinnamic acid and aldehyde group (93.7%). In the *E. sativa* DSM extract, the group of phenyl carboxylic acids was the most representative (58.9%) of the soluble polyphenols fraction, followed

by phenyl cinnamic acids (28.9%) and flavonoids (12.2%). Gallic acid is the main component of soluble phenyl carboxylic acids and aldehydes both in *E. sativa* DSM and in *E. sativa* DSM extract, 84% and 57% respectively, while sinapic acid was the main component within the soluble and insoluble fractions of phenyl cinnamic acids, in *E. sativa* DSM and in the soluble fraction of phenyl cinnamic acids in *E. sativa* DSM extract. Regards to flavonoids group, apigenin and naringenin resulted as the most abundant in soluble and insoluble fraction of DSM, respectively, while luteolin was the most abundant component in *E. sativa* lyophilized extract.

TABLE 1 Soluble conjugated and insoluble bound phenolic acids and aldehyde ($\mu\text{g g}^{-1}$) in *E. sativa* defatted seed meal (DSM) and in *E. sativa* defatted seed meal lyophilized extract (ES). Average values \pm standard deviation ($n = 3$) are shown

| | Gallic acid | Protocatechuic acid | <i>p</i> -hydroxybenzoic acid | Vanillic acid | Syringic acid | Vanillin | Caffeic acid | <i>p</i> -coumaric acid | Sinapic acid | Trans-cinnamic acid |
|--|-----------------|---------------------|-------------------------------|----------------|---------------|-----------------|----------------|-------------------------|-----------------|---------------------|
| Soluble conjugated phenolic acids and aldehyde | | | | | | | | | | |
| DSM | 1,454 \pm 27 | 29.1 \pm 0.2 | 104 \pm 18 | 10.0 \pm 0.9 | 28 \pm 3 | 114 \pm 6 | 22.0 \pm 0.2 | 130 \pm 9 | 965 \pm 59 | 150 \pm 9 |
| ES | 8,309 \pm 426 | 69 \pm 2.4 | 189 \pm 24 | 1,102 \pm 52 | 4.3 \pm 0.6 | 4,822 \pm 122 | 422 \pm 12 | 232 \pm 12 | 6,436 \pm 286 | n.d. |
| Insoluble conjugated phenolic acids and aldehyde | | | | | | | | | | |
| DSM | 30 \pm 24 | n.d. | n.d. | n.d. | 3 \pm 2 | 2.8 \pm 0.2 | 11 \pm 5 | 66 \pm 14 | 1,215 \pm 85 | 49 \pm 12 |

Abbreviation: n.d. not detected.

3.2 | *E. sativa* DSM extract releases H₂S

E. sativa DSM extract incubated at the concentration of 1 mg ml⁻¹, corresponding to 0.4 mM total GSLs, released H₂S with a slow kinetic both in the presence and in the absence of L-cysteine (L-Cys, 4 mM), reaching a stable steady state of about 1,5 μM (Figure 2).

3.3 | *E. sativa* DSM extract shows in vivo anti-hypertensive effects

E. sativa DSM extract administrated in vivo to normotensive rats did not modify the levels of systolic blood pressure; indeed, only a minimal and not significant reduction was observed at the highest dose (100 mg kg⁻¹, Figure 3a). In agreement with literature (Martelli, Piragine, et al., 2020; Talmor et al., 1998), both SHR and PE-treated rats showed Psys values markedly higher if compared with normotensive ones (174 \pm 5 mm Hg and 152 \pm 2 mm Hg, respectively). Interestingly, in such hypertensive conditions, *E. sativa* DSM extract was able to significantly reduce Psys values; indeed, at the end of recording, a clear decrease of Psys was observed in both types of hypertension ($-28 \pm 3\%$ in PE-treated rats and $-18 \pm 4\%$ in SHR; Figure 3b, c). The hypotensive efficacy of *E. sativa* DSM extract was found to be comparable to the isothiocyanate erucin, used as a positive control (Figure S2 in supplementary materials).

3.4 | *E. sativa* DSM extract shows in vivo anti-ischemic cardioprotection

According to literature, 30 min of coronary occlusion followed to 120 minutes of reperfusion produced a significant injury (Testai et al., 2016), indeed ischemic size was 34 \pm 1%; conversely, in the sham group, the amount of injured area was very low ($A_i/A_{LV} = 11 \pm 4\%$). An i.p. injection of *E. sativa* DSM extract (10 mg kg⁻¹) 2 hours before the experimental procedure did not reduce ischemic area ($A_i/A_{LV} = 33 \pm 7\%$), while *E. sativa* DSM extract at the doses 50 and 100 mg kg⁻¹ significantly contained the ischemic damage ($A_i/A_{LV} = 26 \pm 2\%$ and 21 \pm 3%, respectively), similarly to that was observed with IPC procedure (Figure 4a).

The protective effects observed at the highest dose of the extract were almost completely abolished by the pre-treatment with selective mitoKATP and Kv7 channel blockers, 5HD (10 mg kg⁻¹, $A_i/A_{LV} = 39 \pm 2\%$) and XE991 (2 mg kg⁻¹, $A_i/A_{LV} = 31 \pm 2\%$), respectively (Figure 4b).

3.5 | Effects of *E. sativa* DSM on mitochondrial calcium movements

It is well-known that mitochondria greedily uptake calcium ions. The addition of *E. sativa* DSM extract to cardiac mitochondria produced a concentration-dependent reduction in the accumulation of calcium (Figure 5).

TABLE 2 Soluble and insoluble flavonoids ($\mu\text{g g}^{-1}$) in *Eruca sativa* defatted seed meal (DSM) and in *E. sativa* defatted seed meal lyophilized extract (ES). Average values \pm standard deviations ($n = 3$) are shown

| | Luteolin | Vitexin | Apigenin | Naringenin | Rutin |
|---------------------------------|----------------|---------------|----------------|--------------|--------------|
| Soluble conjugated flavonoids | | | | | |
| DSM | 1,135 \pm 81 | 833 \pm 24 | 1838 \pm 188 | 161 \pm 4 | 105 \pm 12 |
| ES | 1,614 \pm 91 | 503 \pm 18 | n.d. | 452 \pm 30 | 426 \pm 18 |
| Insoluble conjugated flavonoids | | | | | |
| DSM | n.d. | 1.4 \pm 0.3 | 17 \pm 6 | 26 \pm 3 | 11 \pm 5 |




FIGURE 2 Amperometric recording of the time-course of H_2S -release from *E. sativa* defatted seed meal (DSM) extract in the presence and in the absence of L-cysteine (Cys 4 mM). The vertical bars indicate SEM

4 | DISCUSSION

Vegetables belonging to *Brassicaceae* family are largely consumed even for possible health benefits. They contain high levels of GSLs, which are sulfur-derivatives able to produce active metabolites, named isothiocyanates, after the hydrolysis mediated by myrosinase enzyme (Martelli, Piragine, et al., 2020). Noteworthy, even if cooking and pH at gastric level can cause denaturation of myrosinase, thioglucosidase activity of gut bacteria can hydrolyze GSLs and ensure the formation of isothiocyanates (Martelli et al., 2014).

To our knowledge, the *E. sativa* seed extract obtained and characterized in this paper is endowed with the highest content of identified molecules and the highest concentration in GSL.

Indeed, other authors reported that *E. sativa* seeds Soxhlet extracted with 95% ethanol contained up to 0.45% glucoerucin and 0.65% total flavonoids (Sarwar et al., 2007). Gugliandolo et al. obtained *E. sativa* seed extract enriched in glucoerucin up to about 4.6% on weight basis, with additional 0.2% (w/w) total flavonoids and 0.016% (w/w) ascorbic acid (Gugliandolo et al., 2018). Recently, an *E. sativa* seed extract obtained in 95% ethanol in an incubator shaker at 30°C for 2 days was reported. GSLs were not analyzed, and probably the extract contained rather high amounts of GSL hydrolysis products, but HPLC analysis of phenolic compounds revealed the presence of 10 phenolic compounds, accounting about 7% on a weight basis, of which the main ones were catechin, caffeic acid, and 3,5-dicaffeoyl quinic acid (Abdel-Rahman et al., 2015). Finally, the LC/MS analysis of

an ethanol extract of *E. sativa* whole seeds revealed the tentative identification of 39 compounds belonging to sulphur-containing compounds, flavonoids, phenolic acids and fatty acids, with no further information on quantitative composition (Abd-Elsalam et al., 2021).

Considering that the isothiocyanate moiety has been recognized as a smart H_2S -donor chemical group (Citi et al., 2014; Lucarini et al., 2018; Martelli, Piragine, et al., 2020; Pagliaro, Santolamazza, Simonelli, & Rubattu, 2015; Ritz, Wan, & Diaz-Sanchez, 2007), we hypothesized that, at least in part, the numerous beneficial effects observed with a *Brassicaceae* enriched diet may be due to the release of this gasotransmitter. In this regards, natural isothiocyanates, that is, erucin and sulforaphane (Citi et al., 2014), can release H_2S only in the presence of endogenous thiols. Surprisingly, *E. sativa* DSM extract shows an L-cysteine-independent H_2S donation. We hypothesize that probably this behavior can be due to the presence of some substances which may favor the H_2S release. Among flavonoids founded in *E. sativa* extract, rutin seemed an attractive candidate for an L-cysteine-independent release of H_2S ; it was in fact associated to an increase of H_2S level, with subsequent up-regulation of the expression of nuclear factor-E2-related factor-2 (Nrf2) and heme oxygenase-1 (HO-1) in dorsal root ganglions of diabetic rats (Tian et al., 2015).

Besides the well-known cancer-preventive properties, vegetables from *Brassicaceae* family show notable beneficial effects on cardiovascular system. Several clinical and pre-clinical evidence point out protection induced by some *Brassica* spp; in particular, sulforaphane (isothiocyanate deriving from the GSL glucoraphanin)-induced expression of detoxification enzymes and several nuclear factors, such as Nrf2, involved in reactive oxygen species elimination and xenobiotic excretion in in vitro models (Ritz et al., 2007). Moreover, rats receiving a supplementation for 14 weeks of 200 mg day⁻¹ of dried broccoli sprouts, showed an improvement of antioxidant reserves and lower blood pressure value in SHR (Wu et al., 2004). Likely, we observed that, in hypertensive conditions, *E. sativa* DSM extract has been able to significantly reduce Psys values; such an effect could be due to the main isothiocyanate erucin, hydrolyzed by glucoerucin. Indeed, Martelli and colleagues recently demonstrated that erucin possesses a H_2S -donor profile in cell-free and in cell-based models and induced clear vasodilating and anti-hypertensive effects (Martelli et al., 2021; Martelli, Piragine, et al., 2020).

In addition, our results demonstrate that *E. sativa* DSM extract possesses anti-ischemic effects, accordingly to other *Brassica* extracts, that have recognized protective effects against oxidative stress on




FIGURE 3 Time-course of changes in systolic blood pressure values (expressed in mmHg, in the left part of the figure) and mean change in Psys (expressed as a % of the basal Psys values, in the right part of the figure). (a) normotensive rats, (b) PE-induced hypertensive rats and (c) spontaneously hypertensive rats after i.p. administration of vehicle or *E. sativa* defatted seed meal (DSM) extract at 10 or 100 mg kg⁻¹. The symbol * indicates significant differences vs vehicle




FIGURE 4 Morphometric quantification of Ai/A_{LV}% observed in ventricular slices of rat hearts, after acute myocardial infarction in vivo. (a) After treatment with vehicle or *E. sativa* defatted seed meal (DSM) extract at different doses (10, 50 and 100 mg kg⁻¹). (b) After the treatment with *E. sativa* DSM extract 100 mg kg⁻¹ in the presence or in the absence of the selective mitoK blockers, mitoKATP and mitoKv7.4, 5HD (10 mg kg⁻¹) and XE991 (2 mg kg⁻¹), respectively. The symbol * indicates significant differences vs vehicle and the symbol # indicates significant differences vs *E. sativa* DSM extract (100 mg kg⁻¹)-treatment




FIGURE 5 Intra-mitochondrial uptake of calcium ions (in % vs. 100 μ M), following the administration of isolated cardiac mitochondria in the presence of vehicle or *E. sativa* defatted seed meal (DSM) extract at different concentrations. The concentration of the extract is related to glucosinolate title. The symbol * indicates a statistically significant difference vs vehicle

cardiomyocytes (Yang, 2018) and in in vivo models of myocardial infarct (Jana et al., 2017).

Moreover, the involvement of mitochondrial potassium (mitoK) channels has been hypothesized, since that specific blockers of mitoKATP and mitoKv7 channels almost completely abolished the cardioprotection. Such a profile further confirms the critical role of mitoK channels and strengthens the hypothesis of the presence of an active compound able to stimulate these ion channels; indeed, a mitoK opener is able to induce a mild depolarization of the cardiac mitochondrial membrane, responsible for a lower driving force for accumulating of calcium ions and then reduces the probability to assemblage and opening of the mitochondrial permeability transition pore (MPTP), preventing ultimately the organelle swelling and cell apoptotic death (Testai, Rapposelli, Martelli, Breschi, & Calderone, 2015).

Although the putative mechanisms of action through which H₂S may play cardioprotective effects are numerous and include the engagement of antioxidant transcription factors (such as Nrf2), anti-inflammatory cytokines and other agents, such as nitric oxide, mitoK channels are recognized as an interesting target. Of note, the cardioprotection promoted by NaHS in a model of myocardial I/R injury was significantly reversed by pharmacological blockage of mitoKATP channels (Citi et al., 2018; Sivarajah et al., 2009). The involvement of mitoKATP channels in the cardioprotective effects of H₂S was also ex vivo and in vivo demonstrated, by using a synthetic H₂S-donor isothiocyanate (Testai et al., 2016). Noteworthy, sulforaphane at micromolar concentrations protected both cultured cardiomyoblasts and adult cardiomyocytes from H₂O₂-induced oxidative stress through the modulation of Nrf2 signaling pathway (Corsac et al., 2018).

5 | CONCLUSIONS

On the basis of the phytochemical characterization of this extract, it is possible to speculate that cardiovascular benefits observed, other than GSL-derived isothiocyanates, may be due to the contribution of flavonoids and phenolics acids found in the *E. sativa* DSM extract; among these luteolin and sinapic acid are particularly relevant. Very recently, it has been demonstrated that luteolin is able to induce cardioprotective effects in vitro and in vivo through mechanisms involving the activation of Nrf2 and the inhibition of the NF- κ B pathways (Vashi & Patel, 2021). Furthermore, luteolin has also been reported to be cardioprotective via the phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K)/Akt pathway and able to increase the phosphorylation of phospholamban (PLB) and sarco/endoplasmic reticulum Ca²⁺-ATPase (SERCA2a) to protect the heart against acute myocardial ischemia-reperfusion injury (Chin, Qin, May, Ritchie, & Woodman, 2017). Moreover, sinapic acid has been recognized in vitro and in vivo as an antioxidant and cardioprotective agent against ischemia-reperfusion injury (Silambarasan et al., 2015).

Furthermore, Olson et al. highlighted that tea polyphenols may autoxidize, thus becoming pro-oxidants and favouring the oxidation of H₂S to polysulfides (H₂S_n; where $n = 2-5$) and other sulfoxides (Olson, Gao, & Straub, 2021).

Taken together with our results, this evidence suggests the possibility of synergistic mechanisms between H₂S released by GSLs and phenolic acids in *E. sativa* extracts, that may contribute to the evidenced benefits on cardiovascular system.

Although an accurate pharmacokinetic analysis will have to be executed, this work demonstrates the benefits of this *E. sativa* DSM extract on the cardiovascular system, suggesting that the daily consumption of this vegetable may be associated to beneficial effects on cardiovascular system and that H₂S may be an end-effector of such a nutraceutical profile. Future studies will be focused on the characterization of the pharmacological profile of main constituents, specially glucoerucin and its derivative erucin.

ACKNOWLEDGMENTS

This work was financially supported by the PRA2017 (grant number PRA2017_26; entitled Botanicals in the Treatment of Ageing-Related Cardiovascular Diseases), PRIN 2017 (protocol 2017XP72RF, entitled Hydrogen Sulfide in the Vascular inflamm-Aging: role, therapeutic Opportunities and development of novel pharmacological tools for age-related cardiovascular diseases (SVAgo) and by the Italian Ministry of Instruction, University and Research (MIUR), PON "Ricerca e Innovazione" 2014-2020-Azione II, by the COMETA research project "Colture autoctone mediterranee e loro valorizzazione con tecnologie avanzate di chimica verde" (Native Mediterranean crops and their enhancement with advanced green chemistry technologies) (ARS01_00606). Grant decree COMETA prot. n. 1741 of 05/07/2018, CUP B26G18000200004 - COR 545910, by the grant "Contratto per l'affidamento del servizio di consulenza farmacologica per la caratterizzazione cinetico-chimica del rilascio di acido solfidrico (H₂S) da farmaci e molecole di origine naturale in soluzione, in vitro e in vivo e per la valutazione degli effetti di una modulazione della

concentrazione di H₂S in modelli in vitro e in vivo di patologie cardiovascolari e tumorali” funded by CREA Council for Agricultural Research and Economics, Italy. MIUR-DAAD Joint Mobility Program (Prog. n. 32678, entitled: “RBC and EC NOS3 in exercise dependent cardioprotection: focus on mitochondria”).

CONFLICTS OF INTEREST

The authors declare no competing financial interest.

AUTHOR CONTRIBUTIONS

Conceptualization, V.Ca., L.T. and E.Pa.; Methodology, E.Pa., E.Pi., L.F., R.M., S.S. A.M.; Data Curation, E.Pi., L.F., A.T., A.M. and V.Ci. Writing – Original Draft Preparation, L.T., A.M., V.Ci., and E.Pa.; Writing – Review & Editing, L.D.M., N.P., C.G., V.Ca., A.T.; Supervision, V.Ca., N.P. Project Administration, V.Ca.; Funding Acquisition, L.T., V.Ca., A.M., N.P.


DATA AVAILABILITY STATEMENT

Data openly available in a public repository that issues datasets with DOIs

ORCID

Lara Testai  <https://orcid.org/0000-0003-2431-6248>

Alma Martelli  <https://orcid.org/0000-0002-2090-7107>

Lorenzo Di Cesare Mannelli  <https://orcid.org/0000-0001-8374-4432>

Vincenzo Calderone  <https://orcid.org/0000-0002-1441-5421>

REFERENCES

- Abdel-Rahman, S., Shaban, N., Haggag, A., Awad, D., Bassiouny, A., & Talaat, I. (2015). Inhibition of NF- κ B, Bcl-2 and COX-2 gene expression by an extract of *Eruca sativa* seeds during rat mammary gland carcinogenesis. *Asian Pacific Journal of Cancer Prevention*, 16(18), 8411–8418. <https://doi.org/10.7314/apjcp.2015.16.18.8411>
- Abd-El salam, R. M., El Badawy, S. A., Ogaly, H. A., Ibrahim, F. M., Farag, O. M., & Ahmed, K. A. (2021). *Eruca sativa* seed extract modulates oxidative stress and apoptosis and up-regulates the expression of Bcl-2 and Bax genes in acrylamide-induced testicular dysfunction in rats. *Environmental Science and Pollution Research*, 28, 53249–53266. <https://doi.org/10.1007/s11356-021-14532-y>
- Barillari, J., Canistro, D., Paolini, M., Ferroni, F., Pedulli, G. F., Iori, R., & Valgimigli, L. (2005). Direct antioxidant activity of purified glucoerucin, the dietary secondary metabolite contained in rocket (*Eruca sativa* mill.) seeds and sprouts. *Journal of Agricultural and Food Chemistry*, 53, 2475–2482. <https://doi.org/10.1021/jf047945a>
- Bell, L., & Wagstaff, C. (2019). Rocket science: A review of phytochemical & health-related research in *Eruca* & *Diplotaxis* species. *Food Chemistry: X*, 30, 100002. <https://doi.org/10.1016/j.fochx.2018.100002>
- Bhandari, D. C., & Chandel, K. P. S. (1996). Status of rocket germplasm in India: Research accomplishment and priorities. In S. Padulosi & D. Pignone (Eds.), *Rocket: A Mediterranean crop for the world*. Legnaro (Padova), Italy: Report of a workshop, 13–14 December.
- Chin, K. Y., Qin, C., May, L., Ritchie, R. H., & Woodman, O. L. (2017). New pharmacological approaches to the prevention of myocardial ischemia-reperfusion injury. *Current Drug Targets*, 18, 1689–1711. <https://doi.org/10.2174/1389450116666151001112020>
- Citi, V., Martelli, A., Testai, L., Marino, A., Breschi, M. C., & Calderone, V. (2014). Hydrogen sulfide releasing capacity of natural isothiocyanates: Is it a reliable explanation for the multiple biological effects of Brassicaceae? *Planta Medica*, 80, 610–613. <https://doi.org/10.1055/s-0034-1368591>
- Citi, V., Piragine, E., Testai, L., Breschi, M. C., Calderone, V., & Martelli, A. (2018). The role of hydrogen sulfide and H₂S-donors in myocardial protection against ischemia/reperfusion injury. *Current Medicinal Chemistry*, 25, 4380–4401. <https://doi.org/10.2174/0929867325666180212120504>
- Corssac, G. B., Campos-Carraro, C., Hickmann, A., da Rosa Araujo, A. S., Fernandes, R. O., & Belló-Klein, A. (2018). Sulforaphane effects on oxidative stress parameters in culture of adult cardiomyocytes. *Biomedicine & Pharmacotherapy*, 104, 165–171. <https://doi.org/10.1016/j.biopha.2018.05.031>
- Di Gioia, F., Avato, P., Serio, F., & Argentieri, M. P. (2018). Glucosinolate profile of *Eruca sativa*, *Diplotaxis tenuifolia* and *Diplotaxis erucoides* grown in soil and soilless systems. *Journal of Food Composition and Analysis*, 69, 197–204. <https://doi.org/10.1016/j.jfca.2018.01.022>
- Franco, P., Spinozzi, S., Pagnotta, E., Lazzeri, L., Ugolini, L., Camborata, C., & Roda, A. (2016). Development of a liquid chromatography-electrospray ionization-tandem mass spectrometry method for the simultaneous analysis of intact glucosinolates and isothiocyanates in brassicaceae seeds and functional foods. *Journal of Chromatography. A*, 1428, 154–161. <https://doi.org/10.1016/j.chroma.2015.09.001>
- Fuentes, E., Alarcón, M., Fuentes, M., Carrasco, G., & Palomo, I. (2014). A novel role of *Eruca sativa* mill. (rocket) extract: Antiplatelet (NF- κ B inhibition) and antithrombotic activities. *Nutrients*, 6, 5839–5852. <https://doi.org/10.3390/nu6125839>
- Gugliandolo, A., Giacoppo, S., Ficcichia, M., Aliquò, A., Bramanti, P., & Mazzone, E. (2018). *Eruca sativa* seed extract: A novel natural product able to counteract neuroinflammation. *Molecular Medicine Reports*, 17, 6235–6244. <https://doi.org/10.3892/mmr.2018.8695>
- Hancock, J. T., & Whiteman, M. (2014). Hydrogen sulfide and cell signaling: Team player or referee? *Plant Physiology and Biochemistry*, 78, 37–42. <https://doi.org/10.1016/j.plaphy.2014.02.012>
- Jana, S., Patel, D., Patel, S., Upadhyay, K., Thadani, J., Mandal, R., & Devkar, R. (2017). Anthocyanin rich extract of *Brassica oleracea* L. alleviates experimentally induced myocardial infarction. *PLoS One*, 12, e0182137. <https://doi.org/10.1371/journal.pone.0182137>
- Khan, H., & Khan, M. A. (2014). Antiulcer effect of extract/fractions of *Eruca sativa*: Attenuation of urease activity. *Journal of Evidence-Based Integrative Medicine*, 19, 176–180. <https://doi.org/10.1177/2156587214527452>
- Lazzeri, L., Errani, M., Leoni, O., & Venturi, G. (2004). *Eruca sativa* spp. oleifera: A new non-food crop. *Industrial Crops and Products*, 20, 67–73. [10.1016/j.indcrop.2002.06.001](https://doi.org/10.1016/j.indcrop.2002.06.001)
- Lucarini, E., Micheli, L., Trallori, E., Citi, V., Martelli, A., Testai, L., & Di Cesare, M. L. (2018). Effect of glucoraphanin and sulforaphane against chemotherapy-induced neuropathic pain: Kv7 potassium channels modulation by H₂S release in vivo. *Phytotherapy Research*, 32, 2226–2234. <https://doi.org/10.1002/ptr.6159>
- Lucarini, E., Pagnotta, E., Micheli, L., Parisio, C., Testai, L., Martelli, A., & Ghelardini, C. (2019). *Eruca sativa* meal against diabetic neuropathic pain: An H₂S-mediated effect of glucoerucin. *Molecules*, 24(16), 3006. <https://doi.org/10.3390/molecules24163006>
- Martelli, A., Piragine, E., Citi, V., Testai, L., Pagnotta, E., Ugolini, L., & Calderone, V. (2020). Erucin exhibits vasorelaxing effects and antihypertensive activity by H₂S-releasing properties. *British Journal of Pharmacology*, 177, 824–835. <https://doi.org/10.1111/bph.14645>
- Martelli, A., Piragine, E., Gorica, E., Citi, V., Testai, L., Pagnotta, E., & Calderone, V. (2021). The H₂S-donor Erucin exhibits protective effects against vascular inflammation in human endothelial and smooth muscle cells. *Antioxidants*, 10, 961. <https://doi.org/10.3390/antiox10060961>
- Martelli, A., Testai, L., Breschi, M. C., Blandizzi, C., Virdis, A., Taddei, S., & Calderone, V. (2012). Hydrogen sulphide: Novel opportunity for drug

- discovery. *Medicinal Research Reviews*, 32, 1093–1130. <https://doi.org/10.1002/med.20234>
- Martelli, A., Testai, L., Citi, V., Marino, A., Bellagambi, F. G., Ghimenti, S., & Calderone, V. (2014). Pharmacological characterization of the vascular effects of aryl isothiocyanates: Is hydrogen sulfide the real player? *Vascular Pharmacology*, 60, 32–41. <https://doi.org/10.1016/j.vph.2013.11.003>
- Mirzabe, A. H., Hajjahmad, A., & Asadollahzadeh, A. H. (2021). Moisture-dependent engineering properties of arugula seed relevant in mechanical processing and bulk handling. *Journal of Food Process Engineering*, 44, e13704. <https://doi.org/10.1111/jfpe.13704>
- Olson, K. R., Gao, Y., & Straub, K. D. (2021). Oxidation of hydrogen sulfide by Quinones: How polyphenols initiate their Cytoprotective effects. *International Journal of Molecular Sciences*, 22, 961. <https://doi.org/10.3390/ijms22020961>
- Pagliaro, B., Santolamazza, C., Simonelli, F., & Rubattu, S. (2015). Phytochemical compounds and protection from cardiovascular diseases: A state of the art. *BioMed Research International*, 2015, 918069. <https://doi.org/10.1155/2015/918069>
- Piragine, E., Flori, L., Di Cesare Mannelli, L., Ghelardini, C., Pagnotta, E., Matteo, R., & Calderone, V. (2021). *Eruca sativa* mill. Seed extract promotes anti-obesity and hypoglycemic effects in mice fed with a high-fat diet. *Phytotherapy Research*, 35, 1983–1990. <https://doi.org/10.1002/ptr.6941>
- Ritz, S. A., Wan, J., & Diaz-Sanchez, D. (2007). Sulforaphane-stimulated phase II enzyme induction inhibits cytokine production by airway epithelial cells stimulated with diesel extract. *American Journal of Physiology*, 292, L33–L39. <https://doi.org/10.1152/ajplung.00170.2006>
- Salma, U., Khan, T., & Shah, A. J. (2018). Antihypertensive effect of the methanolic extract from *Eruca sativa* mill., (Brassicaceae) in rats: Muscarinic receptor-linked vasorelaxant and cardiogenic effects. *Journal of Ethnopharmacology*, 224, 409–420. <https://doi.org/10.1016/j.jep.2018.06.013>
- Sarwar, A. M., Kaur, G., Jabbar, Z., Javed, K., & Athar, M. (2007). *Eruca sativa* seeds possess antioxidant activity and exert a protective effect on mercuric chloride induced renal toxicity. *Food and Chemical Toxicology*, 45(6), 910–920. <https://doi.org/10.1016/j.fct.2006.11.013>
- Silambarasan, T., Manivannan, J., Priya, M. K., Suganya, N., Chatterjee, S., & Raja, B. (2015). Sinapic acid protects heart against ischemia/reperfusion injury and H9c2 cardiomyoblast cells against oxidative stress. *Biochemical and Biophysical Research Communications*, 456, 853–859. <https://doi.org/10.1016/j.bbrc.2014.12.022>
- Sivarajah, A., Collino, M., Yasin, M., Benetti, E., Gallicchio, M., Mazzon, E., & Thiernemann, C. (2009). Anti-apoptotic and anti-inflammatory effects of hydrogen sulfide in a rat model of regional myocardial I/R. *Shock*, 31, 267–274. <https://doi.org/10.1097/SHK.0b013e318180ff89>
- Talmor, D., Roytblat, L., Artru, A. A., Yuri, O., Koefman, L., Katchko, L., & Shapira, Y. (1998). Phenylephrine-induced hypertension does not improve outcome after closed head trauma in rats. *Anesthesia and Analgesia*, 87, 574–578. <https://doi.org/10.1097/0000539-199809000-00014>
- Testai, L., Marino, A., Piano, I., Brancaleone, V., Tomita, K., Di Cesare Mannelli, L., & Calderone, V. (2016). The novel H₂S-donor 4-carboxyphenyl isothiocyanate promotes cardioprotective effects against ischemia/reperfusion injury through activation of mitoKATP channels and reduction of oxidative stress. *Pharmacological Research*, 113, 290–299. <https://doi.org/10.1016/j.phrs.2016.09.006>
- Testai, L., Rapposelli, S., Martelli, A., Breschi, M. C., & Calderone, V. (2015). Mitochondrial potassium channels as pharmacological target for cardioprotective drugs. *Medicinal Research Reviews*, 35, 520–553. <https://doi.org/10.1002/med.21332>
- Tian, R., Yang, W., Xue, Q., Gao, L., Huo, J., Ren, D., & Chen, X. (2015). Rutin ameliorates diabetic neuropathy by lowering plasma glucose and decreasing oxidative stress via Nrf2 signaling pathway in rats. *European Journal of Pharmacology*, 771, 84–92. <https://doi.org/10.1016/j.ejphar.2015.12.021>
- Vashi, R., & Patel, B. M. (2021). NRF2 in cardiovascular diseases: A ray of Hope! *Journal of Cardiovascular Translational Research*, 14(573–586), 573–586. <https://doi.org/10.1007/s12265-020-10083-8>
- Villatoro-Pulido, M., Font, R., Saha, S., Obregon-Cano, S., Anter, J., Munoz-Serrano, A., & Del Rio-Celestino, M. (2012). In vivo biological activity of rocketextracts (*Eruca vesicaria* subsp. *sativa* (miller) Thell) and sulforaphane. *Food and Chemical Toxicology*, 50, 1384–1392. <https://doi.org/10.1016/j.fct.2012.02.017>
- Wathelet, J. P., Iori, R., Leoni, O., Rollin, P., Quinsac, A., & Palmieri, S. (2004). Guidelines for glucosinolate analysis in green tissues used for biofumigation. *Agroindustria*, 3, 257–266. <http://hdl.handle.net/2268/16127>
- Wu, L., Noyan, A. M. H., Facci, M., Wang, R., Paterson, P. G., Ferrie, A., & Juurlink, B. H. (2004). Dietary approach to attenuate oxidative stress, hypertension, and inflammation in the cardiovascular system. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 7094–7099. <https://doi.org/10.1073/pnas.0402004101>
- Yang, D. K. (2018). Cabbage (*Brassica oleracea* var. *capitata*) protects against H₂O₂-induced oxidative stress by preventing mitochondrial dysfunction in H9c2 Cardiomyoblasts. *Evid Based Complement Alternat Med*, 2018, 2179021. <https://doi.org/10.1155/2018/2179021>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Testai, L., Pagnotta, E., Piragine, E., Flori, L., Citi, V., Martelli, A., Mannelli, L. D. C., Ghelardini, C., Matteo, R., Suriano, S., Troccoli, A., Pecchioni, N., & Calderone, V. (2022). Cardiovascular benefits of *Eruca sativa* mill. Defatted seed meal extract: Potential role of hydrogen sulfide. *Phytotherapy Research*, 36(6), 2616–2627. <https://doi.org/10.1002/ptr.7479>