

Editorial

To Do or Not to Do? - The Value of the Preseason Assessment in Sport Injury Prevention

Luciana De Michelis Mendonça^{1 a}

¹ 1. Graduate Program in Rehabilitation and Functional Performance - Physical Therapy Department - Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM) - Diamantina, Minas Gerais, Brazil; President of the International Federation of Sports Physical Therapy (IFSPT)

Keywords: preseason assessment, sport physical therapy, injury prevention

<https://doi.org/10.26603/001c.31871>

International Journal of Sports Physical Therapy

Vol. 17, Issue 2, 2022

In 2011, World Physiotherapy published the Standards of Physical Therapy practice (<https://world.physio/sites/default/files/2020-06/G-2011-Standards-practice.pdf>), which indicates that “the physical therapist performs an initial examination/assessment and evaluation to establish a diagnosis and prognosis/plan of care prior to intervention/treatment.” Since assessment is considered mandatory for a clinical decision-making process in Physical Therapy,^{1,2} it is expected to find sports physical therapists performing preseason assessment with their athletes. A Preseason Assessment (PA) is a battery of tests chosen to identify and characterize the health status of athletes (screening) to prevent injuries and improve performance.³ In addition, the PA might identify athletes with increased likelihood of being injured and guide the initial phase of the preventive program planned by sport physical therapists³.

Screening athletes is mandatory in other professions. For example, the American College of Sports Medicine proposes a preparticipation health screening on athletes to access exercise-related cardiovascular events.⁴ The argument commonly used that “general prevention programs work, so why the concern on assessing and building tailored programs?” is not enough to abandon the standards of our profession. Athletes’ health and safety should be our main concerns and our interventions should be specific for each health condition, each sport injury, each athlete. Therefore, we should deliver our efforts to targeting the best health and safety status. The purpose of this editorial is to discuss the execution of preseason assessment (PA) and planning preventive programs based on the PA results.

To understand injury occurrence, we should know about the sport action and most common movements, collect athlete’s injury history and sport practice, and identify and measure athletes’ needs (physical, psychological, sport performance, etc) to facilitate the outcome measurement (dysfunctions linked to the injury).¹ If injury is an established problem in sport practice, how can we prevent it without knowing/understanding it? An important process that sports physical therapists should do to understand athletic injury is to assess, quantify, define the diagnosis, imple-

ment interventions, follow-up and re-assess. Mehl et al.⁵ indicated that screening, identification, and correction of endangering movement patterns like the dynamic valgus are the first crucial steps in order to prevent knee injuries in athletes. Interestingly, Mendonça et al.¹ developed an international survey and the authors reported a frequency of 75% sports PT performing PA in their athletes.¹ The fact that about one third of these sports physical therapists use the results of the PA to build the prevention program was surprisingly negative.¹

PA would be recognized as mandatory and properly implemented (and even disseminated) if it is validated. To accomplish this, it is necessary to apply the PA results in sport injury prevention program implementation and follow-up injury occurrences to actually validate the prevention program and also the PA itself. Bittencourt et al.⁶ recently published a cohort study which identified that a tailored preventive program reduced the incidence of patellar tendinopathy in elite youth jumping athletes. The necessity of performing this preseason screening has been questioned, mainly based on the statement of lack of strong evidence.⁷

Considering that the pre-competition season usually involves athletes being exposed to frequent training sessions and friendly matches before a break-time. Even a global non-specific prevention program, such as FIFA 11+, could benefit the athlete. However, we might not do all in our power to help our athletes throughout the whole season. For example, Slauterbeck et al.⁸ did not find a reduction in lower extremity injury in schools using the FIFA 11+ program compared with schools using their usual pre-practice warm-up program. In elite athletes, although some studies indicate that FIFA 11+ reduces injury incidence in soccer, Ekstrand et al.⁹ found that hamstring injuries have increased 4% annually, during 13 years follow-up, in elite male soccer teams.

So maybe the problem is not about the PA itself, but how to perform the PA. Which tests to choose? How to apply it? How to do analyze the data? Relative limitations in performing the PA might be the time needed to organize and

^a **Cooresponding author:**

lucianademichelis@gmail.com

President of the International Federation of Sports Physical Therapy (IFSPT)

execute, high-cost equipment and lack of methodological rigor.¹ However, those limitations could be easily addressed with strategies such as substituting tests using expensive equipment for clinical tests, keeping the scientific rigor (i.e. LESS),¹⁰ and possibly involving university students to make the process easier to execute.

The purpose of the PA is not to predict injury, but it to screen our athletes, identify risk profiles, and set specific parameters to improve their capacity to deal with sport demands.⁶ We should use PA results to build a tailored pre-

ventive program to help our athletes achieve the strength and skill to perform.⁶ Considering that PA procedures could be performed on the field using low-cost equipment, these regimens should be promoted and facilitated in sports organizations world-wide, by means of shared consensus amongst the organization's medical and technical staffs.

Submitted: December 01, 2021 CST, Accepted: January 01, 2022 CST



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Mendonça LD, Ley C, Schuermans J, Wezenbeek E, Witvrouw E. How injury registration and preseason assessment are being delivered: An international survey of sports physical therapists. *Phys Ther Sport*. 2021;53:S1466-853X(21)00147-4. doi:10.1016/j.ptsp.2021.08.014
2. Teo PL, Hinman RS, Egerton T, Dziedzic KS, Bennell KL. Identifying and Prioritizing Clinical Guideline Recommendations Most Relevant to Physical Therapy Practice for Hip and/or Knee Osteoarthritis. *J Orthop Sports Phys Ther*. 2019;49(7):501-512. doi:10.2519/jospt.2019.8676
3. Kennedy MD, Fischer R, Fairbanks K, et al. Can pre-season fitness measures predict time to injury in varsity athletes?: a retrospective case control study. *Sports Medicine, Arthroscopy, Rehabilitation, Therapy & Technology*. 2012;4(1):26. doi:10.1186/1758-2555-4-26
4. Riebe D, Franklin BA, Thompson PD, et al. Updating ACSM's Recommendations for Exercise Preparticipation Health Screening. *Med Sci Sports Exerc*. 2015;47(11):2473-2479. doi:10.1249/mss.0000000000000664
5. Mehl J, Diermeier T, Herbst E, et al. Evidence-based concepts for prevention of knee and ACL injuries. 2017 guidelines of the ligament committee of the German Knee Society (DKG). *Arch Orthop Trauma Surg*. 2018;138(1):51-61. doi:10.1007/s00402-017-2809-5
6. Bittencourt NFN, Oliveira RR, Vaz RPM, Silva RS, Mendonça LM. Preventive effect of tailored exercises on patellar tendinopathy in elite youth athletes: A cohort study. *Phys Ther Sport*. 2022;53:60-66. doi:10.1016/j.ptsp.2021.11.006
7. Bahr R. Why screening tests to predict injury do not work—and probably never will...: a critical review. *Br J Sports Med*. 2016;50(13):776-780. doi:10.1136/bjsports-2016-096256
8. Slauterbeck JR, Choquette R, Tourville TW, et al. Implementation of the FIFA 11+ Injury Prevention Program by High School Athletic Teams Did Not Reduce Lower Extremity Injuries: A Cluster Randomized Controlled Trial. *Am J Sports Med*. 2019;47(12):2844-2852. doi:10.1177/0363546519873270
9. Ekstrand J, Waldén M, Häggglund M. Hamstring injuries have increased by 4% annually in men's professional football, since 2001: a 13-year longitudinal analysis of the UEFA Elite Club injury study. *Br J Sports Med*. 2016;50(12):731-737. doi:10.1136/bjsports-2015-095359
10. Padua DA, Marshall SW, Boling MC, Thigpen CA, Garrett WE Jr, Beutler AI. The Landing Error Scoring System (LESS) Is a valid and reliable clinical assessment tool of jump-landing biomechanics: The JUMP-ACL study. *Am J Sports Med*. 2009;37(10):1996-2002. doi:10.1177/0363546509343200