

TRACEBACK: Testing of Historical Tubo-Ovarian Cancer Patients for Hereditary Risk Genes as a Cancer Prevention Strategy in Family Members

Rachel Delahunty, MD^{1,2}; Linh Nguyen, PhD^{1,2}; Stuart Craig, BSc(Hons)¹; Belinda Creighton, GDip³; Dinuka Ariyaratne, BSc(Hons)¹; Dale W. Garsed, PhD^{1,2}; Elizabeth Christie, PhD^{1,2}; Sian Fereday, BSc(Hons)^{1,2}; Lesley Andrews, MMed^{4,5}; Alexandra Lewis, GDip⁶; Shame Limb, MGenCouns^{2,6}; Ahwan Pandey, MSc¹; Joy Hendley¹; Nadia Traficante, BSc(Hons)^{1,2}; Natalia Carvajal, MBiotech¹; Amanda B. Spurdle, PhD⁷; Bryony Thompson, PhD^{8,9}; Michael T. Parsons, BSc(Hons)⁷; Victoria Beshay, PhD¹; Mila Volcheck, MD^{1,10,11}; Timothy Semple, BSc(Hons)¹; Richard Lupat, MPhil¹; Kenneth Doig, PhD^{1,2}; Jiaan Yu, MSc¹; Xiao Qing Chen, MBBS⁷; Anna Marsh, BSc(Hons)⁷; Christopher Love, PhD¹; Sanela Bilic, BSc¹²; Maria Beilin, BSc(Hons)¹²; Cassandra B. Nichols, GDip¹³; Christina Greer, MGenCouns¹³; Yeh Chen Lee^{4,5,14}; Susan Gerty, BSc(Hons)⁴; Lynette Gill, RN⁴; Emma Newton, MPH⁴; Julie Howard, BNurs⁴; Rachel Williams, GDip^{14,15}; Christie Norris, MPH⁴; Andrew N. Stephens, PhD^{16,17,18}; Erin Tutty, MGenCouns⁶; Courtney Smyth, GDip¹⁹; Shona O'Connell, GDip¹⁹; Thomas Jobling, MD²⁰; Colin J.R. Stewart, MD²¹; Adeline Tan, MD²²; Stephen B. Fox, DPhil^{1,2}; Nicholas Pachter^{13,23,24}; Jason Li, PhD^{1,2}; Jason Ellul, BMACompSc(Hons)^{1,2}; Gisela Mir Arnau, PhD^{1,2}; Mary-Anne Young, MHSc²⁵; Louisa Gordon, PhD^{7,26}; Laura Forrest, PhD^{1,2}; Marion Harris, MD²⁰; Karen Livingstone²⁷; Jane Hill³; Georgia Chenevix-Trench, PhD⁷; Paul A. Cohen, MD^{12,28}; Penelope M. Webb, PhD⁷; Michael Friedlander, MD^{4,5,13}; Paul James, MD^{1,2}; David Bowtell, PhD^{1,2}; and Kathryn Alsop, PhD^{1,2}; on behalf of The Australian Ovarian Cancer Study, Ovarian Cancer Prognosis and Lifestyle Study and the TRACEBACK Study

abstract

PURPOSE Tubo-ovarian cancer (TOC) is a sentinel cancer for *BRCA1* and *BRCA2* pathogenic variants (PVs). Identification of a PV in the first member of a family at increased genetic risk (the proband) provides opportunities for cancer prevention in other at-risk family members. Although Australian testing rates are now high, PVs in patients with TOC whose diagnosis predated revised testing guidelines might have been missed. We assessed the feasibility of detecting PVs in this population to enable genetic risk reduction in relatives.

PATIENTS AND METHODS In this pilot study, deceased probands were ascertained from research cohort studies, identification by a relative, and gynecologic oncology clinics. DNA was extracted from archival tissue or stored blood for panel sequencing of 10 risk-associated genes. Testing of deceased probands ascertained through clinic records was performed with a consent waiver.

RESULTS We identified 85 PVs in 84 of 787 (11%) probands. Familial contacts of 39 of 60 (65%) deceased probands with an identified recipient (60 of 84; 71%) have received a written notification of results, with follow-up verbal contact made in 85% (33 of 39). A minority of families ($n = 4$) were already aware of the PV. For many (29 of 33; 88%), the genetic result provided new information and referral to a genetic service was accepted in most cases (66%; 19 of 29). Those who declined referral (4 of 29) were all male next of kin whose family member had died more than 10 years before.

CONCLUSION We overcame ethical and logistic challenges to demonstrate that retrospective genetic testing to identify PVs in previously untested deceased probands with TOC is feasible. Understanding reasons for a family member's decision to accept or decline a referral will be important for guiding future TRACEBACK projects.

ASSOCIATED CONTENT

Data Supplement

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on February 3, 2022 and published at ascopubs.org/journal/jco on March 9, 2022; DOI <https://doi.org/10.1200/JCO.21.02108>

J Clin Oncol 40:2036-2047. © 2022 by American Society of Clinical Oncology

Licensed under the Creative Commons Attribution 4.0 License 

INTRODUCTION

Prevention strategies in individuals at increased genetic risk of cancer offer important opportunities to reduce cancer incidence and mortality. This is particularly relevant for tubo-ovarian cancer (referred to as ovarian cancer). Pathogenic germline variants in *BRCA1* and *BRCA2* (*BRCA1/2*) confer a cumulative risk for ovarian cancer by age 80 years of 44% and 17%, respectively.¹ Other genes, including

RAD51C, *RAD51D*, *PALB2*, *BRIP1*, and genes involved in DNA mismatch repair, are associated with a moderately increased risk of between 5% and 12%.²⁻⁵ In the absence of reliable early detection, prevention of ovarian cancer remains the most effective means to reduce disease impact, as the majority of patients have advanced-stage cancer at diagnosis, which is associated with high rates of relapse and mortality.

CONTEXT

Key Objective

Germline mutations in hereditary cancer risk genes, including *BRCA1* and *BRCA2*, are common in high-grade non-mucinous tubo-ovarian cancers. Retrospective identification of patients who died of their disease without being offered clinical genetic testing, which is now the standard of care, presents an opportunity to identify genetically at-risk family members before further cancer diagnoses.

Knowledge Generated

Our study explored three methods for the identification and genetic testing of deceased patients with ovarian cancer for pathogenic variants in hereditary risk genes. We were able to subsequently contact familial relatives of mutation carriers and offer referral for specialist genetic counseling and follow-up.

Relevance

Retrospective identification of pathogenic germline variants in deceased patients with tubo-ovarian cancer is feasible and importantly, results in uptake of referral to genetic services in family members. Our experiences can guide the development of similar programs internationally and address ethical and logistical challenges.

Over the past decade, two factors have positively influenced genetic testing rates of patients with newly diagnosed ovarian cancer: the demonstrated efficacy of maintenance therapy with poly (ADP-ribose) polymerase inhibitors in germline *BRCA1/2* carriers^{6,7} and recognition that family history–based testing criteria were frequently inadequate.⁸ Subsequently, genetic testing guidelines have been revised in many countries to include all women with ovarian cancer.⁹ There is, however, a legacy of untested, often deceased, patients whose diagnosis predated these changes. In such circumstances, the germline status of untested patients, and therefore familial risk, is likely to remain unknown until subsequent cancer diagnoses in family members, which might have been prevented had they been aware that they had inherited a pathogenic variant (PV).

Pathogenic germline variants in *BRCA1/2* are present in 12%-15% of patients with high-grade nonmucinous ovarian cancer (HGNMOC),^{8,10} the highest prevalence of any malignancy and therefore representing a sentinel cancer for *BRCA1/2* carriers. The identification of a patient with ovarian cancer within a family that is at increased genetic risk (proband) as a strategy for cancer prevention in other family members was explored at a National Cancer Institute workshop in 2016, and a conceptual framework, termed Traceback, developed.¹¹ The subsequent editorial published in this journal¹² highlighted significant ethical, legal, and social implications of retrospective identification and testing of probands and recommended pilot studies.

Here, we provide our experience in establishing and conducting, to our knowledge, the first genetic testing program for deceased patients with ovarian cancer, which we have also termed TRACEBACK. Our pilot demonstrated the feasibility and effectiveness of multiple ascertainment approaches for deceased probands, performed retrospective genetic testing, and returned significant results to

families to facilitate cascade testing. We discuss the ethical and logistic challenges with this type of program.

PATIENTS AND METHODS

This research was approved by the Human Research Ethics Committee (HREC) at the Peter MacCallum Cancer Centre (PMCC, Melbourne, Australia; HREC/17/PMCC/89) and all participating hospitals (including SJOG 1393).

Identification of Probands

Eligible probands were patients diagnosed with high-grade nonmucinous epithelial carcinoma of the ovary, fallopian tube, or peritoneum (HGNMOC) between 2000 and 2016 and were ascertained through existing research studies (cohorts), from specialist gynecologic oncology clinic medical records (clinic), and by a next of kin (NOK) of a deceased HGNMOC proband (NOK referral). Further details of case ascertainment are provided in the Data Supplement (online only). Probands determined to have previously undergone *BRCA1/2* germline testing in a clinical or research setting were excluded.

Vital Status

Although there was provision within the TRACEBACK study to facilitate testing of living probands (Data Supplement), here, we evaluate the implications of genetic testing in the deceased ovarian cancer population.

Genetic Testing

Germline DNA derived from blood was available for most cohort-ascertained probands. If unavailable, we obtained DNA extracted from either frozen or formalin-fixed paraffin-embedded (FFPE) tissue (Fig 1 and Table 1). A gynecologic pathologist reviewed all tissue, and DNA was extracted from areas enriched by needle dissection for normal content where possible. In some instances, only tumor-rich tissue was available (Fig 1 and Table 1).

Genetic testing was performed using a customized Agilent SureSelect^{XT} Low Input capture panel (Design ID 314187) of 79 genes (Data Supplement). Libraries were prepared using the SureSelect^{XT} Low Input Target Enrichment System (Agilent, Santa Clara, CA) and sequenced on an Illumina NextSeq 500 (Data Supplement).

Variant Detection and Curation

Sequence alignment and variant detection are described in the Data Supplement. Ten genes (*BRCA1*, *BRCA2*, *RAD51C*, *RAD51D*, *BRIP1*, *PALB2*, *MLH1*, *MSH2*, *MSH6*, and *PMS2*), regarded as clinically actionable,⁹ were reviewed in PathOS¹³ and single-nucleotide variants and indels filtered to remove synonymous and common variants (global minor allele frequency > 1%). Potentially pathogenic nonsynonymous variants were reviewed manually via Integrative Genomics Viewer.^{14,15} Variant pathogenicity was assigned using databases such as ClinVar,¹⁶ InSiGHT,¹⁷ and literature from expert panels¹⁸ and classified according to the International Agency for Cancer Research (IARC) and American College of Medical Genetics and Genomics (ACMG).^{19,20} All variants of unknown significance (C3), likely pathogenic/pathogenic (C4/C5) variants or any with ambiguous classification, were committee reviewed. Variants determined to be C4/C5 (here after referred to as PVs) were subsequently validated in a National Association of Testing Authorities, Australia (NATA; ISO15189)-accredited clinical laboratory, using independent DNA extractions where available (Data Supplement).

Return of Findings

A familial recipient for the research results was identified from the proband's medical records or the research cohort consent form (Data Supplement). Results were disseminated using a two-step process. This involved delivery of a written notification (Data Supplement) alerting the recipient of genetic information availability without specifying findings, followed by a telephone call from a genetic clinician (counselor/geneticist or a knowledgeable specialist known to the family; GC). During follow-up contact, referral to a local Familial Cancer Clinic (FCC) for predictive testing was offered.

Statistical Analyses

Data were analyzed using GraphPad Prism 8.4 for Windows (GraphPad Software, San Diego, CA) or in the R software. A *P* value of .05 was considered statistically significant (further details are given in the Data Supplement).

RESULTS

A Large Population of Undetected PV Carriers

Approximately 17,000 women were diagnosed with invasive HGNMOC in Australia between 2000 and 2016.^{21,22} On the basis of the reported testing rates from this period in two Australian studies^{8,23} and data captured by participating research studies, we estimated that 12,000 patients missed genetic testing. Incorporating survival data, we anticipated that approximately 60% would have died of

their disease by 2018.²⁴ Previous literature indicates that 15% of Australian patients with HGNMOC carry a pathogenic germline *BRCA1/2* variant.⁸ We therefore estimated that in 2018, approximately 1,100 *BRCA1/2* pathogenic germline variant carriers exist among untested, deceased patients with ovarian cancer in Australia.

Three Approaches to Proband Ascertainment

We used a multimodal approach for proband ascertainment (Data Supplement). The most straightforward involved accessing specimens from national ovarian cancer research studies. TRACEBACK was also open to referral of a deceased HGNMOC proband by a family member (NOK referral), who would otherwise be unable to access Australian government-subsidized genetic testing as it is currently restricted to living probands. We encouraged FCCs and clinicians to refer relatives who may be interested in participating. These individuals provided consent for genetic testing of their family member's tissue.

We recognized that the most efficient way to identify large numbers of eligible probands would be through auditing the medical records of primary treatment centers as most ovarian cancer surgery in Australia is centralized to metropolitan tertiary hospitals. Local clinical investigators reviewed medical records for eligible probands, including determination of previous genetic testing and vital status. In consultation with Institutional Review Boards (IRBs), we carefully considered whether to seek consent from relatives before testing. The Australian National Health and Medical Research Council (NHMRC) National Statement on Ethical Conduct in Human Research²⁵ allows a waiver of the requirement for consent in certain circumstances. The rationale for proceeding with testing under a consent waiver and the processes put in place to minimize harm are detailed in the Data Supplement and Discussion.

PVs

We report results for the first 824 deceased probands, including 512 cohort-ascertained, 80 self-referrals, and 232 ascertained from participating clinics (Fig 1). Blood-derived DNA was available for 88% (451 of 512) of the cohort probands, with normal FFPE tissue used for 85% (265 of 312) of clinic/NOK-referral probands and tumor DNA for the remainder (Fig 1). Clinical features are summarized in Table 1.

After sequencing, 96% (787 of 824) of samples met predefined quality control measures (Data Supplement). Eighty-five PVs were detected in 84 probands (84 of 787; 11%; Fig 2). *BRCA1/2* had the highest frequency of PVs (46 of 85, 54% and 22 of 85, 26%, respectively; Fig 2) with variability depending on the proband source, patient age, and year of diagnosis (Fig 3). Consistent with previous findings,⁸ *BRCA1/2* PVs were predominantly frameshift variants in the largest coding exons (Fig 2 and Data Supplement). Twenty percent (17 of 85) of PVs occurred in genes other than *BRCA1/2*, including *BRIP1* (9 of 85; 11%) and *PALB2* (3 of 85; 4%). Two PVs (2%) were detected in




FIG 1. TRACEBACK probands and samples. Schematic representation of the three ascertainment methods used to identify and test high-grade nonmucinous ovarian cancer probands, diagnosed in Australia between the years 2000 and 2016. Genetic testing of deceased probands ($n = 824$) is reported here, with information available for the living probands ascertained to the study ($n = 265$) in the Data Supplement. ^aSome exceptions were made for deceased probands diagnosed outside 2000-2016 for NOK referral (Data Supplement). ^bPredominately normal tissue was obtained from surgical tumor samples and included both FFPE and fresh-frozen biospecimens (Data Supplement). FFPE, formalin-fixed paraffin-embedded; HGNMOC, high-grade nonmucinous ovarian cancer; NOK-R, Next of Kin-referred; QC, quality control.

Lynch syndrome-associated genes (Fig 2 and Data Supplement).

All PVs identified were independently validated (Data Supplement). Thirteen PVs were initially detected in tumor-derived DNA, including one case with two *BRCA2* PVs (Data Supplement). For the purposes of validation, we were able to enrich the independent DNA for normal tissue content in 5 of 12 cases (Data Supplement). PVs in the remaining seven cases were validated with tumor DNA, and the possibility of somatic occurrence is recognized. Sixteen percent (126 of 787) of probands had a variant of unknown significance detected (Data Supplement), which were not validated or returned because of uncertainty of their clinical significance.

Recipients for Notification of Genetic Test Findings

To date, an appropriate familial recipient (next of kin; NOK) with verified contact details for the notification of a PV was identified for 71% (60 of 84) of the deceased probands

(Fig 4 and Data Supplement). Despite a thorough search, no appropriate contact could be found in six instances (7%) and identification of an appropriate recipient of findings is ongoing for the remainder (18 of 84; 21%).

The study commenced in 2018 and was cross-checking with updated clinical information before notifying results, and 14 of the 60 probands with an identified contact were found to have subsequently received clinical genetic testing since initially identified as eligible (Data Supplement). In each case, the findings of this study were consistent with the result ascertained through clinical testing and notification of research results was not required in these instances. To date, of the remaining 46 of 60 probands with confirmed recipient details for results, 85% (39 of 46) have been sent a written notification (Fig 4 and Data Supplement). This includes seven NOKs that self-referred to TRACEBACK and familial contacts for 32 cohort or clinic-ascertained probands.

TABLE 1. Clinical Characteristics of Deceased TRACEBACK Probands

Characteristic	Cohorts						NOK-R				Clinic		Total	
	ACS	%	AOCS	%	OPAL	%	NOK-R	%	1	%	2	%		%
n	100		265		147		80		153		79		824	
Age, years														
Mean	62	62	63	24	65	44	63	79	67	44	69	87.3	389	47
< 70	75	75	186	70	99	67	56	70	89	58	39	49.4	544	66
≥ 70	25	25	79	30	48	33	24	30	64	42	40	50.6	280	34
Vital status														
Deceased	100	100	265	100	147	100	80	100	153	100	79	100.0	824	100
Year of diagnosis														
Pre-2000	0	0	1	0	0	0	10	13	0	0	0	0.0	11	1
2000-2013	100	100	239	90	51	35	55	69	128	84	79	100.0	652	79
2014-2016	0	0	6	2	96	65	7	9	25	16	0	0.0	134	16
> 2017	0	0	0	0	0	0	8	10	0	0	0	0.0	8	1
Unknown	0	0	19	7	0	0	0	0	0	0	0	0.0	19	2
Histologic type														
HGSOC	78	78	206	78	109	74	64	80	127	83	57	72.2	641	78
HGEn	3	3	6	2	5	3	2	3	4	3	0	0.0	20	2
CC	4	4	12	5	7	5	4	5	7	5	5	6.3	39	5
Others	3	3	13	5	8	5	3	4	10	7	10	12.7	47	6
Undifferentiated	3	3	15	6	0	0	0	0	2	1	0	0.0	20	2
Carcinoma NOS	4	4	6	2	15	10	5	6	0	0	1	1.3	31	4
Mixed	5	5	7	3	3	2	2	3	3	2	6	7.6	26	3
Stage														
I	7	7	6	2	6	4	0	0	6	4	3	3.8	28	3
II	7	7	6	2	5	3	3	4	10	7	3	3.8	34	4
III	70	70	174	66	95	65	15	19	108	71	14	17.7	476	58
IV	13	13	25	9	32	22	3	4	17	11	10	12.7	100	12
Early NOS	0	0	0	0	0	0	6	8	0	0	6	7.6	12	1
Advanced NOS	0	0	0	0	8	5	47	59	12	8	42	53.2	109	13
Unknown	3	3	54	20	1	1	6	8	0	0	1	1.3	65	8
Tissue type tested														
Germline	100	100	204	77	147	100	7	9	0	0	0	0.0	458	56
Normal FFPE	0	0	24	9	0	0	58	73	142	93	65	82.3	289	35
Tumor FFPE	0	0	7	3	0	0	15	19	11	7	14	17.7	47	6
Frozen tumor	0	0	30	11	0	0	0	0	0	0	0	0.0	30	4

NOTE. An overview of the clinical characteristics of the 824 deceased probands enrolled and tested through TRACEBACK.

Abbreviations: ACS, Australian Cancer Study; AOCS, The Australian Ovarian Cancer Study; CC, Clear cell; FFPE, formalin-fixed paraffin-embedded; HGEn, High-Grade Endometrioid; HGSOC, High-Grade Serous ovarian cancer; NOK-R, next of kin-referred; NOS, not otherwise specified; OPAL, Ovarian Cancer Prognosis and Lifestyle Study.

GC Manages Processes After Initial Notification

Approximately two weeks after the notification letter was sent, a GC contacted the identified recipient. To date, verbal contact has been made with 86% (33 of 39) of individuals to whom a notification has been sent (Fig 4 and

Data Supplement). Attempted verbal contact has been made in the remaining six cases.

The feedback of PVs to NOK recruited to TRACEBACK via NOK referral (7 of 33; 21%) was relatively simple and successful, and all accepted an FCC referral (Fig 4 and




FIG 2. Detected C4/C5 variants (PVs). An overview of the clinical and genomic features of 84 of 787 (11%) deceased probands with PVs, including one proband with two PVs, both in *BRCA2* (one frameshift and one nonsense [given in ^a]). ACS, Australian Cancer Study; AOCS, Australian Ovarian Cancer Study; Carcinoma NOS, Carcinoma not otherwise specified; FFPE, Formalin-fixed paraffin-embedded; HGE, High-Grade Endometrioid; HGNMOC, high-grade nonmucinous ovarian cancer; HGS, High-Grade Serous; NOK-R, Next of Kin-referred; Opal, Ovarian Cancer Prognosis and Lifestyle Study; PV, pathogenic variant.

Data Supplement). By contrast, returning results to the NOK of the deceased cohort and clinic-ascertained probands (26 of 33; 79%) was more challenging. Follow-up contact by the GCs to the NOK was unexpected and, in most cases, occurred years after the proband's initial diagnosis, or death. Contact between the GC and NOK was challenging in some instances, for example, to convey to the NOK in a short period of time the significance of an unexpected phone contact. However, in most instances (24 of 26; 92%), a meaningful discussion resulted. Overall, 55% (12 of 22) of NOK presented with new and unsolicited genetic information have already accepted a FCC referral

(Fig 4). Of the remaining NOKs, four were already aware of the familial variant, two declined further clinical support, and six required further input; this included three instances where it was requested that contact is made with an alternative family member and three NOKs requested more time to consider the implications of the findings for their family. Finally, in the remaining two instances (2 of 26; 8%), although an initial verbal contact was made, further communication attempts with the NOK were unsuccessful and interpreted as a decline of the offered information. All declining NOKs were male, and their proband family member had died more than 10 years before.




FIG 3. Frequency of PVs in the 787 deceased probands stratified by (A) method of ascertainment, (B) age at diagnosis, and (C) years of diagnosis. Group differences for categorical variables were examined using the chi-square test (GraphPad Prism 8.4 for Windows). ACS, Australian Cancer Study; AOCS, Australian Ovarian Cancer Study; NOK-R, Next of Kin—referred; OPAL, Ovarian Cancer Prognosis and Lifestyle Study; PV, pathogenic variant.

DISCUSSION

Detection of pathogenic germline variants in *BRCA1/2* or other risk genes is relevant to the treatment of patients with ovarian cancer and provides an opportunity for cancer prevention in family members. Although identifying genetic predisposition in a family can cause distress, for most, it is seen as an opportunity to reduce cancer risk.²⁶ Many women diagnosed with ovarian cancer died before genetic testing criteria became less restrictive, representing missed opportunities for cancer prevention in family members. The value of assessing such patients relies on feasibility, accurate testing, and uptake of risk-reducing strategies in recipients of significant results.

Although conceptually simple, TRACEBACK presented significant ethical and logistic challenges. We explored three different proband ascertainment pathways, recognizing that ascertainment through specialist clinics provided an opportunity to capture a large proband population, independent of socioeconomic and geographical factors, or previous opportunities to participate in research. It was, however, also the most ethically challenging.

TRACEBACK is, to our knowledge, not only the first program of its kind internationally but also the first study to perform research genetic testing for clinical applications under a consent waiver. Although decisions regarding genetic testing should be autonomous, our intention to

focus on deceased probands precluded personal consent, aside from historical unspecified research consent for a proportion of probands. We recognized that contacting family members before the use of their deceased relative’s sample, of which only a minority would be determined to harbor a pathogenic germline variant, would greatly slow progress of the study, reduce the proband population, and likely render the study unfeasible. Instead, we proposed limiting contact only when a PV was identified. This targeted approach reduced the challenge of identifying an appropriate NOK by an order of magnitude and prevented causing unnecessary distress in a majority of families that are not at increased genetic risk of cancer. Working within the Australian NHMRC research guidelines, and in consultation with our IRB, we balanced the risk of causing distress through the receipt of unrequested information against the potential for cancer prevention. In considering these factors, we drew on prior experience of acceptability of genetic testing information within families.^{26,27} In addition, a previous Australian study using immunohistochemistry as a surrogate marker of Lynch syndrome among unconsented patients with colorectal cancer was informative, as a majority of recipients found the unsolicited information valuable.²⁸ We further mitigated the potential of harm caused by the delivery of unwanted information through graded feedback of findings involving a GC,²⁹ designed to highlight the existence of important health




FIG 4. Identification of recipients for research results. A stepwise representation of the feedback of PVs to date: (A) outcome of the feedback process to the NOK of deceased probands who self-referred into the study and (B) outcome for feedback to 77 NOKs of deceased probands ascertained via clinic-based recruitment and existing research studies. Created using BioRender. GC, Genetic clinician; NOK-R, next of kin–referred; PV, pathogenic variant.

information but at the same time providing the recipient some control of how much and what information is disclosed to them.

A positive reception of the research results by family members is critical for the success of TRACEBACK programs. NOKs of deceased probands who self-referred were, as expected, highly engaged with the project, irrespective of their sex, and all accepted referral to an FCC

after notification of clinically relevant results by the TRACEBACK GC. To date, where the TRACEBACK research result was new, 55% (12 of 22) accepted a referral. This favorable rate occurred at a time of added disruption to health services and increased complexity of daily life for many individuals because of the COVID-19 pandemic.

When contacted unexpectedly with new, and unsought, genetic information, female NOKs were more likely to

TABLE 2. TRACEBACK Pilot Program Key Learning Points and Recommendations

Learning points
General
Probands can be ascertained through research cohort studies, referral (by self or family members), and gynecologic clinics
Archival FFPE surgical tissue can be sequenced with a gene panel using capture technology
Germline status can be evaluated by examining DNA extracted from normal tissue isolated in archival surgical samples
Archival FFPE surgical tissues are legally required to be kept for a minimum 10-15 years depending on the Australian state. There is variability in this practice across the country, and many pathology laboratories only keep samples for the minimum required time before discarding
The frequency of pathogenic variants is lower in women age over 70 years and in those diagnosed after the expansion of clinical genetic testing, arising from amended national testing guidelines and increased use of PARP inhibitors
Finding recipients for feedback and verifying that the details obtained are correct and current is labor-intensive and requires several approaches. These can include searching through medical records from clinics where the proband was treated, reaching out to individual treating clinicians known to the proband/family, use of online obituary searches, telephone/address directories, and searches for proband's/NOK's online presence
Considerations when the proband is deceased
Female NOKs contacted with unsolicited and new health information are more likely to accept an FCC/genetics referral than male NOK
Male NOK may benefit from targeted approaches to results notification
When it is identified that the family harbors a pathogenic germline variant, the majority of individuals contacted will accept referral to an FCC, even when the referral is unexpected and unsolicited
A psychosocial investigation into barriers of referral uptake will be informative for future Traceback projects
Considerations when the proband is living
Living probands are more likely to accept an FCC/genetics referral than a NOK; however, the number of untested living probands in the Australian population is relatively low
Recommendations
Use a multimodal approach to proband ascertainment to maximize those tested and to include individuals who might not have previously had an opportunity to participate in research cohort studies or clinical trials
Consider institutional, state, and national policies related to waiver of consent
Take into account logistics, what information is recorded/available in medical records, and how long is this information kept
Where possible use a clinician known to the family, geneticist, and/or experienced genetic counselors to deliver unexpected and unsolicited genetic information, both for the handling of sensitive information and to engage the attention of the NOK to increase the uptake of subsequent testing by family members
Attempt to identify female NOK as recipients of genetic information, especially for cancer conditions that predominately affect females
Future TRACEBACK studies in a research setting should use the clinical approach targeting the years 2000-2016
Consideration should be given to funding clinical genetic testing of a deceased proband for self-referring NOK through health institutions/government or insurance bodies

Abbreviations: FCC, Familial Cancer Clinic; FFPE, formalin-fixed paraffin-embedded; GC, Genetic Clinician; NOK, Next of Kin; PARP, poly (ADP-ribose) polymerase.

accept a FCC referral 63% (5 of 8) compared with male NOK 53% (7 of 14). This is consistent with other studies that demonstrate that men are generally less engaged with genetic research and less likely to have testing.^{30,31}

Further work is planned to understand the factors influencing decision making around the uptake of research-generated genetic information and exploring steps that family members take to reduce their risk. The success of such genetic testing programs, particularly those focused on cancer syndromes that are primarily recognized as affecting females, may be improved by focusing on female family members as the preferred recipient of clinical follow-up. The delivery of unexpected genetic information years after the death of a family member can be confronting, and considerable care is required. The outcome of a potentially difficult interaction may be more successful when conducted by a clinician who is known and trusted by the family.

The molecular findings are similar to previous investigations in HGNMOC,¹⁰ supporting our technical approach. The frequency of *BRCA1/2* PVs in the TRACEBACK cohort is at the lower end of that in other population-based data sets,^{8,10} likely because of an ascertainment bias that excluded probands who had previous genetic testing. For technical reasons, our analysis to date has not included exploration of copy number variants that are expected to contribute a small number of additional PVs.^{8,32} The majority of cohort-ascertained probands had stored blood samples that were able to be accessed for testing purposes and, as expected, performed well in the assay. In the absence of blood-derived DNA, 93% of DNA samples derived from the use of non-neoplastic FFPE tissue passed our quality control measures (Data Supplement). Despite pathologic review and efforts to enrich for non-neoplastic tissue, PV detected and validated in DNA extracted from tumor tissue may indeed have somatic changes detected in contaminating tumor cells. Such results were nevertheless deemed notifiable by the variant review committee and were useful for relatives, who could then undergo funded personal predictive testing.

How far into the past should probands be ascertained for testing? We were successful in achieving a result in the majority of sequenced cases (96%), irrespective of the age of the tissue. However, in general, the longer it has been since a patient has been diagnosed, the more difficult it is to find a NOK and the greater the chance that new cancers have already arisen in the family and genetic risk has become apparent. In this pilot study, we used a nominal cutoff of 17 years and identified 84 at-risk families, in which the PV represented new health information for approximately 80% (66 of 84). Assuming a consistent outcome across the remaining untested Australian population, with upscaling, we estimate that up to 500 more PVs could be found with an expanded program in this country. Given that guidelines for genetic testing in ovarian cancer in most

Western countries have been broadened only within the past decade, there remains a window of opportunity to reduce the risk of cancers in at-risk family members. With a high level of engagement of NOK-referral individuals, consideration should be given to obtaining access to subsidized clinical testing of archival cancer tissue for this

group. We note that the approach used here may also be applied to other cancer types, including triple-negative breast cancer, which has a high rate of *BRCA1/2* germline PVs, and more broadly to solid organ malignancies with a strong association between histology and the presence of inherited susceptibility genes.

AFFILIATIONS

- ¹Peter MacCallum Cancer Centre, Parkville, Victoria, Australia
- ²Sir Peter MacCallum Department of Oncology, The University of Melbourne, Victoria, Australia
- ³Ovarian Cancer Australia, Melbourne, Victoria, Australia
- ⁴Prince of Wales Hospital, Randwick, New South Wales, Australia
- ⁵Royal Hospital for Women, Randwick, New South Wales Australia
- ⁶Parkville Familial Cancer Centre, Peter MacCallum Cancer Centre and Royal Melbourne Hospital, Parkville, Victoria, Australia
- ⁷QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia
- ⁸Department of Pathology, Royal Melbourne Hospital, Parkville, Victoria, Australia
- ⁹Department of Clinical Pathology, University of Melbourne, Parkville, Victoria, Australia
- ¹⁰The Royal Women's Hospital, Melbourne Victoria, Australia
- ¹¹The Royal Children's Hospital, Melbourne Victoria, Australia
- ¹²Department of Gynecological Oncology, St John of God Hospital, Subiaco, Western Australia, Australia
- ¹³Genetic Services of Western Australia, Perth, Western Australia, Australia
- ¹⁴Prince of Wales Clinical School, Faculty of Medicine, University of New South Wales, Sydney, Australia
- ¹⁵Prince of Wales Hereditary Cancer Centre, Prince of Wales Hospital Randwick, New South Wales, Australia
- ¹⁶Hudson Institute of Medical Research, Clayton, Victoria, Australia
- ¹⁷Department of Molecular and Translational Sciences, Monash University, Clayton, Victoria, Australia
- ¹⁸School of Health and Biomedical Sciences, RMIT University, Melbourne, Australia
- ¹⁹Monash Health Familial Cancer Centre, Clayton, Victoria, Australia
- ²⁰Monash Health, Bentleigh East, Melbourne, Australia
- ²¹PathWest, King Edward Memorial Hospital, Subiaco, Western Australia, Australia
- ²²Clinipath Pathology, Osborne Park, Western Australia, Australia
- ²³King Edward Memorial Hospital, Perth, Western Australia, Australia
- ²⁴Faculty of Health and Medical Sciences, University of Western Australia, Perth Australia
- ²⁵Kinghorn Centre for Clinical Genomics, Garvan Institute of Medical Research Darlinghurst, New South Wales, Australia
- ²⁶School of Nursing, Queensland University of Technology, Brisbane, Queensland, Australia
- ²⁷Consumer Advocate, Melbourne, Victoria, Australia
- ²⁸Division of Obstetrics and Gynaecology, Faculty of Health and Medical Sciences, University of Western Australia, Perth, Australia

CORRESPONDING AUTHOR

Kathryn Alsop, PhD, Peter MacCallum Cancer Centre, 305 Grattan St, Melbourne, Victoria 3000, Australia; Twitter: @DrKathAlsop; e-mail: kathryn.alsop@petermac.org.

PRIOR PRESENTATION

Presented in part at World Ovarian Cancer Coalition Initiative: Global Ovarian Cancer Charter Summit Series, online event, November 21, 2021; Agilent Precision Oncology Summit 2021: Towards a Meaningful Clinical Outcome: Discovery and Applications of NGS Singapore, online

event, May 21, 2021; Familial Aspects of Cancer: Research and Practice, Kingscliff, NSW, Australia, August 30-September 1, 2017; poster, September 5, 2018, August 30, 2019, September 2, 2021; Australian and New Zealand Gynecological Oncology Group Annual Scientific Meeting, concept only, Brisbane, Queensland, Australia, April 5, 2018; and The Seventh International Symposium on Hereditary Breast and Ovarian Cancer, poster, Montréal, Québec, Canada, May 8-11, 2018. The basis for the study was published in this journal: "Traceback: A Proposed Framework to Increase Identification and Genetic Counseling of BRCA1 and BRCA2 Mutation Carriers Through Family-Based Outreach"; *J Clin Oncol* 35:2329-2337, 2017.

SUPPORT

Supported by funding from the Australian Government under the Public Health and Chronic Disease Grant Program, The National Health and Medical Research Council/Cancer Australia Postgraduate Scholarship (R.D.: APP1133906), The Donald and Joan Wilson Foundation, and St John of God Subiaco Hospital.

Personnel received additional support from the National Health and Medical Research Council of Australia: APP1117044 (D.B.), APP1092856 (D.B. and M.F.), GNT1173346 (P.M.W.), APP177524 (A.B.S. and M.T.P.), APP1186505 (D.W.G.), and APP1161198 (E.C.). D.W.G. was supported by US Army Medical Research and Materiel Command grant W81XWH-16-2-0010. S.L. was supported by an Australian Government Research Training Program Scholarship and a Peter MacCallum Cancer Foundation Postgraduate Award.

The Australian Ovarian Cancer Study was supported by the US Army Medical Research and Materiel Command under DAMD17-01-1-0729, The Cancer Council Victoria, Queensland Cancer Fund, The Cancer Council New South Wales, The Cancer Council South Australia, The Cancer Council Tasmania, The Cancer Foundation of Western Australia (Multi-State Applications 191, 211, and 182), and the National Health and Medical Research Council of Australia (NHMRC; ID199600, ID400413, and ID400281). The OPAL Study was funded by the National Health and Medical Research Council (NHMRC) of Australia (GNT1025142 and GNT1073898) and Brisbane Women's Club.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.21.02108>.

AUTHOR CONTRIBUTIONS

Conception and design: Rachel Delahunty, Sian Fereday, Alexandra Lewis, Mary-Anne Young, Louisa Gordon, Laura Forrest, Karen Livingstone, Jane Hill, Georgia Chenevix-Trench, Paul A. Cohen, Michael Friedlander, Paul James, David Bowtell, Kathryn Alsop

Financial support: Stephen B. Fox, Jane Hill, David Bowtell

Administrative support: Rachel Delahunty, Sian Fereday, Lesley Andrews, Sanela Bilic, Emma Newton, Christie Norris, Karen Livingstone, Jane Hill, Paul James, David Bowtell

Provision of study materials or patients: Rachel Delahunty, Stuart Craig, Sian Fereday, Lesley Andrews, Nadia Traficante, Xiao Qing Chen, Sanela Bilic, Yeh Chen Lee, Susan Gerty, Lynette Gill, Julie Howard, Andrew N.

Stephens, Courtney Smyth, Stephen B. Fox, Nicholas Pachter, Jane Hill, Penelope M. Webb, Michael Friedlander, Paul James, David Bowtell

Collection and assembly of data: Rachel Delahunty, Stuart Craig, Belinda Creighton, Dinuka Ariyaratne, Sian Fereday, Lesley Andrews, Alexandra Lewis, Sharne Limb, Joy Hendley, Nadia Traficante, Natalia Carvajal, Mila Volcheck, Timothy Semple, Gisela Mir Arnau, Xiao Qing Chen, Anna Marsh, Christopher Love, Sanela Bilic, Maria Beilin, Cassandra B. Nichols, Christina Greer, Yeh Chen Lee, Susan Gerty, Lynette Gill, Emma Newton, Julie Howard, Rachel Williams, Christie Norris, Erin Tutty, Courtney Smyth, Shona O'Connell, Thomas Jobling, Colin J.R. Stewart, Adeline Tan, Stephen B. Fox, Jason Ellul, Laura Forrest, Marion Harris, Penelope M. Webb, Michael Friedlander, Paul James, David Bowtell, Kathryn Alsop

Data analysis and interpretation: Rachel Delahunty, Stuart Craig, Dale W. Garsed, Elizabeth Christie, Sian Fereday, Lesley Andrews, Ahwan Pandey, Amanda B. Spurdle, Bryony Thompson, Michael T. Parsons, Victoria Beshay, Mila Volcheck, Richard Lupat, Kenneth Doig, Jian Yu, Yeh Chen Lee, Andrew N. Stephens, Thomas Jobling, Nicholas Pachter, Jason Li, Jason Ellul, Mary-Anne Young, Marion Harris, Paul A. Cohen, Paul James, David Bowtell, Kathryn Alsop

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

The authors would like to acknowledge and thank Karen Conte, Lisa Webbe, and Vanessa Alford (Ovarian Cancer Australia); Leanne Bowes, Jillian Hung, and Catherine Kennedy (AOCS); Suzanne O'Brien, Hélène O'Neill, and Merran Williams (OPAL); Cassandra Hoskins, Huiling Xu, Steven Macaskill, Anthony Bell, Andrew Fellowes, William Murray, and David Choong; The Bioinformatics, Molecular Pathology, and Molecular Genomics Core Facilities (Peter MacCallum Cancer Centre); Rhiannon Dudley (Hudson Institute of Medical Research); and Daniele Belluoccio (Agilent) and Angela Rutherford for their contributions to the study. The Australian Ovarian Cancer Study (AOCS) gratefully acknowledges support from Ovarian Cancer Australia and the Peter MacCallum Foundation. The AOCS and Ovarian Cancer Prognosis and Lifestyle Study (OPAL) acknowledge all the clinicians and participating institutions who helped make the studies possible. See www.aocstudy.org and opalstudy.qimrberghofer.edu.au for a complete list.

We would like to thank all those women with ovarian cancer and their families that contacted and/or participated in the TRACEBACK study and all women who participated in the ACS, AOCS, and OPAL studies.

REFERENCES

- Kuchenbaecker KB, Hopper JL, Barnes DR, et al: Risks of breast, ovarian, and contralateral breast cancer for BRCA1 and BRCA2 mutation carriers. *JAMA* 317:2402-2416, 2017
- Song H, Dicks E, Ramus SJ, et al: Contribution of germline mutations in the RAD51B, RAD51C, and RAD51D genes to ovarian cancer in the population. *J Clin Oncol* 33:2901-2907, 2015
- Rafnar T, Gudbjartsson DF, Sulem P, et al: Mutations in BRIP1 confer high risk of ovarian cancer. *Nat Genet* 43:1104-1107, 2011
- Yang X, Leslie G, Doroszuk A, et al: Cancer risks associated with germline PALB2 pathogenic variants: An international study of 524 families. *J Clin Oncol* 38:674-685, 2020
- Watson P, Vasen HFA, Mecklin JP, et al: The risk of extra-colonic, extra-endometrial cancer in the Lynch syndrome. *Int J Cancer* 123:444-449, 2008
- Moore K, Colombo N, Scambia G, et al: Maintenance olaparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med* 379:2495-2505, 2018
- Pujade-Lauraine E, Ledermann JA, Selle F, et al: Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): A double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol* 18:1274-1284, 2017
- Alsop K, Fereday S, Meldrum C, et al: BRCA mutation frequency and patterns of treatment response in BRCA mutation-positive women with ovarian cancer: A report from the Australian Ovarian Cancer Study Group. *J Clin Oncol* 30:2654-2663, 2012
- Konstantinopoulos PA, Norquist B, Lacchetti C, et al: Germline and somatic tumor testing in epithelial ovarian cancer: ASCO guideline. *J Clin Oncol* 38:1222-1245, 2020
- Norquist BM, Harrell MI, Brady MF, et al: Inherited mutations in women with ovarian carcinoma. *JAMA Oncol* 2:482-490, 2016
- Samimi G, Bernardini MQ, Brody LC, et al: Traceback: A proposed framework to increase identification and genetic counseling of BRCA1 and BRCA2 mutation carriers through family-based outreach. *J Clin Oncol* 35:2329-2337, 2017
- Schwartz MD: Identification of BRCA1 and BRCA2 mutation carriers through a Traceback framework: Consent, privacy, and autonomy. *J Clin Oncol* 35:2226-2228, 2017
- Doig KD, Fellowes A, Bell AH, et al: PathOS: A decision support system for reporting high throughput sequencing of cancers in clinical diagnostic laboratories. *Genome Med* 9:38, 2017
- Robinson JT, Thorvaldsdóttir H, Winckler W, et al: Integrative genomics viewer. *Nat Biotechnol* 29:24-26, 2011
- Thorvaldsdóttir H, Robinson JT, Mesirov JP: Integrative genomics viewer (IGV): High-performance genomics data visualization and exploration. *Brief Bioinform* 14:178-192, 2013
- Landrum MJ, Lee JM, Benson M, et al: ClinVar: Improving access to variant interpretations and supporting evidence. *Nucleic Acids Res* 46:D1062-d7, 2018
- Thompson BA, Spurdle AB, Plazzer JP, et al: Application of a 5-tiered scheme for standardized classification of 2,360 unique mismatch repair gene variants in the InSIGHT locus-specific database. *Nat Genet* 46:107-115, 2014
- Parsons MT, Tudini E, Li H, et al: Large scale multifactorial likelihood quantitative analysis of BRCA1 and BRCA2 variants: An ENIGMA resource to support clinical variant classification. *Hum Mutat* 40:1557-1578, 2019
- Richards S, Aziz N, Bale S, et al: Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 17:405-424, 2015
- Plon SE, Eccles DM, Easton D, et al: Sequence variant classification and reporting: Recommendations for improving the interpretation of cancer susceptibility genetic test results. *Hum Mutat* 29:1282-1291, 2008
- Australian Institute of Health and Welfare & National Breast and Ovarian Cancer Centre: Ovarian Cancer in Australia: An Overview. Australian Institute of Health and Welfare, Canberra, Australia, 2010
- Ovarian Cancer Statistics in Australia, Cancer Australia, Australian Government. Data sourced from the Australian Institute of Health and Welfare, 2020. <https://ovarian-cancer.cancer australia.gov.au/statistics>
- Cohen PA, Nichols CB, Schofield L, et al: Impact of clinical genetics attendance at a gynecologic oncology tumor board on referrals for genetic counseling and BRCA mutation testing. *Int J Gynecol Cancer* 26:892-897, 2016

24. Surveillance, Epidemiology, and End Results (SEER) Program: SEER* Explorer: Ovary; Survival Rated by Time Since Diagnosis 2000-2017—All Stages, Races and Ages. National Cancer Institute, DCCPS, Surveillance Research Program. <https://seer.cancer.gov/explorer>
 25. National Statement on Ethical Conduct in Human Research 2007, National Health and Medical Research Council, Reference number E72, ISBN 1864962755, 2018. <https://www.nhmrc.gov.au/about-us/publications/national-statement-ethical-conduct-human-research-2007-updated-2018>
 26. Hallowell N, Alsop K, Gleeson M, et al: The responses of research participants and their next of kin to receiving feedback of genetic test results following participation in the Australian Ovarian Cancer Study. *Genet Med* 15:458-465, 2013
 27. Crook A, Plunkett L, Forrest LE, et al: Connecting patients, researchers and clinical genetics services: The experiences of participants in the Australian Ovarian Cancer Study (AOCS). *Eur J Hum Genet* 23:152-158, 2015
 28. Zeps N, Iacopetta BJ, Schofield L, et al: Waiver of individual patient consent in research: When do potential benefits to the community outweigh private rights? *Med J Aust* 186:88-90, 2007
 29. Forrest LE, Young MA: Clinically significant germline mutations in cancer-causing genes identified through research studies should be offered to research participants by genetic counselors. *J Clin Oncol* 34:898-901, 2016
 30. Wakefield CE, Thorne H, Kirk J, et al: Improving mutation notification when new genetic information is identified in research: A trial of two strategies in familial breast cancer. *Genet Med* 15:187-194, 2013
 31. Gauna Cristaldo FB, Touzani R, Apostolidis T, et al: Uptake of genetic counseling among adult children of BRCA1/2 mutation carriers in France. *Psychooncology* 28:1894-1900, 2019
 32. Patch AM, Christie EL, Etemadmoghadam D, et al: Whole-genome characterization of chemoresistant ovarian cancer. *Nature* 521:489-494, 2015
-

ASCO® in Action

Your Source for Cancer Policy and Practice News

ASCO in Action provides the latest news and analysis on cancer policy and practice issues through a frequently updated newsfeed and biweekly newsletter. ASCO in Action provides key details for the cancer community on critical issues affecting the delivery of care, including federal funding for cancer research, the ongoing response to COVID-19, physician reimbursement, and more.

For more information, visit asco.org/ascoaction.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**TRACEBACK: Testing of Historical Tubo-Ovarian Cancer Patients for Hereditary Risk Genes as a Cancer Prevention Strategy in Family Members**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Belinda Creighton

Employment: Cancer Specialists 1/84 Bridge Rd, Richmond VIC 3121 Australia
Consulting or Advisory Role: Bristol Myers Squibb, Eisai

Dinuka Ariyaratne

Research Funding: AstraZeneca

Elizabeth Christie

Honoraria: AstraZeneca

Sian Fereday

Consulting or Advisory Role: Geneseq Biosciences Pty Ltd
Research Funding: AstraZeneca (Inst), AstraZeneca (Inst)

Nadia Traficante

Research Funding: AstraZeneca (Inst)

Bryony Thompson

Consulting or Advisory Role: Genetic Technologies

Kenneth Doig

Stock and Other Ownership Interests: CSL Limited

Christopher Love

Employment: Geneseq Biosciences

Yeh Chen Lee

Consulting or Advisory Role: GlaxoSmithKline

Research Funding: AstraZeneca (Inst)

Travel, Accommodations, Expenses: AstraZeneca

Rachel Williams

Honoraria: AstraZeneca

Andrew N. Stephens

Consulting or Advisory Role: Invion Pty Ltd (Inst)

Research Funding: Invion Pty Ltd (Inst)

Adeline Tan

Employment: Sonic Healthcare

Stock and Other Ownership Interests: Sonic Healthcare

Stephen B. Fox

Consulting or Advisory Role: Novartis (Inst), BMS (Inst), AstraZeneca (Inst), MSD (Inst), Pfizer (Inst), Roche (Inst)

Research Funding: AstraZeneca (Inst), Amgen (Inst), Roche (Inst), BMS (Inst)

Expert Testimony: AstraZeneca (Inst), MSD (Inst)

Travel, Accommodations, Expenses: Ventana Medical Systems, AstraZeneca, MSD

Mary-Anne Young

Uncompensated Relationships: Illumina (Inst)

Jane Hill

Other Relationship: Medicines Australia

Paul A. Cohen

Employment: St John of God Healthcare

Stock and Other Ownership Interests: Clinic IQ

Honoraria: AstraZeneca

Consulting or Advisory Role: Clinic IQ

Research Funding: ANZGOG, St John of God Foundation

Penelope M. Webb

Research Funding: AstraZeneca (Inst)

Michael Friedlander

Honoraria: AstraZeneca, MSD, Lilly, Takeda, Novartis, GlaxoSmithKline

Consulting or Advisory Role: AstraZeneca, MSD, AbbVie, Lilly, Takeda, Novartis, GlaxoSmithKline

Speakers' Bureau: AstraZeneca, ACT Genomics

Research Funding: BeiGene (Inst), AstraZeneca (Inst), Novartis (Inst)

Travel, Accommodations, Expenses: AstraZeneca

David Bowtell

Honoraria: AstraZeneca

Consulting or Advisory Role: Exo Therapeutics

Research Funding: Roche/Genentech, AstraZeneca, BeiGene

Patents, Royalties, Other Intellectual Property: AstraZeneca Genentech Roche

Kathryn Alsop

Research Funding: AstraZeneca (Inst)

No other potential conflicts of interest were reported.