



Species Conservation Profile

Species conservation profile of the stenoendemic cave spider *Pimoa delphinica* (Araneae, Pimoidae) from the Varaita valley (NW-Italy)

Stefano Mammola^{‡,§}, Gustavo Hormiga[‡], Marco Isaia^{‡,§}

[‡] University of Turin, Department of Life Sciences and Systems Biology, Torino, Italy

[§] IUCN SSC Spider & Scorpion Specialist Group, Torino, Italy

| The George Washington University, Washington DC, United States of America

Corresponding author:

Academic editor: Pavel Stoev

Received: 15 Dec 2016 | Accepted: 13 Jan 2017 | Published: 19 Jan 2017

Citation: Mammola S, Hormiga G, Isaia M (2017) Species conservation profile of the stenoendemic cave spider *Pimoa delphinica* (Araneae, Pimoidae) from the Varaita valley (NW-Italy). Biodiversity Data Journal 5: e11509.

<https://doi.org/10.3897/BDJ.5.e11509>

Abstract

Pimoa delphinica Mammola, Hormiga & Isaia, 2016 is a troglophile araneoid spider endemic of the high Varaita valley (Western Alps, Province of Cuneo, NW Italy). In spite of relatively intense field research and examination of museum collections, the species is restricted to ten localities, thus showing a reduced extent of occurrence (EOO; 26 km²) and area of occupancy (AOO; 12 km²). Although the habitat and the subpopulations of *P. delphinica* are not currently threatened, the species is potentially exposed due to its extremely narrow geographic distribution range, subdivision in subpopulations and low dispersal capacity.

Keywords

Western Alps, IUCN, Cave, Military bunker, red list, troglophile species

Contributors

Giulio Gardini, Fulvio Gasparo, Alesandro Girodo, Giulia Marangoni, Paolo Pantini and Alessio Trotta contributed to provide data of occurrence and/or with fieldwork.

Pimoa delphinica Mammola, Hormiga & Isaia, 2016

Species information

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Arachnida	Araneae	Pimoidae

Taxonomic notes

Pimoa delphinica is a medium-sized spider (cephalothorax: ca. 4 mm, abdomen: ca. 7 mm), with slender legs and a brown-reddish coloration (Fig. 1). The species—previously misidentified with *P. rupicola* (Simon, 1884)—was described in 2016. It is readily distinguishable from the other species of European pimoids by morphological characters in male and female genitalia (cf. Mammola et al. 2016b).

Region for assessment:

- Global

Geographic range

Biogeographic realm:

- Palearctic

Countries:

- Italy

Map of records (image):

Fig. 2

Map of records (Google Earth):

Suppl. material 1



Figure 1.

Pimoa delphinica, female from the military bunker near Castelfelfino, Province of Cuneo, Italy.
[Photo credit: Alessandro Girodo]

a: Ventral habitus.

b: Dorsal habitus.




Figure 2.
Known distribution of *Pimoidelphinica*.

Basis of EOO and AOO: Known habitat extent

Basis (narrative)

During relatively intense speleological research studies in the Western Alps, we collected *Pimoidelphinica* in a few caves and military bunkers in the municipalities of Casteldelfino, Sampeyre and Sant'Anna di Bellino (Varaita valley, Western Alps, Province of Cuneo, Italy). Specifically, the species occurred in four small wild caves and a few subterranean military bunkers in Casteldelfino and Sant'Anna di Bellino and an abandoned house in the hamlet of Becetto (Sampeyre). Two additional localities reported in Isaia et al. (2011) and Mammola et al. (2016b)—Buco del Nebin 1 [Speleological Cadastre: Pi 1158] and Buco del Nebin 2 [Pi 1159] caves—are in need of verification, since only juvenile specimens were collected. However, even if confirmed, these additional localities would fall within the known extent of occurrence of the species (cf. Suppl. material 1). Despite our extensive study of *Pimoidelphinica* specimens from Italy and France—including material stored in Museums or private collections and original records—we have not been able to find additional records of *Pimoidelphinica* outside this small area (detail in Mammola et al. 2016b).

We used the verified occurrence records of the species to calculate the extent of occurrence (EOO) as the minimum convex hull and the area of occupancy (AOO) through a 2x2 km² grid, as implemented in the *red R* package (Cardoso 2016).

Min Elevation/Depth (m): 1230

Max Elevation/Depth (m): 2242

Range description

Pimoa delphinica is a stenoendemic species with a restricted Western Alpine distribution. The species is recorded in seven nearby localities, in the high Varaita valley. Its distribution range represents a small enclave (ca. 20 km²) within the range of distribution of the sister species *P. graphitica* Mammola, Hormiga & Isaia, 2016.

New occurrences

Materials

- a. scientificName: *Pimoa delphinica*; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Araneae; family: Pimoidae; genus: *Pimoa*; specificEpithet: *delphinica*; scientificNameAuthorship: Mammola, Hormiga, Isaia, 2016; country: Italy; stateProvince: Piedmont; county: CN; municipality: Casteldelfino; locality: Military bunker 1 near the road to Casteldelfino; verbatimElevation: 1280 m; minimumElevationInMeters: 1280; maximumElevationInMeters: 1280; locationRemarks: Abandoned military bunker (II World War); verbatimCoordinates: 44°35'07.21"N, 7°04'40.02"E; georeferenceProtocol: GPS; samplingProtocol: hand collected; eventDate: 12 Dec 2016; individualCount: 2; sex: female; lifeStage: adult, juvenile; recordedBy: Stefano Mammola, Alessandro Girodo; identifiedBy: Stefano Mammola, Marco Isaia; dateIdentified: 2016; basisOfRecord: PreservedSpecimen
- b. scientificName: *Pimoa delphinica*; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Araneae; family: Pimoidae; genus: *Pimoa*; specificEpithet: *delphinica*; scientificNameAuthorship: Mammola, Hormiga, Isaia, 2016; country: Italy; stateProvince: Piedmont; county: CN; municipality: Casteldelfino; locality: Military bunker 2 near the road to Casteldelfino; verbatimElevation: 1288 m; minimumElevationInMeters: 1288; maximumElevationInMeters: 1288; locationRemarks: Abandoned military bunker (II World War); verbatimCoordinates: 44°35'07.70"N, 7°04'40.15"E; georeferenceProtocol: GPS; samplingProtocol: hand collected; eventDate: 12 Dec 2016; individualCount: 3; sex: female; lifeStage: adult, juvenile; recordedBy: Stefano Mammola, Alessandro Girodo; identifiedBy: Stefano Mammola, Marco Isaia; dateIdentified: 2016; basisOfRecord: PreservedSpecimen
- c. scientificName: *Pimoa delphinica*; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Araneae; family: Pimoidae; genus: *Pimoa*; specificEpithet: *delphinica*; scientificNameAuthorship: Mammola, Hormiga, Isaia, 2016; country: Italy; stateProvince: Piedmont; county: CN; municipality: Casteldelfino; locality: Miniera di Casteldelfino, galleria Auriol (Borgata Ciampanesio); verbatimElevation: ca. 970 m; locationRemarks: Mineshaft; verbatimCoordinates: 44°35'10.3"N, 7°07'06.6"E; verbatimCoordinateSystem: WGS84; decimalLatitude: 44.5862; decimalLongitude: 7.1185; georeferenceProtocol: GPS; samplingProtocol: hand collected; eventDate: 16 Dec 2016; habitat: Subterranean; individualCount: 2; sex: females; lifeStage: adults; recordedBy: Alessandro Girodo, Paolo Bertacco; identifiedBy: Stefano Mammola, Marco Isaia; dateIdentified: 2016; basisOfRecord: PreservedSpecimen

Extent of occurrenceEOO (km²): 26

Trend: Unknown

Justification for trend

The species is troglophile (*sensu* Sket 2008), showing a preference for dark and moist habitats. Yet, the species exhibits a moderate plasticity in its ecological requirements, being able to colonize both cave and extra-cave environments, such as forests in high altitude habitats. The distribution range of *Pimoa delphinica* is enclosed within the range of the more widespread *P. graphitica*, which is apparently competing with the former thus limiting its expansion (Mammola et al. 2016b). The subterranean habitats colonized by *P. delphinica* are as yet not threatened by direct human activities. However, biogeographic studies and genetic data suggested that past climate change determined strong contraction in the distribution ranges of the species of alpine *Pimoa* (Mammola et al. 2016b, Mammola et al. 2015). It is thus plausible that anthropogenic climate change may determine reduction or habitat shift for this species. In order to confirm this hypothesis statistically, a deeper study of the detailed occurrence of this species in its distribution range is required.

Causes ceased?: Unknown

Causes understood?: Unknown

Causes reversible?: Unknown

Area of occupancyAOO (km²): 12

Trend: Unknown

Justification for trend

See paragraph "Extent of Occurrence".

Causes ceased?: Unknown

Causes understood?: Unknown

Causes reversible?: Unknown

Locations

Trend: Stable

Population

Number of individuals: Unknown.

Trend: Unknown

Justification for trend

No information about population size are currently available.

Causes ceased?: Unknown

Causes understood?: Unknown

Causes reversible?: Unknown

Population Information (Narrative)

A census of the population has never been attempted. According to our observations, populations are locally abundant. In two caves in which *Pimoa delphinica* was found in syntopy with the congeneric *P. graphitica*, mixed nuclear alleles between the two species have been found, indicating the existence of unidirectional introgression of males of *P. graphitica* into females of *P. delphinica* (Mammola et al. 2016b).

Subpopulations

Number of subpopulations: 2

Trend: Stable

Justification for trend

Examining the known range of distribution and taking into account habitat connectivity, it is possible to identify two subpopulations within the range. The first subpopulation includes the localities from Casteldelfino and the Bellino valley, which are more connected through alpine scree and larch woods, acting as potential route of dispersal. The other subpopulation is found on the other slope of the Varaita valley, in the nearby of the hamlet of Becetto (municipality of Sampeyre). The subpopulation are as yet not threatened.

Habitat

System: Terrestrial

Habitat specialist: Yes

Habitat (narrative)

The species primarily lives in the twilight zone of wild caves and other similar sheltered habitats, in high alpine environments (Fig. 3a). Healthy populations were also observed in artificial subterranean habitat (military bunkers and mines), offering suitable cool climatic conditions (Fig. 3b, c, d). Further individuals of *P. delphinica* were collected in pitfall traps placed within the rocky debris on the floor of the cellar of an abandoned cottage (Becetto, Sampeyre). During summertime, we observed juveniles—tentatively classified as *P. delphinica*—in an epigeal environment, near the locus typicus (Fig. 3c), in a larch (*Larix decidua*) wood. Accordingly, it is possible that juveniles of *P. delphinica* may be able to disperse through epigeal habitats under suitable climatic conditions. Extra-cave dispersal was also documented for the congeneric alpine species *P. graphitica* and *P. rupicola*, as justified by occasional catches of juveniles and males in pitfall traps placed in the leaf litter of broad-leaved woods at mid-altitudes (e.g., Isaia et al. 2015, Isaia et al. 2014, Jackson 1929, Mammola et al. 2015, Mammola et al. 2016b).



Figure 3.

Habitats of *Pimoa delphinica*. [Photo credit: a) Simone Spedicato; b-d) Alessandro Girodo]

a: The typical alpine environment in the high Varaita valley where the caves colonized by *Pimoa delphinica* are found.

b: An abandoned mineshaft colonized by the species.

c: A disused military bunkers from the Second World War, *locus typicus* of the species.

d: Military bunker, interior.

Trend in extent, area or quality?: Stable

Habitat importance: Major Importance

Habitats:

- 7. Caves and Subterranean Habitats (non-aquatic)
- 7.1. Caves and Subterranean Habitats (non-aquatic) - Caves
- 7.2. Caves and Subterranean Habitats (non-aquatic) - Other Subterranean Habitats

Habitat importance: Marginal

Habitats:

- 1. Forest
- 1.4. Forest - Temperate

Ecology

Size: Total length (leg excluded) = Male: 7 mm, Female: 10.5 mm

Generation length (yr): 1

Dependency of single sp?: No

Ecology and traits (narrative)

Little is known about the ecology of *Pimoa delphinica*. We report the result of our sporadic observations, which are not supported by specific studies or statistical inference. Like other *Pimoa* species (cf. Mammola et al. 2016a, Mammola et al. 2015, Hormiga 1994), *P. delphinica* exhibits a moderate ecological plasticity. In caves, it is found preferentially in the twilight zone. We observed adult males and females during the summertime. Adults display thanatotic behaviour when disturbed, possibly as a protection against predators (cf. Rogers and Simpson 2014, Novak et al. 2016). Depositions of cocoons occurs in July and cocoons are guarded by females. Females affix substrate particles to their cocoons (Mammola et al. 2016b).

Threats

Threat type: Future

Threats:

- 11.1. Climate change & severe weather - Habitat shifting & alteration
- 11.3. Climate change & severe weather - Temperature extremes

Justification for threats

See "Extent of occurrence".

Conservation

Conservation actions

Conservation action type: In Place

Conservation actions:

- 2.1. Land/water management - Site/area management

Conservation action type: Needed

Conservation actions:

- 4. Education & awareness
- 4.3. Education & awareness - Awareness & communications

Justification for conservation actions

A portion of the distribution range of *Pimoa delphinica* falls within the border of the Natural Park "Parco del Monviso".

The installation of information panels educating the visitors about this peculiar endemic species would positively increase the awareness of the caves as a natural heritage deserving protection.

Other

Research needed:

- 1. Research
- 1.2. Research - Population size, distribution & trends
- 1.3. Research - Life history & ecology
- 3. Monitoring
- 3.1. Monitoring - Population trends
- 3.4. Monitoring - Habitat trends

Acknowledgements

We thank Alessandro Girodo and Simone Spedicato for the photos.

References

- Cardoso P (2016) red: IUCN Redlisting Tools. 0.1.0. R. URL: <http://CRAN.R-project.org/package=red>
- Hormiga G (1994) A revision and cladistic analysis of the spider family Pimoidae (Araneioidea: Araneae). *Smithsonian Contributions to Zoology* 549: 1-104. <https://doi.org/10.5479/si.00810282.549>
- Isaia M, Paschetta M, Chiarle A (2015) Annotated checklist of the spiders (Arachnida, Araneae) of the Site of Community Importance and Special Area of Conservation "Alpi Marittime" (NW Italy). *Zoosystema* 37 (1): 57-114. <https://doi.org/10.5252/z2015n1a4>
- Isaia M, Paschetta M, Gobbi M, Zapparoli M, Chiarle A, Taglianti AV (2014) Stand maturity affects positively ground-dwelling arthropods in a protected beech forest. *Annals of Forest Science* 72 (4): 415-424. <https://doi.org/10.1007/s13595-014-0441-x>
- Isaia M, Paschetta M, Lana E, Pantini P, Schönhofer AL, Christian E, Badino G (2011) *Aracnidi sotterranei delle Alpi Occidentali italiane. [Subterranean Arachnids of the Western Italian Alps]*. XLVII. Museo Regionale di Scienze Naturali, Torino, 326 pp. [In Italian and English]. [ISBN 978-88-97189-08-4]
- Jackson AR (1929) A list of spiders found by Mr H. Donisthorpe at Bordighera in Northern Italy. *Entomologist's Record* 38: 26-28.
- Mammola S, Isaia M, Arnedo M (2015) Alpine endemic spiders shed light on the origin and evolution of subterranean species. *PeerJ* 3: e1384. <https://doi.org/10.7717/peerj.1384>
- Mammola S, Piano E, Isaia M (2016a) Step back! Niche dynamics in cave-dwelling predators. *Acta Oecologica* 75: 35-42. <https://doi.org/10.1016/j.actao.2016.06.011>
- Mammola S, Hormiga G, Arnedo MA, Isaia M (2016b) Unexpected diversity in the relictual European spiders of the genus *Pimoa* (Araneae, Pimoidae). *Invertebrate systematics* 30: 566-587. <https://doi.org/10.1071/IS16017>
- Novak T, Novak LS, Janžekovič F, Kozel P (2016) Hot-Dancing Method for Extracting Thanatotic Arachnids from a Substrate. *Entomological News* 126 (2): 121-127. <https://doi.org/10.3157/021.126.0207>
- Rogers S, Simpson S (2014) Thanatosis. *Current Biology* 24 (21): R1031-R1033. <https://doi.org/10.1016/j.cub.2014.08.051>
- Sket B (2008) Can we agree on an ecological classification of subterranean animals? *Journal of Natural History* 42: 1549-1563. <https://doi.org/10.1080/00222930801995762>

Supplementary material

Suppl. material 1: Extent of Occurrence of *Pimoa delphinica*

Authors: Mammola S., Hormiga G., Isaia M.

Data type: Geographic range

Filename: *Pimoa delphinica*.kml - [Download file](#) (675.00 bytes)