

1 **Revisiting the paradigm of anhematophagy in male mosquitoes**

2

3

4 Jovana Bozic<sup>1,2,3¶</sup>, Renuka E. Joseph<sup>1,2,3¶</sup>, Rachel S. Krizek<sup>2,3,4</sup>, Joshua B. Benoit<sup>5</sup>, and  
5 Jason L. Rasgon<sup>1,2,3,4\*</sup>

6

7

8 <sup>1</sup> Department of Entomology, The Pennsylvania State University, University Park, PA,  
9 USA

10

11 <sup>2</sup> The Center for Infectious Disease Dynamics, The Pennsylvania State University,  
12 University Park, PA, USA

13

14 <sup>3</sup> The Huck Institutes of the Life Sciences, The Pennsylvania State University, University  
15 Park, PA, USA

16

17 <sup>4</sup> Department of Biochemistry and Molecular Biology, The Pennsylvania State  
18 University, University Park, PA, USA

19

20 <sup>5</sup> Department of Biological Sciences, University of Cincinnati, Cincinnati, OH, USA

21

22

23

24 ¶ These authors contributed equally to this work

25

26 \* email for correspondence: [jlrs54@psu.edu](mailto:jlrs54@psu.edu)

27

28

29

30

31

32 **Abstract**

33 Female mosquitoes are reproductively obligate bloodfeeders which feed on vertebrate  
34 blood to obtain nutrients required for egg production (driving transmission of vector-  
35 borne pathogens in the process), and which rely on plant sugars for their non-  
36 reproductive energy requirements. Male mosquitoes, on the other hand, are thought to  
37 rely exclusively on plant sugars for their energetic needs; indeed, this dichotomy is one  
38 of the central tenets of medical entomology. Here, we show that male *Culex tarsalis* and  
39 *Aedes aegypti* mosquitoes will readily take blood from a membrane feeder when reared  
40 under dehydration conditions with no toxic effects. Mosquitoes with impaired humidity  
41 detection do not increase their bloodfeeding rates when dehydrated compared to wild-  
42 type controls. While conventionally reared males ignore a human host, dehydrated  
43 males are attracted to and attempt to probe, with some success, although they cannot  
44 access host capillaries. However, they will take blood from a vertebrate host wound.  
45 When fed a blood meal containing West Nile virus, male mosquitoes can become  
46 infected with and orally transmit the pathogen at rates and titers equivalent to females.  
47 These data suggest that under some circumstances male mosquitoes may be able to  
48 probe and/or ingest blood and transmit pathogens to vertebrate hosts, and that their role  
49 in maintaining pathogen transmission cycles should be re-examined.

50  
51 **Key words:** *Culex tarsalis*, *Aedes aegypti*, mosquito, bloodfeeding, sugar feeding,  
52 vector-borne disease

53  
54  
55  
56  
57  
58  
59  
60  
61  
62

## 63 **Introduction**

64 Female mosquitoes (except those which are autogenous) are reproductively obligate  
65 bloodfeeders, feeding on vertebrate blood to obtain nutrients required for egg  
66 production, and relying on plant sugars for their non-reproductive energy requirements.  
67 Male mosquitoes, on the other hand, rely exclusively on plant sugars for their energetic  
68 needs [1]. Indeed, this difference is one of the central tenants of medical entomology;  
69 female mosquitoes bloodfeed, males do not. Female bloodfeeding allows transmission  
70 of blood-borne pathogens, such as viruses or parasites, between vertebrate hosts,  
71 which is why the majority of mosquito research is performed on females rather than  
72 males; even when males are studied, it is usually within the context of how they affect  
73 females (mating behavior and fertility, pathogen transmission modulation etc...) [2-5].

74

75 Female mosquitoes are adapted to feed on blood, and this adaptation is reflected in the  
76 biology of their midgut, where transcripts related to blood digestion are enriched in the  
77 female compared to the male [6]; one would expect that male mosquitoes should not be  
78 attracted to blood as a nutrition source as they are thought to lack the proper physiology  
79 to digest and process it. However, there is one interesting report in the literature where  
80 male mosquitoes were attracted to and fed on blood. Nikbakhtzadeh and colleagues [7]  
81 documented bloodfeeding behavior in a laboratory colony of the mosquito *Culex*  
82 *quinquefasciatus*. When presented with defibrinated sheep blood on a cotton pledget  
83 (and to a much less efficient extent, a Parafilm membrane), male mosquitoes took a  
84 bloodmeal. However, blood was toxic to male mosquitoes, which died in a dose-  
85 dependent manner when blood was mixed with sugar [7], consistent with physiological  
86 adaptations to sugar vs bloodfeeding in males vs. females [6]. Interestingly, males did  
87 not show a preference for sugar compared to blood in a dual-choice assay [7], and the  
88 reason they fed on blood at all, particularly as it was toxic, remains an open question.  
89 As this is (to our knowledge) the only observation of male mosquito bloodfeeding  
90 behavior, it is difficult to speculate. However, there are a multitude of observations that  
91 male mosquitoes are attracted to human host odors and this behavior is suppressed by  
92 mosquito repellants [8], which includes species from arguably the three most important  
93 mosquito genera that act as disease vectors to humans (*Anopheles*, *Culex*, and *Aedes*).

94

95 Here, we present studies on male bloodfeeding behavior in the mosquitoes *Cx. tarsalis*  
96 *and Ae. aegypti*. *Cx. tarsalis* is one of the major West Nile virus (WNV) vectors in North  
97 America, where it is widely distributed across the Western United States [9]. It is  
98 genetically diverse, generally feeds on birds in the wild, and can be facultatively  
99 autogenous [9-10]. After becoming infected with WNV during a bloodfeeding event, it  
100 can also transmit the virus vertically to offspring at relatively high rates [11-12]. *Ae.*  
101 *aegypti* is one of the major invasive arbovirus vectors in the world [13]. We  
102 opportunistically observed *Cx. tarsalis* and *Ae. aegypti* males taking blood during  
103 unrelated laboratory studies, and undertook experiments to document and understand  
104 the behavior. We found that when dehydrated, male *Cx. tarsalis* and *Ae. aegypti* will  
105 predictably take human blood from a membrane feeder, determined the mechanism  
106 driving male bloodfeeding behavior, and present results of experiments examining  
107 potential for male mosquitoes to be involved in pathogen transmission cycles. These  
108 results are a paradigm shift in our understanding of male mosquito biology and suggest  
109 they may be more directly involved in pathogen transmission cycles than previously  
110 recognized.

111

## 112 **Methods**

113 Human subjects: All experiments with a human volunteer used the senior author (JLR)  
114 under PSU IRB Exempt Protocol STUDY00024284.

115

116 Mosquitoes: *Cx. tarsalis* strain (KNWR) and *Aedes aegypti* (Liverpool) were maintained  
117 at 25°C, 16:8 h light:dark diurnal cycle with 80% relative humidity, with 10% sucrose  
118 solution provided at all times through a cotton wick. For general rearing, mosquitoes  
119 were provided with expired anonymous human blood (BioIVT) through a water-jacketed  
120 glass membrane (Parafilm) feeder or a Hemotek feeder for egg development.

121

122 Male dehydration: To stimulate bloodfeeding, male mosquitoes were held at 25°C, 75%  
123 RH without sugar or water for 24 hours [14].

124

125 Survival analysis: Bloodfed male mosquitoes were isolated and placed into cup cages,  
126 held at the previously described standard insectary conditions, and provided with a  
127 cotton ball soaked in 10% sucrose solution. Control males were non-bloodfed and were  
128 maintained under the same conditions. Dead mosquitoes were counted every day and  
129 removed from the cages. Significant differences in survival between mosquito groups  
130 was determined with Kaplan-Meier analysis using GraphPad Prism version 9.0.4.

131  
132 Dehydration and bloodfeeding behavior in male *Cx. tarsalis*: *Cx. tarsalis* males were  
133 reared conventionally (80% RH, with free access to 10% sucrose solution in water), or  
134 under dehydrating conditions (75% RH, 25°C, with no access to water or sugar for 24  
135 hours) [14], then were offered a bloodmeal through a membrane feeder for 30 minutes.  
136 The number of fed and unfed mosquitoes at the end of the feeding period were counted.  
137 Data were analyzed by Fishers Exact test.

138  
139 Ionotropic receptor 93a (Ir93a) mutant mosquito assays: CRISPR protocols have not yet  
140 been developed for *Cx. tarsalis*, but we noted during experiments that males of the  
141 species *Ae. aegypti* (where CRISPR mutagenesis is routine) would also take blood from  
142 a membrane feeder, so we obtained an *Ae. aegypti* line that was a CRISPR knockout  
143 mutant for the Ir93a gene, which inhibits its ability to sense humidity [15]. The mutation  
144 was introgressed into the wild-type Liverpool background for comparison with Liverpool  
145 controls, and both lines reared as described above. For experiments, at 5-6 days post-  
146 emergence, males of each strain were transferred to 10 x 10 x 10 cm cages and  
147 deprived of sucrose and water (or held at normal conditions as controls) for 24 hours  
148 before being offered an anonymous human bloodmeal using an artificial feeding system  
149 (Hemotek). Bloodfeeding rates for each genotype and condition were recorded. Data  
150 were analyzed by Fishers Exact tests and confidence intervals calculated from the  
151 binomial distribution.

152  
153 Landing and probing experiments: Cages of 50 *Cx. tarsalis* males (reared under  
154 standard or dehydrating insectary conditions) were allowed to probe on the hand of the  
155 senior author for five minutes. Mosquito landings (defined as a mosquito alighting on the

156 volunteer hand for any period of time) and probing behavior (defined as exploring and  
157 probing with mouthparts) were counted during the 5 minute interval. The experiment  
158 was repeated 6 times. Data were analyzed by Mann-Whitney U test.

159

160 Host bloodfeeding by male mosquitoes: The senior author had an unrelated small  
161 (3mm) scratch on their hand from obtained from a pet cat a day earlier. A sterile razor  
162 blade was used to pick the scab off the scratch allowing a minimal amount of blood to  
163 be exposed. The wounded hand was placed in a cage of 20 dehydrated male *Cx.*  
164 *tarsalis* mosquitoes and their behavior recorded.

165

166 WNV feeds: Dehydrated *Cx. tarsalis* males and female controls were allowed access to  
167 an infectious blood meal consisting of a 1:1 mix of anonymous human blood (BioIVT)  
168 and  $5.0 \times 10^7$  FFU/ml (focus-forming units/ml) of WN02-1956 (GenBank: AY590222). A  
169 subset of male and females were processed immediately after feeding (“day zero”) to  
170 check for virus viability. Mosquito virus infection and transmission assays were  
171 performed at 7 and 14 days post-blood feeding. Fully engorged mosquitoes were sorted  
172 from non-fed ones for analysis. Mosquitoes were anesthetized with triethylamine  
173 (Sigma, St. Louis, MO), legs/wings from each mosquito were removed and placed  
174 separately in a 2-mL tube filled with 0.5 mL mosquito diluent (MD: 20% heat-inactivated  
175 fetal bovine serum (FBS) in Dulbecco’s phosphate-buffered saline, 50 µg/mL  
176 penicillin/streptomycin, 50 ug/mL gentamicin, and 2.5 µg/mL fungizone, with a sterile 2.0  
177 mm stainless steel bead (Next Advance, Inc. Innovative Lab Products for the Life  
178 Sciences). The proboscis of each mosquito was positioned in a tapered capillary tube  
179 containing approximately 10 µL of a 1:1 solution of 50% sucrose and FBS to induce  
180 salivation. After 30 min, the tube contents were expelled into 0.3 mL MD, and bodies  
181 were placed individually into a 2-mL tube filled with 0.5 mL MD and a stainless steel  
182 bead as described above. Mosquito bodies and legs/wings were homogenized for 30  
183 sec with TissueLyser (QIAGEN, Hilden, Germany) at 24 cycles/sec, followed by  
184 centrifugation for 1 min. Mosquito bodies, legs/wings, and salivary secretion samples  
185 were tested for live, infectious WNV using focus-forming assays (FFAs; see below).

186

187 WNV FFAs: WNV titers were quantified by FFA, which detects live, infectious virus.  
188 C6/36 cells were seeded into 96-well plates at a density of  $1 \times 10^5$  cells/well and  
189 incubated overnight at 28°C in complete RPMI medium without CO<sub>2</sub>. The next day,  
190 medium was removed from the wells. Samples from male and female bodies or  
191 legs/wings were serially diluted in a serum-free RPMI medium; saliva samples were  
192 undiluted. 30 µl of each sample was added in duplicate to the prepared C6/36 cells.  
193 Cells were incubated for 1 hour at 28°C without CO<sub>2</sub>, after which the inoculum was  
194 removed. 100 µl of RPMI containing 0.8% methylcellulose was added to limit viral  
195 spread. Infected cells were incubated for 48 hours at 28°C without CO<sub>2</sub>. At 48 hours  
196 post-infection, infected C6/36 cells were fixed with 50 µl 4% formaldehyde for 30  
197 minutes at room temperature (RT). Cells were washed, permeabilized with 0.2% triton-  
198 X, and blocked with 3% BSA. 30 µl monoclonal flavivirus antibody (Clone D1-4G2-4-15,  
199 BEI-resources, NR-50327) was added and incubated overnight at 4°C. After washing,  
200 30µl of fluorescent secondary antibody (1:1000 dilution; Goat anti-Mouse IgG (H+L)  
201 Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488, Invitrogen/Thermo  
202 Fischer, A-11029) was added and incubated overnight at 4°C. Cells were maintained in  
203 100 µl PBS to prevent drying. West Nile virus foci were imaged using a FITC filter on an  
204 Olympus BX41 microscope with a UPlanFI 4x objective and counted. Infection rate (IR)  
205 was defined as the proportion of mosquitoes exposed to virus that had WNV-positive  
206 bodies. Dissemination rate (DR) was defined as the proportion of mosquitoes with  
207 WNV-positive bodies that had WNV-positive legs/wings. Transmission rate (TR) was  
208 defined as the proportion of mosquitoes with WNV-positive legs/wings that had WNV-  
209 positive saliva. IR, DR, and TR were analyzed with Fisher's exact tests. Viral titers were  
210 analyzed using Mann-Whitney U tests.

211

212

## 213 **Results**

214 Bloodfeeding is not toxic to *Cx. tarsalis* males: While we were bloodfeeding during an  
215 experiment related to relative humidity (e.g. [16]), we noted incidentally that in addition  
216 to females, male mosquitoes were probing the membrane and were taking blood  
217 (Figure 1A,B). As this was a spontaneous occurrence, the total number of males in the



218 cage was not recorded but was on the order of 50-70 based on standard rearing  
219 practices in our lab. Out of this total, we isolated seven blood-engorged males. These  
220 males were placed into a survival cup and a survival experiment conducted, comparing  
221 their survival to 10 non-bloodfed males from the same initial cage. Although  
222 Nikbakhtzadeh et al. [7] demonstrated that blood was highly toxic to male *Cx.*  
223 *quinquefasciatus*, we did not observe any acute toxicity to blood in male *Cx. tarsalis*;  
224 indeed, survival in bloodfed males was marginally (although not statistically) higher than  
225 non-bloodfed (Figure 1C).

226

227 *Cx. tarsalis* male bloodfeeding is driven by dehydration: As we previously demonstrated  
228 that dehydration stimulates elevated bloodfeeding behavior in females [14, 16], we  
229 tested the hypothesis that dehydration was driving bloodfeeding behavior in males.  
230 Cages of male mosquitoes were reared under conventional insectary conditions or  
231 under dehydrating conditions [14]. No conventionally reared male mosquito (N = 64)  
232 took a bloodmeal from the membrane feeder, while 44/163 dehydrated males took a  
233 bloodmeal ( $P < 0.00001$ ).

234

235 Male mosquito bloodfeeding behavior is dependent on their ability to sense humidity:  
236 *Ae. aegypti* and *Anopheles gambiae* mosquitoes sense humidity through ionotropic  
237 receptor Ir93a, by which they locate oviposition sites, and CRISPR Ir93a knock-out  
238 mutants are impaired in this behavior [15]. We noted anecdotally that in our lab, male  
239 *Ae. aegypti* mosquitoes would also take blood from a membrane feeder, and as  
240 CRISPR protocols have not yet been developed for *Cx. tarsalis*, we used an Ir93a *Ae.*  
241 *aegypti* KO mutant for these assays [15]. When reared under standard insectary  
242 conditions, bloodfeeding rates did not differ statistically between wild-type and mutant  
243 mosquitoes. However, when reared under dehydrating conditions, bloodfeeding rates  
244 for the mutant did not increase, while wild-type mosquitoes had significantly elevated  
245 bloodfeeding behavior ( $P = 0.0284$ ) (Figure 2).

246

247 Dehydrated male mosquitoes will probe the hand of a human volunteer: The hand of a  
248 human volunteer was exposed to cages of conventionally reared or dehydrated male



249 *Cx. tarsalis*. Conventionally reared males showed little interest in the host, with  
250 infrequent landings that lasted less than 5 seconds. None demonstrated probing  
251 behavior. In contrast, dehydrated males landed significantly more often on the hand of  
252 the volunteer ( $P = 0.0065$ ), most landings lasted until the end of the time period, and  
253 probing behavior was observed in the majority of landings ( $P = 0.0022$ ) (Figure 3,  
254 Supplementary Videos 1 and 2). One dehydrated male mosquito (out of 6 separate  
255 trials) was able to lightly pierce the skin of the volunteer at the base of the wrist,  
256 although it was unable to reach the capillaries and acquire a bloodmeal (Supplementary  
257 Video 3). The bite resulted in a mild immunogenic reaction that disappeared after  
258 approximately 10 minutes (Supplementary Figure 1). To confirm that only males were in  
259 the cage, after the study was concluded the entire cage was killed by freezing and every  
260 mosquito visually examined for the presence of a female or a gynandromorph; only  
261 males were identified. While this is only a single observation and thus definitive  
262 conclusions cannot be drawn, to our knowledge, this is the first documented case of a  
263 male mosquito biting a vertebrate host.

264

265 Dehydrated male mosquitoes will take blood from a vertebrate host wound: Dehydrated  
266 male *Cx. tarsalis* show keen interest in probing a human host, but were not able to  
267 acquire blood, even from the single observed “successful” probing attempt. We  
268 hypothesized that if blood was made more accessible, male mosquitoes would take a  
269 bloodmeal. The senior author serendipitously had a small scratch on their hand  
270 (acquired from a pet cat a day earlier). The scab was peeled back using a sterile razor  
271 blade, exposing a small amount of blood. The volunteer placed their hand in a cage of  
272 20 dehydrated male mosquitoes. Males were attracted to the wound, and wound  
273 probing behavior was observed by 5 males (see Supplementary Video 4 for example).  
274 One male out of the 5 that probed fed and took a bloodmeal from the wound  
275 (Supplementary Video 5, Figure 4). At the conclusion of the experiment, the fed male  
276 was dissected to confirm the presence of blood in the gut (Figure 4).

277

278 Male *Cx. tarsalis* mosquitoes are competent vectors for West Nile virus: Since we  
279 determined that male *Cx. tarsalis* will probe a human hand or ingest blood from a

280 wound, allowing ingestion of a blood meal from a vertebrate host, we asked the  
281 question: can male mosquitoes become infected with and transmit arboviruses? We  
282 offered dehydrated male mosquitoes a bloodmeal spiked with WNV and assayed their  
283 vector competence at day 7 and day 14 post-exposure. Female *Cx. tarsalis* were  
284 exposed to virus at the same time as a control. We found that both female and male *Cx.*  
285 *tarsalis* were able to become infected with, disseminate, and orally transmit virus; males  
286 transmitted at both day 7 and 14, while females only had detectable virus in their saliva  
287 at day 14. After adjusting for multiple comparisons, infection rates (IR), dissemination  
288 rates (DR), and transmission rates (TR) did not differ statistically between males and  
289 females at either timepoint (Table 1).

290  
291 We quantitated all viral titers using an infectious virus assay. First, a subsample of  
292 males and females were assayed immediately after feeding ("day zero") to confirm virus  
293 viability. All fed males and females had detectable live infectious virus in their bodies,  
294 although females had statistically higher viral titers ( $P = 0.005$ ), likely because they  
295 could physically ingest a larger volume of blood. At day 7 post-exposure, viral titers  
296 were not statistically different between males and females in the bodies, the legs/wings,  
297 or the saliva (Figure 5A). At day 14 post-exposure, females had higher viral titers in their  
298 bodies ( $P = 0.001$ ) and legs/wings ( $P = 0.0083$ ) compared to males, suggesting either  
299 greater viral replication rates, or simply more tissue available for virus replication due to  
300 the larger size of the females. However, viral titers in saliva between males and females  
301 were statistically similar (Figure 5B).

302

303

## 304 **Discussion**

305 Previous work showed that blood was toxic to male *Cx. quinquefasciatus* mosquitoes  
306 [7], suggesting that in this species male bloodfeeding seems to be a maladaptive trait,  
307 perhaps a laboratory artifact. In our study, we demonstrate that males of other species  
308 (*Cx. tarsalis* and *Ae. aegypti*) can tolerate bloodfeeding, and that male bloodfeeding  
309 behavior is driven by water homeostasis during dehydration conditions. When  
310 mosquitoes cannot sense humidity due to *Ir93a* mutagenesis, dehydration does not

311 increase blood seeking behavior. These results are consistent with the role of  
312 dehydration on bloodfeeding behavior in female mosquitoes, where dehydration can  
313 stimulate females to increase their bloodfeeding rates as well [14, 16-17] and thus may  
314 reflect an adaptive trait where mosquitoes (female or male) can maximize their water  
315 intake during drought or periods of low relative humidity if other sources (nectar or free  
316 water) are not available.

317  
318 The mouthparts of male mosquitoes are thought to be physically incapable of  
319 penetrating vertebrate skin; however, in our experiments they were proven adequate to  
320 pierce a Parafilm membrane. Dehydrated *Cx. tarsalis* males were significantly attracted  
321 to and actively probed the hand of a human host, and one individual was even able to  
322 slightly penetrate the outer epidermis, leading to a transitory immune reaction  
323 (Supplementary Video 3 and Supplementary Figure 1). As the saliva of males differs  
324 from that of females, lacking various proteins needed for immunomodulation and  
325 bloodmeal acquisition [18], and it is likely that very little saliva was transferred compared  
326 to the bite of a female mosquito, it is not surprising that the host immune reaction was  
327 mild and rapidly resolving. When allowed access to a wound, dehydrated male  
328 mosquitoes readily probed the wound and one took a bloodmeal. As this experiment  
329 was facilitated by the fortuitous presence of a pre-existing wound on the hand of the  
330 senior author, it could not be deliberately repeated (as we were not allowed to make a  
331 deliberate wound due to IRB concerns). Still, it does suggest that male mosquitoes have  
332 the ability to take blood under specific rare circumstances that require dry periods and,  
333 likely, a host with a wound. According to fossil evidence, male mosquitoes are thought  
334 to once have had the ability to feed on vertebrate blood, and to have lost this ability over  
335 evolutionary time [19]. It is possible that the neural circuitry regulating host seeking and  
336 bloodfeeding behavior may still be conserved among male mosquitoes, or alternatively  
337 that this is simply a unique response to dehydration conditions in the lab.

338  
339 Interestingly our data demonstrate that, if male *Cx. tarsalis* orally acquire a WNV  
340 infection, they are competent vectors and transmit the virus at similar rates and titers  
341 compared to females. In our experiments we explicitly used an assay that quantified

342 live, infectious viral particles rather than quantitative PCR to rule out results that might  
343 be due to carryover of non-infectious viral RNA. Our results suggest that male *Cx.*  
344 *tarsalis* retain the receptors necessary for viral infection on their midgut, salivary glands,  
345 and other body tissues.

346

347 Finally, there is the question “is bloodfeeding behavior by male mosquitoes  
348 epidemiologically significant”? It is already known that male mosquitoes can be  
349 indirectly important for vector-borne disease transmission dynamics. For example,  
350 mating can affect key physiological parameters in females related to pathogen infection  
351 and transmission [2]. More directly, in some species, including *Cx. tarsalis* and WNV,  
352 male mosquitoes can be infected with arboviruses by vertical transmission from infected  
353 mothers [11, 20-21]. Infected males can also transmit some viruses venereally to  
354 females during mating where they can be transmitted to vertebrate hosts during feeding  
355 [20-21]. Consistent collection of males using host-derived attractants suggest that males  
356 are commonly found to move toward hosts [8], increasing the potential of male feeding  
357 on host-derived fluids under specific conditions (dry periods with a lack of sugar and  
358 water resources). Our study suggests the rare possibility of edge cases where male  
359 mosquitoes could be more directly implicated in virus transmission, where males  
360 undergoing dehydrating conditions (for example, during drought) acquire virus through  
361 vertical transmission from infected mothers or by feeding on an open wound of an  
362 infected vertebrate host, then transmit to a naïve host through feeding on an open  
363 wound or by probing the skin, as mosquitoes often transmit the bulk of virus when  
364 probing skin prior to actually taking a bloodmeal [22].

365

366 We must emphasize that while compelling, the results presented in this research are  
367 laboratory-based, and there is no peer-reviewed evidence of male mosquito  
368 bloodfeeding or pathogen transmission in nature (although we suspect that researchers  
369 have not rigorously looked for these phenomena). However, while arbovirus  
370 transmission by males is unlikely to be a major factor in driving disease dynamics,  
371 these data suggest that their canonical role as non-bloodfeeders needs to be re-  
372 examined and their contribution to pathogen transmission explicitly quantified,

373 particularly in light of recent vector-borne disease control strategies that rely on the  
374 mass release of male mosquitoes into natural populations [23-26].

375

## 376 **References**

- 377 1. Clements AN. The Biology of Mosquitoes, Volume 1. CABI Publishers; 1992.  
378
- 379 2. Dahalan FA, Churcher TS, Windbichler N, Lawniczak MKN. The male mosquito  
380 contribution towards malaria transmission: Mating influences the *Anopheles* female  
381 midgut transcriptome and increases female susceptibility to human malaria parasites.  
382 PLoS Pathog. 2019; 15:e1008063.  
383
- 384 3. Marcenac P, Shaw WR, Kakani EG, Mitchell SN, South A, Werling K, Marrogi E,  
385 Abernathy DG, Yerbanga RS, Dabiré RK, Diabaté A, Lefèvre T, Catteruccia F. A  
386 mating-induced reproductive gene promotes *Anopheles* tolerance to *Plasmodium*  
387 *falciparum* infection. PLoS Pathog. 2020;16:e1008908.  
388
- 389 4. Carraretto D, Soresinetti L, Rossi I, Malacrida AR, Gasperi G, Gomulski LM.  
390 Behavioural Responses of male *Aedes albopictus* to different volatile chemical  
391 compounds. Insects. 2022;13:290.  
392
- 393 5. Peng D, Kakani EG, Mamei E, Vidoudez C, Mitchell SN, Merrihew GE, MacCoss  
394 MJ, Adams K, Rinvee TA, Shaw WR, Catteruccia F. A male steroid controls female  
395 sexual behaviour in the malaria mosquito. Nature. 2022;608:93.  
396
- 397 6. Warr E, Aguilar R, Dong Y, Mahairaki V, Dimopoulos G. Spatial and sex-specific  
398 dissection of the *Anopheles gambiae* midgut transcriptome. BMC Genomics. 2007;8:37.  
399
- 400 7. Nikbakhtzadeh MR, Buss GK, Leal WS. Toxic effect of blood feeding in male  
401 mosquitoes. Front Physiol. 2016;7:4.  
402
- 403 8. Paris V, Hardy C, Hoffmann AA, Ross PA. How often are male mosquitoes  
404 attracted to humans? R Soc Open Sci. 2023 Oct 25;10(10):230921.  
405
- 406 9. Venkatesan M, Rasgon JL. Population genetic data suggest a role for mosquito-  
407 mediated dispersal of West Nile virus across the western United States. Mol Ecol.  
408 2010;19:157.  
409
- 410 10. Provost-Javier KN, Chen S, Rasgon JL. Vitellogenin gene expression in  
411 autogenous *Culex tarsalis*. Insect Mol Biol. 2010;19:423.  
412
- 413 11. Goddard LB, Roth AE, Reisen WK, Scott TW. Vertical transmission of West Nile  
414 Virus by three California *Culex* (Diptera: Culicidae) species. J Med Entomol.  
415 2003;40:743.  
416

- 417 12. Nelms BM, Fechter-Leggett E, Carroll BD, Macedo P, Kluh S, Reisen WK.  
418 Experimental and natural vertical transmission of West Nile virus by California *Culex*  
419 (Diptera: Culicidae) mosquitoes. *J Med Entomol.* 2013;50:371.  
420
- 421 13. Brady OJ, Hay SI. The Global Expansion of Dengue: How *Aedes aegypti*  
422 Mosquitoes Enabled the First Pandemic Arbovirus. *Annu Rev Entomol.* 2020 Jan  
423 7;65:191-208.  
424
- 425 14. Hagan RW, Didion EM, Rosselot AE, Holmes CJ, Siler SC, Rosendale AJ,  
426 Hendershot JM, Elliot KSB, Jennings EC, Nine GA, Perez PL, Rizlallah AE, Watanabe  
427 M, Romick-Rosendale LE, Xiao Y, Rasgon JL, Benoit JB. Dehydration prompts  
428 increased activity and blood feeding by mosquitoes. *Sci Rep.* 2018;8:6804.  
429
- 430 15. Laursen WJ, Budelli G, Tang R, Chang EC, Busby R, Shankar S, Gerber R, Greppi  
431 C, Albuquerque R, Garrity PA. Humidity sensors that alert mosquitoes to nearby hosts  
432 ad egg-laying sites. *Neuron.* 2023 Mar 15;111(6):874-887.e8.  
433
- 434 16. Manzano-Alvarez J, Terradas G, Holmes CJ, Benoit JB, Rasgon JL. Dehydration  
435 stress and Mayaro virus vector competence in *Aedes aegypti*. *J Virol.* 2023 Dec  
436 21;97(12):e0069523.  
437
- 438 17. Holmes CJ, Chakraborty S, Ajayi OM, Unran MR, Frigard RA, Stacey CL, Susanto  
439 EE, Chen SC, Rasgon JL, DeGennaro MJ, Xiao Y, Benoit JB. Multiple bouts of blood  
440 feeding in mosquitoes allow prolonged survival and are predicted to increase viral  
441 transmission during drought. *bioRxiv [Preprint].* 2024 Jun 22:2024.05.28.595907.  
442
- 443 18. Lu S, Martin-Martin I, Ribeiro JM, Calvo E. A deeper insight into the sialome of male  
444 and female *Ochlerotatus triseriatus* mosquitoes. *Insect Biochem Mol Biol.* 2022  
445 Aug;147:103800.  
446
- 447 19. Azar D, Nel A, Huang D, Engel MS. The earliest fossil mosquito. *Curr Biol.* 2023  
448 Dec 4;33(23):5240-5246.e2.  
449
- 450 20. Schopen S, Labuda M, Beaty B. Vertical and venereal transmission of California  
451 group viruses by *Aedes triseriatus* and *Culiseta inornata* mosquitoes. *Acta Virol.*  
452 1991:35:37.  
453
- 454 21. Patrican LA, DeFoliart GR. *Aedes triseriatus* and La Crosse virus: similar venereal  
455 infection rates in females given the first bloodmeal immediately before mating or several  
456 days after mating. *Am J Trop Med Hyg.* 1987 May;36(3):648-52  
457
- 458 22. Visser I, Koenraad CJM, Koopmans MPG, Rockx B. The significance of mosquito  
459 saliva in arbovirus transmission and pathogenesis in the vertebrate host. *One Health.*  
460 2023 Feb 12;16:100506.  
461



462 23. Bansal S, Lim JT, Chong CS, Dickens B, Ng Y, Deng L, Lee C, Tan LY, Kakani EG,  
463 Yoong Y, Du Yu D, Chain G, Ma P, Sim S, Ng LC, Tan CH. Effectiveness of *Wolbachia*-  
464 mediated sterility coupled with sterile insect technique to suppress adult *Aedes aegypti*  
465 populations in Singapore: a synthetic control study. *Lancet Planet Health*. 2024  
466 Sep;8(9):e617-e628.

467  
468 24. Carvalho DO, McKemey AR, Garziera L, Lacroix R, Donnelly CA, Alphey L,  
469 Malavasi A, Capurro ML. Suppression of a Field Population of *Aedes aegypti* in Brazil  
470 by Sustained Release of Transgenic Male Mosquitoes. *PLoS Negl Trop Dis*. 2015 Jul  
471 2;9(7):e0003864.

472  
473 25. Apte RA, Smidler AL, Pai JJ, Chow ML, Chen S, Mondal A, Sánchez C HM,  
474 Antoshechkin I, Marshall JM, Akbari OS. Eliminating malaria vectors with precision-  
475 guided sterile males. *Proc Natl Acad Sci U S A*. 2024 Jul 2;121(27):e2312456121.

476  
477 26. Rasgon JL. Precision-guided tools for malaria control. *Proc Natl Acad Sci U S A*.  
478 2024 Aug 6;121(32):e2411587121.

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498



499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528

**Declarations**

Ethics approval and consent to participate: All experiments with a human volunteer used the senior author (JLR) under PSU IRB Exempt Protocol STUDY00024284.

Acknowledgements: This research was supported by NIH/NIAID grant R01AI150251, USDA Hatch Project 4769, a grant with the Pennsylvania Department of Health using Tobacco Settlement Funds, and funds from the Dorothy Foehr Huck and J. Lloyd Huck endowment to JLR, and NIH/NIAID grant R01AI148551 to JBB and JLR. We thank Ms. Amelia Romo and Ms. Heather Engler for assistance with mosquito rearing and Paul Garrity for providing the transgenic mosquitoes used in this study (Ir93a).

Authors' contributions: JB, REJ, RSK, JBB, and JLR conducted the research, JBB contributed materials and reagents, JLR analyzed the data, JB, REJ, JBB, and JLR wrote the manuscript.

529

530

531

532

### 533 **Figure legends**

534 **Figure 1. Male *Cx. tarsalis* bloodfeeding behavior and survival.** A) mosquitoes  
535 congregating at and bloodfeeding from a paraffin membrane. Arrow points to male  
536 orienting toward the membrane. B) Blood-engorged male mosquito. Un-engorged male  
537 can be seen in-frame. C) Survival curve of bloodfed vs. non-bloodfed male *Cx. tarsalis*  
538 mosquitoes. No significant difference was observed between treatments.

539

540 **Figure 2. CRISPR deletion of *Ir93a* ablates male mosquito bloodfeeding behavior**  
541 **under dehydration conditions.** When reared conventionally, both wild-type and  
542 mutant *Ae. aegypti* exhibit baseline levels of bloodfeeding behavior. When reared under  
543 dehydration conditions, wild-type males significantly increase bloodfeeding behavior but  
544 humidity-insensitive mutant mosquitoes do not. Confidence intervals were calculated  
545 from the binomial distribution. WT = wild-type.

546

547 **Figure 3. Host probing behavior of dehydrated male mosquitoes.** A) Control cage  
548 of conventionally reared male *Cx. tarsalis*. Mosquitoes ignore the human host. B) Cage  
549 of dehydrated male *Cx. tarsalis*. Dehydrated males land on and probe the human host.  
550 C, D) Stills from Supplementary Videos 1 and 2 showing dehydrated male mosquito  
551 probing behavior. See videos for complete behavioral responses. E) Landing responses  
552 for dehydrated vs. conventionally reared (“standard”) male *Cx. tarsalis*. F) Probing  
553 behavior for dehydrated vs. conventionally reared (“standard”) male *Cx. tarsalis*. Error  
554 bars = SEM. \*\* =  $P < 0.01$

555

556 **Figure 4. Male *Cx. tarsalis* taking a human bloodmeal from a wound.** A) Male *Cx.*  
557 *tarsalis* feeding on an open wound. B) Close-up of feeding behavior. C) Blood is  
558 observable in the male mosquito gut. D) Blood in the male mosquito gut was confirmed  
559 by dissection. See Supplementary Video 3 for complete behavioral response.

560

561 **Figure 5. West Nile virus vector competence for male and female *Cx. tarsalis*.** A) 7  
562 days post-exposure. B) 14 days post-exposure. Red = bodies (infection); blue =  
563 legs/wings (dissemination); black = saliva (transmission). Males = closed circles,  
564 females = open circles. Zero values had 0.01 added purely for log-scale plotting  
565 purposes ( $10^{-2}$  = uninfected); analysis was performed on untransformed data. \*\* =  $P <$   
566 0.01.



567

568

Table 1. WNV Infection rate (IR), dissemination rate (DR) and transmission rate (TR) of dehydrated males and females at 7 and 14 days post-infection. No comparisons were statistically significant after correcting for multiple comparisons.

	N	IR	DR	TR
<b>Day 7</b>				
Males	43	0.53	0.70	0.56
Females	10	0.50	0.60	0.00
<b>Day 14</b>				
Males	14	0.50	0.86	0.50
Females	12	0.92	0.91	0.30

Figure 1



**Figure 2**





Figure 3





**Figure 4**






Figure 5



## Supplementary material

**Supplementary Figure 1. Host immune reaction to probing of dehydrated male *Culex tarsalis* mosquito.** A) Bite reaction 2 minutes post-probing (arrow). B) Immune reaction resolved by 10 minutes post-probing.



**Supplementary Video 1.** Probing behavior of dehydrated male *Cx. tarsalis* mosquito on the thumb of a human volunteer.

**Supplementary Video 2.** Probing behavior of dehydrated male *Cx. tarsalis* mosquito on the index finger of a human volunteer.

**Supplementary Video 3.** Probing behavior of dehydrated *Cx. tarsalis* male mosquito on the wrist of a human volunteer. This mosquito succeeded in slightly penetrating the outer epidermis (see Supplementary Figure 1).

**Supplementary Video 4.** Male *Cx. tarsalis* probing a human host wound.

**Supplementary Video 5.** Male *Cx. tarsalis* feeding from a human host wound.