

# Slumber in a cell: honeycomb used by honey bees for food, brood, heating... and sleeping

Barrett A. Klein<sup>1</sup> and M. Kathryn Busby<sup>2</sup>

<sup>1</sup> Biology Department, University of Wisconsin-La Crosse, La Crosse, WI, USA

<sup>2</sup> Graduate Interdisciplinary Program in Entomology and Insect Science, University of Arizona, Tucson, AZ, USA

## ABSTRACT

Sleep appears to play an important role in the lives of honey bees, but to understand how and why, it is essential to accurately identify sleep, and to know when and where it occurs. Viewing normally obscured honey bees in their nests would be necessary to calculate the total quantity and quality of sleep and sleep's relevance to the health and dynamics of a honey bee and its colony. Western honey bees (*Apis mellifera*) spend much of their time inside cells, and are visible only by the tips of their abdomens when viewed through the walls of an observation hive, or on frames pulled from a typical beehive. Prior studies have suggested that honey bees spend some of their time inside cells resting or sleeping, with ventilatory movements of the abdomen serving as a telltale sign distinguishing sleep from other behaviors. Bouts of abdominal pulses broken by extended pauses (discontinuous ventilation) in an otherwise relatively immobile bee appears to indicate sleep. Can viewing the tips of abdomens consistently and predictably indicate what is happening with the rest of a bee's body when inserted deep inside a honeycomb cell? To distinguish a sleeping bee from a bee maintaining cells, eating, or heating developing brood, we used a miniature observation hive with slices of honeycomb turned in cross-section, and filmed the exposed cells with an infrared-sensitive video camera and a thermal camera. Thermal imaging helped us identify heating bees, but simply observing ventilatory movements, as well as larger motions of the posterior tip of a bee's abdomen was sufficient to noninvasively and predictably distinguish heating and sleeping inside comb cells. Neither behavior is associated with large motions of the abdomen, but heating demands continuous (vs. discontinuous) ventilatory pulsing. Among the four behaviors observed inside cells, sleeping constituted 16.9% of observations. Accuracy of identifying sleep when restricted to viewing only the tip of an abdomen was 86.6%, and heating was 73.0%. Monitoring abdominal movements of honey bees offers anyone with a view of honeycomb the ability to more fully monitor when and where behaviors of interest are exhibited in a bustling nest.

Submitted 17 March 2020

Accepted 30 June 2020

Published 5 August 2020

Corresponding author

Barrett A. Klein,  
barrett@pupating.org

Academic editor

Joseph Gillespie

Additional Information and  
Declarations can be found on  
page 21

DOI 10.7717/peerj.9583

© Copyright

2020 Klein and Busby

Distributed under

Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Animal Behavior, Ecology, Entomology

**Keywords** Sleep, Honeycomb, Heating bees, Thermography, Discontinuous ventilation

## INTRODUCTION

Sleep is a behavior steeped in mystery, yet it appears to offer essential benefits (*Rattenborg et al., 2007; Cirelli & Tononi, 2008*). Sleep may specifically assist with honey bee communication (*Klein et al., 2010, 2018*) and memory (*Zwaka et al., 2015*), so accurately identifying sleep and knowing when and where it occurs is essential for further investigating sleep's role in honey bee ecology. To better understand sleep's benefits, or the detriments that come with sleep loss, it is essential to monitor sleep, including when it occurs in dark, hidden places. Several species of honey bees (*Apis* spp.) nest inside cavities, and all species of honey bees spend periods of their lives concealed inside honeycomb cells, within which much of the colony's behaviors occur and without which the colony would inevitably perish. Honeycomb is where honey bees store honey and pollen, rear brood, and where some appear to sleep. Young adult workers (callows/cell cleaners) appear to sleep almost exclusively inside cells (*Klein et al., 2008, 2014*), but workers spend less and less time sleeping inside cells as they age and change tasks, with foragers spending none of their time asleep inside cells (*Kaiser, 1988; Klein et al., 2008*). If significant periods of sleep occur within honeycomb cells, it would be wise to take inside-cell behavior into account.

Looking for sleep within the dark confines of a honey bee nest requires close examination or monitoring of bees, and modifying a nest to expose the inner workings of a colony has a long and curious history (*Crane, 1983*), including displaying comb by adding transparent glass jars to hives (*Kritsky, 2010*), and designing research-friendly observation hives (*Von Frisch, 1967; Seeley, 1995*). Observation hives increase visibility by encapsulating frames of honeycomb between panes of glass, and this innovation has led to revolutionary discoveries in animal behavior (*Von Frisch, 1967; Seeley, 2019*). *Lindauer (1952)* cleverly and patiently recorded the behaviors of individual honey bee workers in a modified observation hive that allowed him to observe activities within a subset of honeycomb cells. Lindauer made a point of recording when a bee was "Müssig" (idle), and concluded that time spent exhibiting this behavior, a portion of which occurred inside cells, far outweighed the time spent performing other tasks (e.g., bee #107: 68 h 53 min out of the total 176 h 45 min observed). He referred to bees seeking out undisturbed resting places, like empty or egg-containing cells, and spending long, calm periods inside these cells. Lindauer used an icon of a couch or bed to symbolize this behavior, despite including cleaning movements and other somewhat superficially immobile states in his calculations. We now know that a portion of this relatively immobile time in cells is devoted to heating brood in adjacent cells (*Kleinhenz et al., 2003*), but to what extent the remaining time is spent sleeping has never been rigorously established.

Few studies following Lindauer's observations have addressed immobility or potentially sleep-like states in cells. These include reports of honey bees exhibiting a resting state ("Ruhezustand", *Sakagami, 1953*), a "motionless" state (*Moore et al., 1998*), "rest" (*Kaiser, 1988*), or ventilatory signs indicative of sleep (*Sauer, Menna-Barreto & Kaiser, 1998; Kleinhenz et al., 2003*). Sleep is typically defined by several behavioral criteria, most important of which is an increased threshold of responsiveness to a stimulus. An animal with

an increased response threshold exhibits a specific posture during states of relative immobility that are reversible (Flanigan, Wilcox & Rechtschaffen, 1973). This suite of sleep signs is internally controlled (Tobler, 1985), meaning that if deprived of the state, the organism will respond with an increased expression of the behavior. Kaiser (1988) and Sauer, Herrmann & Kaiser (2004) confirmed that these coincident behavioral traits exist in honey bees.

Not all sleep signs can be observed simultaneously, so a dependable indicator of sleep would be of great value. Antennal immobility is a feature that has been used as a proxy for sleep (Eban-Rothschild & Bloch, 2008; Hussaini et al., 2009; Zwaka et al., 2015; Vázquez et al., 2020) because the amount of antennal immobility per unit time correlates with an increased response threshold (Kaiser, 1988). Another feature that covaries with antennal immobility (Sauer et al., 2003) and could, therefore, be an alternative proxy for sleep, is discontinuous ventilation. The honey bee's metasoma (hereafter referred to as "abdomen") moves in anterior-posterior pumping motions (pulses) at various rates and degrees of continuity. Easily observed extremes include "continuous" and "discontinuous" ventilation, in which the interim between anterior-posterior abdominal motions is consistently brief (continuous) or occasionally broken by extended pauses of at least 10 s (discontinuous; Kleinhenz et al., 2003). Honey bees exhibit continuous and discontinuous ventilation inside cells (Kleinhenz et al., 2003) and outside cells (Kaiser, 1988), suggesting higher and lower rates of respiration (Bailey, 1954). A discontinuously ventilating honey bee appears to almost always have a higher response threshold, a hallmark of sleep (B. Klein, 2020, in preparation). Because it can be difficult or impossible to gauge antennal movement in a nest, especially if a bee is inserted in a honeycomb cell, ventilatory activity holds promise as a more suitable indicator of sleep under natural or close-to-natural conditions. It is worth noting that antennal immobility may even be a misleading indicator of sleep because brood-incubating (heater) bees, which may appear to be asleep on the comb surface because of an absence of large body movements, also exhibit slow to no antennal movement (Bujok et al., 2002).

Kleinhenz et al. (2003) modified Lindauer's (1952) hive manipulation and used ventilatory rates, in part, to distinguish resting versus heating honey bees. We adopted this approach to peer at the undisturbed activities of worker bees in comb cells to see if what can be seen outside a cell (tip of abdomen; Fig. 1A) can serve as a reliable indication of what is going on with the rest of a bee's body inside the cell (Figs. 1B–1D). We hypothesized that ventilatory rate (continuous vs. discontinuous ventilation) and the presence/absence of larger movements of the abdomen can be used to predict behavior of bees inside cells. If correlations are robust between behavior of honey bees inside cells with behavior that is observable outside cells, someone observing only the posterior tips of honey bees when bees are inserted in honeycomb should be able to identify the bees' behaviors, including sleep.

## MATERIALS AND METHODS

We set up a small colony of honey bees in an observation hive with honeycomb positioned so that the interiors of some cells were visible. We recorded bees' behaviors inside the



**Figure 1** Visibility of honey bee workers deep inside cells in observation hives. (A) Clear view of posterior end of worker abdomen, center. This is the typical view of an adult bee inside a cell when removing frames from a beehive, or, in this case, when viewed through the glass window of an observation hive (Würzburg, Germany, 2006). (B–D) Unusually clear views of bees sleeping inside cells built on windows of an observation hive (USA, 2019). Sometimes bees construct comb against the glass of an observation hive, exposing the occasional bee's activities within comb cells. Photos by Barrett Klein.

Full-size  DOI: [10.7717/peerj.9583/fig-1](https://doi.org/10.7717/peerj.9583/fig-1)

visible cells using an infrared-sensitive camera and a thermographic camera, first by surveying all of the visible cells, then by zooming into and recording examples of behaviors for later analysis. We classified behaviors into four categories based on body movement, ventilatory rate and surface temperatures. To test the viability of identifying behaviors based solely on viewing the portion of a bee that is visible when honeycomb is exposed in a

more conventional hive, we asked naïve viewers to identify behaviors from a subset of the videos. The viewers used the same four behavioral categories, but the videos were modified so only the posteriors of the bees were visible. Their identifications, made under limited-visibility conditions, were compared to our identifications, which benefitted from careful examination of behavior visible only inside the cells, and surface temperatures visible using the thermographic camera.

### Study organisms and hive

We collected one queen, two frames of honeycomb, and 800–1,000 Carniolan worker honey bees (*A. mellifera carnica* Pollman, 1879) from a bee yard hive, with permission from Dr. Jürgen Tautz and the University of Würzburg (Würzburg, Germany; 49°46'47" N, 9°58'31" E). We cut out three sections of comb to fit within a honey bee mating cage (Begattungskästchen; see [Kleinhenz et al., 2003](#)) and cleaned out most of the cells along the edges to increase our likelihood of viewing visiting workers. The interiors of 93 empty cells and 22 cells with food were visible along the edges of the hive ([Fig. 2](#)). The sections of comb included brood cells, pollen (at least 10 cells), uncapped honey and empty cells. The comb slice on the left side of the hive contained only uncapped honey and empty cells. The middle slice contained 50 capped brood on the left side (nine were one-cell deep from the empty edge cells) and 27 capped brood on the right side (two were one-cell deep). The right slice contained 40 capped brood on the left side (two were one-cell deep) and 41 capped brood on the right side (two were edge cells and five were one-cell deep, with at least six uncapped larval cells toward the back of the comb). Twenty-one hours after inserting the queen, followed by the workers, we introduced 49 uniquely paint-marked callows using nontoxic, oil-based markers (Sharpie, Oak Brook, IL, USA). Marking did not noticeably affect temperature readings in preliminary tests. The intent of introducing individually-marked callows was to increase our likelihood of observing sleep inside cells, because young adults appear to sleep more inside cells than older adults ([Klein et al., 2008, 2014](#)). The callows had been incubated at 36 °C and collected within 24 h of emergence, marked on dorsal side of mesosoma (hereafter referred to as “thorax”) and abdomen, and placed in a small cage on top of the new hive, separated from the hive by a screen. After 5 h of callows being exposed to nest odor, the screen was removed and newly marked bees were accepted without any sign of aggression. Ultimately, only a subset of the data recorded came from these introduced, marked workers (see ventilatory and thermal methods, below).

The hive allowed for unrestricted access to the outdoors for the duration of the study (20–24 August 2008, with data collected from 23–24 August) via an entrance tunnel. The hive window was replaced with a sheet of transparent polypropylene giftwrap (pbsfactory, Artikel 00347, Rheinland-Pfalz, Germany) that remained in place for the entire study to allow for thermographic recordings. The ambient temperature of the small room was maintained high enough by using a space heater that insulation was not used to cover the hive during any portion of the short study. Diet was supplemented with honey and sugar water ad libitum.



**Figure 2** Observation hive with slices of honeycomb and exposed cells. (A) Still image from infrared video of inhabited hive, with hive entrance leading to tubular tunnel at lower left. (B) Slices of comb were held in place by nails inserted through wood frame and comb slices were aligned at natural distances from each other. (C) Comb on right is angled prior to study to show width and some of cell contents, including uncapped and capped brood. Bottom, back corner of each comb was removed to allow easy travel by workers through the hive. Photos by Barrett Klein. [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90\_img.jpg\) DOI: 10.7717/peerj.9583/fig-2](https://doi.org/10.7717/peerj.9583/fig-2)

**Table 1** Criteria used to define behaviors of honey bees when visible in comb cells, or when visibility was limited to posterior ends of abdomens. Honey bees were observed inside exposed comb cells (i.e., in cells on edge of comb, visible through plastic window), or visibility was limited by digitally obscuring cell interiors (for testing predictability of behavior from observations of abdominal tip alone; Fig. 4). Ventilatory movements appear as anterior-posterior abdominal pulses, occasionally consolidated into “bouts.” Continuous ventilation = respiratory pulses separated by <10 s of immobility (rarely, if ever, >10 s); discontinuous ventilation = respiratory pulses in bouts, separated by >10 s of immobility (Kleinhenz *et al.*, 2003). Sample size refers to bees for which we collected respiratory rate data.


Behavior	Criteria used to identify behavior when cell interior was visible	Criteria used to identify behavior when cell interior was digitally obscured (test videos)	<i>n</i> (bees)
Sleeping	Discontinuously ventilating, otherwise relatively immobile (Klein <i>et al.</i> , 2008)	Discontinuously ventilating, otherwise immobile	12
Maintaining cells	Body active in empty cell, often obscuring continuous ventilation; mandibular or antennal movement commonly observed (i.e., cleaning or building cells)	Continuously ventilating, often coupled with larger body movements (in and out, or rotating in cell; Sakagami, 1953)	10
Eating	Tongue extended in cell containing liquid; continuously ventilating	Continuously ventilating with possible body movement, and only partially in cell (cell contents prevent bee from going deeper)	3
Heating	Continuously ventilating, otherwise immobile while deep in cell; thorax obviously hotter than surroundings (when viewed using thermal camera)	Continuously ventilating, otherwise immobile	12

### Behaviors of interest

Four categories of behaviors were recorded: sleeping, maintaining cells, eating, and heating (Table 1). Sleeping was identified by a bee’s discontinuous ventilation and otherwise relative immobility (see description, above). Maintaining cells (i.e., cleaning or building) was identified by occasional large body movements, or obvious mandibular activity while continuously ventilating (although ventilation could be difficult to assess during large body movement episodes) in a cell devoid of food. Sakagami (1953) identified cleaners as externally quiet or irregularly moving, rotating once in a while in a cell. Eating was rarely observed, but obvious when it did occur; a continuously ventilating bee extended her tongue into a cell containing liquid. Heating was identified when a bee with a relatively hot thorax was deep in a cell, continuously ventilating and otherwise immobile. We have no data for bees packing pollen and, because we had no uncapped brood in exposed cells, we have no data involving development or direct tending of brood.

We conducted three sets of analyses: (1) We surveyed behaviors of bees visible inside cells across multiple time points, and after zooming in with the video camera to record exemplars of the different behaviors, we (2) analyzed a subset of the surveyed bees for ventilatory rates, then (3) used the thermography to measure surface temperatures associated with a subset of the bees that had been analyzed for ventilatory rates. By restricting thermal analyses to only those bees for which we acquired ventilatory rate data, we could test whether heating bees could be identified by ventilatory rates (and relative immobility) alone.

For survey data (Dataset S1 and S2), we scanned the cells with visible interiors ( $n = 115$ ) at 49 discrete time points, and recorded behavior for all bees inside cells. Surveys were separated by at least 10 min, and, because cell maintenance was so commonly observed (considered the default behavior when not explicitly announced by B.A.K.), surveys



**Figure 3** Still images from infrared-sensitive videos of behaviors, with all honey bees head-first inside cells. (A) Sleeping bee, center cell, is facing left, venter up. (B) Bee is eating, with mouthparts extended and body less fully inserted in cell; facing left, venter down. (C) Heating bee is facing left, venter facing observer (sideways). (D) Sleeping bee, center, is facing left, venter down, and is to be compared with heating bee, at right, facing right, dorsum facing observer (sideways). All other bees inside cells are maintaining (cleaning or building) cells, including two visible in (A) and three in (B), indicated by black circles superimposed on the middle of their thoraces. Contrast and brightness alterations of image serve to highlight sleeping, eating, and heating bees. See [Movies S1–S4](#), from which these images were taken. Images by Barrett Klein. [Full-size](#) [DOI: 10.7717/peerj.9583/fig-3](https://doi.org/10.7717/peerj.9583/fig-3)

sometimes started when a behavior other than cell maintenance was detected to ensure sampling of these other behaviors. Each survey involved examining every bee inserted at least partially inside comb cells for at least three to five seconds if obviously maintaining cells (i.e., cleaning or building), or eating, and for longer (>10 s, and sometimes for several minutes) if a worker appeared to be sleeping or heating (Table 1; Fig. 3; Movies S1–S4). Surveys stopped immediately after identifications of behaviors, or after a few minutes of close-up filming for subsequent ventilatory analysis. B.A.K. identified behaviors in real time and identified individually paint-marked bees by briefly shining a tiny white light on the abdomen. Each behavioral count represented a unique bee within each survey, but some bees were undoubtedly repeatedly measured across surveys.


We collected ventilatory rate data (Dataset S3) from a subset of the surveyed bees. Sleeping and heating bees are relatively immobile (no major head, wing, leg, or body movements), except for ventilatory motions of the abdomen, described above (for more details describing relative immobility during sleep, see Klein et al., 2008). Discontinuous ventilation, identified by bouts of abdominal pulses separated by pauses of stillness exceeding 10 s, occasionally included a single, isolated, apparently spontaneous abdominal



jerk during one of these pauses. We excluded a pulse (jerk) if isolated from other pulses by  $>5$  s before and after. We recorded ventilatory rates using JWatcher, an event recorder and analytical software package designed for study of behavior (version 1.0, <http://www.jwatcher.ucla.edu/>). Recording events with JWatcher entails pressing keys assigned to represent behaviors of interest on a keyboard, with event times automatically recorded. M.K.B. manually pressed one key in time with a pulsing abdomen replayed at  $0.3\times$  the normal speed, to increase accuracy and consistency of data transcription. Although we cannot be certain that each behavioral recording represents a unique bee, three steps were taken to increase the likelihood: (1) 12 of the 37 bees were individually marked, and individually-marked bees were analyzed only once; (2) some of the recordings of unmarked bees captured several unmarked bees concurrently, so each was unique within those recordings; and (3) surveys were separated by at least 10 min, and sometimes by several hours.

In addition to behaviors exhibited in exposed cells, we recorded mean surface temperature of a bee's thorax ( $T_{th}$ ) and mean surface temperature of her surroundings ( $T_{surr}$ ) using FLIR's analysis software package (ResearchIR Max version 4, FLIR Systems, Inc.) from a subset of the bees for which we analyzed ventilatory rates, above (Dataset S4). To calculate the mean temperature of a bee's thorax ( $T_{th}$ ), we drew a circle (within which a mean temperature could be automatically generated) over the region of interest (thorax) in an image taken at the beginning of a thermal recording (several seconds after entering cell), the middle of the recording, and the end (several seconds before exiting cell). To calculate the mean surface temperature of her surroundings ( $T_{surr}$ ) at identical time points, we dragged the same ellipse over three regions bordering the bee's thorax: above and below thorax, and anterior to head. These regions of interest surrounding the bee's thorax included almost exclusively cells and cell walls and, unlike a previous study by Klein *et al.* (2014), did not include any portion of the bee herself. This updated method avoids problems of the bee's body contributing to the measurement of  $T_{surr}$ . We report the difference of  $T_{surr}$  from  $T_{th}$  to indicate the surface temperature of the bee relative to the surface temperature of her surroundings ( $T_{diff} = T_{th} - T_{surr}$ ) (Klein *et al.*, 2014). There was no statistically meaningful difference between using the mean body temperature from the middle time point versus the mean of means across all three time points for any behavior, so we use the middle point when reporting  $T_{th}$  ( $W = 27, 52, 4, 89$ ;  $P = 0.65, 0.94, 1.00, 0.07$  for bees sleeping, maintaining cells, eating, and heating, respectively) and  $T_{diff}$  ( $W = 26, 52, 3, 69$ ;  $P = 0.58, 0.91, 0.70, 0.61$  for bees sleeping, maintaining cells, eating and heating, respectively). Because these bees represent a subset of the bees analyzed above (32 of the 37 bees for which we analyzed ventilatory rates; 11 of the 32 bees were individually marked), the same discussion of unique sampling applies here as well.

To test how predictable a behavior is from observing tips of abdomens alone, we first edited video clips so that they were without sound and a dark gray bar concealed cell contents, revealing only what extended beyond each cell (tip of abdomen and, sometimes, distal portions of hindlegs). We placed a tiny digital mark to indicate the bee(s) of interest in each video (Fig. 4; Movies S5–S8). B.A.K. trained 54 students for 20–30 min by showing



**Figure 4** Still image from infrared test video. Gray boxes obscure cell interiors, and small light gray rectangle within box on right marks bee of interest. Test yourself with [Movies S5–S8](#), a sample of video clips we used to test the reliability of identifying inside-cell behavior from what is visible outside the cell. (Answers included.) Photo by Barrett Klein. [Full-size](#) DOI: [10.7717/peerj.9583/fig-4](https://doi.org/10.7717/peerj.9583/fig-4)

and describing behaviors (criteria: [Table 1](#)) in 12 video clips (four sleeping, five maintaining cells, two heating, and one showing food in a cell; eating was described but not shown due to lack of additional examples from our recordings). Videos used during training were not used during testing, but were made available to students, had they wished to continue training on their own. Once trained, students independently watched 30 video clips of bees with digitally obscured cell contents—a subset of the 37 bee recordings used in our ventilatory rate analyses (11 sleeping, six maintaining cells, two eating and 11 heating)—and recorded what they believed to be each bee’s behavior.

### Recording equipment

We eliminated outdoor light and lit the room with a single desk lamp covered with a red acetate filter (#27 Medium Red, transparency = 4%, peak at 670 nm, Supergel by Rosco, Stamford, CT, USA), selected because honey bees may be less sensitive to frequencies beyond 600 nm ([Von Frisch & Lindauer, 1977](#)) or 650 nm ([Dustmann & Geffcken, 2000](#)). The same filter was applied to a headlamp, used to facilitate observations. The warm lights were kept away from the hive, and angled to minimize glare that would otherwise affect thermal measures. We filmed under the low, red light, and with an infrared spotlight by using an infrared-sensitive video camera (AGDVC 30, Panasonic, Japan) side-by-side with a thermal camera (FLIR SC660, FLIR Systems Inc., Boston, MA, USA; accuracy 1 °C or 1% of reading, according to FLIR manual and FLIR technical support). We adjusted thermal camera settings to match the emissivity value of a honey bee’s thorax (0.97; [Stabentheiner & Schmaranzer, 1987](#)), although wax and other surface temperatures were recorded for  $T_{\text{surr}}$ , and set the transmissivity to that of polypropylene (0.89).

The giftwrap used as the observation hive's window produced a nonlinear error when recording temperature as temperature increased, so we adjusted absolute temperature measurements (Dataset S5; see Klein et al., 2014 for details). Some data were taken using an audio recorder (Olympus VN-4100PC Digital Voice Recorder) and later transcribed. Audio was synchronized with video recordings by the researcher making a noise, followed by announcing the exact time as was recorded on video when the noise was made. Bees were often pointed out when announced, and this served to synchronize thermal imagery with video and audio.

## Statistical analysis

### *Behavior surveys*

To determine how prevalent each behavior was within the hive, we compared total counts of bees performing each behavior using a Kruskal–Wallis Rank Sum test. We then conducted post-hoc pairwise tests using six two-sided, non-paired Wilcoxon–Mann–Whitney tests. To avoid multiple testing problems, we corrected resulting *P*-values using the Holm method in the R function `p.adjust()`. To account for day/night differences between behaviors, we conducted three Kolmogorov–Smirnov tests using the R function `ks.test()`. This two-sided test's null hypothesis states that two sets of data, *x* and *y*, were drawn from the same continuous distribution. Therefore, we set our *x* and *y* to be the day and night distributions of each behavior, respectively. We performed three such tests to include sleeping, heating, and cell-maintaining behaviors, each time testing that each behavior count distribution remained the same from day to night. We did not perform this test on eating behavior because we did not have a large enough sample. We used local sunrise/sunset times to distinguish day and night (<https://www.gaisma.com/en/location/wurzburg.html>).

### *Ventilatory rates*

To address whether different behaviors could be distinguished using the time separations between their individual within-bout abdominal ventilation pulses (“pulse separations”), we performed a Kruskal–Wallis Rank Sum test. To test specifically for differences in pulse separations among behaviors, we filtered data to include only those abdominal pulses separated by <1 s, and repeated the aforementioned Kruskal–Wallis Rank Sum test. We then conducted pairwise Wilcoxon–Mann–Whitney tests comparing pulse separations across behavior groups. We applied a Holm correction for multiple testing. Since individual bees were monitored for different durations while performing behaviors inside cells, some bees could have disproportionately influenced the separation interval of the behavioral category to which they belonged. To account for any effect of individual bees on the timing between abdominal pulses, we performed a linear mixed effects logistic regression analysis using the R library `lme4` (in package `lmer` test), with bee ID as random factor and behavior as fixed effect. Because our residuals were initially not normally distributed, we performed a rank transformation before conducting the regression analysis.

### **Thermal measures**


We measured  $T_{th}$  and  $T_{surr}$  across three timepoints (beginning, middle and end of each bee's behavior duration). If we were to use all temperatures in our analyses, then we would have three  $T_{th}$  and nine  $T_{surr}$  per bee. If we were to use only the middle temperature measurements (which might avoid behaviorally transitional complications), then we would have one  $T_{th}$  and three  $T_{surr}$  per bee. To determine whether either method would affect behavior mean  $T_{th}$  or  $T_{diff}$ , we compared mean  $T_{th}$ , then mean  $T_{diff}$  of each behavior between the two methods using four Wilcoxon-Mann-Whitney tests. Corrections for multiple testing were not necessary. Heating behavior was confirmed by comparing a bee's thoracic temperature to that of the surrounding region. For temperature difference analyses, because the data were not normally distributed and the sample size was relatively small, we applied the Wilcoxon-Mann-Whitney test with Holm correction. To see if the temperature associated with behavior differed across behaviors, we applied a Kruskal-Wallis Chi-square test using the  $T_{th}$  and  $T_{surr}$  from the middle timepoint thermal measurements. We then conducted post-hoc pairwise Wilcoxon-Mann-Whitney tests with Holm correction on each of six combinations of behavior pairs. We repeated these methods for  $T_{diff}$ . To determine whether  $T_{diff}$  changed over the duration of a bee's behavior in a cell, we used the R package nparLD (for nonparametric longitudinal data; [Noguchi et al., 2012](#)) to conduct a non-parametric ANOVA-type test. We applied the formula F1-LD-F1, which tests for group (behavior) differences, change over time, and the interaction between group and time.  $T_{diff}$  was compared across temperature measurement periods 1, 2 and 3 for all bees, grouped by behavior. Before analyzing any thermal data, we corrected for the thermal signature of the thin film of giftwrap that functioned in enclosing the observation colony. To do this, we used thermographic measurements of the same neutral surface with and without the giftwrap film covering at a range of room temperatures from 26.5–43.6 °C. Differences between the two measurements at the same nominal temperatures were used to generate a set of correction values, which were then added as offsets to all thermal measurements of the colony behind the giftwrap ([Dataset S5](#)).

### **Limited-visibility test**

To calculate the reliability of identifying behavior based on observing only the posterior tip of a bee's abdomen, we applied a binomial test (Binomial Test Calculator, <https://www.socscistatistics.com/tests/binomial/default2.aspx>), with the null hypothesis that determination of behavior is random and not related to the actual correct behaviors. We then corrected for multiple testing using the Holm method. We set alpha at 0.05, report two-tailed  $P$ -values for all tests, and report errors as standard deviations. M.K.B. performed all statistical tests using R ([R Core Team, 2019](#)), except for binomial test on limited-visibility experiment.

## **RESULTS**

We conducted 49 surveys (21 nighttime, 28 daytime) of behaviors exhibited inside comb cells across 34.5 h. Absolute counts of each behavior differed across the surveys (Kruskal-Wallis rank sum test  $\chi^2 = 123.3$ ,  $df = 3$ ,  $P < 2.2 \times 10^{-16}$ ). Of the 455 behavioral




**Figure 5** Number of observations spent sleeping, maintaining cells, eating, or heating inside cells across time. Note that number of observations during any given survey ranged from 1 to 17 bees. Sleeping data are presented top-down so comparisons of number of occurrences can more easily be compared within this behavior across time. Two eating occurrences are presented here, but a third eating event was recorded for ventilatory and thermal data between surveys (Figs. 6 and 8).  
Full-size [DOI: 10.7717/peerj.9583/fig-5](https://doi.org/10.7717/peerj.9583/fig-5)

events monitored inside cells, bees spent 16.9% sleeping ( $n = 63$ ), 76.4% maintaining cells ( $n = 362$ ), 0.4% eating ( $n = 2$ ) and 6.4% heating ( $n = 28$ ). Bees slept for bouts of  $1316 \pm 1038$  s (range: 257–3,346 s), maintained cells for  $237 \pm 257$  s (range: 61–845 s), ate for  $447 \pm 233$  s (range: 197–659 s), and heated for  $956 \pm 509$  s (range: 452–2,214 s) ( $n = 7, 9, 3, 10$  recordings of entire duration in cell for each behavior category, respectively). Behaviors were exhibited day and night, with no evidence of day-night bias for any behavior (2-sample Kolmogorov–Smirnov tests,  $D = 0.13, 0.26, 0.09$ ;  $P = 0.99, 0.40, 1.00$  for sleeping, maintaining cells and heating, respectively; eating sample size was too low for test to be meaningful; Fig. 5). None of the discontinuously ventilating bees exhibited visible signs of wakeful activity (larger movements of body, antennal movement, chewing, etc.), and because discontinuous ventilation covaries with other sleep signs (see above), “sleeping” is used as a shorthand for discontinuous ventilation + relative immobility inside cells, below.

### Ventilatory signatures as indicators of behavior

Ventilatory patterns differed among behaviors exhibited inside cells, as evident when plotting abdominal pulses (Fig. 6), and time between pulses (Fig. 7) (Kruskal–Wallis rank




**Figure 6** Ventilatory pulses of abdomen over time by worker bees that were sleeping, maintaining cells, eating, or heating. Gray areas to the right of each pulse sequence signify post-recording periods (no data). Bees (y-axis) included 12 uniquely marked individuals (total  $n = 37$  bees; 12 sleeping, 10 cleaning, three eating, 12 heating). [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90\_img.jpg\) DOI: 10.7717/peerj.9583/fig-6](https://doi.org/10.7717/peerj.9583/fig-6)

sum test,  $\chi^2 = 185.2$ ,  $df = 3$ ,  $P = 2.2 \times 10^{-16}$ ). Discontinuous ventilation associated with sleep was identified by having discrete bouts of abdominal pulses, with the bouts separated by at least 10 s. Bouts of pulses were separated by  $34.5 \text{ s} \pm 12.7 \text{ s}$  (range: 10.1–336.6 s,  $n = 179$  bout separations with a mean of 10 bout separations per bee across 12 bees). Pulses within bouts were separated by  $0.27 \text{ s} \pm 0.06 \text{ s}$ , when excluding pulse separations  $>1 \text{ s}$ , which helped to exclude possible spontaneous abdominal jerks that appeared distinct from bouts of pulses ( $n = 1166$  pulse separations with a mean of 97 pulse separations per bee across 12 bees).

Continuous ventilation (by bees maintaining cells, eating, or heating) rarely included separation of pulses by greater than 10 s ( $n = 28$  out of 490 pulse separations when maintaining cells, 14 out of 394 when eating, and only 16 out of 5,525 when heating; Fig. 6), and instead featured relatively continuous abdominal pulses, which were separated by the same amount of time as sleeping bees, above ( $0.33 \text{ s} \pm 0.08 \text{ s}$  for pulse separations  $<1 \text{ s}$ ,  $n = 5701$  pulse separations with a mean of 328 pulses and a mean of 78 pulse separations per bee across 25 bees; linear mixed model after rank transformation,  $F_{3,34} = 1.19$ ;  $P = 0.33$ ).


Abdominal pulses can be difficult to discern when bees are very active (maintaining cells or eating) in cells, so ventilatory rates should be viewed in the context of whether or not a bee is exhibiting larger body motions.



**Figure 7** Time between pulses of the abdomen (= pulse separations) when exhibiting different behaviors. Ventilatory behavior is commonly distinguished as either continuous or discontinuous, depending on pattern of pulse separations. (A–D) Histograms displaying frequency of pulse separations when bees were sleeping, maintaining cells, eating, or heating. Y-axes break at frequency = 14 to show spread of data along x-axes; maximum y-values (1,166, 335, 328, 5,039) are superimposed above 14 in each plot. Pulse separations that were (E) long (>10 s) and typically associated with discontinuous ventilation, or (F) short (<10 s), by behavior. Pulses separated by <1 s are typical within pulse bouts that are separated by long pauses during discontinuous ventilation (associated with sleep), and are common throughout continuous ventilation. Note different y-axis scales in E and F. ( $n_{\text{pulse separations}} = 1,342$  during sleeping, 490 maintaining cells, 394 eating and 5,525 heating;  $n_{\text{bees analyzed}} = 12$  sleeping, 10 maintaining cells, three eating, 12 heating). [Full-size DOI: 10.7717/peerj.9583/fig-7](https://doi.org/10.7717/peerj.9583/fig-7)

### Thermal measures as indicators of behavior

Body temperatures ( $T_{\text{th}}$ ) differed from surrounding temperatures ( $T_{\text{surr}}$ ) (Kruskal–Wallis  $\chi^2 = 21.2$ ,  $df = 3$ ,  $P = 9.5 \times 10^{-5}$ ), but only when bees were heating.  $T_{\text{th}}$  did not differ from  $T_{\text{surr}}$  when bees were sleeping, maintaining cells, or eating ( $T_{\text{diff}} = 0.20 \pm 0.23$  °C when sleeping,  $0.20 \pm 0.33$  °C when maintaining cells, and  $0.86 \pm 0.31$  °C when eating;  $n = 8, 10, 3$  bees and  $W = 37.5, 58.5, 7$ , respectively; corrected  $P$ -values using Holm method = 1.0 in each case) (Fig. 8).  $T_{\text{th}}$  was only statistically different from  $T_{\text{surr}}$  in heating bees




**Figure 8** Temperatures of bees and of bees relative to their surroundings when exhibiting different behaviors. Individual bees' thorax temperatures relative to surrounding temperatures ( $T_{\text{diff}} = T_{\text{th}} - T_{\text{surr}}$ , left axis; colored circles), and absolute temperature of thorax (right axis; + signs) when sleeping, maintaining cells, eating, or heating. Solid lines represent mean temperature of  $T_{\text{diff}}$  across bees for each behavior, and dashed lines represent mean temperature of thorax ( $T_{\text{th}}$ ) across bees for each behavior. Temperatures are all surface measurements taken remotely with a thermal camera. Bees (x-axis) included 11 uniquely marked individuals (total  $n = 32$  bees; eight sleeping, 10 cleaning, three eating, 11 heating).

Full-size [DOI: 10.7717/peerj.9583/fig-8](https://doi.org/10.7717/peerj.9583/fig-8)

( $T_{\text{diff}} = 2.62 \pm 1.37$  °C;  $n = 11$  bees,  $W = 112$ ,  $P = 0.0032$ ). Heating bees'  $T_{\text{diff}}$  was greater than other bees'  $T_{\text{diff}}$  (vs. sleeping:  $W = 88$ ,  $P = 0.0002$ ; vs. maintaining cells:  $W = 110$ ,  $P = 0.0006$ ; vs. eating:  $W = 32$ ,  $P = 0.044$ ).  $T_{\text{diff}}$  did not differ between sleeping and maintaining cells ( $W = 39$ ,  $P = 0.96$ ), nor did  $T_{\text{diff}}$  differ between maintaining cells and eating ( $W = 2$ ,  $P = 0.068$ ), but eating bees'  $T_{\text{diff}}$  was greater than sleeping bees' ( $W = 0$ ,  $P = 0.044$ ). A heating bee's body temperature visibly differed from her surrounding temperature when using thermal imagery (Figs. 8, 9A and 9B; Movies S9 and S10), and we used this visible difference to initially identify heating bees, prior to analyzing ventilatory rates. Our aim here is to quantitatively confirm this difference ( $T_{\text{diff}}$ ) so that we can confidently associate heaters' telltale heat emission with complementary behaviors (immobility + continuous ventilation) to confirm that the complementary behaviors alone can be used to distinguish heating bees from bees exhibiting the other behaviors. Ventilatory rates are important because a heating bee's thoracic temperature fluctuates over time (Kleinhenz et al., 2003;





**Figure 9** Still images from thermal imaging videos. (A) A worker heating while inside a cell, head facing left and part of abdomen, wings and hind leg extending outside cell, to the right. Image taken from [Movie S9](#). (B) The two brightest spots, at left, each show the relatively hot thorax of a heating bee inside a cell, one immediately behind the plastic window of the hive (top arrow), and one that is one cell deep, seen through the wax wall of a cell (bottom arrow). These workers spent a total of ca. 5 min and 25 min heating inside these cells, respectively. Image taken from [Movie S10](#) at 35 s after 21:41 h. (C) The arrows point to a heating bee with fluctuating body temperature. Images taken from [Movie S11](#) at 10 s after 07:58 h and 2 min 48 s later. Total heating time exceeded 11 min. (D) Each bright spot is a relatively hot thorax, but of a bee maintaining cells, not heating. Image taken from [Movie S12](#).

Full-size [DOI: 10.7717/peerj.9583/fig-9](#)

[Fig. 9C](#); [Movie S11](#)), and a relatively hot thorax does not necessarily mean a worker is actively performing as a heater, but could instead be transitioning into another behavioral state ([Fig. 9D](#); [Movie S12](#)).

Time spent exhibiting a behavior inside cells (beginning, middle and end of stay) did not affect relative body temperature ( $T_{diff}$ ; ANOVA-Type statistic = 0.48,  $df = 1.7$ ,  $P = 0.58$ ).

### Reliability of observing posterior tip of abdomen for identifying behaviors

We tested how reliable watching only the posterior tip of a bee's abdomen is for identifying a behavior when a bee is inside a cell. Fifty-four human subjects correctly identified when honey bee workers were sleeping 86.6% of the time ( $n = 461$  of 540 observations of 11 bees; binomial test, expected = 0.25;  $z = 32.3$ ,  $P = 4.0 \times 10^{-5}$ ; 13.5% misidentifications, with 8.1% identified as heating), maintaining cells 50.1% ( $n = 174$  of 353 observations of seven bees;  $z = 10.5$ ,  $P = 4.0 \times 10^{-5}$ ; most common misidentification: 49.2% eating), eating 70.4% ( $n = 76$  of 108 observations of two bees;  $z = 10.8$ ,  $P = 4.0 \times 10^{-5}$ ; most

common misidentification: 31.5% maintaining cells), and heating 73.0% ( $n = 446$  of 617 observations of 12 bees;  $z = 27.1$ ,  $P = 4.0 \times 10^{-5}$ ; most common misidentification: 18.5% maintaining cells). Participants typically reported difficulty determining behavior due to blurriness of abdomen (one video) or jostling of bee by other bees (1–2 videos). All percentages are means of percentages across bees to address effect of bee, some behaviors of which were more difficult than others to identify. For this reason, percentages may not sum perfectly to 100.

## DISCUSSION

Of the behaviors we recorded inside comb cells, sleep made up 16.9% of the observations, second only to maintaining cells. Maintaining cells and eating were easily identified when observing movement of body or mouthparts, and contents of the cell. Sleeping and heating bees lacked large movements of body or head, and were distinguished from each other using ventilatory rates (discontinuous vs. continuous pulses of the abdomen, respectively) and body surface temperature (relative to surrounding surface temperature). When visibility was restricted (i.e., when the contents of cells were obscured and only the posterior tips of honey bees' abdomens were visible), maintaining cells and eating were difficult to distinguish from each other, but sleeping and heating were identifiable based on ventilatory rates and lack of major body motions alone (86.6% and 73.0% of observations were correctly identified, respectively). We used these two indicators to initially identify sleeping bees and, despite the relative ease of using thermography to distinguish heating bees, the same two indicators (ventilatory rate and lack of major body motions) appear most reliable to identify heating as well. We base this on the fact that a heating bee's temperature can fluctuate, or confusion can arise when bees transition from one behavior to the next (Figs. 9C and 9D; Movies S11 and S12). We also base this on the high reliability of identifying heating bees in our limited-visibility reliability test, which we expect would increase by training observers for longer than 20–30 min.

This study's findings match or differ from other studies in revealing ways. Foragers sleep more during the night than during the day (Kaiser, 1988; Sauer et al., 2003; Sauer, Herrmann & Kaiser, 2004; Eban-Rothschild & Bloch, 2008; Klein et al., 2008), but in this study sleep did not occur more at night (Fig. 5), suggesting that we were likely observing younger workers (e.g., cell cleaners and nurse bees). These younger "hive" bees sleep primarily in cells, and behave arrhythmically (Sauer, Menna-Barreto & Kaiser, 1998; Sauer et al., 1999; Eban-Rothschild & Bloch, 2008; Klein et al., 2008, 2014; but see a report of day-night differences inside cells in Moore et al. (1998)). Bees were sleeping in 16.9% of observations, which falls within the wide range of caste-dependent sleep observed inside cells by Klein et al. (2008) (1.6% observations of foragers—39.4% of cell cleaners). Comparisons with Lindauer (1952) are not feasible because he recorded data from only two individuals under relatively normal conditions, did not distinguish discontinuously ventilating or restful states from superficially similar behavioral states, and did not specify whether calculations were based on idleness exhibited within versus outside cells. Sleeping bees' surface temperatures did not differ from their surroundings, and were slightly higher ( $T_{th} = 34.7 \pm 0.8$  °C,  $n = 8$  bees) than were reported in "resting" bees by

*Kleinhenz et al. (2003)*, which were also measured inside cells ( $32.7 \pm 0.1$  °C– $33.4 \pm 0.3$  °C,  $n = 5$  bees). These resting bees exhibited discontinuous ventilation, with inter-bout durations lasting up to 58 s (vs.  $34.5 \pm 12.7$  s, lasting up to 337 s in this study). Heating bees are typically notably hotter than their surroundings, but the contrast was not as extreme in this study ( $T_{\text{diff}} = 2.6 \pm 1.4$  °C;  $n = 11$  bees) as it was in *Kleinhenz et al. (2003)* ( $4.2 \pm 1.6$  °C;  $n = 8$  bees), but the body temperatures were equally high in both studies ( $T_{\text{th}} = 38.7 \pm 1.6$  °C,  $n = 11$  bees;  $38.3 \pm 1.6$  °C,  $n = 8$  bees).

### Limitations of study

Our observation hive approximated natural conditions in that it was kept in a relatively dark and warm room, featured combs spaced natural distances apart, contained food and brood, the queen was free-roaming, and an entrance allowed full access outdoors. Despite these similarities to natural nests, we supplied the colony with food ad libitum, and comb was limited to narrow slices attached on one side to a plastic window. Reports by Gontarski and Geschke (as communicated by *Von Frisch (1967)*, p. 7), suggest that 500 or 500–1,000 members are sufficient for developing the same division of labor as in normal colonies, but we cannot know if our tiny colony (800–1,000 bees) developed a natural division of labor during this short study. It is important to note that the cells visible along one edge of each comb from which we collected our data may present behavioral biases, which would affect results related to the proportions of behaviors exhibited in cells reported in our surveys. Contents removed from edge cells to increase visibility of comb cells could have caused increased cell cleaning and building activity. We wanted to increase the likelihood of observing sleep in the visible cells, so our emptying of edge cells could have caused a higher rate of discontinuous ventilation within these edge cells. Small numbers of brood or small size of comb could have resulted in unnatural rates of heating, as well. Our limited-visibility reliability test for predicting behaviors featured a lateral view of abdominal tips (*Fig. 4*) when the typical view would be posterior view of abdominal tips (*Fig. 1A*). The limited-visibility test included only three bees eating, and the sole training video devoted to eating did not include eating behavior, only presence of food with description of behavior.

### Why sleep inside cells?

Accounting for sleep inside honeycomb may help to resolve contradictory or confusing evidence reported under less natural conditions (*Sauer, Menna-Barreto & Kaiser, 1998*; *Eban-Rothschild & Bloch, 2008*). If we can rely on discontinuous ventilation + absence of major body motions as markers of sleep in limited-visibility, in-cell situations, the youngest adult workers spend more time asleep than later in life (*Klein et al., 2008*). Sleeping more earlier in life is normal across animals, and much research has considered the current utility of this standard feature of sleep ontogeny. Cell cleaners and nurse bees sleep primarily inside cells that are located in or close to brood comb, and this could be for a variety of functionally interesting reasons (see *Klein et al., 2008*). Comb cells may protect sleeping adults from being disturbed, which could reduce the damaging effects associated with sleep fragmentation. Comb cells could provide warmth for regenerative or

cognitive processes, or serve as a site that reduces sleepers' interference with other workers bustling about the comb. Alternatively, sleeping in cells could be a nonadaptive behavior, during which honey bees simply use comb cells as a default site between acts of cell maintenance, nursing, or heating.

Poets, philosophers and scientists have long pondered the societal marvels of honey bee colonies (*Preston, 2006*), and making visible the bees' activities is a pursuit that has changed our understanding of what nonhuman animals are capable of. Activities, like sleeping, can be difficult to access, particularly when performed inside honeycomb hidden within a dark tree hollow. What specific benefits are conferred by bees sleeping inside cells awaits further investigation, and likely will depend on technical innovations involving noninvasive imaging of standard hives or natural nests, or testing sleep and sleep loss in noncircadian subjects.

## CONCLUSIONS

The best view of a honey bee inside a honeycomb cell is typically restricted to the tip of its abdomen, under the best of circumstances. We hypothesized that even with such constraints, the capacity to identify sleep and other behaviors can be high, based on brief observations of the ventilatory rates (discernible by timing of abdominal pulsing motions) combined with the presence or absence of major body movements. Viewing bees inside cells using a special hive and filming with an infrared-sensitive camera and thermal camera made identifying all behaviors relatively easy in this study, but identifying sleeping or heating bees was also reliable with the limited visibility available to an observer without this special hive or thermal camera. Simply observing ventilatory movements, as well as larger motions evident in the tip of a bee's abdomen was sufficient to noninvasively identify sleeping or heating inside comb cells. Cell maintenance was frequently confused with eating under limited visibility conditions, but both were clearly distinguishable from sleeping and heating. Sleeping and heating were accurately identified (86.6% and 73.0% of observations, respectively) by observing ventilatory rates (discontinuous versus continuous, respectively), combined with a lack of major body movements. Although reliability of identifying behaviors was high, the specialized hive we used may have biased proportions of time bees slept, heated, ate, or maintained cells. Sleep appeared frequently enough to suggest that it is an important behavior experienced within honeycomb cells, supporting previous examinations of sleep inside comb cells, and lending credibility to future ventures, which can rely on similarly less invasive manipulations to reveal the dynamics and functions related to sleep in nature.

## ACKNOWLEDGEMENTS

Jürgen Tautz, Hartmut Vierle, and the entire BEEgroup generously hosted B.A.K.'s research at the University of Würzburg. Sven Mayer and Dirk Ahrens-Lagast ably and graciously assisted with innovative beekeeping. Markus Mika helped to translate a portion of *Lindauer's (1952)* manuscript for us. Alistair Alesch edited video clips for testing predictability of behaviors based on visibility of abdominal tips alone, and University of Wisconsin—La Crosse students took the limited-visibility test. David Reineke

(University of Wisconsin—La Crosse) and Jeff Oliver and Keaton Wilson (University of Arizona Library) contributed statistical and R coding consultation. Logan Schoolcraft provided help in data preparation. Bryant Tran assisted with video editing and file processing. Laura Still and Melody Latronico assisted with some preliminary data transcription. Two anonymous reviewers and Stephen Pratt provided profoundly helpful critiques and suggestions. As always, we thank our sleeping, cell-maintaining, eating and heating honey bees.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the Deutscher Akademischer Austausch Dienst (DAAD #A0670415). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
Deutscher Akademischer Austausch Dienst: DAAD #A0670415.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Barrett A. Klein conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- M. Kathryn Busby analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Honey bee research was conducted using bees from an on-campus bee yard with approval by the University of Würzburg (Würzburg, Germany).

### Data Availability

The following information was supplied regarding data availability:

Data are available in the [Supplemental Files](#). All Supplemental Information, plus additional files, are also available through Open Science Framework: Klein, Barrett A., and M.K. Busby. 2020. “Slumber in a cell: honeycomb used by honey bees for food, brood, heating... and sleeping.” OSF. July 3. [osf.io/eux5y](https://osf.io/eux5y).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9583#supplemental-information>.

## REFERENCES

- Bailey L. 1954. The respiratory currents in the tracheal system of the adult honey-bee. *Journal of Experimental Biology* **31**:589–593.
- Bujok B, Kleinhenz M, Fuchs S, Tautz J. 2002. Hot spots in the bee hive. *Naturwissenschaften* **89**(7):299–301 DOI [10.1007/s00114-002-0338-7](https://doi.org/10.1007/s00114-002-0338-7).
- Cirelli C, Tononi G. 2008. Is sleep essential? *PLOS Biology* **6**(8):1605–1611 DOI [10.1371/journal.pbio.0060216](https://doi.org/10.1371/journal.pbio.0060216).
- Crane E. 1983. *The archaeology of beekeeping*. Ithaca: Cornell University Press.
- Dustmann JH, Geffcken H. 2000. *Bienen können Farben unterscheiden*. Celle: Landesinstitut für Bienenkunde.
- Eban-Rothschild AD, Bloch G. 2008. Differences in the sleep architecture of forager and young honeybees (*Apis mellifera*). *Journal of Experimental Biology* **211**(15):2408–2416 DOI [10.1242/jeb.016915](https://doi.org/10.1242/jeb.016915).
- Flanigan WF, Wilcox RH, Rechtschaffen A. 1973. The EEG and behavioral continuum of the crocodylian, *Caimen sclerops*. *Electroencephalography and Clinical Neurophysiology* **34**(5):521–538 DOI [10.1016/0013-4694\(73\)90069-2](https://doi.org/10.1016/0013-4694(73)90069-2).
- Hussaini SA, Bogusch L, Landgraf T, Menzel R. 2009. Sleep deprivation affects extinction but not acquisition memory in honeybees. *Learning & Memory* **16**(11):698–705 DOI [10.1101/lm.1578409](https://doi.org/10.1101/lm.1578409).
- Kaiser W. 1988. Busy bees need rest, too: behavioural and electromyographical sleep signs in honeybees. *Journal of Comparative Physiology A* **163**(5):565–584 DOI [10.1007/BF00603841](https://doi.org/10.1007/BF00603841).
- Klein BA, Klein A, Wray MK, Mueller UG, Seeley TD. 2010. Sleep deprivation impairs precision of waggle dance signaling in honey bees. *Proceedings of the National Academy of Sciences of the United States of America* **107**(52):22705–22709 DOI [10.1073/pnas.1009439108](https://doi.org/10.1073/pnas.1009439108).
- Klein BA, Olzowy KM, Klein A, Saunders KM, Seeley TD. 2008. Caste-dependent sleep of worker honey bees. *Journal of Experimental Biology* **211**(18):3028–3040 DOI [10.1242/jeb.017426](https://doi.org/10.1242/jeb.017426).
- Klein BA, Stiegler M, Klein A, Tautz J. 2014. Mapping sleeping bees within their nest: spatial and temporal analysis of worker honey bee sleep. *PLOS ONE* **9**(7):e102316 DOI [10.1371/journal.pone.0102316](https://doi.org/10.1371/journal.pone.0102316).
- Klein BA, Vogt M, Unrein K, Reineke DM. 2018. Followers of honey bee waggle dancers change their behaviour when dancers are sleep-restricted or perform imprecise dances. *Animal Behaviour* **146**:71–77 DOI [10.1016/j.anbehav.2018.10.010](https://doi.org/10.1016/j.anbehav.2018.10.010).
- Kleinhenz M, Bujok B, Fuchs S, Tautz J. 2003. Hot bees in empty broodnest cells: heating from within. *Journal of Experimental Biology* **206**(23):4217–4231 DOI [10.1242/jeb.00680](https://doi.org/10.1242/jeb.00680).
- Kritsky G. 2010. *The quest for the perfect hive: a history of innovation in bee culture*. New York: Oxford University Press.
- Lindauer M. 1952. Ein Beitrag zur Frage der Arbeitsteilung im Bienenstaat. *Zeitschrift für Vergleichende Physiologie* **34**(4):299–345 DOI [10.1007/BF00298048](https://doi.org/10.1007/BF00298048).
- Moore D, Angel JE, Cheeseman IM, Fahrbach SE, Robinson GE. 1998. Timekeeping in the honey bee colony: integration of circadian rhythms and division of labor. *Behavioral Ecology and Sociobiology* **43**(3):147–160 DOI [10.1007/s002650050476](https://doi.org/10.1007/s002650050476).
- Noguchi K, Gel YR, Brunner E, Konietzschke F. 2012. nparLD: an R software package for the nonparametric analysis of longitudinal data in factorial experiments. *Journal of Statistical Software* **50**(12):12 DOI [10.18637/jss.v050.i12](https://doi.org/10.18637/jss.v050.i12).
- Preston C. 2006. *Bee*. London: Reaktion Books.

- R Core Team. 2019.** *R: a language and environment for statistical computing*. Vienna: The R Foundation for Statistical Computing. Available at <http://www.R-project.org/>.
- Rattenborg NC, Lesku JA, Martinez-Gonzalez D, Lima SL. 2007.** The non-trivial functions of sleep. *Sleep Medicine Reviews* **11**(5):405–409 DOI [10.1016/j.smrv.2007.04.003](https://doi.org/10.1016/j.smrv.2007.04.003).
- Sakagami SF. 1953.** Untersuchungen über die Arbeitsteilung in einem Zwergvolk der Honigbiene: Beiträge zur Biologie des Bienenvolkes, *Apis mellifera* L.I. *Japanese Journal of Zoology* **11**:117–185.
- Sauer S, Happel U, Neubecker R, Menna-Barreto L, Herrmann E, Kaiser W. 1999.** Ontogeny of the circadian rest-activity cycle in honey bees kept in isolation. In: Elsner N, Eysel U, eds. *Proceedings of the 1st Göttingen Conference of the German Neuroscience Society, Vol. II: 27th Göttingen Neurobiology Conference*. Stuttgart: Georg Thieme Verlag, 241.
- Sauer S, Herrmann E, Kaiser W. 2004.** Sleep deprivation in honey bees. *Journal of Sleep Research* **13**(2):145–152 DOI [10.1111/j.1365-2869.2004.00393.x](https://doi.org/10.1111/j.1365-2869.2004.00393.x).
- Sauer S, Kinkelin M, Herrmann E, Kaiser W. 2003.** The dynamics of sleep-like behaviour in honey bees. *Journal of Comparative Physiology A: Sensory, Neural, and Behavioral Physiology* **189**(8):599–607 DOI [10.1007/s00359-003-0436-9](https://doi.org/10.1007/s00359-003-0436-9).
- Sauer S, Menna-Barreto L, Kaiser W. 1998.** The temporal organization of rest and activity in newly emerged honey bees kept in isolation—initial results. *Apidologie* **29**:445–447.
- Seeley TD. 1995.** *The wisdom of the hive: the social physiology of honey bee colonies*. Cambridge: Harvard University Press.
- Seeley TD. 2019.** *The lives of bees: the untold story of the honey bee in the wild*. Princeton: Princeton University Press.
- Stabentheiner A, Schmaranzer S. 1987.** Thermographic determination of body temperatures in honey bees and hornets: calibration and applications. *Thermology* **2**:563–572.
- Tobler I. 1985.** Deprivation of sleep and rest in vertebrates and invertebrates. In: Inoue S, Borbely AA, eds. *Endogenous Sleep Substances and Sleep Regulation (Taniguchi Symposia, Series no. 8)*. Utrecht: VNU Science Press, 57–66.
- Vázquez DE, Balbuena MS, Chaves F, Gora J, Menzel R, Farina WM. 2020.** Sleep in honey bees is affected by the herbicide glyphosate. *Scientific Reports* **10**:10516 DOI [10.1038/s41598-020-67477-6](https://doi.org/10.1038/s41598-020-67477-6).
- Von Frisch K. 1967.** *The dance language and orientation of bees*. Cambridge: The Belknap Press.
- Von Frisch K, Lindauer M. 1977.** *Nachweis des Farbsehens bei der Honigbiene*. Göttingen: IWF.
- Zwaka H, Bartels R, Gora J, Franck V, Culo A, Götsch M, Menzel R. 2015.** Context odor presentation during sleep enhances memory in honeybees. *Current Biology* **25**(21):2869–2874 DOI [10.1016/j.cub.2015.09.069](https://doi.org/10.1016/j.cub.2015.09.069).