



CORRECTION

## Correction to: A Dalbavancin Lock Solution Can Reduce Enterococcal Biofilms After Freezing

Marta Díaz-Navarro · Rama Hafian · Irene Manzano · María J. Pérez-Granda ·  
Emilia Cercenado · Cristina Pascual · Carmen Rodríguez ·  
Patricia Muñoz · María Guembe

Published online: April 13, 2022  
© The Author(s) 2022

Correction to: Infect Dis Ther  
<https://doi.org/10.1007/s40121-021-00579-4>

---

The original article can be found online at <https://doi.org/10.1007/s40121-021-00579-4>.

---

M. Díaz-Navarro · M. J. Pérez-Granda ·  
E. Cercenado · P. Muñoz · M. Guembe (✉)  
Department of Clinical Microbiology and Infectious  
Diseases, Hospital General Universitario Gregorio  
Marañón, C/Dr. Esquerdo, 46, 28007 Madrid, Spain  
e-mail: mariaguembe@hotmail.com;  
maria.guembe@iisgm.com

M. Díaz-Navarro · M. J. Pérez-Granda · P. Muñoz ·  
M. Guembe  
Instituto de Investigación Sanitaria Gregorio  
Marañón, Madrid, Spain

R. Hafian · I. Manzano  
Biology Department, School of Biology, Universidad  
Complutense de Madrid, Madrid, Spain

M. J. Pérez-Granda · P. Muñoz  
CIBER Enfermedades Respiratorias-CIBERES (CB06/  
06/0058), Madrid, Spain

E. Cercenado · P. Muñoz  
Medicine Department, School of Medicine,  
Universidad Complutense de Madrid, Madrid, Spain

C. Pascual  
Hematology Department, Hospital General  
Universitario Gregorio Marañón, Madrid, Spain

C. Rodríguez  
Pharmacy Department, Hospital General  
Universitario Gregorio Marañón, Madrid, Spain

In the original publication of the article, Funding must be changed and figures files are wrong labeled, so both issues has been addressed.

1. **Funding:** The study was partially financed by grants from the ISCIII (PI18/00045), the European Regional Development Fund (FEDER) “A way of making Europe” and from Fundación Mutua Madrileña (FMM21/01).
2. **Results: last paragraph:**

In contrast, in *E. faecium* biofilms, the only statistically and clinically significant difference between DH and dalbavancin was observed for the median (IQR) percentage reduction in log cfu/ml at month 6 (13.6% [13.0–18.9%] vs. 20.4% [14.6–26.6%],  $p = 0.01$ , 25% of dalbavancin value: 15.3%) (Table 3, Fig. 3C). No significant reduction in metabolic activity was detected during either of the 2 periods (months 3 and 6) (Table 3, Fig. 3D).

3. **Figures files** (you can find them attached correctly labeled).

Fig. 1A 1B must be 3A 3B.




Fig. 2A 2B must be 3C 3D.





Fig. 3A 3B must be 1A 1B.






Fig. 3C 3D must be 2A 2B.



This has been corrected in this paper.

**Open Access.** This article is licensed under a Creative Commons Attribution-Non-Commercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were

made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view

a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

#### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.