

COMMENTARY

Open Access

The summer of hydrogen sulfide: highlights from two international conferences

John W Calvert

Abstract

A great deal of interest has been paid recently to the hydrogen sulfide, the newest member of the gasotransmitter family. With the growing interest in the biology of H₂S, the need for meetings and conferences dedicated solely to the field of H₂S has also grown. In 2009, scientist from around the world met in Shanghai, China for the first time to discuss the physiological relevance of H₂S. In 2012, two conferences were organized to bring scientists, clinicians, and industry representatives together to discuss the latest breakthroughs concerning the emergent field of H₂S. The following is a summary report of The First European Conference on the Biology of Hydrogen Sulfide and the Second International Conference on Hydrogen Sulfide Biology and Medicine.

Keywords: Gasotransmitter, Hydrogen sulfide, Nitric oxide, Carbon monoxide

Introduction

Gasotransmitters are small, labile, endogenously-produced gaseous molecules that play important roles in cellular signaling [1]. Hydrogen sulfide (H₂S) is the newest member of the gasotransmitter family. Historically, it was considered to be highly toxic and hazardous to the environment [2]. However, H₂S has recently emerged as a critical physiological gaseous signaling molecule that is produced enzymatically in all mammalian species at low micromolar levels via the action of cysteine metabolic enzymes: cystathionine-γ-lyase (CSE), cystathionine-β-synthase (CBS), and 3-mercaptopyruvate sulfutransferase (3-MST) [3]. Because of this discovery, the biological and physiological role of H₂S has been re-evaluated. As such, an extensive amount of work has been conducted over the last several decades and has led to the discovery that H₂S possesses a number of physiological actions.

In 1989, while investigating the effects of acute H₂S poisoning in the brain, Warenycia *et al.* [4] first reported that H₂S was produced endogenously. Soon after this discovery, other investigators started to take a closer look at H₂S and evidence soon emerged to demonstrate its physiological importance. For instance, Skrajny *et al.* [5] found increased levels of serotonin and reduced levels of norepinephrine in the frontal cortex of a rat

when chronically exposed to 20 ppm of H₂S and in 1996, Abe *et al.* [6] suggested that H₂S was an endogenous neuromodulator, as they showed that physiological concentrations of H₂S enhanced *N*-methyl-D-aspartate (NMDA) receptor-mediated responses and aided in the induction of hippocampal long-term potentiation. Shortly after, Hosoki *et al.* [7] reported that an H₂S-producing enzyme was present in the vasculature and proposed that H₂S may be an endogenous smooth muscle relaxant. Since these initial offerings, studies have begun to characterize its biological profile and therapeutic potential. Over the last several years, a great amount of work by investigators all over the world have led to the suggestion that H₂S is a powerful signaling molecule with a diverse physiological profile. Moreover, because of its physiological actions, H₂S has been reported to protect the heart, brain, liver, kidney, and lungs against a variety of pathological stimuli, including ischemia/reperfusion (I/R) injury [8-17].

With the growing interest in the biology of H₂S, the need for meetings and conferences dedicated solely to the field of H₂S has also grown. In 2009, scientists from around the world met in Shanghai, China for the first time to discuss the physiological relevance of H₂S. In 2012, two conferences were organized to bring scientists, clinicians, and industry representatives together to discuss the latest breakthroughs concerning the emergent

Correspondence: jcalver@emory.edu
Department of Surgery, Division of Cardiothoracic Surgery, Carlyle Fraser Heart Center, Emory University of Medicine, Atlanta, GA 30308, USA

field of H₂S. The following is a summary of the highlights from those two meetings.

First European conference on the biology of hydrogen sulfide

The First European Conference on the Biology of Hydrogen Sulfide was held at Smolenice Castle on the outskirts of Bratislava, Slovakia from June 15 to 18, 2012. The Institute of Molecular Physiology and Genetics from the Slovak Academy of Sciences sponsored this conference. Dr Karol Ondrias from the Slovak Academy of Sciences served as the Conference Chairman and Dr Matthew Whiteman from the University of Exeter served as Honorary Chairman and Chairman of the International Advisory Committee. Together with the rest of the members of the International Advisory Committee, Drs Ondrias and Whiteman organized an outstanding scientific program consisting of 12 plenary lectures, 34 short oral presentations, and 15 poster presentations.

The meeting started with a Keynote Lecture delivered by Dr Rui Wang from Lakehead University in Thunder Bay, Ontario, Canada. In his lecture, Dr Wang discussed several recent findings regarding where inside the eukaryotic cell H₂S is produced and how this intracellular compartmentalization is realized. He also discussed the functional consequences of H₂S production in response to difference stimuli. Dr Wang's lecture was followed by a very nice welcome reception where the attendees were inundated with local Slovakian cuisine.

Day 2 of the conference began with a session focused on the effects of H₂S on heart and vascular physiology. Dr Y.Z. Zhu from Fudan University in Shanghai China began the session by detailing recent work on the involvement of H₂S in the cardiovascular disorder processes and the prospects of new therapeutic strategies based on the regulation of endogenous H₂S production [18]. This session was highlighted by Dr David Lefer from Emory University in Atlanta GA who presented recent work investigating the role of cystathionine gamma lyase derived H₂S in the development of pressure-induced heart failure. He also presented novel and exciting data regarding the effects of oral H₂S therapy on preventing cardiac dysfunction following the induction of heart failure.

There is emerging evidence that members of the gasotransmitter family can interact with each other. For instance, NO and H₂S have been suggested to collaborate in regulating vascular homeostasis and vasodilation. Additional evidence suggests that NO can increase CGL activity acutely, and that chronic exposure to NO up-regulates CGL expression. Moreover, at low concentrations H₂S has been shown to enhance the release of NO from vascular endothelium and increase the vasorelaxant effect of the NO donor sodium nitroprusside [19]. Furthermore, a 10

fold increase in H₂S levels have been reported with administration of NO donors [20], suggesting a synergistic relationship between NO and H₂S. Dr Andreas Papapetropoulos from the University of Patras in Greece detailed very exciting and new results that further demonstrate the interdependence of the gasotransmitters on each other. Specifically, he provided the audience with compelling evidence that angiogenesis and vasodilatation require the simultaneous production of H₂S and NO, suggesting that both are mutually required for the control of vascular function [21].

Day 2 concluded with two afternoon sessions. The first session focused on the ability of H₂S to regulate respiratory and gastrointestinal function. Dr John Wallace from McMaster University in Hamilton, Ontario, Canada discussed his group's latest findings regarding the ability of novel H₂S releasing anti-inflammatory drugs to alleviate the development of gastrointestinal (GI) ulcers [22]. The last session of Day 2 focused on H₂S donors and was highlighted by Dr Mathew Whiteman's presentation on the use of H₂S donors to treat inflammatory joint disease.

Dr Csabo Szabo from the University of Texas Medical Branch in Galveston, TX opened the morning session on Day 3 by discussing the role of H₂S deficiency in the pathogenesis of diabetic vascular complications. In his presentation he provided the audience with compelling data suggesting that H₂S deficiency is emerging as a significant pathogenic mechanism in diabetic complication, while H₂S supplementation has the potential to be a novel therapeutic approach [23]. The session on H₂S and diabetes was followed by a session regarding the interaction of H₂S with the mitochondria. This session was highlighted by Dr Katalin Modis' (University of Texas Medical Branch in Galveston, TX) presentation detailed her exciting work demonstrating that H₂S has a physiological role as a positive modulator of mitochondrial electron transport that is dependent on a basal activity of the Krebs cycle [24].

In the afternoon, Dr Hideo Kimura from the National Institute of Neuroscience in Tokyo, Japan led off a very interesting and stimulating session on the synthesis of hydrogen sulfide. In his talk, Dr Kimura detailed his group's efforts to identify the endogenous reducing substances (thioredoxin and dihydrolipoic acid) that facilitate the production of H₂S from 3-mercaptopyruvate sulfurtransferase [25]. This was followed by a very stimulating presentation from the Ondrias group detailing their work suggesting that H₂S induces the release of NO from nitroso groups. Day 3 ended with a very informative session regarding the interaction of H₂S with different ion channels.

The Final Day of the conference ended with several talks discussing the latest findings regarding the role of H₂S in the setting of cellular stress and cell proliferation.

Additionally, awards were handed out at the conclusion of the conference to several young investigators.

The second international conference on H₂S biology and medicine

With the momentum gained at Smolenic Castle in Bratislava, scientists, clinicians, and industry representatives from around the world descended on Atlanta, GA from September 20 – 22, 2012 to attend the Second International Conference on H₂S Biology and Medicine. Dr David Lefer from Emory University in Atlanta, GA organized this conference. Together with the rest of the member of the Organizing Committee, Dr Lefer planned a well-balanced scientific program detailing the latest findings concerning the chemistry and biology of H₂S. The final program consisted of 30 plenary lectures, 4 short oral presentations, and over 75 poster presentations.

Dr Solomon Snyder from Johns Hopkins University in Baltimore, MD delivered the keynote lecture to start the conference. In his lecture, Dr Solomon eloquently discussed his group's efforts to characterize sulfhydration, which is the mechanism by which H₂S modifies reactive cysteine residues on targeted proteins. Although analogous to protein nitrosylation, sulfhydration is substantially more prevalent and usually increases the catalytic activity of targeted proteins [26].

Dr Snyder's lecture was followed by several presentations on the biology and chemistry of H₂S. Highlights from this session included Dr Nicholas Tonk's (Cold Spring Harbor Laboratory) presentation on his recent work detailing of the effects of H₂S on the endoplasmic reticulum stress response. Specifically, he presented data demonstrating that H₂S reversibly inhibits the activity of protein tyrosine phosphatase 1B, which promotes the activity of protein kinase-like endoplasmic reticulum (ER) kinase (PERK) during the response to ER stress [27]. With the growing interest in the physiological and biological role of H₂S, there is a need for better detection methods for H₂S levels and signaling. This was a constant point of discussion throughout the meeting. As such, Dr Ming Xian's (Washington State University in Pullman, WA) presentation on new chemical tools for H₂S research was of particular interest. In his talk, Dr Xian provided evidence that newly developed fluorescent probes provide good selectivity and sensitivity for H₂S [28].

The afternoon session of Day 1 was dedicated to discussion on the role that H₂S plays in the development and treatment of cardiovascular disease. Dr Frank Sellke from Brown University in Providence, RI led off this session by discussing his labs previous work investigating the impact of H₂S on myocardium in the setting of cold crystalloid cardioplegia and cardiopulmonary bypass [29]. In his talk, Dr Sellke provided compelling evidence that treatment with H₂S offered myocardial protection

via attenuation of caspase-independent apoptosis and autophagy in a porcine model of cardiopulmonary bypass. Later in this session, Dr Suresh Tyagi from the University of Louisville in Louisville, KY summarized his groups' findings regarding the effects of H₂S on cardiovascular remodeling in response to heart failure [30]. Day 1 concluded with a reception and presentation of posters.

Day 2 of the conference featured three sessions. The first session of the morning was focused on H₂S and the vasculature. Dr Rui Wang from Lakehead University in Thunder Bay, Ontario, Canada started the session with a nice introduction to the effects of H₂S on the vascular endothelium. This was followed by Dr Guiseppe Cirino's (University of Naples, Naples, Italy) presentation on H₂S as an endogenous inhibitor of phosphodiesterases and how this affects vascular tone.

The second session of Day 2 was focused on the H₂S and gastrointestinal diseases. Presentations in this session from Drs. John Wallace (McMaster University in Hamilton, Ontario, Canada), Mauro Perretti (Queen Mary University), Nathalie Vergnolle (INSERM), and Andre Buret (University of Calgary) detailed the latest findings regarding the role of H₂S in the resolution and repair of the GI tract and detailed the latest developments in use of novel H₂S releasing drugs to treat GI disorders.

The final session of Day 2 was concentrated on the role of H₂S in shock and sepsis. This session began with a very informative presentation by Dr Konstantin Shatalin (New York University) in which he discussed the role H₂S plays in mediating bacteria defense against antibiotics [31]. Specifically, Dr Shatalin provided evidence that inactivation of putative cystathionine β-synthase, cystathionine γ-lyase, or 3-mercaptopyruvate sulfurtransferase in *Bacillus anthracis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Escherichia coli* suppresses H₂S production, rendering these pathogens highly sensitive to a multitude of antibiotics. Dr Fumito Ichinose from Harvard University in Boston, MA then detailed recent findings from his lab in which they demonstrated that inhaled H₂S prevents inflammation and improves survival in the setting of endotoxemia by altering sulfide metabolism in mice [32].

The final day of the conference began with presentations selected from submitted abstracts. This forum gave deserving young investigators and trainees the opportunity to present their latest findings in a more formal setting. This session was followed by a second session on H₂S and the vasculature. Dr Christopher Peers from the University of Leeds in Leeds, England highlighted this session with his presentation on the regulation of ion channels by H₂S. Dr Chris Kevil from LSU-Health Sciences Center in Shreveport, LA delivered

the last talk of the conference detailing his groups latest work on the actions of H₂S in the setting of high limb ischemia.

Summary and comments

In summary, the First European Conference on the Biology of H₂S and the Second International Conference on H₂S Biology and Medicine were both a tremendous successes. Each conference provided an excellent platform for the presentation and discussion of the latest findings regarding the biology and chemistry of H₂S. In addition, both of these conferences served as a reminder that that is still so much about H₂S that we do not know. In this regard, a number of important issues/questions were raised at each conference that will certainly drive research in the years to come. These include (1) the need to develop more sensitive ways to measure H₂S, (2) the need to develop new drugs or delivery mechanisms to deliver sustained levels of H₂S, and (3) the need to understand the molecular targets of H₂S and the manner in which H₂S modifies these cellular targets.

It was recently announced that there will be a Second European Conference in Exeter England from September 8–11, 2013. Dr Matthew Whiteman will serve as the Organizer. In addition, Dr Hideo Kimura will serve as the organizer for the next International H₂S meeting. He announced at the conclusion of the Atlanta meeting that the next International conference on H₂S will be held in Japan in 2014. For more information about this conference and other conferences related to H₂S please log on to the European Network on Gasotransmitters website for the latest news and developments (<http://www.gasotransmitters.eu/>).

Competing interest

All authors declare that they have no competing interest.

Disclosures

John Calvert served as Vice Chairman of the Organizing committee for the Second International Conference on H₂S Biology and Medicine.

Sources of funding

Supported by grants from the American Diabetes Association (7-09-BS-26) and the National Institutes of Health National Heart Lung and Blood Institute (NHLBI) (5R01HL098481-03) to J.W.C. This work was also supported by funding from the Carlyle Fraser Heart Center (CFHC) of Emory University Hospital Midtown.

Received: 23 December 2012 Accepted: 15 February 2013

Published: 25 February 2013

References

1. Nicholson CK, Calvert JW: **Hydrogen sulfide and ischemia-reperfusion injury.** *Pharmacol Res* 2010, **62**:289–297.
2. Ramazzini B: *Diseases of Workers - De Morbis Artificum Diatriba - 1713.* New York: C. Hafner Publishing Co Inc; 1964.
3. Li L, Rose P, Moore PK: **Hydrogen sulfide and cell signaling.** *Annu Rev Pharmacol Toxicol* 2011, **51**:169–187.
4. Warencya MW, Goodwin LR, Benishin CG, Reiffenstein RJ, Francom DM, Taylor JD, Dieken FP: **Acute hydrogen sulfide poisoning. Demonstration**

- of selective uptake of sulfide by the brainstem by measurement of brain sulfide levels. *Biochem Pharmacol* 1989, **38**:973–981.
5. Skrajny B, Hannah RS, Roth SH: **Low concentrations of hydrogen sulphide alter monoamine levels in the developing rat central nervous system.** *Can J Physiol Pharmacol* 1992, **70**:1515–1518.
6. Abe K, Kimura H: **The possible role of hydrogen sulfide as an endogenous neuromodulator.** *J Neurosci* 1996, **16**:1066–1071.
7. Hosoki R, Matsuki N, Kimura H: **The possible role of hydrogen sulfide as an endogenous smooth muscle relaxant in synergy with nitric oxide.** *Biochem Biophys Res Commun* 1997, **237**:527–531.
8. Bian JS, Yong QC, Pan TT, Feng ZN, Ali MY, Zhou S, Moore PK: **Role of hydrogen sulfide in the cardioprotection caused by ischemic preconditioning in the rat heart and cardiac myocytes.** *J Pharmacol Exp Ther* 2006, **316**:670–678.
9. Kimura Y, Kimura H: **Hydrogen sulfide protects neurons from oxidative stress.** *FASEB J* 2004, **18**:1165–1167.
10. Elrod JW, Calvert JW, Morrison J, Doeller JE, Kraus DW, Tao L, Jiao X, Scalia R, Kiss L, Szabo C, et al: **Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function.** *Proc Natl Acad Sci U S A* 2007, **104**:15560–15565.
11. Sivarajah A, McDonald MC, Thiernemann C: **The production of hydrogen sulfide limits myocardial ischemia and reperfusion injury and contributes to the cardioprotective effects of preconditioning with endotoxin, but not ischemia in the rat.** *Shock* 2006, **26**:154–161.
12. Bliksoen M, Kaljusto ML, Vaage J, Stenslokken KO: **Effects of hydrogen sulphide on ischaemia-reperfusion injury and ischaemic preconditioning in the isolated, perfused rat heart.** *Eur J Cardiothorac Surg* 2008, **34**:344–349.
13. Florian B, Vintilescu R, Balseanu AT, Buga AM, Grisk O, Walker LC, Kessler C, Popa-Wagner A: **Long-term hypothermia reduces infarct volume in aged rats after focal ischemia.** *Neurosci Lett* 2008, **438**:180–185.
14. Jha S, Calvert JW, Duranski MR, Ramachandran A, Lefer DJ: **Hydrogen sulfide attenuates hepatic ischemia-reperfusion injury: role of antioxidant and antiapoptotic signaling.** *Am J Physiol Heart Circ Physiol* 2008, **295**:H801–H806.
15. Tripatara P, Patel NS, Collino M, Gallicchio M, Kieswich J, Castiglia S, Benetti E, Stewart KN, Brown PA, Yaqoob MM, et al: **Generation of endogenous hydrogen sulfide by cystathionine gamma-lyase limits renal ischemia/reperfusion injury and dysfunction.** *Lab Invest* 2008, **88**:1038–1048.
16. Xu Z, Prathapasinghe G, Wu N, Hwang SY, Siow YLOK: **Ischemia-reperfusion reduces cystathionine-beta-synthase-mediated hydrogen sulfide generation in the kidney.** *Am J Physiol Renal Physiol* 2009, **297**:F27–F35.
17. Fu Z, Liu X, Geng B, Fang L, Tang C: **Hydrogen sulfide protects rat lung from ischemia-reperfusion injury.** *Life Sci* 2008, **82**:1196–1202.
18. Gu X, Zhu YZ: **Therapeutic applications of organosulfur compounds as novel hydrogen sulfide donors and/or mediators.** *Expert Rev Clin Pharmacol* 2011, **4**:123–133.
19. Zhao W, Zhang J, Lu Y, Wang R: **The vasorelaxant effect of H(2)S as a novel endogenous gaseous K(ATP) channel opener.** *EMBO J* 2001, **20**:6008–6016.
20. Eto K, Kimura H: **The production of hydrogen sulfide is regulated by testosterone and S-adenosyl-L-methionine in mouse brain.** *J Neurochem* 2002, **93**:80–86.
21. Coletta C, Papapetropoulos A, Erdelyi K, Olah G, Modis K, Panopoulos P, Asimakopoulou A, Gero D, Sharina I, Martin E, Szabo C: **Hydrogen sulfide and nitric oxide are mutually dependent in the regulation of angiogenesis and endothelium-dependent vasorelaxation.** *Proc Natl Acad Sci U S A* 2012, **109**:9161–9166.
22. Wallace JL: **Hydrogen sulfide: a rescue molecule for mucosal defence and repair.** *Dig Dis Sci* 2012, **57**:1432–1434.
23. Szabo C: **Roles of hydrogen sulfide in the pathogenesis of diabetes mellitus and its complications.** *Antioxid Redox Signal* 2012, **17**:68–80.
24. Modis K, Coletta C, Erdelyi K, Papapetropoulos A, Szabo C: **Intramitochondrial hydrogen sulfide production by 3-mercaptopyruvate sulfurtransferase maintains mitochondrial electron flow and supports cellular bioenergetics.** *Faseb J* 2012, **27**:601–611.
25. Mikami Y, Shibuya N, Kimura Y, Nagahara N, Ogasawara Y, Kimura H: **Thioredoxin and dihydrolipoic acid are required for 3-mercaptopyruvate sulfurtransferase to produce hydrogen sulfide.** *Biochem J* 2011, **439**:479–485.

26. Ambrus GG, Al-Moyed H, Chaieb L, Sarp L, Antal A, Paulus W: **The fade-in–short stimulation–fade out approach to sham tDCS–reliable at 1 mA for naive and experienced subjects, but not investigators.** *Brain Stimul* 2012, **5**:499–504.
27. Krishnan N, Fu C, Pappin DJ, Tonks NK: **H₂S-Induced sulfhydration of the phosphatase PTP1B and its role in the endoplasmic reticulum stress response.** *Sci Signal* 2011, **4**:ra86.
28. Liu C, Peng B, Li S, Park CM, Whorton AR, Xian M: **Reaction based fluorescent probes for hydrogen sulfide.** *Org Lett* 2012, **14**:2184–2187.
29. Osipov RM, Robich MP, Feng J, Chan V, Clements RT, Deyo RJ, Szabo C, Sellke FW: **Effect of hydrogen sulfide on myocardial protection in the setting of cardioplegia and cardiopulmonary bypass.** *Interact Cardiovasc Thorac Surg* 2010, **10**:506–512.
30. Givvimani S, Munjal C, Gargoum R, Sen U, Tyagi N, Vacek JC, Tyagi SC: **Hydrogen sulfide mitigates transition from compensatory hypertrophy to heart failure.** *J Appl Physiol* 2011, **110**:1093–1100.
31. Shatalin K, Shatalina E, Mironov A, Nudler E: **H₂S: a universal defense against antibiotics in bacteria.** *Science* 2011, **334**:986–990.
32. Tokuda K, Kida K, Marutani E, Crimi E, Bougaki M, Khatri A, Kimura H, Ichinose F: **Inhaled hydrogen sulfide prevents endotoxin-induced systemic inflammation and improves survival by altering sulfide metabolism in mice.** *Antioxid Redox Signal* 2012, **17**:11–21.

doi:10.1186/2045-9912-3-5

Cite this article as: Calvert: The summer of hydrogen sulfide: highlights from two international conferences. *Medical Gas Research* 2013 **3**:5.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

