

## The essential role of prevention in reducing the cancer burden in Europe: a commentary from Cancer Prevention **Europe**

2020, Vol. 106(5) NP2-NP4 © Fondazione IRCCS Istituto Nazionale dei Tumori 2019



Article reuse guidelines: sagepub.com/journals-permissions DOI: 10.1177/0300891619851865 journals.sagepub.com/home/tmj

(\$)SAGE

Carolina Espina D. Linda Bauld<sup>2,3</sup>, Bernardo Bonanni<sup>4</sup>, Hermann Brenner<sup>5</sup>, Karen B Joakim Dillner<sup>7</sup>, Ellen Kampman<sup>8</sup>, Mef Nilbert<sup>9</sup>, Paolo Vineis<sup>10</sup>, Matty P. Weijenker Alison Cox<sup>3</sup>, Theo M. de Kok<sup>11</sup>, Daniela Fecht<sup>10</sup>, Giota Mitrou<sup>12</sup>, David C Muller zive Duplicate Pi Davide Serrano<sup>4</sup>, Karen Steindorf<sup>5</sup>, Hans Storm<sup>9</sup>, Mangesh A. Thorat<sup>6,13</sup> Fränzel van Duijnhoven<sup>8</sup>, Elisabete Weiderpass<sup>1</sup> and Joachim Schüz<sup>1</sup>

#### **Keywords**

Epidemiology and prevention, Oncology, cancer prevention

Date received: 28 February 2019; accepted: 19 April 2019

The potential of cancer prevention

In 28 out of 40 United Nations-defined European tries, cancer is now the leading cause of prematical leath, and the second most common in the other 12 with a total of 1.93 million deaths and 3.91 million new resident cases in 2018. Those numbers are projected to use to 2.55 million deaths and 4.75 million incident cases in Europe by 2040, as a result of population agi g and growth, representing an overall increase in number of deaths by 32%.2 Notably, however, these estites are based on prediction models taking current in electronic rates and time trends into consideration, so that preventive actions taken today could change this forecast, V.e., lead to lower than those expected numbers. The sancer cost in 27 countries in the European Union in 2009 was €126 billion, 60% incurred in nonhealthcare areas, with almost €43 billion in lost productivity due to early death. Of the 4 cancers studied, lung cancer had the highest economic burden.3

Sancer prevention has a broad scope encompassing primary, secondary, and tertiary prevention, and research in this domain ranges from the submicroscopic study of the mechanisms of carcinogenesis to the supramacroscopic analysis of the "causes of the causes," also known as the social determinants of health.4 Recent studies from France, the United Kingdom, and Germany have estimated that around 40% of cancers in these countries could be prevented.<sup>5–9</sup> Established means of primary prevention include legislation and policies (e.g., on tobacco, alcohol, or hazardous agents), vaccination programs (e.g., human papilloma

virus or hepatitis B), and education about healthy lifestyles and behaviors (e.g., tobacco, alcohol, diet, and UV exposure), as well as other risk-adapted medical prevention strategies (e.g., tamoxifen or aspirin). By cancer type, a

International Agency for Research on Cancer (IARC/WHO), 150 cours Albert Thomas, 69372 Lyon CEDEX 08, France <sup>2</sup>Usher Institute, College of Medicine and Veterinary Medicine, University of Edinburgh, Edinburgh EH8 9AG, United Kingdom, UK <sup>3</sup>Cancer Research UK, 407 St John Street, London ECIV 4AD, UK <sup>4</sup>European Institute of Oncology, Via Ripamonti, 435 - 20141 Milano, Italy <sup>5</sup>German Cancer Research Center (DKFZ), Im Neuenheimer Feld 280, 69120 Heidelberg, Germany

<sup>6</sup>UK Therapeutic Cancer Prevention Network, Leicester Cancer Research Centre, University of Leicester, Leicester, LE2 7LX, UK <sup>7</sup>Karolinska University Laboratory, Karolinska University Hospital, 14186 Stockholm, Sweden

<sup>8</sup>Division of Human Nutrition and Health, Wageningen University, PO Box 17, 6700 AA Wageningen, the Netherlands

<sup>9</sup>Danish Cancer Society, Strandboulevarden 49, 2100 Copenhagen, Denmark

<sup>10</sup>MRC-PHE Centre for Environment and Health, School of Public Health, Imperial College, Norfolk Place, London W2 IPG, UK

11 GROW School for Oncology and Developmental Biology, Maastricht University, P.O. Box 616, 6200 MD Maastricht, the Netherlands

<sup>12</sup>World Cancer Research Fund International, London, UK

<sup>13</sup>UK Therapeutic Cancer Prevention Network, Centre for Cancer Prevention, Wolfson Institute of Preventive Medicine, Queen Mary University of London, London, UK

#### Corresponding author:

Carolina Espina, International Agency for Research on Cancer (IARC/ WHO), 150 cours Albert Thomas, 69372 Lyon Cedex 08, France. Email: cpe@iarc.fr

NP3 Tumori Journal 106(5)

proportion of 75% to 100% of all cancers of the lung, cervix, esophagus, oral cavity, melanoma, and stomach in Europe are suggested to be preventable as a consequence of changes in established risk and protective factors; similarly, 25% to 74% of all colorectal, bladder, kidney, liver, uterus, pancreas, and breast cancers are potentially preventable.<sup>10</sup> Primary prevention synergistically benefits other noncommunicable diseases by reducing exposures to shared risk factors, creating healthy environments for current and future generations. 11,12 Secondary prevention or early detection through organized screening programs can significantly reduce mortality from breast, cervical, and colorectal cancers, as well as incidence of cervical and colorectal cancers<sup>13</sup>; however, implementation of secondary prevention in Europe is scattered, as recently assessed.<sup>14</sup> Tertiary prevention refers to care aimed at reducing morbidity, disability, and risk of second primary cancer, as well as at restoring function and improving quality of life and participation in society in people diagnosed with and being treated for cancer. The potential to expand preventive interventions remains large and the evidence base for a number of measures has been summarized in the 4th edition of the European Code against Cancer (ECAC) (http://cancer-code-europe. iarc.fr/index.php/en/).15 The ECAC is an integrated instrument for cancer prevention that informs the general public how to avoid or reduce exposures to established causes of cancers, to adopt behaviors to reduce cancer risk, and to participate in vaccination and screening programs under the appropriate national guidelines. It also acts as a guide to all development of national health policies in cancer pontion, as has been shown by the adoption of the ECAN posed structure in the comprehensive National of several European countries.<sup>16</sup>

### **Cancer Prevention Europ9**

Research is required to provide new evidence-based preventive interventions (including the continued search for additional causes of cancer and to understand the factors that hamper their implementation within health care systems and in the community. In 2018, the international and multidisciplinary consortium Cancer Prevention Europe (CPE) was created to de elop world-class prevention research to be translated into effective cancer prevention guidelines and t the national and international level. 10,17 CPE is a consortium of leading European research institutions committed to prioritizing cancer prevention through cooperation between countries and programs in order to ensure a dedicated research agenda with long-term vision, and sustainable funding and infrastructure for such research. CPE will focus on expanding preventive interventions and strengthening cancer prevention in Europe by increasing awareness of the needs, the associated required resources, and reducing inequalities in access to cancer prevention across Europe. 18 CPE will be broad in scope, covering a spectrum of research

from human studies and laboratory science to policy research, as well as dissemination of the best evidence, quality indicators, and practices used. Assessment of the cost-effectiveness of different interventions, in relation to costs of treatment, care, and productivity loss, will be a core component of the initiative. Emphasis will also be placed on the evaluation of the impact of preventive interventions (primary, secondary, and tertiary) and advocacy dimensions of the prevention agenda. The agenda for CPE includes (1) research into optimizing the implementation of known preventive strategies (implementation research), (2 dissemination and research translation to inform policy and practice (advocacy), and (3) the identification of fixed risk factors and targets for prevention (improvement and expansion). This agenda will be supported by a range of platforms, networks, and infrastructures and will draw together a wide network of partners. Training and capacity building will be integral to the initiative.

# Cancer Mission for Europe: Now!

fragmented in all aspects of cancer bur-Reidence varies considerably across European Partially due to differing risk factor prevalence and stages of implementation of primary prevention<sup>18</sup>; this is specially evident in the case of tobacco smoking. 19 Even in countries at the forefront of cancer awareness, around 40% of cancers—as shown—could be prevented; instead, numbers continue to rise.<sup>20</sup> Cancer survival and mortality differs between European countries, 21,22 as well as within countries across socioeconomic groups,<sup>23</sup> even in a country of high social welfare such as Denmark.<sup>24</sup> Cancer is not just a common cause of death but often comes with physical suffering for the patient and psychological suffering for patients and their families, not counting the economic burden of €102 per citizen, as estimated in 2009 for the European Union.3 A Europe-wide cancer prevention mission to strengthen and join forces in primary and secondary prevention, as well as tertiary prevention, diagnosis, and treatment, and palliative care, well-resourced for significant improvements in all regards, is timely and urgently needed to reduce the growing cancer burden in Europe. CPE has been founded to be one of the major drivers of such a mission.

#### **Acknowledgements**

Secretariat of and projects conducted by Cancer Prevention Europe are at present exclusively supported by the Cancer Prevention Europe member institutions. The authors thank Drs. Christopher P. Wild (former Director of IARC) and David Forman (IARC) for their roles in launching Cancer Prevention Europe.

#### **Author contributions**

The article was drafted by C.E. and J.S. All authors contributed to the writing of this review article and approved its final version.

Espina et al. NP4

#### **Declaration of conflicting interest**

The authors declare that there is no conflict of interest.

#### **Disclaimer**

Where authors are identified as personnel of the International Agency for Research on Cancer / World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer / World Health Organization.

#### **Funding**

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

#### **ORCID iDs**

Carolina Espina Dhttps://orcid.org/0000-0001-6848-4687

Mangesh A. Thorat https://orcid.org/0000-0001-8673-5320

Elisabete Weiderpass https://orcid.org/0000-0003-2237-0128

#### References

- Ferlay J, Colombet M, Soerjomataram I, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries and 25 major cancers in 2018. Eur J Cancer 2018; 103: 356–387.
- Global Cancer Observatory, International Agency f Research on Cancer, http://gco.iarc.fr/ (2018, accessed February 2019).
- February 2019).
  Luengo-Fernandez R, Leal J, Gray A, et al. Economic burden of cancer across the European Union: a topolation-based cost analysis. *Lancet Oncol* 2013; 14: 163-1174.
- 4. World Health Organisation Commission on the Social Determinants of Health. Claring the Gap in a Generation: Health Equity Through Action on the Social Determinants of Health: Final Refort of the Commission on Social Determinants of Health. Geneva: World Health Organisation; 2008.
- Soerjomataram I, Shied K, Marant Micallef C, et al. Cancers related to instyle and environmental factors in France in 2015. Bar Cancer 2018; 105: 103–113.
- Brown KF, Runday H, Dunlop C, et al. The fraction of cancer attributable to modifiable risk factors in England, Wales, Scotland, Northern Ireland, and the United Kingdom in 20 K. Br. J Cancer 2018; 118: 1130–1141.
- 7. Rehrans G, Gredner T, Stock C, et al. Cancers due to excess verght, low physical activity, and unhealthy diet. *Dtsch Arztebl Int* 2018; 115: 578–585.
- Gredner T, Behrens G, Stock C, et al. Cancers due to infection and selected environmental factors. *Dtsch Arztebl Int* 2018; 115: 586–593.
- Mons U, Gredner T, Behrens G, et al. Cancers due to smoking and high alcohol consumption. *Dtsch Arztebl Int* 2018; 115: 571–577.

 Forman D, Bauld L, Bonanni B, et al. Time for a European initiative for research to prevent cancer: a manifesto for Cancer Prevention Europe (CPE). J Cancer Policy 2018; 17: 15–23.

- United Nations General Assembly. United Nations General Assembly resolution 66.2, Sixty-sixth session, New York, 24 January 2012, Political Declaration of the High-Level Meeting of the General Assembly on the Prevention and Control of Non-Communicable Diseases (A/RES/66/2).
- 12. World Cancer Research Fund: American Institute for Cancer Research. Diet, Nutrition, Physical Activity and Cancer: A Global Perspective. Continuous Update Project Expert Report 2018. Washington, DC: World Cancer Research Fund/American Institute for Cancer Research; 2018.
- Armaroli P, Villain P, Suonio E, et M. European Code Against Cancer, 4th edition: center screening. *Cancer Epidemiol* 2015; 39 (Suppl N: 32)–S152.
- Basu P, Ponti A, Anttila Co. I. Status of implementation and organization of cancer screening in the European Union Member States: surfacery results from the second European screening report. *Int J Cancer* 2018; 142: 44–56.
- Schüz J, Espine Villain P, et al. European Code Against Cancer, 4th Option: 12 ways to reduce your cancer risk. Cancer Epidymiol 2015; 39 (Suppl 1): S1–S10.
- 16. Espined Soerjomataram I, Forman D, et al. Cancer prevention pency in the EU: best practices are now well recognist; no reason for countries to lag behind. *J Cancer Policy* 2018; 18: 40–51.
- Wild CP, Espina C, Bauld L, et al. Cancer Prevention Europe. *Mol Oncol*. Epub ahead of print 22 Jan 2019. doi: 10.1002/1878-0261.12455.
- Schüz J, Espina C and Wild CP. Primary prevention: a need for concerted action. *Mol Oncol*. Epub ahead of print 24 Dec 2018. doi: 10.1002/1878–0261.12432.
- Leon ME, Peruga A, McNeill A, et al. European Code against Cancer, 4th edition: tobacco and cancer. *Cancer Epidemiol* 2015; 39 (Suppl 1): S20–S33.
- 20. Forbes LJ, Simon AE, Warburton F, et al. Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *Br J Cancer* 2013; 108: 292–300.
- Allemani C, Weir HK, Carreira H, et al. Global surveillance of cancer survival 1995–2009: analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet* 2014; 385: 977–1010.
- 22. Munro AJ. Comparative cancer survival in European countries. *Br Med Bull* 2014; 110: 5–22.
- Vaccarella S, Lortet-Tieulent J, Saracci R, et al. Reducing social inequalities in cancer: setting priorities for research. CA Cancer J Clin 2018; 68: 324–326.
- Dalton SO, Schüz J, Engholm G, et al. Social inequality in incidence of and survival from cancer in a population-based study in Denmark, 1994–2003: summary of findings. *Eur J Cancer* 2008; 44: 2074–2085.