

RESEARCH ARTICLE

Open Access



# Variability in newborn telomere length is explained by inheritance and intrauterine environment

Li Chen<sup>1\*</sup> , Karen Tan Mei Ling<sup>1</sup>, Min Gong<sup>1</sup>, Mary F. F. Chong<sup>1,2</sup>, Kok Hian Tan<sup>3</sup>, Yap Seng Chong<sup>1,4</sup>, Michael J. Meaney<sup>1,5,6</sup>, Peter D. Gluckman<sup>1,7</sup>, Johan G. Eriksson<sup>1,6,8,9</sup> and Neerja Karnani<sup>1,10,11\*</sup>

## Abstract

**Background:** Telomere length (TL) and its attrition are important indicators of physiological stress and biological aging and hence may vary among individuals of the same age. This variation is apparent even in newborns, suggesting potential effects of parental factors and the intrauterine environment on TL of the growing fetus.

**Methods:** Average relative TLs of newborns (cord tissue,  $N = 950$ ) and mothers (buffy coat collected at 26–28 weeks of gestation,  $N = 892$ ) were measured in a birth cohort. This study provides a comprehensive analysis of the effects of heritable factors, socioeconomic status, and in utero exposures linked with maternal nutrition, cardiometabolic health, and mental well-being on the newborn TL. The association between maternal TL and antenatal maternal health was also studied.

**Results:** Longer maternal TL ( $\beta = 0.14$ ,  $P = 1.99\text{E}-05$ ) and higher paternal age ( $\beta = 0.10$ ,  $P = 3.73\text{E}-03$ ) were positively associated with newborn TL. Genome-wide association studies on newborn and maternal TLs identified 6 genetic variants in a strong linkage disequilibrium on chromosome 3q26.2 (Tag SNP-*LRRC34*-rs10936600:  $P_{\text{meta}} = 5.95\text{E}-08$ ). Mothers with higher anxiety scores, elevated fasting blood glucose, lower plasma insulin-like growth factor-binding protein 3 and vitamin B12 levels, and active smoking status during pregnancy showed a higher risk of giving birth to offspring with shorter TL. There were sex-related differences in the factors explaining newborn TL variation. Variation in female newborn TL was best explained by maternal TL, mental health, and plasma vitamin B12 levels, while that in male newborn TL was best explained by paternal age, maternal education, and metabolic health. Mother's TL was associated with her own metabolic health and nutrient status, which may have transgenerational effects on offspring TL.

**Conclusions:** Our findings provide a comprehensive understanding of the heritable and environmental factors and their relative contributions to the initial setting of TL and programming of longevity in early life. This study provides valuable insights for preventing in utero telomere attrition by improving the antenatal health of mothers via targeting the modifiable factors.

**Trial registration:** [ClinicalTrials.gov](https://clinicaltrials.gov), NCT01174875. Registered on 1 July 2010

**Keywords:** Newborn, Telomere length, Intrauterine exposures, Inheritance, Sex differences

\* Correspondence: [chen\\_li@sics.a-star.edu.sg](mailto:chen_li@sics.a-star.edu.sg); [neerja\\_karnani@sics.a-star.edu.sg](mailto:neerja_karnani@sics.a-star.edu.sg)

<sup>1</sup>Singapore Institute for Clinical Sciences, A\*STAR, Singapore, Singapore  
Full list of author information is available at the end of the article



© The Author(s). 2022 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Telomeres are nucleoprotein structures formed of non-coding tandem repeats of hexamer TTAGGG at the end of the chromosomes. Functionally, telomeres play an important role in maintaining genomic stability and protect chromosomes from end-to-end fusion and degradation [1]. Telomeres shorten with each cell division, reflecting the age of a cell and the time until senescence [1]. Telomere (TL) is a biomarker of biological aging and has been associated with physiological and environmental stress, age-related diseases, and early mortality [2, 3]. Sex differences have also been reported in multiple studies with females having longer TL than males [4]. Shortened leukocyte TL has been associated with smoking [5], type 2 diabetes [6], higher 2-h post-load glucose concentration [7], lower insulin-like growth factor 1 [8], and higher leptin [9] and homocysteine levels [10]. Multiple studies reported telomere shortening to be associated with hypertension [11], cardiovascular disease [12], and mental disorders [13]. Some studies reported diet [14], parity [15, 16], and educational attainment [17] to explain inter-individual variation in TL.

In addition to diet, lifestyle, and other environmental exposures, heritable factors such as genetic variants have also been recognized to play a significant role in determining an individual's TL. Genome-wide association studies (GWAS) have been performed on TL in many adult studies, which have identified TL-linked genetic variants at multiple loci in the human genome: *TERC* (3q26.2) [18–22], *TERT* (5q15.33) [20, 21, 23], *RTEL1* (20q13.33) [21, 24], *OBFC1* (10q24.3) [20, 21], *NAF1* (4q32.2) [21], *ZNF208* (19p12) [21], and *ACY2* (2p16.2) [21].

Newborn TL is the initial setting of TL and highly variable among individuals [25]. It has important implications on telomere dynamics and molecular longevity over the lifespan [26, 27]. Current advances in newborn TL research indicate plasticity in the programming of telomere biology and the initial setting of TL [28]. The in utero environment has been suggested to be a significant contributor to this effect. Cord blood TL association studies with specific in utero exposures such as maternal nutrient status (e.g., folate [29] and vitamin D [30]), smoking status [31], educational attainment [32], or metabolic [33–35] and mental health [36] have provided valuable insights. Offspring TL has also been reported to associate with parental age and TL [37–41]. However, as the existing studies have primarily focused on individual exposures, a comprehensive understanding of the magnitude of their independent effects on offspring TL is lacking in the field. Also, though leukocyte TL in newborn is known to differ by sex, the sexual dimorphism in the factors explaining newborn TL variation is underexplored.

This study provides the comprehensive analysis of the effects of heritable factors, socioeconomic status, and in utero exposures linked with maternal nutrition, cardio-metabolic health, and mental well-being on the newborn TL. Sex-specific effects and the contributions of factors to variation in newborn TL were assessed. This comprehensive analysis was feasible due to the availability of a large sample size ( $N = 950$ ) with deep phenotyping and genomics data generated in the Growing Up in Singapore Towards healthy Outcomes (GUSTO) birth cohort.

## Methods

### Study population

We used data from the Growing Up in Singapore Towards healthy Outcomes (GUSTO) study, which is a prospective mother-offspring birth cohort study designed for investigating developmental origins of health and disease (DOHaD). The GUSTO study recruited 1247 pregnant women in their first trimester of pregnancy from two major public hospitals in Singapore, KK Women's and Children's Hospital (KKH) and National University Hospital (NUH), between June 2009 and September 2010 [42]. It conducted extensive maternal assessments at 26–28 weeks of gestation and assessments of offspring development and behavior from birth onwards. Participants could be of Chinese, Malay, or Indian ethnicity, but with homogeneous parental ethnic backgrounds.

### Tissue collection and DNA extraction

#### Umbilical cord

Detailed information of collection and processing for the cord tissue has been described previously [43]. Briefly, umbilical cord tissue samples were collected after the extraction of cord blood and then cleaned with phosphate-buffered saline solution. The cord samples were snap-frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until subsequent DNA extraction. Before DNA extraction, frozen umbilical cords were pulverized and treated with 10 U/mL hyaluronidase enzyme and then incubated at  $37^{\circ}\text{C}$  for 30 min. The tissue was homogenized using a Xiril Dispo-mix homogenizer after adding 250  $\mu\text{L}$  of Tris-NaCl-EDTA-SDS solution. Samples were pulse spun to pellet the tissue prior to adding proteinase K, and incubated overnight at  $55^{\circ}\text{C}$ . DNA extraction from the lysates was performed using the QIASymphony Midi DNA kit (QIAGEN), as per the manufacturer's instructions. Genomic DNA integrity of DNA samples was evaluated by agarose gel electrophoresis to ensure no apparent DNA degradation (Fig. S1A).

#### Maternal blood

Up to 20 mL of blood was collected from the peripheral vein of pregnant women at 26–28 weeks of gestation

into EDTA tubes. Blood samples were then centrifuged within 4 h at 1600 g at 4 °C for 10 min to separate the blood into three distinct layers—plasma, buffy coat, and erythrocytes. The top plasma layer was then carefully extracted (without disturbing the buffy coat), followed by extraction of the buffy coat layer. The buffy coat was stored at −80 °C. DNA extraction from the buffy coat was performed using either the QIAasyphony Midi DNA kit (QIAGEN) as per the manufacturer's instructions or phenol/chloroform extraction. Briefly, an equal volume of phenol/chloroform was added to the buffy coat and mixed, followed by centrifugation at 13,200 rpm for 10 min. DNA was precipitated from the top aqueous layer using 1/10th volume of 3 M NaAc and 2 volumes of ice-cold 100% ethanol. The DNA pellet was obtained by centrifugation at 13,200 rpm for 10 min and air-dried at 60 °C before dissolving in double-distilled water. Genomic DNA integrity of DNA samples was evaluated by agarose gel electrophoresis to ensure no apparent DNA degradation (Fig. S1B).

#### **Sample/DNA procedures**

Cord tissue and maternal blood samples were collected within 1.5 years. Sample collection month was divided into four quarters in cord tissue samples (January to March, 33.8%; April to June, 16.3%; July to September, 21.6%; and October to December, 28.3%) and maternal blood samples (January to March, 18.5%; April to June, 20.3%; July to September, 27.0%; and October to December, 34.2%). Sample storage time was varied from 4~6 years for cord tissue samples and 3~4 years for maternal blood samples. DNA storage time (−80 °C freezer) was varied from 0~2 years (81.3%, < 1 year, and 18.7%, ~ 2 years) for cord tissue and 3~4 years (17.8%, ~ 4 years, and 82.2%, ~ 3 years) for maternal blood samples.

#### **Average relative telomere length measurement**

Average relative TLs of cord tissue and maternal buffy coat samples were measured by the Blackburn Laboratory (<https://blackburnlab.ucsf.edu>). Cord tissue and maternal samples were measured separately and run as two batches, using the same reagent lots for each batch. They were measured by a modified quantitative real-time PCR (qPCR) protocol which has been described previously in detail [44]. Briefly, this method generates a measure of the average TL of each DNA sample as a ratio (T/S) of telomere repeat length (T) to the copy number of a single-copy gene (S, human beta-globin). All PCRs were performed on a Roche LightCycler 480 real-time PCR machine with 384-well capacity. Eight control DNA samples were included in each run to account for inter-assay variability. The T/S ratio of each control DNA was divided by the average T/S for the same DNA from 10 runs to obtain a normalizing factor. This

procedure was implemented for all eight control samples, and then the average normalizing factor was used to correct the participant DNA samples to obtain the final T/S ratio. Each sample was measured twice for the T/S ratio. But if the difference between the two measurements varied by more than 7%, the sample was run a third time and then the average of the two closest values was reported. The above method was used for measuring TLs of newborn and maternal DNA samples. Nine assay plates were used for cord tissue samples, and ten assay plates were used for maternal samples. The inter-assay coefficient of variant (CV) for newborn TL was 2.5%, and the CV for maternal TL was 2.7%. The intra-class correlation coefficient (ICC) [45] was 0.924 (95%CI 0.913–0.934) for newborn TL and 0.978 (95%CI 0.975–0.980) for maternal TL (Additional file 1: Table S1).

A pilot study was conducted for the comparison of the average relative TL of cord blood and cord tissue using 20 subjects.

#### **Genotype data**

Both newborn and maternal DNA samples were genotyped on Illumina HumanOmniExpressExome arrays. Samples with a call rate less than 97%, cryptic relatedness, or sex/ethnic discrepancies were excluded. SNPs with call rates less than 95% or minor allele frequency less than 5% or Hardy-Weinberg equilibrium *P* value less than 1.00E−06 and non-autosomal SNPs were excluded.

#### **Newborn sex, birth weight, gestational age, and paternal age**

Newborn sex and birth weight were extracted from the medical records. Gestational age (GA) was assessed by ultrasonography by trained ultrasonographers. GA was first assessed during the first trimester of pregnancy and reported in completed weeks. Paternal age was collected at recruitment by interviewer-administered questionnaires. Five paternal subjects with age > 55 years were excluded from the analysis.

#### **Maternal characteristics during pregnancy Demographic and anthropometric data**

At enrollment, interviewer-administered questionnaires were used to collect information on age, ethnicity (Chinese, Malay, and Indian), educational attainment (secondary and below, post-secondary, and university), monthly household income (≤ S\$1999, 2000–5999, and ≥ 6000), and pre-pregnancy weight. Parity (primiparous and multiparous) was extracted from hospital medical records.

Weight and height were measured at 26–28 weeks of gestation. Gestational weight gain (GWG) was calculated as the difference between pre-pregnancy weight and weight at 26–28 weeks' gestation. Pre-pregnancy body

mass index (ppBMI) was calculated as pre-pregnancy weight divided by height squared. Peripheral systolic pressure and diastolic blood pressure were measured from the brachial artery at 26–28 weeks' gestation.

#### **Maternal mental health**

Information on depressive and anxiety symptoms was obtained by questionnaires at 26–28 weeks of gestation. The Edinburgh Postnatal Depression Scale (EPDS) was used to assess the depressive symptoms by 10 items of common depressive symptoms over the past week. Anxiety was assessed by the Spielberger State-Trait Anxiety Inventory (STAI), which consists of 40 items with a 4-point Likert scale. Twenty items assess the state measure which reflects transient characteristics of anxiety (i.e., anxiety disorders), while the other twenty items were used to assess the trait measure, reflecting a more stable personality characteristic, such as an anxious personality.

#### **Maternal plasma glucose concentration**

The participants underwent an oral glucose tolerance test (OGTT) at the 26th–28th week of pregnancy visit [46]. Fasting plasma glucose concentrations (FPG) were measured after an overnight fasting (8–14 h), and 2-h post-load glucose concentrations (2-h PG) were measured at 2 h after taking 75 g of glucose. Gestational diabetes mellitus (GDM) status was diagnosed on the basis of the World Health Organization 1999 criteria:  $\geq 7.0$  mmol/L for FPG and/or  $\geq 7.8$  mmol/L for 2-h PG [47]. Three subjects were excluded with  $FPG > 7.0$  mmol/L since they were diabetic. In this study, all GDM cases are diagnosed based on the cutoff of 2-h post-load glucose concentrations.

#### **Maternal plasma fatty acids, vitamins, metabolites, and biomarkers**

Blood was drawn at 26–28 weeks' gestation into EDTA tubes. Blood samples were centrifuged at 1600 g at 4 °C for 10 min within 4 h and stored at  $-80$  °C prior to analysis. Plasma fatty acids, vitamins, metabolites, and biomarkers were measured as reported previously [48–52].

Plasma phosphatidylcholine (PC) fatty acids were measured, and the fatty acids were expressed in percentage of total fatty acids in plasma PC. We investigated the total saturated fatty acids (SFA), the total mono-unsaturated fatty acids (MUFA), the total n-3 poly-unsaturated fatty acids (PUFA), the total n-6 PUFA, and the ratio of them (n-6:n-3 PUFA). We also investigated the top three abundant fatty acids in n-3 PUFA (docosa-hexaenoic acid (DHA), docosapentaenoic acid (DPA), and eicosapentaenoic acid (EPA)) and n-6 PUFA (linoleic acid (LA), dihomo-gamma-linolenic acid (DGLA), and arachidonic acid (AA)).

Vitamin B6, betaine, choline, and total homocysteine concentrations were measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS) at BEVITAL AS (<http://www.bevital.no>). Folate and vitamin B12 concentrations were measured using the ADVIA Centaur Immunoassay System. 25-Hydroxy vitamin D<sub>3</sub> was analyzed by isotope-dilution liquid chromatography-tandem mass spectrometry (ID-LC-MS/MS).

Protein biomarkers were analyzed by MILLIPLEX® Multiplex Assays Using Luminex®. We investigated adiponectin, leptin, C-reactive protein (CRP), plasminogen activator inhibitor-1 (PAI-1), insulin-like growth factors (IGF1 and IGF2), and insulin-like growth factor binding proteins (IGFBP1, IGFBP3, IGFBP4, and IGFBP7).

#### **Maternal smoking and drinking**

Maternal alcohol consumption before and during pregnancy and maternal smoking status before and during pregnancy were extracted from an interviewer-administered questionnaire at 26–28 weeks' gestation. Plasma cotinine concentration was measured by LC-MS/MS from blood samples drawn at 26–28 weeks' gestation. In order to avoid underreported smoking during pregnancy, a combined method was applied using self-reported smoking status and plasma cotinine level. Smoking status was derived when self-reported status is smoking or cotinine level is greater than 10 ng/mL [53]. Non-smoking status was identified by self-reported non-smoking and cotinine level  $\leq 10$  ng/mL.

#### **Statistical analysis**

After evaluating DNA quality and PCR performances of TL measurement, the selection criteria for newborns (APGAR  $\geq 9$  and singleton), and the availability of genotype data, 950 subjects were included in this study as they had their newborn TL measured with complete information on sex, ethnicity, maternal age, gestational age, and genotype data. The basic characteristics of these 950 newborns were not significantly different from those of a total of 1177 live singleton newborns in the cohort (Additional file 1: Table S2). A subset of participants had available paternal age, maternal TL, and maternal characteristics during pregnancy as shown in Additional file 1: Table S3. Normality was checked for all the continuous variables. As the distributions of some continuous variables were skewed, we did log<sub>10</sub> transformation and then the values were truncated to the nearest possible value if they were  $> 4$  SDs from the mean.

#### **Genome-wide association studies (GWAS)**

Trans-ethnic GWAS were applied on newborn TL ( $N = 950$ ) and maternal TL ( $N = 892$ ). For newborn or maternal genotype data, we did SNP quality control processing in each ethnicity. The SNPs that passed the quality

control in at least one ethnicity were used for trans-ethnic GWAS on newborn TL (615,105 SNPs) and maternal TL (613,953 SNPs). The genotypes of each genetic variant were coded as 0-AA, 1-Aa, and 2-aa using an additive model (A, reference allele; a, effect allele). The coded genotype data and  $z$ -scores of newborn/maternal TL were used in regression analysis. The effect size indicates the SD difference in TL per dosage change of effect allele. Sex and ethnicity were adjusted in newborn trans-ethnic GWAS ( $N = 950$ ), while ethnicity and maternal age were adjusted in maternal trans-ethnic GWAS ( $N = 892$ ). A meta-analysis of two studies was carried out using the inverse-variance-based approach (609,579 SNPs). As a measure for heterogeneity between studies, the fraction of variation across studies ( $I^2$ ) that is due to heterogeneity rather than chance and  $P$  value for the heterogeneity test ( $P_{\text{het}}$ ) were calculated. GWAS and meta-analysis were performed in PLINK 1.9.

#### **Association with newborn TL**

Linear regression was used to study the association between newborn TL (outcome) and clinical variables (predictors) using univariate and multivariate analyses. For continuous predictors,  $z$ -scores of outcome and predictor variables were used in regression analysis, and then the standardized effect sizes  $\beta$  (SD/SD) were reported in the results. For categorical (i.e., ethnicity, GDM status, smoking status, alcohol assumption, and parity) and ordinal (i.e., education and household income) predictors, only  $z$ -scores of the outcome variable were used in regression analysis, and then the effect sizes  $\beta$  (SD difference between the groups) were reported in the results. DNA extraction for all umbilical cord samples was performed using the same method. Sex and ethnicity were adjusted in multivariate analysis. Furthermore, sex stratification analysis was performed after adjustment for ethnicity. The association between newborn TL and influencing factors was examined separately in male and female newborns. The subjects with missing values were excluded in the analysis of each factor.

#### **Association with maternal TL**

The association study between maternal TL and antenatal maternal health was examined by linear regression. As maternal age, ethnicity, and DNA extraction methods of maternal blood samples were significantly associated with maternal TL, they were adjusted in the regression models. Two multivariate models were implemented to assess the significance of each factor after accounting for the effects of (1) maternal age, ethnicity, and DNA extraction method and (2) maternal age, ethnicity, DNA extraction method, and GDM status. The subjects with missing values were excluded in the analysis of each factor. The calculation of effect sizes was the same as

above. The mediating effect of maternal TL was studied for the factors that were significantly associated with both newborn and maternal TLs.

#### **Comparative assessment factors associated with newborn TL**

In order to check for independent significances of influencing factors and their contributions to variance of newborn TL, twelve significant factors listed in Table 2 were considered in this analysis. Representative variables were selected from parental age and mental health since variables are highly correlated within the same category. For parental ages, paternal age was selected for its dominant effect on newborn TL. For mental health, STAI trait score was selected for representing this category because it exhibited much stronger association than STAI state and EPDS. In consideration of the GWAS result, ten selected factors were examined in three multivariate models using all, female-only, and male-only subjects. They are newborn genetic variant *LRRC34*-rs10936600, maternal TL, paternal age, STAI trait score, fasting glucose, DGLA%, IGFBP3, vitamin B12, maternal education, and smoking status during pregnancy. In the process of model selection, sex and ethnicity were included in the basic model for all subjects, and ethnicity only was used in the basic models for male-only and female-only subjects. The forward selection method was applied to add predictors, and the Akaike information criterion was used to select the best multivariate model, in which sample size was reduced due to the missing values of required predictors.

#### **Sensitivity analysis**

As sample collection, DNA storage procedures, and seasonality have been reported to be associated with telomere length measurement [54], additional adjustment of these factors in the best multivariate models of newborn TL and the association studies between maternal TL and antenatal factors were performed. Sample storage time is highly correlated to DNA storage time in our study. Hence, only DNA storage time (two groups) and sample collection month (four quarters) were studied in the sensitivity analysis.

The association analyses with newborn TL and maternal TL were performed in MATLAB R2019b. The calculation of intra-class correlation coefficients (rptR package) and mediation analysis of maternal TL (mediation package) were carried out in R 3.6.3.

## **Results**

### **Participant characteristics and TL measures**

Depending on the availability of key variables required for this study, data from 950 of 1247 mother-offspring dyads were analyzed from the GUSTO cohort. A flow chart of sample selection and data analysis is provided in

Fig. S2. Clinical characteristics and TL measurement summary of all participants in this study are shown in Additional file 1: Table S3. These include newborn TL, maternal TL, infant sex, ethnicity, parental age, gestational age, birth weight, maternal smoking and alcohol consumption status, and a wide range of antenatal maternal cardiometabolic, mental health, and plasma nutrient measures. Both newborn and maternal TLs showed normal distribution (Figs. 1A and 2A). The average newborn TL is 2.00 with a standard deviation of 0.25, and the average maternal TL is 1.02 with a standard deviation of 0.23.

#### Sex difference and ethnic diversity in newborn TL

Newborns in the study comprised 52.6% males and 47.4% females. Notably, sex was the most significant factor ( $\beta = 0.40$ ,  $P = 4.45E-10$ ) associated with newborn TL in univariate analysis (Additional file 1: Table S3). The average TL at birth in females (mean = 2.05, SD = 0.24) was longer than that in males (mean = 1.96, SD = 0.24) (Fig. 1B). All mother-offspring dyads in this study were of Asian ethnic origin, i.e., Chinese (58.4%), Malay (24.6%), and Indian (17.0%). Compared to the TL of newborn of Chinese ethnic origin (mean = 2.02, SD = 0.25), TLs of Malay (mean = 1.97, SD = 0.24;  $\beta = -0.21$ ,  $P = 6.63E-03$ ) and Indian (mean = 1.97, SD = 0.24;  $\beta = -0.21$ ,  $P = 1.67E-02$ ) newborns were slightly shorter (Fig. 1B).

#### Paternal age is positively associated with newborn TL

Newborn TL was not associated with gestational age and birth weight (Additional file 1: Table S3). As is evident from Fig. 1C, maternal age ( $\beta = 0.07$ ,  $P = 2.78E-02$ ) and paternal age ( $\beta = 0.10$ ,  $P = 3.73E-03$ ) showed a significant positive association with the newborn TL. Since maternal age is highly correlated with the paternal age ( $R^2 = 0.50$ ) in the GUSTO cohort, we also analyzed them in the same linear regression model with sex and ethnicity as covariates. Interestingly, the association between maternal age and newborn TL was lost in such an analysis, but paternal age remained significant ( $\beta = 0.14$ ,  $P = 4.22E-03$ ). This finding indicates that among the parents, paternal age has a dominant effect on newborn TL, and hence, offsprings of older fathers are born with longer TL.

#### Inheritance/heritability of TL in newborn

To elucidate the factor of heritability in newborn TL, we first examined if maternal TL was a predictor of newborn TL. The correlation analysis confirmed this to be true as the mothers with longer TL gave birth to offsprings with longer TL (Fig. 1D;  $\beta = 0.14$ ,  $P = 1.99E-05$ ).

We next examined if the heritability of TL was influenced by genetics via performing genome-wide association studies (GWAS) of newborn and maternal

TLs. Though no SNPs passed the genome-wide significance cutoff ( $P = 5.00E-08$ ) in the individual GWASs (Fig. S3A-B), a meta-analysis of these two studies (Fig. S3C) revealed top six genetic variants (*LRRC34*-rs10936600, rs13069553, *LRRC34*-rs7621631, *MYNN*-rs1317082, *MYNN*-rs10936599, and rs12638862) within the same genomic region 3q26.2 (Table 1 and Additional file 1: Table S4). The top variant *LRRC34*-rs10936600 (missense) showed a borderline genome-wide significance ( $P = 5.95E-08$ ;  $\beta = -0.18$ ). Locus zoom plot using the 1000 genome ASN population (Fig. S3D) identified these six genetic variants to be in a strong linkage disequilibrium (LD) with each other. Their pairwise  $R^2$  in three ethnic groups are provided in Additional file 1: Table S5 ( $R^2 \geq 0.82$ ). Among these six SNPs, *LRRC34*-rs10936600 constituted the tag SNP of this LD block. This 3q26.2 genomic region has previously been reported to be associated with TL in adults [18, 19]. Notably, rs12638862 is located 4891 bp downstream of the *TERC* (telomerase RNA component) gene, which is known to be essential for telomere length maintenance. All six variants showed no observed heterogeneity between maternal and newborn genotype data ( $P_{\text{het}} > 0.1$  and  $I^2 = 0$ ).

Ethnicity-specific box plots for each of the SNPs and their allele frequencies are provided in Fig. S4-S5. It was noted that the dosage of A allele of the top variant *LRRC34*-rs10936600 was positively associated with TL. Furthermore, sex stratification results on newborn *LRRC34*-rs10936600 showed that its effect on newborn TL was much stronger in males ( $\beta = -0.23$ ,  $P = 4.46E-04$ ) than in females ( $\beta = -0.16$ ,  $P = 1.59E-02$ ).

#### Maternal antenatal health, smoking, and nutrient status have a significant impact on newborn TL

Since the variability in TL can arise from both genetic and environmental factors, we next studied the effects of maternal antenatal health, blood nutrient levels, and socioeconomic status on newborn TL. These variables included mother's mental health, adiposity, blood pressure, plasma glucose concentration, plasma fatty acids and vitamins, plasma protein biomarkers, educational attainment and household income, smoking status, alcohol consumption, and parity. Heat map of pairwise Pearson correlation coefficients between these clinical variables (continuous measures) is provided in Fig. S6. The results of univariate and multivariate regression analyses for these variables are provided in Additional file 1: Table S3.

The link between newborn TL and antenatal mental health was studied using the Edinburgh Postnatal Depression Scale (EPDS) and the State-Trait Anxiety Inventory (STAI) scores collected during mid-pregnancy (26–28 weeks). While EPDS scores showed no significant




Fig. 1 (See legend on next page.)


(See figure on previous page.)

**Fig. 1** Association studies of newborn telomere length. **A** Histogram of newborn telomere length. **B** Boxplots of sex and ethnicity (univariate *P* values). **C, D** Scatter plots of gestational age, maternal age, paternal age, and maternal telomere length (*P* values after adjusting for sex and ethnicity). **E, F** Effect size plot of factors significantly associated with newborn telomere length using all, female-only, and male-only subjects (error bar, 95%CI). Related to Table 2. **E** Categorical/ordinal variables. **F** Continuous variables

association with newborn TL, STAI scores showed strong negative correlation with newborn TL, indicating higher antenatal anxiety to be associated with shorter newborn TL. Among STAI measures, STAI trait scores ( $\beta = -0.09, P = 3.61E-03$ ) were more significantly

associated with newborn TL than STAI state scores ( $\beta = -0.07, P = 2.93E-02$ ).

For cardiometabolic health measures, maternal adiposity (ppBMI, GWG, and height), blood pressure, plasma protein biomarkers, and antenatal glycemia (fasting glucose,





**Table 1** Top six genetic variants (3q26.2) in the meta-analysis results of genome-wide association studies on newborn telomere length and maternal telomere length

SNP	Chr: position <sup>#</sup>	Allele R/E	Gene	Newborn telomere length (N = 950)		Maternal telomere length (N = 892)		Meta-analysis			
				$\beta$ (SE)	P value	$\beta$ (SE)	P value	$\beta$ (SE)	P value	$P_{het}$	$I^2$
rs10936600	3:169514585	A/T <sup>*</sup>	<i>LRRC34</i> <sup>a</sup>	-0.20 (0.05)	2.35E-05	-0.16 (0.05)	6.79E-04	-0.18 (0.03)	5.95E-08	0.58	0
rs13069553	3:169508272	A/G	Intergenic <sup>b</sup>	-0.20 (0.05)	1.57E-05	-0.16 (0.05)	9.54E-04	-0.18 (0.03)	6.00E-08	0.50	0
rs7621631	3:169512145	C/A	<i>LRRC34</i> <sup>c</sup>	-0.20 (0.05)	2.00E-05	-0.16 (0.05)	8.30E-04	-0.18 (0.03)	6.35E-08	0.54	0
rs1317082	3:169497585	A/G	<i>MYNN</i> <sup>d</sup>	-0.20 (0.05)	1.76E-05	-0.16 (0.05)	9.69E-04	-0.18 (0.03)	6.79E-08	0.50	0
rs10936599	3:169492101	G/A	<i>MYNN</i> <sup>e</sup>	-0.19 (0.05)	3.93E-05	-0.16 (0.05)	9.15E-04	-0.18 (0.03)	1.32E-07	0.59	0
rs12638862	3:169477506	G/A	Intergenic <sup>f</sup>	0.16 (0.05)	5.86E-04	0.15 (0.05)	1.18E-03	0.16 (0.03)	2.14E-06	0.91	0

$P_{het} > 0.1$  and  $I^2 = 0$  showed no observed heterogeneity

R/E reference/effect allele,  $\beta$  (SE) effect size (standard error)

<sup>#</sup>hg19 genome build

<sup>\*</sup>Forward strand

<sup>a</sup>Missense L286I

<sup>b</sup>767 bp downstream of *MYNN*

<sup>c</sup>Intron variant, 3' UTR variant

<sup>d</sup>Intron variant

<sup>e</sup>Synonymous codon

<sup>f</sup>4891 bp downstream of *TERC*

2-h post-load glucose concentrations, and GDM status) were investigated for their associations with newborn TL. Only fasting glucose ( $\beta = -0.08$ ,  $P = 1.60E-02$ ) and insulin-like growth factor-binding protein 3 (IGFBP3) ( $\beta = -0.08$ ,  $P = 7.62E-03$ ) demonstrated significant negative associations with newborn TL.

As maternal nutrition plays a critical role in fetal development, we also investigated the effects of antenatal plasma fatty acids and vitamin levels on newborn TL. Percentages of ten fatty acids and n-6:n-3 PUFA ratio in antenatal plasma were assessed, but none of them showed significant association. Among the vitamins tested, vitamin B12 level in gestation showed a significant positive association with newborn TL ( $\beta = 0.08$ ,  $P = 2.87E-02$ ), and vitamin D<sub>3</sub> level was significant only in the univariate analysis.

Two factors describing the socioeconomic status (SES) were evaluated, i.e., household income and maternal education. Both were significant only in the univariate analysis. As smoking is a well-known factor in shortening of TL, maternal smoking before and during pregnancy was assessed. Smoking status before pregnancy was not significant while maternal smoking during pregnancy (39 of 827 mothers are smokers) showed stronger effects on the newborn TL ( $\beta = -0.35$ ,  $P = 3.06E-02$ ). The effects of alcohol consumption before and during pregnancy on newborn TL were not significant; however, the outcomes from such an analysis should be interpreted with caution as the sample size ( $N = 21$ ) of mothers consuming alcohol during pregnancy was very small in the GUSTO cohort. Lastly, parity (primiparous and multiparous) was explored, but it had no significant effects on newborn TL.

From the above analyses, we concluded that a women's mental well-being, metabolic health, and nutrient and smoking status during pregnancy have significant effects on newborn TL. Figure 1E, F provides a comparison of the effect sizes of these factors.

#### Sex-specific effects of factors influencing newborn TL

In order to study the sex difference in newborn TL, both stratification and interaction analyses were performed. Stratification analysis studied sex-specific effects of factors influencing newborn TL. Table 2 summarizes the significant factors derived from multivariate regression analyses using either all, or female-only, or male-only newborns.

Interestingly, maternal TL was the only factor that showed significant association with TL in both male ( $\beta = 0.11$ ,  $P = 1.67E-02$ ) and female ( $\beta = 0.17$ ,  $P = 3.09E-04$ ) newborns, but the effects were much stronger in females. Maternal mental health showed significant negative effects on female newborn TL (EPDS:  $\beta = -0.10$ ,  $P = 4.23E-02$ ; STAI state:  $\beta = -0.13$ ,  $P = 3.79E-03$ ; STAI trait:  $\beta = -0.14$ ,  $P = 2.22E-03$ ) but no significant effects on male newborn TL. Female newborn TL-specific associations were also observed with maternal antenatal plasma vitamin B12 levels ( $\beta = 0.09$ ,  $P = 5.39E-02$ ). Overall, TL in female newborn was more susceptible to variation in maternal TL, mental health, and vitamin B12 levels (Fig. 1E, F and Fig. S7).

For male newborn TL-specific factors, parental age showed significant positive associations (paternal age:  $\beta = 0.13$ ,  $P = 3.53E-03$ ; maternal age:  $\beta = 0.11$ ,  $P = 1.60E-02$ ). Likewise, it was noted that plasma fasting glucose concentration ( $\beta = -0.12$ ,  $P = 1.39E-02$ ), DGLA% ( $\beta = -0.11$ ,  $P = 3.04E-02$ ) and IGFBP3 level ( $\beta = -0.15$ ,  $P =$

**Table 2** Linear regression results between newborn telomere length and significant factors using all, male-only, and female-only subjects

Variable	Multivariate model (adjusted for sex and ethnicity), all			Multivariate model (adjusted for ethnicity), male			Multivariate model (adjusted for ethnicity), female		
	N	$\beta$ (95%CI)	P value	N	$\beta$ (95%CI)	P value	N	$\beta$ (95%CI)	P value
Maternal telomere length (T/S)	892	0.14 (0.08, 0.20)	<b>1.99E-05**</b>	465	0.11 (0.02, 0.2)	<b>1.67E-02*</b>	427	0.17 (0.08, 0.26)	<b>3.09E-04**</b>
Paternal age (years)	805	0.10 (0.03, 0.17)	<b>3.73E-03*</b>	427	0.13 (0.04, 0.22)	<b>3.53E-03*</b>	378	0.06 (-0.05, 0.16)	2.83E-01
Maternal age (years)	950	0.07 (0.01, 0.14)	<b>2.78E-02*</b>	500	0.11 (0.02, 0.20)	<b>1.60E-02*</b>	450	0.03 (-0.06, 0.12)	5.36E-01
EPDS	918	-0.05 (-0.11, 0.02)	1.40E-01	484	-0.01 (-0.1, 0.07)	7.77E-01	434	-0.10 (-0.19, 0.00)	<b>4.23E-02*</b>
STAI State Score	895	-0.07 (-0.14, -0.01)	<b>2.93E-02*</b>	470	-0.02 (-0.11, 0.08)	7.37E-01	425	-0.13 (-0.22, -0.04)	<b>3.79E-03*</b>
STAI Trait Score	891	-0.09 (-0.16, -0.03)	<b>3.61E-03*</b>	468	-0.05 (-0.15, 0.04)	2.54E-01	423	-0.14 (-0.22, -0.05)	<b>2.22E-03*</b>
Plasma fasting glucose (mmol/L)	905	-0.08 (-0.14, -0.01)	<b>1.60E-02*</b>	476	-0.12 (-0.22, -0.02)	<b>1.39E-02*</b>	429	-0.04 (-0.13, 0.04)	3.21E-01
Plasma DGLA%	831	-0.03 (-0.09, 0.04)	4.65E-01	432	-0.11 (-0.22, -0.01)	<b>3.04E-02*</b>	399	0.04 (-0.05, 0.13)	3.28E-01
Plasma IGFBP3 (ng/mL), log <sub>10</sub>	938	-0.08 (-0.15, -0.02)	<b>7.62E-03*</b>	493	-0.15 (-0.23, -0.06)	<b>8.29E-04**</b>	445	-0.02 (-0.11, 0.07)	7.09E-01
Plasma vitamin B12 (pg/mL), log <sub>10</sub>	834	0.08 (0.01, 0.14)	<b>2.87E-02*</b>	435	0.05 (-0.04, 0.15)	2.80E-01	399	0.09 (0.00, 0.19)	<u>5.39E-02</u>
Maternal education									
1: Secondary and below	283	0.07 (-0.01, 0.16)	7.79E-02	154	0.11 (0.00, 0.23)	<u>5.28E-02</u>	129	0.03 (-0.09, 0.15)	6.55E-01
2: Post-secondary	336			172			164		
3: University	319			166			153		
Maternal Smoking during Pregnancy									
0: no	788	-0.35 (-0.67, -0.03)	<b>3.06E-02*</b>	-	-	-	-	-	-
1: yes	39								

The details of characteristics and the univariate and multivariate analysis results of all factors were provided in Additional file 1: Table S3

Underlined data indicates borderline P value

$\beta$  effect size, CI confidence interval, EPDS Edinburgh Postnatal Depression Scale, STAI State-Trait Anxiety Inventory, DGLA dihydro-gamma-linolenic acid, IGFBP3 insulin-like growth factor-binding protein 3

\* $p < 0.05$  and \*\* $p < 0.001$

8.29E-04) exhibited significant negative association with male newborn TL. A borderline significant association with male newborn TL was also found with maternal educational attainment ( $\beta = 0.11$ ,  $P = 5.28E-02$ ). Taken together, variation in newborn male TL was more explained by their parental age, maternal education, plasma fasting glucose, DGLA%, and IGFBP3 levels (Fig. 1 E, F and Fig. S7).

Association of newborn TL with smoking status before pregnancy did not explain sex differences. We were unable to perform a similar analysis for smoking status during pregnancy, as the sample size was extremely small.

Furthermore, interaction analysis was applied to study the interaction between sex and exposure factors after adjusting for ethnicity. The results revealed that only two factors (plasma DGLA%:  $P = 3.22E-02$  and plasma IGFBP3 level:  $P = 3.87E-02$ ) were significant for sex interaction.

#### Maternal TL is associated with antenatal metabolic health and the nutrient status


As significant inheritance effect from maternal TL was found on newborn TL, the association studies were performed between maternal TL and antenatal maternal factors. It was observed that maternal age ( $\beta = -0.14$ ,  $P = 2.73E-05$ ), ethnicity (Indian vs Chinese:  $\beta = -0.31$ ,  $P = 6.04E-04$ ; Malay vs Chinese:  $\beta = 0.06$ ,  $P = 0.44$ ), and DNA extraction method ( $P < 1.00E-05$ ) strongly affected maternal TL measurement, so they were adjusted in the multivariate analysis (Fig. 2B).

We found nine factors to be significantly associated with maternal TL (Additional file 1: Table S6 and Fig. S8). For antenatal glycemia, plasma 2-h post-load

glucose concentration ( $\beta = -0.09$ ,  $P = 6.62E-03$ ) and GDM status ( $\beta = -0.27$ ,  $P = 1.64E-03$ ) identified significant negative association. For maternal nutrients, total n-3 PUFA% ( $\beta = 0.08$ ,  $P = 3.23E-02$ ), n-6:n-3 PUFA ratio ( $\beta = -0.08$ ,  $P = 2.19E-02$ ), and folate level ( $\beta = -0.08$ ,  $P = 2.27E-02$ ) showed strong association with maternal TL. Among plasma protein biomarkers, PAI-1 ( $\beta = -0.07$ ,  $P = 2.40E-02$ ) and IGF2 ( $\beta = 0.07$ ,  $P = 3.25E-02$ ) levels presented significant associations with maternal TL. In addition, maternal educational attainment ( $\beta = 0.09$ ,  $P = 2.73E-02$ ) and maternal smoking during pregnancy ( $\beta = -0.38$ ,  $P = 2.42E-02$ ) were also significantly associated with maternal TL. A comparison of effect sizes for all the antenatal factors associated with maternal TL is shown in Fig. 2C, D.

As GDM status showed the strongest association with maternal TL, an additional model was used to identify the interdependencies between GDM status and other significant factors by adjusting for maternal age, ethnicity, DNA extraction method, and GDM status (Additional file 1: Table S6). Interestingly, most of factors showed consistent significance except plasma folate level. Plasma folate level has been reported to be positively associated with 2-h post-load glucose concentration in the same cohort [55], which may explain the negative association between plasma folate level and maternal TL.

Overall, maternal TL is strongly associated with antenatal factors, especially metabolic health and nutrient status, and consequently, these may have a transgenerational bearing on the offspring's TL.



**Fig. 3** Schematic diagram summarizing the factors contributing to newborn TL variation

### Comparative assessment factors associated with newborn TL

A schematic summary of all the heritable factors and the exposures contributing to the variation in newborn TL is shown in Fig. 3. In order to compare the contributions of individual factors in influencing the newborn TL variation, ten selected factors were studied by the model selection method using either all, male-only, or female-only subjects (the “Methods” section). Table 3 shows the results of the multivariate models that best explained newborn TL.

Seven factors were found to best explain newborn TL variation in all subjects ( $N = 721$ ). Among these, sex was the most significant factor ( $\beta = 0.44$ ,  $P = 7.42E-10$ ), followed by *LRRC34*-rs10936600 ( $\beta = -0.20$ ,  $P = 1.70E-04$ ), paternal age ( $\beta = 0.13$ ,  $P = 2.16E-04$ ), maternal TL ( $\beta = 0.12$ ,  $P = 4.92E-04$ ), plasma IGFBP3 level ( $\beta = -0.08$ ,  $P = 1.45E-02$ ), plasma fasting glucose concentration ( $\beta = -0.08$ ,  $P = 2.42E-02$ ), and ethnicity (Indian vs Chinese:  $\beta = -0.22$ ,  $P = 3.73E-02$ ). This model explained 12.4% of variance in newborn TL (Fig. S9A).

For male newborns, five key factors were identified in the best multivariate model ( $N = 378$ ) that explained 12.6% of variance in TL (Fig. S9B). Paternal age was the most significant factor ( $\beta = 0.20$ ,  $P = 3.43E-05$ ), followed by plasma IGFBP3 level ( $\beta = -0.18$ ,  $P = 1.41E-04$ ), *LRRC34*-rs10936600 ( $\beta = -0.23$ ,  $P = 1.53E-03$ ), plasma fasting glucose concentration ( $\beta = -0.15$ ,  $P = 3.91E-03$ ), and maternal TL ( $\beta = 0.11$ ,  $P = 3.16E-02$ ). Notably, paternal age, fasting glucose, and IGFBP3 levels were male TL-specific factors.

In female newborns, four factors were identified in the best multivariate model ( $N = 405$ ) that explained 7.91% of variance in TL (Fig. S9C). Maternal TL showed the strongest association ( $\beta = 0.16$ ,  $P = 5.37E-04$ ). STAI trait score ( $\beta = -0.13$ ,  $P = 4.12E-03$ ) ranked second, followed by ethnicity (Malay vs Chinese:  $\beta = -0.33$ ,  $P = 4.16E-03$ ) and *LRRC34*-rs10936600 ( $\beta = -0.15$ ,  $P = 2.86E-02$ ). STAI trait score was a female TL-specific factor.

### Sensitivity analysis

For sensitivity analysis, DNA storage time and sample collection month were further adjusted in the association studies with newborn and maternal TLs, respectively. For the best multivariate models of newborn TL (Table 3), these two factors did not show significant association after adding them in the models (Additional file 1: Table S7). For maternal TL, the association results were slightly affected (Additional file 1: Table S8). The associations of maternal TL with plasma PAI-1 and IGF2 levels lost significance in the supplementary model.

### Discussion

To the best of our knowledge, this is the first study to investigate newborn TL variation in umbilical cord tissue and report a comprehensive analysis of the effects of heritable factors, socioeconomic status, antenatal maternal health, and nutrition on this variation. It is known that TL is maximal at birth and decreases progressively with advancing age, and thus is considered a marker of biological aging [26]. In our study, we found relative

**Table 3** The contributing factors that best explained newborn telomere length using all, male-only, and female-only subjects by model selection method. In each multivariate model, the factors were analyzed simultaneously for comparing their contributions to variation in newborn telomere length

Variable	All ( $N = 721$ )		Male ( $N = 378$ )		Female ( $N = 405$ )	
	$\beta$ (95%CI)	P value	$\beta$ (95%CI)	P value	$\beta$ (95%CI)	P value
Sex						
Male	Ref	Ref	–	–	–	–
Female	0.44 (0.30,0.58)	<b>7.42E-10***</b>				
Ethnicity						
Chinese	Ref	Ref	Ref	Ref	Ref	Ref
Malay	-0.17 (-0.34, 0.00)	5.65E-02	-0.10 (-0.32, 0.13)	4.01E-01	-0.33 (-0.56, -0.11)	<b>4.16E-03*</b>
Indian	-0.22 (-0.43, -0.01)	<b>3.73E-02*</b>	-0.20 (-0.50, 0.09)	1.83E-01	-0.21 (-0.48,0.05)	1.19E-01
Newborn <i>LRRC34</i> -rs10936600 (0-AA 1-AT 2-TT)	-0.20 (-0.30, -0.10)	<b>1.70E-04**</b>	-0.23 (-0.37, -0.09)	<b>1.53E-03*</b>	-0.15 (-0.28,-0.02)	<b>2.86E-02*</b>
Maternal telomere length (T/S)	0.12 (0.05, 0.19)	<b>4.92E-04**</b>	0.11 (0.01, 0.21)	<b>3.16E-02*</b>	0.16 (0.07, 0.25)	<b>5.37E-04**</b>
Paternal age (years)	0.13 (0.06, 0.20)	<b>2.16E-04**</b>	0.20 (0.10, 0.29)	<b>3.43E-05**</b>	–	–
Plasma fasting glucose (mmol/L)	-0.08 (-0.14, -0.01)	<b>2.42E-02*</b>	-0.15 (-0.24, -0.05)	<b>3.91E-03*</b>	–	–
Plasma IGFBP3 (ng/mL), log <sub>10</sub>	-0.08 (-0.15, -0.02)	<b>1.45E-02*</b>	-0.18 (-0.27, -0.09)	<b>1.41E-04**</b>	–	–
STAI Trait Score	–	–	–	–	-0.13 (-0.22, -0.04)	<b>4.12E-03*</b>

\* $P < 0.05$ ; \*\* $P < 0.001$ ; \*\*\* $P < 1.00E-06$

average TL of newborns to be longer than the length of TL observed in their mothers (average maternal age, 31 years old). Maternal TL negatively correlated with maternal age ranging from 18 to 46 years. Newborn TL was not associated with gestational age but showed positive association with parental age. Comparison of the paternal and maternal age effects in the same regression model showed paternal age to have a dominant effect. This finding suggests that offsprings of older mothers could have longer TL in the analysis simply because the offsprings' fathers were also older. Supporting this hypothesis, we indeed found a strong correlation between the parents' age in our cohort. Previous studies [37–39] have reported increase in sperm TL with age as a potential reason for offsprings of older fathers to inherit longer telomeres. As oocytes are produced prenatally, while sperm are continually produced throughout life, it is believed that there is greater potential for TL plasticity with age in sperms than in oocytes. Effect of paternal age on newborn TL is intriguing as it potentiates a scenario of intergenerational genetic plasticity in which the DNA passed on to the offspring is systematically changed based upon the reproductive age of one's father.

Genetic variants are additional factors known to have heritable effects on TL. The genome-wide association analysis in this study identified a LD block within the 3q26.2 region to be significantly associated with both maternal TL and newborn TL. *LRRC34*-rs10936600 was the top variant in this LD block. By using LDproxy and CHB population data [<https://analysistools.nci.nih.gov/LDlink>], we were able to extract the full list of genetic variants in a strong LD ( $R^2 > 0.85$ ) with rs10936600 (Additional file 1: Table S9). This analysis identified *TERC* variant rs2293607 ( $R^2 = 0.98$ ) to also be a member of this LD block. *TERC* encodes a long non-coding RNA found in eukaryotes that is a component of telomerase, the enzyme used to extend telomeres. The T allele of this variant (correlated to A allele in rs10936600) has been reported to be associated with an increase in *TERC* expression and TL [56]. Our finding is consistent with this reported trend as the dosage of A allele in rs10936600 is positively associated with TL (Fig. S4A). Further, the association results of previously reported genetic variants (*TERT*, *RTEL1*, *OBFC1*, *NAF1*, *ZNF208*, and *ACYP2*) are shown in Additional file 1: Table S10. They showed a weak association with TL in our study. Among them, *TERT*-rs2853677 (telomerase reverse transcriptase) showed the most significant association with maternal TL ( $\beta = 0.18$ ,  $P = 1.85E-4$ ).

For the previously reported factors associated with cord blood TL [25–36], we similarly found significant associations for sex, parental age, maternal TL, antenatal stress, maternal educational attainment, maternal smoking during pregnancy, and genomic region at 3q26.2, but

the associations with maternal pre-pregnancy BMI, GDM status, hypertension, plasma folate, and vitamin D concentrations were not found in our study. As our pilot study (the “Methods” section) showed cord blood and cord tissue TLs were highly correlated ( $R^2 = 0.64$ , Fig. S10), it is not surprising that we find similar associations with many factors. However, the discrepancy between our findings with the previous reports could be due to the sample size, population effects, and tissue-specific differences. In our study, we identified five new significant factors influencing newborn TL: ethnicity (Asian population), plasma fasting glucose concentration, plasma IGFBP3 level, plasma DGLA%, and vitamin B12 level (Table 2).

Sex-specific analysis identified distinct male vs female effects of factors influencing newborn TL. Interestingly, heritable factors such as maternal TL and newborn *LRRC34*-rs10936600 showed significant effects on both male and female newborn TLs with different effect sizes. *LRRC34*-rs10936600 showed a stronger effect on male newborns TL, while maternal TL had more pronounced effects on the female newborn TL. For other influencing factors, sex-specific effects were also observed. Female newborn TLs were more susceptible to the variation in maternal mental health (depression/anxiety) and vitamin B12 levels, while male newborn TLs were strongly affected by the variation of paternal age, maternal educational attainment, plasma fasting glucose concentration, plasma DGLA%, and IGFBP3 level. Although the negative impact of antenatal maternal distress on newborn TL has been widely reported, this study is the first to note that anxiety scores (STAI state and trait) have much stronger effects than depression scores (EPDS) on TL, and all scores exhibit female-specific effects. Low vitamin B12 level was linked to negative impacts on cognitive, motor, and growth outcomes for fetal development and was related to depression in mothers [57]. Interestingly, vitamin B12 level showed similar female-specific effects as did antenatal depression/anxiety. The effect of paternal age on offspring's TL was linked to telomere elongation in sperm from older men as reported previously [39]. It is interesting to observe the male-specific effect of paternal age on newborn TL. For maternal educational attainment, the male-specific effect is consistent with a recent report that highlighted the male-specific effect of parental SES on newborn TL [58]. Impaired glucose metabolism and the related hormonal imbalance may generate physiological stress for the growing fetus [59] and may lead to TL attrition, especially for the male offspring. IGFBP3 is the main insulin-like growth factor transport protein in the bloodstream and plays an important role in senescence as an aging marker [60]. We observed a negative effect on male newborn TL. For all the TL-associated factors, we

identified eight independent predictors of newborn TL (Table 3). These include sex, ethnicity, newborn *LRR34*-rs10936600, maternal TL, paternal age, antenatal anxiety, plasma fasting glucose, and IGF1. The contributions of these factors influencing the variation in newborn TL were compared and ranked in three best multivariate models using all, female-only, and male-only subjects.

In the association study between maternal TL and antenatal maternal health, our results were similar to the findings from previous studies on TL associations with age [1], diabetes [6], 2-h post-load glucose concentration [7], educational attainment [17], and smoking status during pregnancy [5]. Although n-3 PUFA supplementation has been reported to affect TL [14], our study for the first time showed that higher plasma n-3 PUFA% and lower n-6:n-3 PUFA ratio are associated with longer maternal TLs. In addition, previous reports showed telomere shortening was associated with cardiovascular disease [12]. Elevated plasminogen activator inhibitor-1 (PAI-1) is a risk factor for thrombosis and atherosclerosis and associated with major adverse cardiovascular events (MACE) [61]. Notably, our study showed that shorter maternal TL was associated with higher plasma PAI-1 level. Finally, plasma IGF2 levels were positively associated with maternal TL. Higher circulating levels of IGF1, a related protein of IGF2, are known to be associated with longer TL in healthy subjects [62]. IGFs are known to play an essential role in the pathogenesis of several age-related diseases, including dementia, cardiovascular, and metabolic diseases; hence, their levels in circulation play a significant role in predicting healthspan and biological aging. The association of maternal TL with plasma PAI-1 and IGF2 levels should be interpreted with caution as their associations lost significance in the sensitivity analysis.

While comparing the factors influencing the newborn and maternal TLs, we found some consistent trends. For example, higher maternal educational attainment showed a positive association, and smoking during pregnancy showed a negative association with TL in both the mother and the offspring. The mediation analysis showed these two factors affected newborn TL through the mediating effect of maternal TL (Fig. S11). We also observed both mother and newborn TL to be associated with antenatal glycemia. As fasting and 2-h post-load glucose concentrations have different underlying etiologies and pathophysiologies, different effects on newborn/maternal TL are expected. It was noteworthy that maternal GDM or higher 2-h post-load glucose concentration was significantly associated with reduced TL only in mothers, while fasting glucose concentration had an adverse impact only on offsprings. Impaired glucose tolerance has been previously shown to associate with

impaired telomerase activity [63] which may explain the shortening of TL in mothers. Although both impaired glucose tolerance and impaired fasting glucose can be harmful to the growing fetus [59], our findings showed that the latter had a more detrimental effect on fetal TL. Finally, we found that the association between antenatal depression/anxiety and TL was observed in the offspring but not in the mothers. This may be attributed to a high vulnerability of the fetus to stress during intrauterine development.

Our study has some limitations. First, the GUSTO cohort had a disproportionate number of participants within the Chinese, Malay, and Indian groups. Therefore, analyses of ethnic-specific effects of TL associated factors were limited by sample size. Second, due to the absence of paternal TL data in the cohort, we were unable to compare it with the effects of maternal TL on newborn TL. Finally, pre-pregnancy maternal weight, smoking status and alcohol consumption before/during pregnancy, and SES (maternal education and household income) were drawn from self-reported questionnaires. Bias may exist in the self-reported variables. Pre-pregnancy weight was assessed to be reliable as it highly correlated with the maternal weight at the first-trimester visit. In order to avoid underreported smoking, maternal smoking status during pregnancy was determined by the combination results of self-reported questionnaires and plasma cotinine level. As the DNA from maternal blood was extracted using two different methods (manual and automated), this technical variation could have impacted the TL measures. Hence, we adjusted the analysis models for this technical variation.

Telomeres serve as the biological timekeepers of cellular health, and hence, their attrition relative to chronological age is an indication of advanced biological aging, existence of a stressed environment, and potential risk to disease. Our study found antenatal maternal health to be a crucial determinant of TL programming in utero, which could potentially impact subsequent offspring health outcomes over the life span, including aging and longevity. Hence, improving antenatal health of mothers by targeting modifiable factors can help prevent in utero telomere attrition and enhance cellular longevity. As evident from the findings of this study, smoking during pregnancy, maternal nutrient insufficiency (i.e., vitamin B12), and cardiometabolic and mental health adversities are suboptimal conditions for fetal programming of telomere biology. In addition, our findings in sex-specific associations suggest different focus of antenatal care may be required for mothers carrying babies of different sex.

## Conclusions

We found evidence that the genetic variants at 3q26.2, paternal age, maternal TL, and antenatal maternal health have a significant impact on newborn TL. The

comparative analysis of these factors identified the differences in the magnitude of their effects. Sex stratification analysis provided new insights into the factors explaining the male vs female TL variation. We also found that mother's TL was significantly associated with her own metabolic health and nutrient status, which may have transgenerational effects on offspring TL. Our findings provide a comprehensive understanding of the heritable and environmental factors and their relative contributions to the initial setting of TL and programming of longevity in early life.

#### Abbreviations

AA: Arachidonic acid; APGAR: A score is determined by evaluating the newborn baby on five simple criteria on a scale from 0 to 2; DGLA: Dihomo-gamma-linolenic acid; DHA: Docosahexaenoic acid; DOHaD: Developmental origins of health and disease; DPA: Docosapentaenoic acid; EPA: Eicosapentaenoic acid; EPDS: Edinburgh Postnatal Depression Scale; GA: Gestational age; GDM: Gestational diabetes mellitus; GUSTO: Growing Up in Singapore Towards healthy Outcomes; GWAS: Genome-wide association study; GWG: Gestational weight gain; IGF: Insulin-like growth factor; IGFBP: Insulin-like growth factor-binding protein; LA: Linoleic acid; LD: Linkage disequilibrium; MUFA: Mono-unsaturated fatty acid; PAI-1: Plasminogen activator inhibitor-1; ppBMI: Pre-pregnancy body mass index; PUFA: Poly-unsaturated fatty acid; SD: Standard deviation; SFA: Saturated fatty acid; SNP: Single-nucleotide polymorphism; STAI: State-Trait Anxiety Inventory; TL: Telomere length

#### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12916-021-02217-9>.

**Additional file 1: Figure S1.** DNA quality analysis by agarose gel electrophoresis. **Figure S2.** Flowchart of sample selection and analysis steps. **Figure S3.** Trans-ethnic genome-wide association studies on telomere length. **Figure S4.** Boxplots of the top six genetic variants. **Figure S5.** Effect allele frequencies of the top six genetic variants. **Figure S6.** Heat map of pairwise Pearson correlation coefficients between clinical variables. **Figure S7.** Significant sex-specific effects of the selected factors on newborn telomere length. **Figure S8.** Association between maternal telomere length and antenatal maternal factors. **Figure S9.** The variance percentage explained by each factor. **Figure S10.** Scatter plot of average relative telomere length of cord blood and cord tissue. **Figure S11.** Mediation analysis of maternal telomere length. **Table S1.** Intra-class correlation coefficient of intra-assay and inter-assay for telomere length measurements. **Table S2.** Comparison of the basic characteristics of 950 subjects and the full cohort. **Table S3.** Clinical characteristics of maternal-offspring subjects in this study and linear regression results for newborn TL. **Table S4.** The association of SNPs at 3q26.2 in the GWAS results of newborn and maternal telomere lengths and the meta-analysis results. **Table S5.** Pairwise Linkage Disequilibrium measures between the top six genetic variants. **Table S6.** Linear regression results between maternal telomere length and antenatal maternal factors. **Table S7.** The results of sensitivity analysis after adding DNA storage time and sample collection month in the best multivariate models of newborn telomere length. **Table S8.** The results of sensitivity analysis after further adjustment for DNA storage time and sample collection month in the association studies between maternal telomere length and antenatal maternal factors. **Table S9.** The genetic variants in a strong Linkage Disequilibrium with rs10936600. **Table S10.** The association of candidate genes in the GWAS results of newborn and maternal telomere lengths and the meta-analysis results.

#### Acknowledgements

We would like to thank the participants and the GUSTO study group which includes Allan Sheppard, Amutha Chinnadurai, Anne Eng Neo Goh, Anne

Rifkin-Graboi, Anqi Qiu, Arijit Biswas, Bee Wah Lee, Birit F.P. Broekman, Boon Long Quah, Borys Shuter, Chai Kiat Chng, Cheryl Ngo, Choon Looi Bong, Christiani Jeyakumar Henry, Cornelia Yin Ing Chee, Yam Thiam Daniel Goh, Doris Fok, Fabian Yap, George Seow Heong Yeo, Helen Chen, Hugo P S van Bever, Iliana Magiati, Inez Bik Yun Wong, Ivy Yee-Man Lau, Jeevesh Kapur, Jenny L. Richmond, Jerry Kok Yen Chan, Joanna D. Holbrook, Johan G. Eriksson, Joshua J. Gooley, Keith M. Godfrey, Kenneth Kwek, Kok Hian Tan, Krishnamoorthy Niduvaje, Leher Singh, Lin Lin Su, Lourdes Mary Daniel, Lynette P Shek, Marielle V. Fortier, Mark Hanson, Mary Foong-Fong Chong, Mary Rauff, Mei Chien Chua, Michael Meaney, Mya Thway Tint, Neeraja Karnani, Ngee Lek, Oon Hoe Teoh, P. C. Wong, Peter D. Gluckman, Pratibha Agarwal, Rob M. van Dam, Salome A. Rebello, Seang-Mei Saw, Shang Chee Chong, Shirong Cai, Shu-E Soh, Sok Bee Lim, Chin-Ying Stephen Hsu, Victor Samuel Rajadurai, Walter Stunkel, Wee Meng Han, Wei Wei Pang, Yap-Seng Chong, Yin Bun Cheung, Yiong Huak Chan, and Yung Seng Lee.

We thank the Blackburn Laboratory for the telomere length measurement and Dr. John Connolly and Dr. Gerard Wong for the protein biomarker measurement and preprocessing.

#### Authors' contributions

LC and NK conceived and supervised the study. JGE, YSC, NK, and PDG contributed to the data and sample collection in the GUSTO cohort. MJM, MFFC, and KHT contributed to the acquisition of the phenotypic data. KTM and MG contributed to the interpretation of the data. LC carried out the statistical analysis and association studies. LC and NK interpreted the data and wrote the manuscript. All authors critically read and approved the final manuscript content.

#### Funding

This research is supported by the Singapore National Research Foundation under its Translational and Clinical Research (TCR) Flagship Program and administered by the Singapore Ministry of Health's National Medical Research Council (NMRC), Singapore—NMRC/TCR/004-NUS/2008; NMRC/TCR/012-NUHS/2014. Additional funding is provided by the Singapore Institute for Clinical Sciences (SICS) – Agency for Science, Technology and Research (A\*STAR), Singapore.

#### Availability of data and materials

Data are not publicly available due to ethical restrictions but can be obtained from the authors upon reasonable request and subject to appropriate approvals, including from the GUSTO cohort's Executive Committee.

#### Declarations

##### Ethics approval and consent to participate

Written informed consent was obtained from all women who participated in the study. Approval for the study was granted by the ethics boards of both KK Women's and Children's Hospital and National University Hospital, which are the Centralised Institute Review Board (reference 2009/280/D) and the Domain Specific Review Board (reference D/09/021), respectively.

##### Consent for publication

Not applicable.

##### Competing interests

NK and YSC are part of an academic consortium that has received research funding from Abbott Nutrition, Nestec, EVOLVE Biosystems, DSM, and Danone. All other authors declare that they have no competing interests.

##### Author details

<sup>1</sup>Singapore Institute for Clinical Sciences, A\*STAR, Singapore, Singapore. <sup>2</sup>Saw Swee Hock School of Public Health, National University of Singapore (NUS), Singapore, Singapore. <sup>3</sup>KK Women's and Children's Hospital, Singapore, Singapore. <sup>4</sup>Department of Obstetrics and Gynaecology and Human Potential Translational Research Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore. <sup>5</sup>Sackler Program for Epigenetics & Psychobiology at McGill University, Montréal, Canada. <sup>6</sup>Ludmer Centre for Neuroinformatics and Mental Health, Douglas Mental Health University Institute, McGill University, Montréal, Canada. <sup>7</sup>Centre for Human Evolution, Adaptation and Disease, Liggins Institute,

University of Auckland, Auckland, New Zealand. <sup>8</sup>Folkhalsan Research Center, Helsinki, Finland. <sup>9</sup>Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland. <sup>10</sup>Bioinformatics Institute, A\*STAR, Singapore, Singapore. <sup>11</sup>Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore.

Received: 7 July 2021 Accepted: 14 December 2021

Published online: 25 January 2022

## References

- Blackburn EH. Structure and function of telomeres. *Nature*. 1991;350(6319):569–73. <https://doi.org/10.1038/350569a0>.
- Sanders JL, Newman AB. Telomere length in epidemiology: a biomarker of aging, age-related disease, both, or neither? *Epidemiol Rev*. 2013;35(1):112–31. <https://doi.org/10.1093/epirev/mxs008>.
- Wang Q, Zhan Y, Pedersen NL, Fang F, Hagg S. Telomere length and all-cause mortality: a meta-analysis. *Ageing Res Rev*. 2018;48:11–20. <https://doi.org/10.1016/j.arr.2018.09.002>.
- Gardner M, Bann D, Wiley L, Cooper R, Hardy R, Nitsch D, et al. Gender and telomere length: systematic review and meta-analysis. *Exp Gerontol*. 2014;51:15–27. <https://doi.org/10.1016/j.exger.2013.12.004>.
- Astuti Y, Wardhana A, Watkins J, Wulaningsih W, Network PR. Cigarette smoking and telomere length: a systematic review of 84 studies and meta-analysis. *Environ Res*. 2017;158:480–9. <https://doi.org/10.1016/j.envres.2017.06.038>.
- Wang J, Dong X, Cao L, Sun Y, Qiu Y, Zhang Y, et al. Association between telomere length and diabetes mellitus: a meta-analysis. *J Int Med Res*. 2016;44(6):1156–73. <https://doi.org/10.1177/0300060516667132>.
- Khalangot M, Krasnienkov D, Vaiserman A, Avilov I, Kovtun V, Okhrimenko N, et al. Leukocyte telomere length is inversely associated with post-load but not with fasting plasma glucose levels. *Exp Biol Med (Maywood)*. 2017;242(7):700–8. <https://doi.org/10.1177/1535370217694096>.
- Kaplan RC, Fitzpatrick AL, Pollak MN, Gardner JP, Jenny NS, McGinn AP, et al. Insulin-like growth factors and leukocyte telomere length: the cardiovascular health study. *J Gerontol A Biol Sci Med Sci*. 2009;64(11):1103–6. <https://doi.org/10.1093/gerona/glp036>.
- Broer L, Raschenberger J, Deelen J, Mangino M, Codd V, Pietilainen KH, et al. Association of adiponectin and leptin with relative telomere length in seven independent cohorts including 11,448 participants. *Eur J Epidemiol*. 2014;29(9):629–38. <https://doi.org/10.1007/s10654-014-9940-1>.
- Richards JB, Valdes AM, Gardner JP, Kato BS, Siva A, Kimura M, et al. Homocysteine levels and leukocyte telomere length. *Atherosclerosis*. 2008;200(2):271–7. <https://doi.org/10.1016/j.atherosclerosis.2007.12.035>.
- Tellechea ML, Pirola CJ. The impact of hypertension on leukocyte telomere length: a systematic review and meta-analysis of human studies. *J Hum Hypertens*. 2017;31(2):99–105. <https://doi.org/10.1038/jhh.2016.45>.
- Mwasongwe S, Gao Y, Griswold M, Wilson JG, Aviv A, Reiner AP, et al. Leukocyte telomere length and cardiovascular disease in African Americans: the Jackson Heart Study. *Atherosclerosis*. 2017;266:41–7. <https://doi.org/10.1016/j.atherosclerosis.2017.09.016>.
- Vakonaki E, Tsiminikaki K, Plaitis S, Fragkiadaki P, Tsoukalas D, Katsikantami I, et al. Common mental disorders and association with telomere length. *Biomed Rep*. 2018;8(2):111–6. <https://doi.org/10.3892/br.2018.1040>.
- Kang JX. Differential effects of omega-6 and omega-3 fatty acids on telomere length. *Am J Clin Nutr*. 2010;92(5):1276–7; author reply 1277. <https://doi.org/10.3945/ajcn.110.000463>.
- Pollack AZ, Rivers K, Ahrens KA. Parity associated with telomere length among US reproductive age women. *Hum Reprod*. 2018;33(4):736–44. <https://doi.org/10.1093/humrep/dey024>.
- Barha CK, Hanna CW, Salvante KG, Wilson SL, Robinson WP, Altman RM, et al. Number of children and telomere length in women: a prospective, longitudinal evaluation. *PLoS One*. 2016;11(1):e0146424. <https://doi.org/10.1371/journal.pone.0146424>.
- Stephoe A, Hamer M, Butcher L, Lin J, Brydon L, Kivimaki M, et al. Educational attainment but not measures of current socioeconomic circumstances are associated with leukocyte telomere length in healthy older men and women. *Brain Behav Immun*. 2011;25(7):1292–8. <https://doi.org/10.1016/j.bbi.2011.04.010>.
- Codd V, Mangino M, van der Harst P, Braund PS, Kaiser M, Beveridge AJ, et al. Common variants near TERC are associated with mean telomere length. *Nat Genet*. 2010;42(3):197–9. <https://doi.org/10.1038/ng.532>.
- Shen Q, Zhang Z, Yu L, Cao L, Zhou D, Kan M, et al. Common variants near TERC are associated with leukocyte telomere length in the Chinese Han population. *Eur J Hum Genet*. 2011;19(6):721–3. <https://doi.org/10.1038/ejhg.2011.4>.
- Pooley KA, Bojesen SE, Weischer M, Nielsen SF, Thompson D, Amin Al Olama A, et al. A genome-wide association scan (GWAS) for mean telomere length within the COGS project: identified loci show little association with hormone-related cancer risk. *Hum Mol Genet*. 2013;22(24):5056–64. <https://doi.org/10.1093/hmg/ddt355>.
- Codd V, Nelson CP, Albrecht E, Mangino M, Deelen J, Buxton JL, et al. Identification of seven loci affecting mean telomere length and their association with disease. *Nat Genet*. 2013;45(4):422–7. <https://doi.org/10.1038/ng.2528>.
- Prescott J, Kraft P, Chasman DI, Savage SA, Mirabello L, Berndt SI, et al. Genome-wide association study of relative telomere length. *PLoS One*. 2011;6(5):e19635. <https://doi.org/10.1371/journal.pone.0019635>.
- Liu Y, Cao L, Li Z, Zhou D, Liu W, Shen Q, et al. A genome-wide association study identifies a locus on TERT for mean telomere length in Han Chinese. *PLoS One*. 2014;9(1):e85043. <https://doi.org/10.1371/journal.pone.0085043>.
- Delgado DA, Zhang C, Chen LS, Gao J, Roy S, Shinkle J, et al. Genome-wide association study of telomere length among South Asians identifies a second RTEL1 association signal. *J Med Genet*. 2018;55(1):64–71. <https://doi.org/10.1136/jmedgenet-2017-104922>.
- Factor-Litvak P, Susser E, Kezios K, McKeague I, Kark JD, Hoffman M, et al. Leukocyte telomere length in newborns: implications for the role of telomeres in human disease. *Pediatrics*. 2016;137(4). <https://doi.org/10.1542/peds.2015-3927>.
- Martens DS, Van Der Stukken C, Derom C, Thiery E, Bijmens EM, Nawrot TS. Newborn telomere length predicts later life telomere length: tracking telomere length from birth to child- and adulthood. *EBioMedicine*. 2021;63:103164. <https://doi.org/10.1016/j.ebiom.2020.103164>.
- Benetos A, Verhulst S, Labat C, Lai TP, Girerd N, Toupance S, et al. Telomere length tracking in children and their parents: implications for adult onset diseases. *FASEB J*. 2019;33(12):14248–53. <https://doi.org/10.1096/fj.201901275R>.
- Entringer S, de Punder K, Buss C, Wadhwa PD. The fetal programming of telomere biology hypothesis: an update. *Philos Trans R Soc Lond B Biol Sci*. 2018;373(1741). <https://doi.org/10.1098/rstb.2017.0151>.
- Entringer S, Epel ES, Lin J, Blackburn EH, Buss C, Shahbaba B, et al. Maternal folate concentration in early pregnancy and newborn telomere length. *Ann Nutr Metab*. 2015;66(4):202–8. <https://doi.org/10.1159/000381925>.
- Kim JH, Kim GJ, Lee D, Ko JH, Lim I, Bang H, et al. Higher maternal vitamin D concentrations are associated with longer leukocyte telomeres in newborns. *Matern Child Nutr*. 2018;14(1). <https://doi.org/10.1111/mcn.12475>.
- Mirzakhani H, De Vivo I, Leeder JS, Gaedigk R, Vyhldal CA, Weiss ST, et al. Early pregnancy intrauterine fetal exposure to maternal smoking and impact on fetal telomere length. *Eur J Obstet Gynecol Reprod Biol*. 2017;218:27–32. <https://doi.org/10.1016/j.ejogrb.2017.09.013>.
- Wojcicki JM, Olveda R, Heyman MB, Elwan D, Lin J, Blackburn E, et al. Cord blood telomere length in Latino infants: relation with maternal education and infant sex. *J Perinatol*. 2016;36(3):235–41. <https://doi.org/10.1038/jp.2015.178>.
- Tellechea M, Gianotti TF, Alvarinas J, Gonzalez CD, Sookoian S, Pirola CJ. Telomere length in the two extremes of abnormal fetal growth and the programming effect of maternal arterial hypertension. *Sci Rep*. 2015;5(1):7869. <https://doi.org/10.1038/srep07869>.
- Martens DS, Plusquin M, Gyselaers W, De Vivo I, Nawrot TS. Maternal pre-pregnancy body mass index and newborn telomere length. *BMC Med*. 2016;14(1):148. <https://doi.org/10.1186/s12916-016-0689-0>.
- Xu J, Ye J, Wu Y, Zhang H, Luo Q, Han C, et al. Reduced fetal telomere length in gestational diabetes. *PLoS One*. 2014;9(1):e86161. <https://doi.org/10.1371/journal.pone.0086161>.
- Send TS, Gilles M, Codd V, Wolf I, Bardtke S, Streit F, et al. Telomere length in newborns is related to maternal stress during pregnancy. *Neuropsychopharmacology*. 2017;42(12):2407–13. <https://doi.org/10.1038/npp.2017.73>.
- Prescott J, Du M, Wong JY, Han J, De Vivo I. Paternal age at birth is associated with offspring leukocyte telomere length in the nurses' health study. *Hum Reprod*. 2012;27(12):3622–31. <https://doi.org/10.1093/humrep/des314>.



38. Broer L, Codd V, Nyholt DR, Deelen J, Mangino M, Willemsen G, et al. Meta-analysis of telomere length in 19,713 subjects reveals high heritability, stronger maternal inheritance and a paternal age effect. *Eur J Hum Genet.* 2013;21(10):1163–8. <https://doi.org/10.1038/ejhg.2012.303>.
39. Kimura M, Cherkas LF, Kato BS, Demissie S, Hjelmborg JB, Brimacombe M, et al. Offspring's leukocyte telomere length, paternal age, and telomere elongation in sperm. *PLoS Genet.* 2008;4(2):e37. <https://doi.org/10.1371/journal.pgen.0040037>.
40. Nordfjall K, Larefalk A, Lindgren P, Holmberg D, Roos G. Telomere length and heredity: indications of paternal inheritance. *Proc Natl Acad Sci U S A.* 2005;102(45):16374–8. <https://doi.org/10.1073/pnas.0501724102>.
41. Eisenberg DT. Inconsistent inheritance of telomere length (TL): is offspring TL more strongly correlated with maternal or paternal TL? *Eur J Hum Genet.* 2014;22(1):8–9. <https://doi.org/10.1038/ejhg.2013.202>.
42. Soh SE, Tint MT, Gluckman PD, Godfrey KM, Rifkin-Graboi A, Chan YH, et al. Cohort profile: Growing Up in Singapore Towards healthy Outcomes (GUSTO) birth cohort study. *Int J Epidemiol.* 2013.
43. Lin X, Teh AL, Chen L, Lim IY, Tan PF, Maclsaac JL, et al. Choice of surrogate tissue influences neonatal EWAS findings. *BMC Med.* 2017;15(1):211. <https://doi.org/10.1186/s12916-017-0970-x>.
44. Lin J, Epel E, Cheon J, Kroenke C, Sinclair E, Bigos M, et al. Analyses and comparisons of telomerase activity and telomere length in human T and B cells: insights for epidemiology of telomere maintenance. *J Immunol Methods.* 2010;352(1-2):71–80. <https://doi.org/10.1016/j.jim.2009.09.012>.
45. Verhulst S, Susser E, Factor-Litvak PR, Simons M, Benetos A, Steenstrup T, et al. Response to: Reliability and validity of telomere length measurements. *Int J Epidemiol.* 2016;45(4):1298–301. <https://doi.org/10.1093/ije/dyw194>.
46. Chong YS, Cai S, Lin H, Soh SE, Lee YS, Leow MK, et al. Ethnic differences translate to inadequacy of high-risk screening for gestational diabetes mellitus in an Asian population: a cohort study. *BMC Pregnancy Childbirth.* 2014;14(1):345. <https://doi.org/10.1186/1471-2393-14-345>.
47. Wendland EM, Torloni MR, Falavigna M, Trujillo J, Dode MA, Campos MA, et al. Gestational diabetes and pregnancy outcomes—a systematic review of the World Health Organization (WHO) and the International Association of Diabetes in Pregnancy Study Groups (IADPSG) diagnostic criteria. *BMC Pregnancy Childbirth.* 2012;12(1):23. <https://doi.org/10.1186/1471-2393-12-23>.
48. van Lee L, Tint MT, Aris IM, Quah PL, Fortier MV, Lee YS, et al. Prospective associations of maternal betaine status with offspring weight and body composition at birth: the Growing Up in Singapore Towards healthy Outcomes (GUSTO) cohort study. *Am J Clin Nutr.* 2016;104(5):1327–33. <https://doi.org/10.3945/ajcn.116.138818>.
49. Ong YL, Quah PL, Tint MT, Aris IM, Chen LW, van Dam RM, et al. The association of maternal vitamin D status with infant birth outcomes, postnatal growth and adiposity in the first 2 years of life in a multi-ethnic Asian population: the Growing Up in Singapore Towards healthy Outcomes (GUSTO) cohort study. *Br J Nutr.* 2016;116(4):621–31. <https://doi.org/10.1017/S0007114516000623>.
50. Chong MF, Wong JX, Colega M, Chen LW, van Dam RM, Tan CS, et al. Relationships of maternal folate and vitamin B12 status during pregnancy with perinatal depression: the GUSTO study. *J Psychiatr Res.* 2014;55:110–6. <https://doi.org/10.1016/j.jpsychires.2014.04.006>.
51. Chong MF, Ong YL, Calder PC, Colega M, Wong JX, Tan CS, et al. Long-chain polyunsaturated fatty acid status during pregnancy and maternal mental health in pregnancy and the postpartum period: results from the GUSTO study. *J Clin Psychiatry.* 2015;76(7):e848–56. <https://doi.org/10.4088/JCP.14m09191>.
52. Bernard JY, Tint MT, Aris IM, Chen LW, Quah PL, Tan KH, et al. Maternal plasma phosphatidylcholine polyunsaturated fatty acids during pregnancy and offspring growth and adiposity. *Prostaglandins Leukot Essent Fatty Acids.* 2017;121:21–9. <https://doi.org/10.1016/j.plefa.2017.05.006>.
53. Reese SE, Zhao S, Wu MC, Joubert BR, Parr CL, Haberg SE, et al. DNA methylation score as a biomarker in newborns for sustained maternal smoking during pregnancy. *Environ Health Perspect.* 2017;125(4):760–6. <https://doi.org/10.1289/EHP333>.
54. Rosero-Bixby L, Rehkopf DH, Dow WH, Lin J, Epel ES, Azofeifa J, et al. Correlates of longitudinal leukocyte telomere length in the Costa Rican Longevity Study of Healthy Aging (CRELES): on the importance of DNA collection and storage procedures. *PLoS One.* 2019;14(10):e0223766. <https://doi.org/10.1371/journal.pone.0223766>.
55. Lai JS, Pang WW, Cai S, Lee YS, Chan JKY, Shek LPC, et al. High folate and low vitamin B12 status during pregnancy is associated with gestational diabetes mellitus. *Clin Nutr.* 2018;37(3):940–7. <https://doi.org/10.1016/j.clnu.2017.03.022>.
56. Jones AM, Beggs AD, Carvajal-Carmona L, Farrington S, Tenesa A, Walker M, et al. TERC polymorphisms are associated both with susceptibility to colorectal cancer and with longer telomeres. *Gut.* 2012;61(2):248–54. <https://doi.org/10.1136/gut.2011.239772>.
57. Pepper MR, Black MM. B12 in fetal development. *Semin Cell Dev Biol.* 2011;22(6):619–23. <https://doi.org/10.1016/j.semcdb.2011.05.005>.
58. Martens DS, Janssen BG, Bijns EM, Clemente DBP, Vineis P, Plusquin M, et al. Association of parental socioeconomic status and newborn telomere length. *JAMA Netw Open.* 2020;3(5):e204057. <https://doi.org/10.1001/jamanetworkopen.2020.4057>.
59. Liu B, Chen H, Xu Y, An C, Zhong L, Wang X, et al. Fetal growth is associated with maternal fasting plasma glucose at first prenatal visit. *PLoS One.* 2014;9(12):e116352. <https://doi.org/10.1371/journal.pone.0116352>.
60. Hong S, Kim MM. IGFBP-3 plays an important role in senescence as an aging marker. *Environ Toxicol Pharmacol.* 2018;59:138–45. <https://doi.org/10.1016/j.etap.2018.03.014>.
61. Jung RG, Motazedian P, Ramirez FD, Simard T, Di Santo P, Visintini S, et al. Association between plasminogen activator inhibitor-1 and cardiovascular events: a systematic review and meta-analysis. *Thromb J.* 2018;16(1):12. <https://doi.org/10.1186/s12959-018-0166-4>.
62. Barbieri M, Paolisso G, Kimura M, Gardner JP, Boccardi V, Papa M, et al. Higher circulating levels of IGF-1 are associated with longer leukocyte telomere length in healthy subjects. *Mech Ageing Dev.* 2009;130(11-12):771–6. <https://doi.org/10.1016/j.mad.2009.10.002>.
63. Kuhlow D, Florian S, von Figura G, Weimer S, Schulz N, Petzke KJ, et al. Telomerase deficiency impairs glucose metabolism and insulin secretion. *Aging (Albany NY).* 2010;2(10):650–8. <https://doi.org/10.18632/aging.100200>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

