

A Genome-Wide Screen in Yeast Identifies Specific Oxidative Stress Genes Required for the Maintenance of Sub-Cellular Redox Homeostasis

Anita Ayer¹, Sina Fellermeier¹, Christopher Fife¹, Simone S. Li¹, Gertien Smits², Andreas J. Meyer³, Ian W. Dawes^{1*}, Gabriel G. Perrone⁴

1 University of New South Wales, Sydney, Australia, **2** Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands, **3** INRES- Chemical Signalling, University of Bonn, Bonn, Germany, **4** University of Western Sydney, Penrith, Australia

Abstract

Maintenance of an optimal redox environment is critical for appropriate functioning of cellular processes and cell survival. Despite the importance of maintaining redox homeostasis, it is not clear how the optimal redox potential is sensed and set, and the processes that impact redox on a cellular/organellar level are poorly understood. The genetic bases of cellular redox homeostasis were investigated using a green fluorescent protein (GFP) based redox probe, roGFP2 and a pH sensitive GFP-based probe, pHluorin. The use of roGFP2, in conjunction with pHluorin, enabled determination of pH-adjusted sub-cellular redox potential in a non-invasive and real-time manner. A genome-wide screen using both the non-essential and essential gene collections was carried out in *Saccharomyces cerevisiae* using cytosolic-roGFP2 to identify factors essential for maintenance of cytosolic redox state under steady-state conditions. 102 genes of diverse function were identified that are required for maintenance of cytosolic redox state. Mutations in these genes led to shifts in the half-cell glutathione redox potential by 75–10 mV. Interestingly, some specific oxidative stress-response processes were identified as over-represented in the data set. Further investigation of the role of oxidative stress-responsive systems in sub-cellular redox homeostasis was conducted using roGFP2 constructs targeted to the mitochondrial matrix and peroxisome and E_{GSH} was measured in cells in exponential and stationary phase. Analyses allowed for the identification of key redox systems on a sub-cellular level and the identification of novel genes involved in the regulation of cellular redox homeostasis.

Citation: Ayer A, Fellermeier S, Fife C, Li SS, Smits G, et al. (2012) A Genome-Wide Screen in Yeast Identifies Specific Oxidative Stress Genes Required for the Maintenance of Sub-Cellular Redox Homeostasis. PLoS ONE 7(9): e44278. doi:10.1371/journal.pone.0044278

Editor: Mick F. Tuite, University of Kent, United Kingdom

Received: May 31, 2012; **Accepted:** July 31, 2012; **Published:** September 6, 2012

Copyright: © 2012 Ayer et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was funded by Australian Postgraduate Award (to AA) and a Discovery grant from the Australian Research Council (to IWD and GGP). Work in GJS's lab was partially funded by Unilever and work in AJM's laboratory was funded by the Deutsche Forschungsgemeinschaft (grant ME1567/5-1). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have read the journal's policy and have the following interests: One of the authors (GJS) has received funding from Unilever. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLoS ONE policies on sharing data and materials.

* E-mail: i.dawes@unsw.edu.au

Introduction

The maintenance of an optimal redox environment is critical to the functioning of many biological processes. The majority of intracellular functions require a strongly reducing environment, which maintains key sulfhydryl groups in a reduced and active form and other key elements of proteins and enzymes to remain functional. In eukaryotic cells, highly reducing environments are found in compartments including the cytosol, mitochondrial matrix and peroxisome. Conversely, optimal oxidative protein folding and maturation requires disulfide bond formation and a more oxidative redox environment, found in the endoplasmic reticulum [1,2] and mitochondrial inter-membrane space [3–5]. Maintenance of appropriate cellular and sub-cellular redox environments is crucial since alterations in redox can adversely affect biological processes such as signal transduction [6], protein structure and function, DNA and RNA synthesis [7], enzyme activation and regulation of the cell cycle [8,9]. Moreover, altered cellular redox homeostasis has been associated with neurodegen-

erative disorders including Alzheimer's and Parkinson's diseases [10] and metabolic syndromes such as diabetes [11].

Despite the importance of redox homeostasis for fundamental cell biology and many disease states, our current understanding has been limited by available assay methods relying on whole-cell lysate analysis and the use of altered resistance to oxidants as a phenotypic indicator of redox imbalance or dysfunction. While the thioredoxin (TrxSS/Trx(SH)₂) system, glutathione system (GSSG/2GSH), the NADP⁺/NADPH couple, protein thiols and antioxidant defence systems have all been observed to impact on redox homeostasis on a cell-wide level [9,12,13], the exact genetic mechanisms in place for the maintenance of redox homeostasis are still unknown. Studying how cells regulate redox state on a sub-cellular level could lead to a better understanding of how aberrant redox biology may impact on cellular health in a variety of settings.

In order to gain a better insight into the genetic organization underpinning cellular redox homeostasis, a genome-wide screen of cytosolic redox state was carried out using the *Saccharomyces cerevisiae* tetO₇ promoter replacement collection (essential genes)

[14] and the homozygous diploid non-essential gene deletion collection [15]. Due to its relatively high concentration (in the millimolar range) and low half-cell reduction potential at pH 7 (-240 mV), the glutathione couple is considered the primary cellular redox buffer and its redox state (E_{GSH}) is used as an indicator of the state of both cellular and/or sub-cellular redox environment. Cytosolic redox potential (E_{GSH}) was systematically analyzed to identify genes affecting cytosolic redox homeostasis. Redox measurements were determined using the roGFP2 probe, which predominantly equilibrates with the GSSG/2GSH redox couple [16,17] and acts as an *in vivo* sensor of this couple. 102 genes were identified as essential for maintenance of cytosolic redox state, the majority of which were previously not known to be important for redox homeostasis. Specific oxidative stress genes were over-represented in the genes identified and by taking advantage of the ability of roGFP2 to be targeted to individual sub-cellular compartments, a targeted screen of 42 mutants affected in oxidative stress pathways in the cytosol, mitochondrial matrix and peroxisome was conducted to better understand the systems required to maintain redox homeostasis on a sub-cellular level.

Results

Verification of Redox-responsiveness of roGFP2-based Probes

Probes used to measure the redox state of the GSSG/2GSH couple based on the roGFP2 probe that were targeted to the cytosol and mitochondrial matrix were generated and verified as described in [18] and localization of peroxisome-targeted roGFP2 was analyzed by confocal microscopy in this study (Figure S1). In yeast cells expressing roGFP2-SKL a punctate fluorescent pattern consistent with peroxisomal localization was observed. Additionally, we also expressed roGFP2-SKL in *pex5* cells, which are defective in the import of peroxisomal proteins with a peroxisomal targeting signal 1 (PTS1) such as -SKL. In *pex5* cells, a cytosolic fluorescent pattern was observed indicative that roGFP2-SKL construct was correctly targeted and is imported via the PTS1-mediated import pathway into the peroxisome.

The roGFP2 probes response to redox state has been calibrated *in vitro* (18). Additionally, the responsiveness to *in vivo* redox state changes. To verify that roGFP2 in organelle-targeted plasmids was redox responsive *in vivo*, wild-type cells expressing organelle-targeted roGFP2 were grown to exponential phase in SC_{URA} and treated with various concentrations of dithiothreitol, a strong thiol-containing reductant or diamide a known GSH oxidizing agent. Wild-type cells treated with various concentrations of DTT, resulted in only a small change in E_{GSH} indicating that in the cytosol, peroxisome and mitochondrial matrix the roGFP2 probe was almost entirely in a fully reduced state (Figure S2A). Conversely, diamide-treated cells displayed progressively more oxidized E_{GSH} values that reached a plateau when the reporter was at its oxidized limit (Figure S2A). This verified that the organelle-targeted roGFP2 probes were able to respond dynamically from its most reduced to most oxidized state to intracellular redox state changes. To verify that changes observed via flow cytometry were due to changes in the redox-state of the roGFP2 protein *in vivo*, redox western blots were carried out to determine the redox status of the roGFP2 probe in untreated cells and cells treated with diamide (10 mM; 20 min) or DTT (10 mM; 20 min). Proteins were harvested by trichloroacetic acid precipitation and protein concentration determined by a detergent-compatible Bradford protein assay (Figure S2B). The western blots indicate that in the cytosol, mitochondrial matrix and peroxisome the

roGFP2 probe was almost completely reduced in untreated conditions. The roGFP2 protein extracted from untreated cells had mobility identical to the protein extracted from cells treated with 10 mM DTT. Treatment with diamide resulted in a shift in the protein migration corresponding to that of the oxidized roGFP2 protein. Conversely, wild-type GFP (non-redox sensitive GFP) did not respond to treatment with either diamide or DTT. Together these data indicate that the organelle-targeted roGFP2 probes can sense changes in the intracellular redox environment and are responsive to changes to redox changes.

High Throughput Screen of the Yeast Essential and Non-essential Deletion Libraries Identifies 102 Genes Required for Cytosolic Redox State Maintenance under Steady-state Conditions

A genome-wide screen was carried out to identify mutants affected in cytosolic redox environment using the redox-sensitive green fluorescent protein (roGFP2). The collection of 4,800 homozygous diploid mutants, each deleted for one non-essential gene [15] was transformed with the roGFP2 construct. Cells were grown to exponential phase at 25°C in SC_{URA} in 96-well plate format and the ratio of the intensity of emission after excitation at 405 nm and 488 nm ($R_{405/488}$) was determined for each mutant. The primary round of screening was conducted using a 96-well plate method and a fluorescent spectrophotometer to estimate the emission ratio. All potential mutants showing a shift in $R_{405/488}$ and any decrease in emission after excitation at 488 nm and/or any increase in emission after excitation at 405 nm were subsequently rescreened using flow cytometry, which was found to be more sensitive. An increase in the $R_{405/488}$ indicated a shift in the cytosolic redox state towards a more oxidized environment. Mutants which displayed statistically different $R_{405/488}$ values compared to wild type using a p-value of 0.001 were then identified. To rule out false positives, mutants were treated with 10 mM dithiothreitol (DTT) as a strong reductant. If the change in $R_{405/488}$ was not redox dependent, for example due to the accumulation of autofluorescent metabolites or cell-wall components, treatment with DTT would not significantly alter the $R_{405/488}$ in a pronounced way. However, if the change in $R_{405/488}$ was redox-dependent, DTT treatment would restore the $R_{405/488}$ to that of the wild type. After this stringent testing process, 89 mutants with a significantly higher $R_{405/488}$ compared to wild type were identified. To allow for pH-corrected E_{GSH} values, the cytosolic pH of the wild type was determined to be pH 7.5 using the cytosolic pHluorin probe. This value was used for the determination of cytosolic redox potential (E_{GSH}) and the cytosolic pH of identified mutants was verified to be similar to the wild type. Table 1 lists the top 15 mutants showing a significantly more oxidized cytosol than the wild type with the full list of mutants identified given in Dataset S1.

The screen of essential gene mutations was carried out using the tetO₇ promoter replacement (tetO₇) collection [14], which contains ~ 800 mutants, each mutant has a Kan^R-tetO₇-TATA_{CYC1} cassette integrated into the promoter of an essential gene and the tetracycline transactivator (tTA) integrated at the *URA3* locus to form a “tet-off” system. Addition of doxycycline to the medium represses expression of genes under the control of the tetO₇-CYC1 promoter. Mutants of the tetO₇ collection were separately transformed with the cytosolic roGFP2 construct (leucine-based selection) and grown at 25°C in SC_{LEU} in 96-well plate format in the presence of doxycycline. The ratio of the intensity of emission after excitation at 405 nm and 488 nm ($R_{405/488}$) was determined for each mutant grown after 24 hours and

Table 1. List of non-essential *S. cerevisiae* genes that when deleted significantly affect cytosolic redox state.

Mutant	Function	$E_{GSH} \pm SD$ (-mV)
Wild type	-	349 \pm 3
<i>glr1</i>	Cytosolic/mitochondrial glutathione reductase	275 \pm 4
<i>yap1</i>	Oxidative stress responsive transcription factor	319 \pm 4
<i>skn7</i>	Oxidative stress responsive transcription factor	320 \pm 5
<i>dog2</i>	2-deoxyglucose-6-phosphate phosphatase	325 \pm 4
<i>dig1</i>	MAP kinase-responsive inhibitor of the Ste12p	327 \pm 6
<i>dgr2</i>	Protein of unknown function	330 \pm 4
<i>dit1</i>	Sporulation-specific enzyme	330 \pm 3
<i>tkl1</i>	Transketolase	330 \pm 4
<i>ipk1</i>	Inositol 1,3,4,5,6-pentakisphosphate 2-kinase	332 \pm 3
<i>ysp1</i>	Mitochondrial protein	332 \pm 4
<i>nvj1</i>	Nuclear envelope protein	332 \pm 3
<i>rpl26b</i>	Component of the large (60S) ribosomal subunit	332 \pm 6
<i>inp2</i>	Peroxisome-specific receptor	332 \pm 6
<i>yju3</i>	Monoglyceride lipase	332 \pm 5
<i>ubi4</i>	Ubiquitin	332 \pm 4

Mutants are listed from most to least oxidized.
doi:10.1371/journal.pone.0044278.t001

$R_{405/488}$ calculated. As with the non-essential homozygous diploid collection, mutants from the tetO₇ collection having an $R_{405/488}$ significantly greater than the wild type were treated with DTT to identify redox-dependent changes and cytosolic pH was measured. 13 genes were identified with a significantly higher $R_{405/488}$ compared to the wild type and the cytosolic redox potential (E_{GSH}) was calculated (Table 2).

Functional Analysis of Genes Identified

The mutants identified in this screen were analyzed to identify significantly enriched ($p < 0.01$) cellular processes and functions using the Gene Ontology and Munich Information Center for Protein Sequences (MIPS) databases via FunSpec [19] as described in the Materials and Methods. Importantly, FunSpec analysis showed no enrichment for slow growing mutants, indicating that altered growth alone does not affect cytosolic redox state. As seen in Figure 1, the significantly enriched terms generated from these databases represent a range of cellular processes and functions with the two databases categorizing the data in appreciably different ways. From this analysis, genes involved in endoplasmic reticulum protein transport and translocation including components of the signal recognition particle (SRP) complex were enriched in the data set. While we could not study the redox state of the endoplasmic reticulum since it is too oxidizing to be measured by roGFP2, it is evident that defects in endoplasmic reticulum protein translocation affect cytosolic redox state. Mutations in SRP and downstream effects could be through the increased cytosolic accumulation of secretory protein precursor and chaperones or through the accumulation of misfolded proteins in the endoplasmic reticulum (as suggested by [20]) leading to oxidative stress through rounds of operation of the protein folding machinery.

Many of the genes identified possibly represent new functions important in redox homeostasis. *DOG2*, which encodes 2-deoxyglucose-6-phosphatase, is induced by oxidative and osmotic stress [20] and the results point to it having a significant role in regulating redox homeostasis, more so than its current functional description indicates. Additionally, the identification of genes that have near identical duplicates (for example *HTA1* and *RPL26*) or part of a greater protein complex (*LEA1*; required for U2 snRNP) suggest that there are a sub-set of genes that moonlight as redox sensors. There are two genes encoding *RPL26* in *S. cerevisiae*, *RPL26A* and *RPL26B* which are nearly identical to *HTA2*, yet only mutation to *RPL26B* and *HTA1* caused changes in cytosolic redox state. The fact that only one mutation has such a profound effect on redox homeostasis indicates that genes such as *RPL26B* and *HTA1* may have a specific function in redox homeostasis aside

Table 2. Mutants from the tetO₇ promoter replacement collection that exhibited a significantly more oxidized cytosol than wild type after treatment with doxycycline.

Gene	Function	$E_{GSH} \pm SD$ (-mV)
Wild type		322
tet-O ₇ - <i>STT4</i>	Phosphatidylinositol-4-kinase	272 \pm 6
tet-O ₇ - <i>SRP14</i>	Signal recognition particle complex subunit	282 \pm 16
tet-O ₇ - <i>yor146w</i>	Dubious opening reading frame	284 \pm 6
tet-O ₇ - <i>CDC34</i>	Ubiquitin-conjugating enzyme	291 \pm 7
tet-O ₇ - <i>YPT1</i>	Rab family GTPase	298 \pm 14
tet-O ₇ - <i>SPC98</i>	Component of the microtubule-nucleating Tub4p complex	298 \pm 3
tet-O ₇ - <i>SAH1</i>	S-adenosyl-L-homocysteine hydrolase	298 \pm 6
tet-O ₇ - <i>SEC53</i>	Phosphomannomutase	299 \pm 5
tet-O ₇ - <i>RIB5</i>	Riboflavin synthase	299 \pm 6
tet-O ₇ - <i>TRR1</i>	Cytoplasmic thioredoxin reductase	300 \pm 4
tet-O ₇ - <i>GPI16</i>	Transmembrane protein subunit of the glycosylphosphatidylinositol transamidase complex	307 \pm 6
tet-O ₇ - <i>NUP159</i>	Nucleoporin subunit of the nuclear pore complex	310 \pm 10
tet-O ₇ - <i>NAB3</i>	Single-stranded RNA binding protein	310 \pm 3

doi:10.1371/journal.pone.0044278.t002

A

GO Database		
Molecular Function	Biological Process	Cellular Component
*Transcriptional regulatory region RNA binding ($p=0.0003$) *Flavin adenine nucleotide binding ($p=0.0005$) *Signal sequence binding ($p=0.004$) *Manganese ion binding ($p=0.007$) *7S RNA binding ($p=0.007$)	*Regulation of transcription from RNA polymerase II promoter in response to oxidative stress ($p=0.001$) *Response to singlet oxygen ($p=0.001$) *Protein targeting to ER ($p=0.003$) *Termination of RNA polymerase II transcription (exosome dependent) ($p=0.004$) *SRP dependent cotranslational protein targeting to membrane, signal sequence recognition ($p=0.005$)	*Ndr1 complex ($p=0.0008$) *Signal recognition particle ($p=0.002$)

B

MIPS Database	
Functional Classification	Phenotypes
*Protein targeting, sorting and translocation ($p=0.0007$) *Oxidative stress response ($p=0.002$) *Protein Transport ($p=0.008$)	*Other or general oxidising agent sensitivity ($p=0.001$) *Hydrogen peroxide sensitivity ($p=0.003$) *Secretory mutants ($p=0.003$) *Sucrose fermentation ($p=0.003$)

Figure 1. Characterization of mutants displaying a significantly more oxidized cytosol relative to wild type. All mutants that exhibited a significantly more oxidized cytosol compared to wild type were analyzed for enriched terms using the Gene Ontology (GO) and Munich Information Center for Protein Sequences (MIPS) databases via FunSpec [18]. p -values (Bonferroni corrected) represent the probability that the intersection of a given list within any functional category occurs by chance. The terms identified that are related to oxidative stress are highlighted in bold. doi:10.1371/journal.pone.0044278.g001

from their main function. Alternatively, the gene identified may be expressed more highly than its homologue leading to its deletion having a greater effect on redox homeostasis. In the same vein, the identification of only one gene in a protein complex such as *LEA1* (component of the U2 snRNP complex) or *PXA1* (subunit of the peroxisomal ATP-binding cassette transporter complex; Pxa1p-Pxa2p), may indicate a specialized role for these genes in redox homeostasis.

While functional categorization of gene lists using GO terms is especially useful when working with large amounts of data, often such terms are not sufficiently fine or descriptive to identify functional groups or allocate genes whose products have pleiotropic effects or are less studied, into established groups. Closer inspection of the data revealed that processes affecting cytosolic redox homeostasis such as mitochondrial function (*YPS1*, *NDE1*, *MRPL25*, *MRPS16*, *MRPL35*, *MRPL24*, *ICP55*, *MNE1*, *MSF1*, *MRPL40*, *COQ10*, *MSW1*, *TIM8*, *AIM38*) and the ubiquitin system (*UBI4*, *UIP4*, *UBP13*, *CDC34*) were not detected by either the GO or MIPS databases. These represent new pathways somehow involved in redox homeostasis. Respiratory-incompetent cells have increased levels of GSSG in stationary phase relative to exponential phase and mitochondrial function is required for resistance to oxidative stress [21]. Ccp1p is also involved in oxidative stress signaling leading to activation of the Skn7p transcription factor [22]. This could explain the larger shift in

matrix redox state observed in the *ccp1* mutant compared to the *skn7* mutant. Deletion of *CCP1* may cause increased levels of hydrogen peroxide to accumulate and decreased activation of Skn7p doubly affecting matrix redox state. Taken together, it is apparent that there is a connection between mitochondrial function and redox homeostasis. The link between mitochondrial function and redox homeostasis may be due to several reasons. Many mitochondrial processes are integral to the generation of redox-active species such as NADH and NADPH, they involve redox-dependent electron transfer reactions and produce reactive oxygen species. Consequently, the redox-association of many mitochondrial functions may indicate they have a potential role in the setting, sensing or maintenance of the cellular redox environment.

Four mutants (*ubi4*, *uip4*, *ubp13* and *cdc34-tetO7*) involved in ubiquitin-dependent protein degradation were identified as having a more oxidized cytosol than the wild type. Moreover, decreased expression of *CDC34*, an essential gene which encodes a ubiquitin-conjugating enzyme [23], led to a more oxidized cytosol. Mutants of the ubiquitin system were previously identified in a screen for mutants that over-excrete glutathione [24] and *CDC34* was identified as a high-copy suppressor of the poor growth phenotype of *gsh2* cells, which lack the second step of glutathione synthesis [25]). *CDC34* was shown to activate *GSH1* (encoding γ -glutamylcysteine synthetase) expression by modulating the activity

of Met4p, the transcriptional activator of the sulphur assimilation pathway [25], providing a link between the GSH and ubiquitin systems. Given that overexpression of *CDC34* increases GSH levels and repression of *CDC34* leads to a more oxidized cytosolic redox state, tight control of Cdc34p is required for optimal intracellular glutathione concentrations, and alterations in the activity of Cdc43p would affect glutathione homeostasis. Disruption to the ubiquitin system and to protein degradation processes on a broad level not only via Cdc34p, may affect *GSH1* gene expression and regulation of the glutathione couple and redox state.

Specific Oxidative Stress Responsive Genes are Essential for Cytosolic Redox State Homeostasis

Interestingly, both databases highlighted the enrichment of genes involved in oxidative stress and response to reactive oxygen species as shown in bold in Figure 1. Some oxidative-stress responsive genes identified showed the most extensive shift in E_{GSH} compared to the wild type (ΔE_{GSH}), with the *glr1*, *yap1*, *skn7* and *tkl1* mutants being in the top 10 mutants with the largest ΔE_{GSH} values. Genes involved in the detoxification of reactive oxygen species have conventionally been regarded as response systems rather than maintenance systems. However the results of the screen indicate that some key oxidative stress responsive systems serve a dual role. This challenges the paradigm that maintenance and response systems are distinct elements in redox regulation and supports the view that some of the systems that are needed in the response to counteract deleterious oxidative changes are required to maintain redox homeostasis under steady state conditions. Clearly, some genes identified in this screen will indirectly affect the redox environment rather than being direct modulators of cytosolic redox state. For instance it is possible that mutants may have increased levels of oxidative stress, which could affect the roGFP2 readout rather than directly affecting the glutathione couple. However, delineating between primary and secondary redox effects is complex and not the focus of this present study. Additionally, it is interesting that ~10% of the genes identified in this screen have no known or putative function. These now can be seen to have some involvement, directly or indirectly in modulating cytosolic redox state.

Targeted Cross-compartmental Screen to Understand the Role of Oxidative-stress Systems in Redox Homeostasis

From the mutants identified in the screen, only five (*glr1*, *yap1*, *skn7*, *tkl1* and *trr1*; [12,26]) have been previously reported to be involved in redox homeostasis. Some genes previously identified as affecting cellular redox state were not identified through this screen possibly due to the differences in methods used to estimate E_{GSH} and redundancy of some known antioxidant functions. Given that genes involved in antioxidant function and resistance to oxidative stress were identified as important to cytosolic redox maintenance through the genome-wide screen, we then looked more closely at known pathways involved in antioxidant function and oxidative stress responses to understand better their role in setting, sensing and maintenance of redox state. Moreover, these analyses were carried out on a compartmental level (in the cytosol, mitochondrial matrix and peroxisome) to understand the role of these functions in these organelles and the nature of systems maintaining redox state at a sub-cellular level.

For this study, cells transformed with compartment-targeted roGFP2 constructs were grown in SC_{URA} to exponential ($A_{600} \sim 0.5$) and stationary phase ($A_{600} = 5-6$) and analyzed via flow cytometry. E_{GSH} was measured in the cytosol, mitochondrial matrix and peroxisome of 42 mutants affected in the transcrip-

tional response to oxidative stress, NADPH generation, the thioredoxin and glutathione systems and antioxidant defences. A full list of the pathways and mutants is given in Table 3. Compartmental E_{GSH} was subsequently estimated for all mutants with the experiment carried out in triplicate. For the calculation of E_{GSH} the pH values of the cytosol, mitochondrial matrix and peroxisome using targeted pHluorin probes were calculated at 7.5, 7.7 and 8 respectively and were used for E_{GSH} determination. Data obtained for cells in both exponential phase ($A_{600} \sim 0.5$) and stationary phase ($A_{600} = 5-6$) are presented in Dataset S2.

From this targeted screen, genes affecting redox state in all organelles tested and genes unique for maintenance of redox state in a particular compartment were identified in both exponential and stationary phase. These are outlined in Figure 2A and 2B. Only 12 mutants (exponential phase) and 10 mutants (stationary phase) out of the 42 tested had altered redox in one or more of the compartments. Given the fact that a large number of genes and systems are known to deal with ROS and major perturbations of redox homeostasis, it is significant that mutations in most of these systems had no influence on redox state measured using roGFP2. This indicates that cellular redox homeostasis is tightly regulated and that only a subset of key oxidative stress proteins modulate E_{GSH} .

The most prominent result was the requirement for glutathione reductase (Glr1p), NADPH regeneration systems and the transcription factors Yap1p and Skn7p for the maintenance of redox state across compartments, highlighting the fundamental role of these systems in redox homeostasis. The requirement for glutathione reductase was independent of growth phase unlike that of Yap1p and Skn7p, which appeared important only during exponential phase in all compartments. The identification of Yap1p and Skn7p as essential for cytosolic redox and response to stress indicate that both transcription factors sit at the boundary of maintenance-response systems. Basal levels of expression of the genes they regulate are important for steady-state redox homeostasis, while induction and activation of target genes appears important for mounting appropriate responses to oxidative stress.

From the data it appears that both the cytosol and mitochondrial matrix contain systems required to meet the unique redox and metabolic requirements of each organelle. The mitochondrial redox state was only affected by deletion of oxidative stress transcription factors and mitochondrial NADPH generating enzymes such as the NADH kinase encoded by *POS5* and the NADP⁺-dependent isocitrate dehydrogenase encoded by *IDP1* or mitochondrial ROS detoxifiers Mn-superoxide dismutase and cytochrome c peroxidase encoded by *SOD2* and *CCP1* respectively. The *ccp1* mutant had an altered matrix redox state in both growth phases, while *sod2* mutants only exhibited altered matrix redox in stationary phase, probably due to the increased requirement for Sod2p in stationary phase [27]. The significant role that these two genes play in maintaining mitochondrial redox state reflects the higher burden this compartment faces in terms of endogenous ROS production and subsequent detoxification as part of normal metabolism compared to other compartments. The effect of deleting *CCP1* is particularly interesting because in no other compartment did deletion of a peroxide-detoxifying gene result in alteration to redox state. Given that the mitochondrial matrix has other peroxide-detoxifying enzymes, for example Prx1p [28] and Cta1p [29], the data suggest a separate and significant role for Ccp1p in mitochondrial redox regulation.

In contrast, it was observed that peroxisomal redox was highly reliant on cytosolic rather than peroxisomal redox systems, with glutathione reductase activity, the operation of the pentose phosphate pathway and Cu,Zn superoxide dismutase all critical

Table 3. Genes and known pathways important antioxidant function and resistance to oxidative stress [53].

Process	Genes involved
Transcription factors	<i>msn2, msn4, yap1, skn7 gnd1, gnd2, tal1,</i>
NADPH regeneration	<i>sol3, sol4,rpe1, tk11, ald6, idp1, pos5, idp2, idp3</i>
GSH system	<i>glr1, gpx1, gp2, gpx3, grx1, grx2, grx3, grx4, grx5, grx6, grx7, grx8</i>
Thioredoxin system	<i>trx1, trx2, trx3, trr2</i>
Peroxioredoxins	<i>prx1, tsa1, tsa2, ahp1, dot5</i>
Antioxidants	<i>ctt1, cta1, sod1, sod2, ccp1</i>

Description and gene information were obtained from the *Saccharomyces Genome Database* (SGD).
doi:10.1371/journal.pone.0044278.t003

for maintenance of the redox environment in the peroxisome. Since very little is known about the redox systems in place in the peroxisome, the data may help to understand the manner in which the peroxisome regulates redox homeostasis and its relationship with other compartments. The dependence of the peroxisome on cytosolic pathways indicates a clear inter-relationship between peroxisomal and cytosolic redox homeostasis.

One outcome of this study has been the identification of a function for genes that are annotated in the *Saccharomyces Genome Database* as being “of unknown or putative function”. This has led to new or improved annotations of 10 genes (Table 4) on the basis of their involvement in redox homeostasis and potentially in oxidative stress responses. A good example is *HMS2*, which is a paralogue of *SKN7* arising from the ancient genome duplication [30]. *HMS2* is a putative transcription factor that is annotated by SGD as a “protein with similarity to heat shock transcription factors” involved in pseudohyphal growth under nitrogen depletion. Despite rapid protein divergence between *HMS2* and *SKN7*, *HMS2* shares the heat-shock factor and DNA-binding domains of Skn7p and has been predicted on the basis of microarray data to regulate a subset of responses to reducing conditions, heat shock and hydrogen peroxide [30]. Moreover, Hms2p is structurally very similar not only to Skn7p but also to Sfl1p, a transcriptional activator and repressor involved in the activation of several stress-response genes including Hsp30, which is strongly induced in response to various environmental stress conditions [31]. *HMS2* therefore, has a greater role in redox and oxidative stress

regulation than previously thought, and may represent a transcriptional system acting in concert with *YAPI/SKN7* transcription factors to regulate redox state.

Discussion

The present study was undertaken with the aim of better understanding the genetic basis of the setting, sensing and maintenance of cellular redox homeostasis. The development of redox-sensitive fluorescent probes such as roGFP2 has enabled the high throughput study of redox state in an *in vivo* and real-time manner within distinct cellular compartments, circumventing many of the issues associated with traditional techniques for estimating redox state, such as mixing of compartmental glutathione pools and cell volume estimation. This study represents the first systematic genome-wide study analyzing cellular redox state and is the first to use genome-wide redox potential data to comprehensively understand the bases of redox homeostasis at a compartmental level. The combination of a genome-wide screen with a targeted sub-screen to identify factors contributing to sub-cellular redox homeostasis, highlights the value of *in vivo* redox probes in examining redox homeostasis and broader cellular issues involving redox perturbations. While roGFP2 predominantly equilibrates with the glutathione couple, its redox state also acts as a reflection of that of the protein-sulfhydryl redox buffering system. Protein-thiols contribute significantly to the cellular redox state [13] and measurements using




Figure 2. Functional overlap of mutants that have a more oxidizing cytosol and/or mitochondrial matrix and/or peroxisome. Mutants of known pathways involved in antioxidant defence and oxidative stress responses were analyzed in exponential ($A_{600} \sim 0.5$) and stationary phase ($A_{600} = 5-6$) in the cytosol, mitochondrial matrix and peroxisome. Analysis was carried out using flow cytometry and E_{GSH} estimated. 10,000 cells were counted in each conditions and each experiment was conducted in triplicate. Mutants with a ΔE_{GSH} compared to wild type of 10 mV or more in A) exponential and B) stationary phase in each compartment tested are shown. Data taken Dataset S2.
doi:10.1371/journal.pone.0044278.g002

Table 4. Genes annotated as “putative or unknown function” identified as contributing to cytosolic redox homeostasis in this study and their oxidative stress phenotype(s).

Gene	Phenotype	Reference
<i>DOG2</i>	Induced by oxidative stress	[20]
<i>DGR2</i>	Implicated in stress response signalling	[54]
<i>RPI1</i>	Inhibitor of Ras-cAMP pathway	[55]
<i>PMP3</i>	Increased glutathione excretion	[24]
<i>HMS2</i>	Predicted to regulated some responses to reducing conditions, heat shock and hydrogen peroxide. Homolog of <i>SKN7</i>	[56]
<i>SVF1</i>	Inhibits reactive oxygen species generation and promotes cell survival under oxidative stress conditions	[57]
<i>ECM25</i>	Increased glutathione secretion; decreased oxidative stress resistance	[24,58]
<i>ISF1</i>	Similar to Mbr1p, a protein involved in mitochondrial function and stress responses	[59]

doi:10.1371/journal.pone.0044278.t004

roGFP2 are likely to reflect both the glutathione redox state and the protein-thiol network allowing the reporter to be a multi-dimensional read-out of redox conditions in the cell. Since the roGFP2 probe in the wild-type strain is in an almost fully reduced state across the three compartments studied, it could not be used to identify factors that resulted in more reducing cellular conditions. Re-engineered roGFP2 probes with various mid-point potentials [16,32] could be used to study hyper-reducing conditions or investigate the redox state of the endoplasmic reticulum, which is too oxidizing to be studied using roGFP2. It must be kept in mind, however that there is not one definitive cellular redox state. Glutathione is present at a high concentration, which is an important feature of a buffer. Many other important redox couples exist in cells with varying interactions such as the $\text{NADP}^+/\text{NADPH}$ and $\text{Trx}_{\text{OX}}/\text{Trx}_{\text{RED}}$ couples. An increasing body of evidence is pointing to a more complex relationship between the glutathione and thioredoxin systems with a functional overlap between them. The thioredoxin system is important in preventing over-accumulation of GSSG in cells lacking glutathione reductase [33,34]. Additionally, it has been found that the thioredoxin system is specifically involved in regulating glutathione redox potential in the nucleus even in the presence of glutathione reductase [35]. Future studies would be greatly enhanced by the use of a reporter constructs that equilibrate with the thioredoxin system and $\text{NADP}^+/\text{NADPH}$ couple to get a broader understanding of redox cellular dynamics.

From the data obtained, redox homeostasis in the cytosol, mitochondrial matrix and peroxisome is primarily controlled by regulated glutathione cycling in a growth-phase-independent manner since the *glr1* mutant exhibited the highest ΔE_{GSH} . In exponential phase only the Yap1p and Skn7p transcription factors affected sub-cellular redox. The lack of redox shift in the *msn2* and *msn4* mutants indicates that while the Msn2p and Msn4p transcription factors are important for the general stress response [36], individually they play little or no role in maintenance of redox state. Therefore, it is unlikely that genes whose basal levels are highly regulated by Msn2p/Msn4p such as *GRX2* [37] would be required for steady state redox homeostasis and this is reflected in the results. The reduced need during stationary phase for Yap1p and Skn7p in all compartments is probably due to the fact that, during exponential phase, cells are actively growing and require the transcription of redox maintenance genes however this diminishes in stationary phase when active metabolism and protein synthesis [38] decline. In stationary phase, the need to regenerate existing reduced glutathione is more important and the

activity of Glr1p is therefore predominant. Similarly, NADPH regeneration systems were important for maintaining the redox state in exponential phase with multiple NADPH regeneration mutants identified (cytosol - *tkl1*, *gnd1*; matrix - *tkl1*, *gnd1*, *idp1*, *pos5*; peroxisome - *rpe1*, *gnd1*, *tkl1*).

The finding that key oxidative stress response elements such as the transcription factors Yap1p and Skn7p, and GSH regeneration are essential for redox maintenance under steady state conditions alter the concept that systems involved in the maintenance of redox homeostasis and systems for response to ROS are distinct from each other. These data support the view that some oxidative stress response systems evolved from mechanisms that were important for maintenance of redox homeostasis. It is evident from this study that redox homeostasis is tightly regulated, with redundant systems in place to ensure the maintenance of the appropriate redox environment in each compartment. For the most part, several mutations would be needed to cause large shifts in the redox environment, highlighting the robustness of the cell in terms of redox control. The phenotype of single mutations does not always reflect the complete network of cell functions. We did not observe a significant redox shift in the *trx1* or *trx2* mutants. However, Drakulic and colleagues (2005) showed that the presence of either *TRX1* or *TRX2* was required for maintenance of E_{GSH} [12], despite the redundancy between cytosolic thioredoxins and glutaredoxins for cell viability [12]. In that study, the greatest shift in E_{GSH} from the wild-type in exponential phase was seen in the *trx1 trx2* double mutant. Therefore, while cellular redox systems are redundant and robust, complex redox relationships still do exist. Multiple mutations that have additive effects most likely represent pathways acting in concert for correct sensing and regulating of particular redox couples and redox homeostasis in general, for example the glutathione and thioredoxin systems. As state previously, there are numerous connections between thioredoxin and glutathione systems. The thioredoxin reductase-glutathione reductase double mutant (*trr1 glr1*) is inviable [39] and the *trr1 glr1* double mutant is inviable [39] and only one single thioredoxin or glutaredoxin is essential for viability in *S. cerevisiae* [40]. Deletion of either or both thioredoxin reductases results in increased intracellular concentrations of oxidized and reduced glutathione [41,42] and recently, it was found that the thioredoxin/thioredoxins reductase system could function as an alternative system to reduce oxidized glutathione *in vivo* in *S. cerevisiae* [33] and *Arabidopsis thaliana* [34].

There is a strong connection between mitochondrial function and redox homeostasis. Deletion of *CCPI* encoding cytochrome

c peroxidase affected matrix redox state in both growth phases. As Ccp1p detoxifies hydrogen peroxide and other peroxides, it appears that detoxification of endogenous hydrogen peroxide is required to maintain mitochondrial matrix redox state. Interestingly, Ccp1p is also involved in oxidative stress signaling leading to activation of the Skn7p transcription factor [43]. This could explain the larger shift in matrix redox state observed in the *ccp1* mutant compared to the *skn7* mutant. Deletion of *CCP1* may simultaneously cause increased hydrogen peroxide accumulation and decreased activation of Skn7p, doubly affecting matrix redox state. The signaling function of Ccp1p provides a clear example of the existence of mitochondrial signaling molecules and of cross-compartmental redox signaling. Despite mitochondrial function being absent from those functions significantly enriched in the dataset using the GO and MIPS databases, it is clear that a relationship exists between some mitochondrial functions and maintenance of cytosolic redox state with many mutations affecting mitochondria identified as affecting cytosolic redox state. From previous work it is known that respiratory-incompetent cells have increased levels of GSSG in stationary phase relative to exponential phase and that mitochondrial function is required for resistance to oxidative stress [44]. Key mitochondrial proteins may act in the cellular signaling and sensing of redox change through retrograde redox signaling pathway(s). Since mitochondria are the largest site of endogenous ROS production, redox state changes in mitochondria may be an initial indicator for overall cellular redox dysfunction.

Little is known about peroxisomal redox despite the peroxisome being a significant site of endogenous ROS production. Despite this, the peroxisome exhibited distinctive redox characteristics, with a curious dependence on cytosolic proteins such as glutathione reductase (Glr1p), enzymes involved in the pentose phosphate pathway (Tk1p, Rpe1p) and Cu, Zn superoxide dismutase (Sod1p) rather than peroxisomal proteins for the maintenance of its redox state in both exponential and stationary phase. These results raise some interesting possibilities regarding the nature of peroxisomal redox maintenance and its relationship with other compartments, the cytosol in particular. Free diffusion of small molecules between the cytosol and peroxisome has been hypothesized to account for the pool of GSH and GSSG and hence the redox state of the peroxisome [45] and the results are suggestive of peroxisomal redox state being reliant on cytosolic redox maintenance systems. However, recently, an isoform of glutathione reductase has been found to be dually targeted to the cytosol and peroxisome in *Arabidopsis thaliana* by a novel peroxisomal targeting sequence [46] raising the possibility that glutathione reductase may be localized to the peroxisome in other organisms and that the peroxisome may itself contain a discrete glutathione system.

This study identified approximately 100 genes involved in setting cytosolic redox state using the roGFP2 redox reporter construct. The processes identified highlight that certain redox-active systems traditionally thought of as stress response systems are critical for steady state redox homeostasis, and indicate that redox maintenance systems may have evolved to also be important for responding to reactive oxygen species. The study also found a phenotype of redox dysfunction in 12 genes of unknown or putative function and provides a rich source for further analysis of their cellular role and function in redox homeostasis.

Materials and Methods

Strains, Media and Growth Conditions

For genome-wide screening, the homozygous diploid knock-out mutant collection (Euroscarf) [15] (*MAT α /MAT α his3 Δ 1/his3 Δ 1 leu2 Δ 0/leu2 Δ 0 met15 Δ 0/MET15 LYS2/lys Δ 0 ura3 Δ 0/ura3 Δ 0, *KanMX4::target* gene) and the tetO₇ promoter replacement collection (OpenBiosystems) [14] (*MAT α ; his3 Δ 1; leu2 Δ 0; met15 Δ 0; URA::CMV-*tTA*, kanR-tetO₇-TATA::target* gene) were used. Strains were grown in synthetic complete (SC) medium (0.17% yeast nitrogen base without amino acids, 0.5% ammonium sulfate, 2% w/v glucose) supplemented with appropriate amino acids and bases (Table S1). Dropout medium with the appropriate supplement omitted to provide plasmid selection is denoted by a subscript, for example SC medium lacking leucine is denoted as SC_{LEU}. Media were solidified by the addition of 2% (w/v) agar. Strains were transformed with plasmids using a 96-well plate transformation protocol [47].*

Organelle-targeted roGFP2 Construction

Details of organellar-targeted roGFP2 constructs - pAG416-roGFP2 and pAG416-COX4roGFP2 are described in [18]. The peroxisome-targeted roGFP2 probe, pAG416-roGFP2-SKL was generated using PCR amplification and the Invitrogen Gateway System[®]. Briefly roGFP2 was PCR amplified from the plasmid pBinCM-TKTP-*GRX*/roGFP2 and the peroxisomal C-terminal targeting sequence seryl-lysyl-leucine (SKL) was added to the primer used for amplification. *attb1* or *attb2* sites were added to each primer as appropriate so that the resulting PCR products were flanked by a 5' *attb1* site and a 3' *attb2* site. The resulting PCR fragments were used for the Gateway[®] BP clonase[®] reaction using the destination vector pDonr221. The pDonr221-roGFP2-SKL plasmid was generated using the Invitrogen Gateway[®] system according to manufacturer's instructions and the resulting plasmid pDonr221roGFP2-SKL was used for the Gateway LR[®] clonase[®] reaction using the destination vector pAG416GPD-ccdB [48]. pHluorin organelle-targeted probes were made in an identical manner to the roGFP2 based probes using pCB901YpHc as the starting vector encoding pHluorin a kind gift from R. Rao (Johns Hopkins University School of Medicine, Baltimore).

Organelle-targeted roGFP2 Localization and Redox-sensitivity Verification

Correct localization of mitochondrial matrix-targeted roGFP2 was confirmed by confocal co-localization [18]. Correct localization of all plasmids used in the study was done via confocal microscopy (Figure S1). The redox responsiveness of cells expressing roGFP2-based plasmids was validated via flow cytometry and western blot. Cells were grown in SC_{URA} for 48 h (30°C; 600 rpm) and inoculated into SC_{URA} (A_{600} = 0.001) and grown to exponential phase. Cells were treated with various concentrations of the strong reductant dithiothreitol and the oxidant diamide for 30 min and redox state of roGFP2 determined (data not shown).

Analysis of Compartmental Redox Potential

Cells transformed with the roGFP2 constructs were grown in SC_{URA} for 2 d (30°C; 700 rpm), inoculated in SC_{URA} medium and grown with shaking at 25°C until appropriate cell yield had been reached. All analyses were carried out in 96-well plate format. For preliminary analyses, cytosolic redox state of the cells was analyzed in a fluorescent spectrophotometer (SpectraMax M2, Molecular Devices) with excitation at 405 nm and 488 nm. The emission after excitation at each wavelength was measured and

then the $R_{405/488}$ ratio calculated, with $R_{405/488}$ representing the ratio of emission after excitation at 405 nm divided by the emission after excitation at 488 nm. Given that this was the first round of screening a lenient cut-off were used to identify mutants with altered cytosolic E_{GSH} . In this preliminary screening we identified mutants that had any change in $R_{405/488}$ and any change in either or both the emission after excitation at 405 nm and the emission after excitation at 488 nm. These were then further screened. Subsequent sub-screening and organellar redox analyses were conducted using a FACS Canto II flow cytometer (Becton Dickinson, Australia). For non-essential mutants, cells transformed with roGFP2 were grown to exponential phase (A_{600} 0.5–2.0) in SC_{URA} . For screening of the tetO₇ collection, cells transformed with roGFP2 (leucine selection) were grown in SC_{LEU} for 2 d (25°C; 700 rpm) and inoculated into $SC_{LEU}+10$ µg/ml doxycycline and grown for 24 or 36 h (25°C; 700 rpm). Cells were then analyzed using the FACS Canto II (BD Bioscience) equipped with a laser that excites at 405 nm and 488 nm and emission was detected at 512 nm. The ratio of the emission after excitation at 405 and 488 nm was then calculated. The $R_{405/488}$ and E_{GSH} calculated as previously reported [18,49]. For each sample 10,000 cells were analyzed in at least three biological replicates. For analysis of significance differences in $R_{405/488}$ and E_{GSH} values t-tests were conducted and all E_{GSH} values reported as significant were calculated using a p -value of 0.001. In each case the entire experiment was repeated at least twice. Data obtained were analyzed with FloJo version 7.4 (Tree Star Inc.).

E_{GSH} Calculation

The following modified Nernst equation [49] was used to determine E_{GSH} from the roGFP2 405/488 ratio. Cells were also treated for 20 min with DTT (10 mM) or diamide (10 mM) to obtain measurements of the fully reduced or oxidized (respectively) forms of the roGFP2 probe that are necessary to calculate E_{GSH} using the following formulae derived from the Nernst equation.

Briefly, the degree of oxidation (OxD) of the roGFP2 sensor was calculated from

$$OxD = \frac{R - R_{red}}{\frac{I_{488ox}}{I_{488red}}(R_{ox} - R) + (R - R_{red})}$$

where R is the ratio of emission after excitation at 405/488 nm, R_{red} is the emission ratio of the fully reduced form following incubation with 10 mM DTT (20 min), R_{ox} is the emission ratio of the fully oxidized form following incubation with 10 mM diamide (20 min) and I_{488ox} and I_{488red} are the respective emission intensities at 488 nm for the fully oxidized and fully reduced forms of a given sample. The redox potential was then estimated using:

$$E_{GSH} = E_{0(roGFP2)}^{pH} - \frac{2.303RT}{zF} \log_{10} \frac{1 - OxD_{roGFP2}}{OxD_{roGFP2}}$$

where R is the gas constant (8.315 J K⁻¹ mol⁻¹), T is the absolute temperature (298.15 K), z is the number of transferred electrons (2), F is the Faraday constant (9.648 10⁴ C mol⁻¹) and $E_{0(roGFP2)}^{pH}$ is the mid-point redox potential based on the standard mid-point potential E_0 at 30°C and pH 7, adjusted for the estimated compartment pH and experimental temperature (20–25°C) according to:

$$E_{0(roGFP2)}^{pH} = E_{0(roGFP2)}' - \frac{2.303RT}{zF} (pH - 7)$$

with the mid-point redox potential for roGFP2 ($E_{roGFP2}^{0'}$) estimated as -272 mV.

For analysis of significance differences, t-tests were conducted on samples using $\alpha = 0.001$.

Analysis of Compartmental pH

Wild-type (BY4743) cells transformed with the pHluorin constructs were grown in SC_{URA} for 2 d (30°C; 700 rpm), inoculated in SC_{URA} medium and grown with shaking at 25°C until appropriate cell yield had been reached. Cells were analyzed using the FACS Canto II (BD Bioscience) with excitation at 405 nm and 488 nm and emission was detected at 512 nm. The ratio of the emission after excitation at 405 nm and 488 nm was then calculated and used for pH estimation. 10,000 cells were counted for each sample. Data obtained were analyzed with FlowJo version 7.4 software. To obtain pH values using the pHluorin probe in each compartment, a standard curve was established to allow estimation of intracellular pH by determining the fluorescence of digitonin-permeabilized cells incubated in buffers with pH values of 5 to 8 [50–52]. For calibration of the cytosolic-, mitochondrial matrix-, and peroxisomal-localized pHluorin, cells were suspended in 0.65 M sorbitol containing 100 mg/ml digitonin and incubated (10 min; 600 rpm; 24°C). Cells were pelleted, washed and resuspended in calibration buffer containing 50 mM 2-(*N*-morpholino)ethanesulfonic acid (MES), 50 mM *N*-(2-hydroxyethyl)piperazine-*N*-(2-ethanesulfonic acid) (HEPES), 50 mM KCl, 50 mM NaCl, 200 mM ammonium acetate, 10 mM Na₃N, 10 mM 2-deoxyglucose, 75 µM monensin and 10 µM nigericin, titrated to different pH values using NaOH within the range of pH 6.0–8.0. A calibration curve was plotted using the ratio of fluorescence intensity at 405 nm and 488 nm (after background subtraction) against pH.

Analysis of Genome-wide Data

Cellular processes over-represented in the genome-wide screen data were identified using the Gene Ontology (GO) and Munich Information Centre for Protein Sequences (MIPS) databases via FunSpec [19] with the p -value threshold set at 0.01. FunSpec calculated p -values using a hypergeometric distribution, and the p -value represents the probability that the intersection of a given list with any given functional category occurs by chance. The p -values were Bonferroni corrected to account for spurious significance due to multiple testing over the categories of a database.

Supporting Information

Figure S1 Confocal microscopy images of yeast cells expressing peroxisomal roGFP2.
(EPS)

Figure S2 Verification of organelle-targeted roGFP2 responsiveness to redox state.
(EPS)

Table S1 Supplements for synthetic complete (SC) medium.
(DOC)

Dataset S1 Non-essential mutants identified as having a significantly more oxidized cytosol than wild type.
(XLS)

Dataset S2 E_{GSH} in the cytosol, mitochondrial matrix and peroxisome of cells lacking genes involved in redox homeostasis and antioxidant defence in exponential and stationary phase.
(DOC)

References

- Tu BP, Ho-Schleyer SC, Travers KJ, Weissman JS (2000) Biochemical basis of oxidative protein folding in the endoplasmic reticulum. *Science* 290: 1571–1574.
- Tu BP, Weissman JS (2004) Oxidative protein folding in eukaryotes: mechanisms and consequences. *J Cell Biol* 164: 341–346.
- Riemer J, Bulleid N, Herrmann JM (2009) Disulfide formation in the ER and mitochondria: two solutions to a common process. *Science* 324: 1284–1287.
- Tokatlidis K (2005) A disulfide relay system in mitochondria. *Cell* 121: 965–967.
- Mesecke N, Terziyska N, Kozany C, Baumann F, Neupert W, et al. (2005) A disulfide relay system in the intermembrane space of mitochondria that mediates protein import. *Cell* 121: 1059–1069.
- Janssen-Heininger YM, Mossman BT, Heintz NH, Forman HJ, Kalyanaraman B, et al. (2008) Redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radic Biol Med* 45: 1–17.
- Muller EG (1995) A redox-dependent function of thioredoxin is necessary to sustain a rapid rate of DNA synthesis in yeast. *Arch Biochem Biophys* 318: 356–361.
- Burhans WC, Heintz NH (2009) The cell cycle is a redox cycle: linking phase-specific targets to cell fate. *Free Radic Biol Med* 47: 1282–1293.
- Schafer FQ, Buettner GR (2001) Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med* 30: 1191–1212.
- Barnham KJ, Masters CL, Bush AI (2004) Neurodegenerative diseases and oxidative stress. *Nat Rev Drug Discov* 3: 205–214.
- Kaneto H, Katakami N, Kawamori D, Miyatsuka T, Sakamoto K, et al. (2007) Involvement of oxidative stress in the pathogenesis of diabetes. *Antioxid Redox Signal* 9: 355–366.
- Drakulic T, Temple MD, Guido R, Jarolim S, Breitenbach M, et al. (2005) Involvement of oxidative stress response genes in redox homeostasis, the level of reactive oxygen species, and ageing in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 5: 1215–1228.
- Hansen RE, Roth D, Winther JR (2009) Quantifying the global cellular thiol-disulfide status. *Proc Natl Acad Sci USA* 106: 422–427.
- Mnaimneh S, Davierwala AP, Haynes J, Moffat J, Peng WT, et al. (2004) Exploration of essential gene functions via titratable promoter alleles. *Cell* 118: 31–44.
- Winzler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, et al. (1999) Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* 285: 901–906.
- Meyer AJ, Brach T, Marty L, Kreye S, Rouhier N, et al. (2007) Redox-sensitive GFP in *Arabidopsis thaliana* is a quantitative biosensor for the redox potential of the cellular glutathione redox buffer. *Plant J* 52: 973–986.
- Gutscher M, Pauleau AL, Marty L, Brach T, Wabnitz GH, et al. (2008) Real-time imaging of the intracellular glutathione redox potential. *Nat Meth* 5: 553–559.
- Ayer A, Tan SX, Grant CM, Meyer AJ, Dawes IW, et al. (2010) The critical role of glutathione in maintenance of the mitochondrial genome. *Free Radic Biol Med* 49: 1956–1968.
- Robinson M, Grigull J, Mohammad N, Hughes T (2002) FunSpec: a web-based cluster interpreter for yeast. *BMC bioinformatics* 3: 35.
- Tsujimoto Y, Izawa S, Inoue Y (2000) Cooperative regulation of *DOG2*, encoding 2-deoxyglucose-6-phosphate phosphatase, by Snf1 kinase and the high-osmolarity glycerol-mitogen-activated protein kinase cascade in stress responses of *Saccharomyces cerevisiae*. *J Bact* 182: 5121–50418.
- Grant C, MacIver FH, Dawes IW (1997) Glutathione synthetase is dispensable for growth under both normal and oxidative stress conditions in the yeast *Saccharomyces cerevisiae* due to an accumulation of the dipeptide gamma-glutamylcysteine. *Mol Biol Cell* 8: 1699–1707.
- Charizanis C, Juhnke H, Krems B, Entian KD (1999) The mitochondrial cytochrome c peroxidase Ccp1 of *Saccharomyces cerevisiae* is involved in conveying an oxidative stress signal to the transcription factor Pos9 (Skn7). *Mol Gen Genet* 262: 437–447.
- Goebel MG, Yochem J, Jentsch S, McGrath JP, Varshavsky A, et al. (1988) The yeast cell cycle gene *CDC34* encodes a ubiquitin-conjugating enzyme. *Science* 241: 1331–1335.
- Perrone GG, Grant CM, Dawes IW (2005) Genetic and environmental factors influencing glutathione homeostasis in *Saccharomyces cerevisiae*. *Mol Biol Cell* 16: 218–230.
- Wheeler GL, Quinn KA, Perrone G, Dawes IW, Grant CM (2002) Glutathione regulates the expression of glutamylcysteine synthetase via the Met4 transcription factor. *Mol Micro* 46: 545–556.
- Tan SX, Teo M, Lam YT, Dawes IW, Perrone GG (2009) Cu, Zn superoxide dismutase and NADP(H) homeostasis are required for tolerance of endoplasmic reticulum stress in *Saccharomyces cerevisiae*. *Mol Biol Cell* 20: 1493–1508.
- Flattery-O'Brien JA, Grant CM, Dawes IW (1997) Stationary-phase regulation of the *Saccharomyces cerevisiae SOD2* gene is dependent on additive effects of HAP2/3/4/5- and STRE-binding elements. *Mol Micro* 23: 303–312.
- Pedrajas JR, Miranda-Vizuete A, Javanmardy N, Gustafsson J, Spyrou G (2000) Mitochondria of *Saccharomyces cerevisiae* contain one-conserved cysteine type peroxiredoxin with thioredoxin peroxidase activity. *J Biol Chem* 275: 16296–16301.
- Petrova VY, Drescher D, Kujumdzieva AV, Schmitt MJ (2004) Dual targeting of yeast catalase A to peroxisomes and mitochondria. *Biochem J* 380: 393–400.
- Kellis M, Birren BW, Lander ES (2004) Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature* 428: 617–624.
- Ansanay Galeote V, Alexandre H, Bach B, Delobel P, Dequin S, et al. (2007) Sfl1p acts as an activator of the *HSP30* gene in *Saccharomyces cerevisiae*. *Curr Genet* 52: 55–63.
- Lohman JR, Remington SJ (2008) Development of a family of redox-sensitive green fluorescent protein indicators for use in relatively oxidizing subcellular environments. *Biochem* 47: 8678–8688.
- Tan SX, Greetham D, Raeth S, Grant CM, Dawes IW, et al. (2010) The thioredoxin-thioredoxin reductase system can function *in vivo* as an alternative system to reduce oxidized glutathione in *Saccharomyces cerevisiae*. *J Biol Chem* 285: 6118–6126.
- Marty L, Siala W, Schwarzlander M, Fricker MD, Wirtz M, et al. (2009) The NADPH-dependent thioredoxin system constitutes a functional backup for cytosolic glutathione reductase in *Arabidopsis*. *Proc Natl Acad Sci USA* 106: 9109–9114.
- Dardalhon M, Kumar C, Iraqi I, Vernis L, Kienda G, et al. (2012) Redox-sensitive YFP sensors monitor dynamic nuclear and cytosolic glutathione redox changes. *Free Radic Biol Med* 52: 2254–2265.
- Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, et al. (2000) Genomic expression programs in the response of yeast cells to environmental changes. *Mol Biol Cell* 11: 4241–4257.
- Grant CM, Luikenhuis S, Beckhouse A, Soderbergh M, Dawes IW (2000) Differential regulation of glutaredoxin gene expression in response to stress conditions in the yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta-Gene Structure and Expression* 1490: 33–42.
- Fuge EK, Braun EL, Werner-Washburne M (1994) Protein synthesis in long-term stationary-phase cultures of *Saccharomyces cerevisiae*. *J Bacteriol* 176: 5802–5813.
- Trotter EW, Grant CM (2003) Non-reciprocal regulation of the redox state of the glutathione, glutaredoxin and thioredoxin systems. *EMBO reports* 4: 184–188.
- Draculic T, Dawes IW, Grant CM (2000) A single glutaredoxin or thioredoxin gene is essential for viability in the yeast *Saccharomyces cerevisiae*. *Mol Micro* 36: 1167–1174.
- Muller E (1996) A glutathione reductase mutant of yeast accumulates high levels of oxidized glutathione and requires thioredoxin for growth. *Mol Biol Cell* 7: 1805–1813.
- Garrido EO, Grant CM (2002) Role of thioredoxins in the response of *Saccharomyces cerevisiae* to oxidative stress induced by hydroperoxides. *Mol Micro* 43: 993–1003.
- Charizanis C, Juhnke H, Krems B, Entian KD (1999) The mitochondrial cytochrome c peroxidase Ccp1 of *Saccharomyces cerevisiae* is involved in conveying an oxidative stress signal to the transcription factor Pos9 (Skn7). *Mol Gen Genet* 262: 437–447.
- Grant CM, MacIver FH, Dawes IW (1997) Mitochondrial function is required for resistance to oxidative stress in the yeast *Saccharomyces cerevisiae*. *FEBS Lett* 410: 219–222.
- Antonovkov VD, Grunau S, Ohlmeier S, Hiltunen JK (2010) Peroxisomes are oxidative organelles. *Antioxid Redox Signal* 13: 525–537.
- Kataya AR, Reumann S (2010) *Arabidopsis* glutathione reductase 1 is dually targeted to peroxisomes and the cytosol. *Plant Signal Behav* 5: 171–175.
- Gietz RD, Schiestl RH (2007) Microtiter plate transformation using the LiAc/SS carrier DNA/PEG method. *Nature Protoc* 2: 5–8.
- Alberti S, Gitler AD, Lindquist S (2007) A suite of Gateway cloning vectors for high-throughput genetic analysis in *Saccharomyces cerevisiae*. *Yeast* 24: 913–919.
- Schwarzlander M, Fricker MD, Muller C, Marty L, Brach T, et al. (2008) Confocal imaging of glutathione redox potential in living plant cells. *J Microsc* 231: 299–316.
- Brett CL, Tukaye DN, Mukherjee S, Rao R (2005) The yeast endosomal Na⁺K⁺/H⁺ exchanger Nhx1 regulates cellular pH to control vesicle trafficking. *Mol Biol Cell* 16: 1396–1405.

Author Contributions

Conceived and designed the experiments: AA GGP IWD. Performed the experiments: AA SF CF SSL GJS. Analyzed the data: AA GGP IWD. Contributed reagents/materials/analysis tools: AJM. Wrote the paper: AA GGP IWD.

51. Ali R, Brett CL, Mukherjee S, Rao R (2004) Inhibition of sodium/proton exchange by a Rab-GTPase-activating protein regulates endosomal traffic in yeast. *J Biol Chem* 279: 4498–4506.
52. van Roermund CWT, de Jong M, Ijlst L, van Marle J, Dansen TB, et al. (2004) The peroxisomal lumen in *Saccharomyces cerevisiae* is alkaline. *Journal of cell science* 117: 4231–4237.
53. Jamieson DJ (1998) Oxidative stress responses of the yeast *Saccharomyces cerevisiae*. *Yeast* 14: 1511–1527.
54. Villa-Garcia MJ, Choi MS, Hinz FI, Gaspar ML, Jesch SA, et al. (2010) Genome-wide screen for inositol auxotrophy in *Saccharomyces cerevisiae* implicates lipid metabolism in stress response signaling. *Mol Genet and Genom* 285: 125–149.
55. Kim J, Powers S (1991) Overexpression of *RP11*, a novel inhibitor of the yeast Ras-cyclic AMP pathway, down-regulates normal but not mutationally activated ras function. *Mol Cell Biol* 11: 3894–3904.
56. Segal E, Shapira M, Regev A, Peier D, Botstein D, et al. (2003) Module networks: Discovering regulatory modules and their condition specific regulators from gene expression data. *Nature Genet* 34: 166–176.
57. Brace JL, VanderWeele DJ, Rudin CM (2005) Svf1 inhibits reactive oxygen species generation and promotes survival under conditions of oxidative stress in *Saccharomyces cerevisiae*. *Yeast* 22: 641–652.
58. Higgins VJ, Alic N, Thorpe GW, Breitenbach M, Larsson V, et al. (2002) Phenotypic analysis of gene deletion strains for sensitivity to oxidative stress. *Yeast* 19: 203–214.
59. Daignan-Fornier B, Nguyen CC, Reisdorf P, Lemeignan B, Bolotin-Fukuhara M (1994) *MBR1* and *MBR3*, two related yeast genes that can suppress the growth defect of hap2, hap3 and hap4 mutants. *Mol Gen Genet MGG* 243: 575–583.