

Association of Thymidylate Synthase Polymorphisms with Acute Pancreatitis and/or Peripheral Neuropathy in HIV-Infected Patients on Stavudine-Based Therapy

Pere Domingo^{1*}, Maria del Carmen Cabeza¹, Ferran Torres², Juliana Salazar³, Maria del Mar Gutierrez¹, Maria Gracia Mateo¹, Esteban Martínez⁴, Joan Carles Domingo⁵, Irene Fernandez¹, Francesc Villarroya⁵, Esteban Ribera⁶, Francesc Vidal⁷, Montserrat Baiget⁸

1 Infectious Diseases Unit, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain, **2** Biostatistics and Data Management Platform, IDIBAPS, Hospital Clinic; Biostatistics Unit, School of Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain, **3** CIBERER (U-705), Barcelona, Spain, **4** Infectious Diseases Service, Hospital Clínic Universitari, Universitat de Barcelona, Barcelona, Spain, **5** Department of Biochemistry and Molecular Biology, University of Barcelona, Barcelona, Spain, **6** Infectious Diseases Service, Hospital Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain, **7** Hospital Universitari Joan XXIII, Universitat Rovira i Virgili, Tarragona, Spain, **8** Department of Genetics, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain

Abstract

Background: Low expression thymidylate synthase (TS) polymorphism has been associated with increased stavudine triphosphate intracellular (d4T-TP) levels and the lipodystrophy syndrome. The use of d4T has been associated with acute pancreatitis and peripheral neuropathy. However, no relationship has ever been proved between TS polymorphisms and pancreatitis and/or peripheral neuropathy.

Methods: We performed a case-control study to assess the relationship of TS and methylene-tetrahydrofolate reductase (MTHFR) gene polymorphisms with acute pancreatitis and/or peripheral neuropathy in patients exposed to d4T. Student's t test, Pearson's correlations, one-way ANOVA with Bonferroni correction and stepwise logistic regression analyses were done.

Results: Forty-three cases and 129 controls were studied. Eight patients (18.6%) had acute pancreatitis, and 35 (81.4%) had peripheral neuropathy. Prior AIDS was more frequent in cases than in controls (OR = 2.36; 95%CI 1.10–5.07, P = 0.0247). Low expression TS and MTHFR genotype associated with increased activity were more frequent in patients with acute pancreatitis and/or peripheral neuropathy than in controls (72.1% vs. 46.5%, OR = 2.97; 95%CI: 1.33–6.90, P = 0.0062, and 79.1% vs. 56.6%, OR = 2.90, 95%CI: 1.23–7.41, P = 0.0142, respectively). Independent positive or negative predictors for the development of d4T-associated pancreatitis and/or peripheral neuropathy were: combined TS and MTHFR genotypes (reference: A+A; P = 0.002; OR_{A+B} = 0.34 [95%CI: 0.08 to 1.44], OR_{B+A} = 3.38 [95%CI: 1.33 to 8.57], OR_{B+B} = 1.13 [95%CI: 0.34 to 3.71]), nadir CD4 cell count >200 cells/mm³ (OR = 0.38; 95%CI: 0.17–0.86, P = 0.021), and HALS (OR = 0.39 95%CI: 0.18–0.85, P = 0.018).

Conclusions: Low expression TS plus a MTHFR genotype associated with increased activity is associated with the development of peripheral neuropathy in d4T-exposed patients.

Citation: Domingo P, Cabeza MdC, Torres F, Salazar J, Gutierrez MdM, et al. (2013) Association of Thymidylate Synthase Polymorphisms with Acute Pancreatitis and/or Peripheral Neuropathy in HIV-Infected Patients on Stavudine-Based Therapy. PLoS ONE 8(2): e57347. doi:10.1371/journal.pone.0057347

Editor: Rui Medeiros, IPO, Inst Port Oncology, Portugal

Received: October 9, 2012; **Accepted:** January 21, 2013; **Published:** February 28, 2013

Copyright: © 2013 Domingo et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work has been partially funded by Fondo de Investigaciones Sanitarias (FIS 02/1280, 05/1591, 07/0976, 08/00256), Fundación para la Prevención del SIDA en España (FIPSE 36610, 36572/06), and Red de Investigación en SIDA (RIS RD06/006/0022). Pere Domingo is supported by Programa de Intensificación de Investigadores del Instituto de Salud Carlos III (INT12/383). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: pdomingo@santpau.cat

These authors contributed equally to this work.

Introduction

The doubtless efficacy of highly active antiretroviral therapy (HAART) is still shadowed by drug toxicity, specially that appearing in the long-term [1]. Some of the most important toxic effects experienced by patients on HAART are caused by the nucleoside reverse transcriptase inhibitor (NRTI) part of the regime [2]. Among NRTI-associated toxicity, that caused by

thymidine analogues, namely zidovudine (AZT) and stavudine (d4T) stands out. The common mechanism underlying thymidine analogues' toxic effects is mitochondrial toxicity [3]. The ability of the triphosphate active NRTI to inhibit mitochondrial gamma polymerase causes mitochondrial DNA depletion that may eventually lead to mitochondrial respiratory chain protein de-

pletion which in turn leads to mitochondrial dysfunction [4]. This can ultimately cause cellular dysfunction or even cellular death [5].

The ability of thymidine analogues to inhibit mitochondrial gamma polymerase will depend on the intracellular concentrations of their triphosphate metabolites [6]. Therefore, all the circumstances increasing the levels of these metabolites may be accompanied by a higher degree of mitochondrial toxicity [7]. We recently described how thymidylate synthase (TS) polymorphisms may modify d4T triphosphate (d4T-TP) intracellular concentrations, and how higher intracellular levels of d4T-TP are associated with the HIV/HAART-associated lipodystrophy syndrome (HALS) [7,8].

Our working hypothesis was that TS and MTHFR polymorphisms could be associated with other manifestations of mitochondrial toxicity, such as d4T-associated pancreatitis and d4T-associated peripheral neuropathy. To test this hypothesis we performed a case-control study in d4T-exposed patients who developed pancreatitis or peripheral neuropathy while on d4T-based therapy.

Patients and Methods

Population Studied

All patients and controls were recruited at the same HIV-1 infection clinic at the *Hospital de la Santa Creu i Sant Pau*, which attended a population of 3355 HIV-1-infected patients between 1994 and 2010. All were patients on active follow up with an established diagnosis of HIV-1 infection on d4T-based antiretroviral therapy. Patients were eligible if they had developed pancreatitis and/or peripheral neuropathy while receiving therapy with d4T as part of their antiretroviral regimens. Controls were subjects followed in the same cohort who did not develop pancreatitis and/or neuropathy while on d4T therapy. Controls were selected in a 3:1 ratio with respect to cases. They were matched to patients with respect to age (± 5 years), sex, and exposure to d4T (but not time on d4T). A normal dose of d4T was 40 mg twice daily if the patient weighed >60 kg, and 30 mg twice daily if the patient weighed <60 kg, whereas a reduced dose was 30 mg/12 h and 20 mg/12 h for patients with a weight of ≥ 60 or <60 kg, respectively. Subjects who were hospitalized or had a frank cognitive impairment such as delirium or dementia on enrolment were not eligible. Patients with opportunistic infections, neoplasms or fever of undetermined origin were excluded from the study too. Written informed consent was obtained from the patients at study entry. The study was approved by the Ethics Committee of the *Hospital de la Santa Creu i Sant Pau*.

Definition of Clinical Endpoints

A case of acute pancreatitis was defined as a clinical history consistent with pancreatitis (i.e. abdominal pain with or without findings of shock and hypotension where other causes of abdominal pain have been excluded) with supporting biochemical evidence of pancreatitis: elevated lipase (3 times the upper normal limit) or amylase (3 times the upper normal limit) and evidence of pancreatitis from radiological investigation or hemorrhagic pancreatitis at laparotomy or post-mortem exam [9,10]. Asymptomatic elevations of pancreatic enzymes were not considered to be cases of acute pancreatitis. Chronic pancreatitis cases were excluded from the study.

The diagnosis of neuropathy (sensory or mixed) required the presence of numbness, paresthesias, or dysesthesias in the patient's lower or lower and upper extremities with onset after starting a d4T-based antiretroviral regime [11,12]. A confirmatory electromyogram or a nerve conduction analysis was not a com-

pulsory criterion for the diagnosis of neuropathy. The diagnosis of AIDS was based on the 1993 revised case definition of the Centers for Disease Control and Prevention [13].

The presence or absence of lipodystrophy, lipohypertrophy, and mixed syndrome was determined as previously described [7]. The metabolic syndrome was defined according to the U.S. National Cholesterol Education Program (NCEP) Adult Treatment Panel III Guidelines [14] and modified as recommended in the latest American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement [15]. Alcohol abuse was defined when alcohol intake was >40 g daily [16].

Biochemistry Laboratory Measurements

All laboratory investigations were performed after a 12 h overnight fast and at least 15 minutes after the placement of a peripheral intravenous catheter, as previously described [7,8].

Genotyping Analyses

The genomic DNA was extracted from the peripheral leucocytes by the salting-out procedure [17]. In the TS gene, the variable number tandem repeat (VNTR) of 28 bp polymorphism and the G \rightarrow C SNP in the first and second repeat were analyzed. A DNA fragment was amplified using previously described PCR conditions and primers [7,8], and directly sequenced using an ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA). This G to C substitution changes a critical residue in the USF E-box consensus element, abolishes USF-1 binding, and alters transcriptional activity. TS genotypes of the patients were classified according to Kawakami and Watanabe into two groups: high expression type (2/3G, 3C/3G and 3G/3G) and low expression type (2/2, 2/3C and 3C/3C) [18].

The MTHFR gene polymorphisms (677C \rightarrow T [rs1801133] and 1298 A \rightarrow C [rs1801131]) were determined. These two polymorphisms were analyzed using Fluidigm's Biomark system. This technology is designed for the allelic discrimination 5' nuclease assay. The samples and the TaqMan Gene expression assays (Applied Biosystems, Foster City, CA, USA) were prepared following manufacturer's instructions. The 48.48 dynamic arrays used were automatically loaded using an IFC Controller (Fluidigm Corporation), and real-time reactions were performed and analyzed using BioMark Real-Time PCR System and Analysis software (Fluidigm Corporation), respectively. As a quality control, normal, heterozygote and homozygote sequenced samples were included on every array for each genotype. MTHFR genotypes were classified also into two groups: those associated with a decreased enzymatic activity (homozygous 677T, homozygous 1298C and compound heterozygous patients), and genotypes associated with an increased enzymatic activity (heterozygous and wild-type patients) [19]. Combined TS and MTHFR genotypes with respect to enzyme activities produced four groups: low TS expression and increased MTHFR activity (B+A), low TS expression and decreased MTHFR activity (B+B), high TS expression and increased MTHFR activity (A+A), and high TS expression and decreased MTHFR activity (A+B).

Statistical Analyses

Data are expressed as median with interquartile range (IQR) or as otherwise specified. Continuous variables were assessed with the nonparametric Mann-Whitney test and categorical data such as genotype and allele frequencies were compared by use of the Fisher's exact test. The level of significance was established at the 0.05 level and all reported P values are two-sided. A logistic regression analysis was used to examine the association of peripheral neuropathy and/or pancreatitis with TS and MTHFR

polymorphisms and other parameters; variables associated with a $P < 0.1$ in the univariate analyses were included in the multivariate stepwise analysis. All analyses were performed with the SAS version 9.1.3 software (SAS Institute Inc., Cary, NC).

Results

Population Studied

From January 1994 to December 2010, there were 102 cases of pancreatitis and/or peripheral neuropathy in our cohort; 34 cases of pancreatitis, 62 peripheral neuropathies and 6 cases with both pancreatitis and neuropathy. Of them, 63 (61.7%) were associated with d4T exposure, which had been taken at any time by 899 patients for a total exposure of 39,684 months. The incidence of d4T-associated pancreatitis and/or neuropathy was 19.05 (95% CI: 14.28–24.39) cases per 1000 patient-years (PY) in those exposed to d4T and 4.28 (3.13–5.86) in those not exposed ($P < 0.0001$). For peripheral neuropathy the incidences were 18.75 (14.62–24.05) and 1.98 (1.25–3.14), respectively ($P < 0.0001$), whereas for pancreatitis they were 10.28 (7.35–14.39) and 2.3 (1.50–3.54), respectively ($P < 0.0001$). We studied 172 HIV-1-infected, Caucasian patients, exposed to a d4T-based therapy (43 cases and 129 controls). There were no differences between the 43 cases studied and the 20 patients excluded because of lack of DNA. Demographic, means of acquiring HIV-1 infection and the immunovirological status of patients and controls are shown in table 1. There were 127 men (73.8%) and 45 women (26.2%), with a mean age of 51.1 ± 10.9 years (median: 49.0 [IQR: 42.0–58.0] years). Sixty-nine patients (40.1%) had had a prior AIDS-defining condition. Eight patients (18.6%) had pancreatitis, and 35 (81.4%) had peripheral neuropathy. Prior AIDS was significantly more frequent in cases than in controls (OR = 2.36; 95%CI: 1.10–5.07, $P = 0.0247$). No patient was diabetic or was using insulin or hypoglycemic agents. Alcohol abuse was documented in 11 patients (6.3%). The median daily alcohol consumption for alcohol abusers was 63.0 (IQR: 55.0–94.5) grams, without statistically significant differences between cases and controls ($P = 0.8738$).

Antiretroviral Drug Exposure and Immunovirological Status

Most of the patients (124, 72.9%) had undetectable viral load at the time of the study. The median viral load for those who had it detectable was 2.0 (IQR: 1.68–3.06) \log_{10} copies/ml. The mean CD4 count was 589 ± 291 cells/ mm^3 (median: 539 [IQR: 383–784]). Nadir CD4 cell count was < 200 cells/ mm^3 in 109 patients (60.5%) and < 100 cells/ mm^3 in 75 (41.7%). A CD4 cell count nadir < 200 cells/ mm^3 was significantly more frequent in cases (OR = 2.34; 95%CI: 1.03–5.58, $P = 0.0437$) (Table 1). The maximum viral load was above 5 \log_{10} copies/ml in 106 patients (61.6%). Four cases (9.3%) and 31 controls (24.0%) were receiving reduced d4T doses (OR = 0.32; 95%CI: 0.08–1.01, $P = 0.0630$). The cumulated exposure to antiretroviral drugs is shown in table 2. More cases than controls were receiving d4T plus ddI at event (53.5% vs. 34.1%, OR = 2.22; 95%CI: 1.04–4.76, $P = 0.0379$). None of the patients was receiving hydroxyurea.

Metabolic Data, Metabolic Syndrome and HALS

Metabolic and fat data are shown in table 3. There were no differences between cases and controls with respect to anthropometric and metabolic parameters. Sixty-nine patients (38.3%) had metabolic syndrome, without differences between cases and controls (OR = 1.59; 0.75–3.32, $P = 0.2498$). Ninety-three patients (54.1%) had HALS. HALS was less frequent in cases than in

controls (41.9% vs. 58.1%, OR = 0.52; 95%CI: 0.24–1.10, $P = 0.0932$).

TS and MTHFR Polymorphisms, and Development of Pancreatitis and/or Peripheral Neuropathy

The distribution of the different genotypes is shown in table 4. Ninety-one patients (52.9%) had a low expression TS genotype. This genotype was more frequent in patients with pancreatitis and/or peripheral neuropathy than in controls (72.1% vs. 46.5%, OR = 2.97; 95%CI: 1.33–6.90, $P = 0.0062$). The association remained statistically significant when peripheral neuropathy alone was considered (68.6% vs. 46.5%, OR = 2.51; 95%CI: 1.07–6.14, $P = 0.0336$), whereas there was a trend to significance for pancreatitis (87.5% vs. 46.5%, OR = 8.05; 95%CI: 0.98–367.94, $P = 0.0592$). Neither 677C→T ($P = 0.1119$) or 1298A→C ($P = 0.0708$) MTHFR polymorphisms were associated with the development of pancreatitis and/or peripheral neuropathy. However, when functional translation was taken into account MTHFR genotypes associated with increased enzymatic activity were more frequent in cases than in controls (79.1% vs. 56.6%, OR = 2.90, 95%CI: 1.23–7.41, $P = 0.01423$). The association remained statistically significant for peripheral neuropathy (77.1% vs. 58.4%, OR = 2.40, 95%CI: 0.97–6.55, $P = 0.0417$), but not for pancreatitis (87.5% vs. 60.9%, OR = 4.48, 95%CI: 0.55–205.51, $P = 0.2553$).

Combined TS and MTHFR genotypes with respect to peripheral neuropathy or pancreatitis are shown in table 4, where it is apparent that those combinations with a greater negative impact on TS activity were more frequently seen in cases.

Factors Associated with Development of Pancreatitis and/or Peripheral Neuropathy

A multivariable analysis was performed taking as the dependent variable the development of pancreatitis and/or peripheral neuropathy and as independent variables, age, sex, AIDS, CD4 count nadir $< 200/\text{mm}^3$, CD4 nadir $< 100/\text{mm}^3$, d4T exposure (m), d4T exposure (g), d4T exposure (mg/kg), AZT exposure (m), AZT exposure (g), ABC exposure (m), d4T plus ddI at event, EFV exposure (m), PI exposure (m), NRTI exposure (m), HALS, TS genotype (low vs. high expression), MTHFR genotype (increased vs. decreased enzymatic activity), and combined TS and MTHFR genotypes, all variables associated with a P value < 0.1 in the univariate analysis. Independent positive or negative predictors for the development of d4T-associated pancreatitis and/or peripheral neuropathy were: combined TS and MTHFR genotypes (reference: A+A; $P = 0.002$; OR_{A+B} = 0.34 [95%CI: 0.08 to 1.44], OR_{B+A} = 3.38 [95%CI: 1.33 to 8.57], OR_{B+B} = 1.13 [95%CI: 0.34 to 3.71]), nadir CD4 cell count > 200 cells/ mm^3 (OR = 0.38; 95%CI: 0.17–0.86, $P = 0.021$), and HALS (OR = 0.39 95%CI: 0.18–0.85, $P = 0.018$).

Discussion

Our study suggests an association between TS and MTHFR polymorphisms and the appearance of d4T-related toxicity in the form of acute pancreatitis or peripheral neuropathy. However, our work has inherent limitations. First, this is a case-control study and therefore no causal relationships should or must be drawn. Second, in case-control studies endpoint verification is of paramount importance. It is known that peripheral neuropathy or acute pancreatitis in the setting of HIV-1 infection treated with HAART may be caused not only by antiretroviral drugs but also by other drugs, alcohol abuse, or biliary stones and hypertriglyceridemia in the case of pancreatitis [20]. On the other hand,

Table 1. Demographics and immunovirological status of the population studied.

	Cases (N = 43)	Controls (N = 129)	P value
Age (yrs.)	48.0 (43.0–57.7)	50.0 (42.0–58.2)	0.9042
Males, n (%)	32 (74.4)	95 (73.6)	0.9999
Risk group			
Msm, n (%)	17 (39.5)	45 (34.9)	0.3366
Htsx, n (%)	14 (32.5)	55 (42.6)	
IDU, n (%)	11 (23.2)	30 (21.7)	
Other*, n (%)	2 (4.6)	1 (0.8)	
Years since diagnosis	13.0 (9.0–15.7)	13.0 (10.0–17.0)	0.4065
AIDS, n (%)	24 (55.8)	45 (34.9)	0.0195
Smokers, n (%)	17 (39.5)	70 (54.3)	0.1138
Alcohol abuse, n (%)	1 (2.3)	10 (7.7)	0.2952
HBV co-infection, n (%)	6 (13.9)	11 (8.5)	0.3751
HCV co-infection, n (%)	13 (30.2)	40 (31.0)	0.9999
Current CD4 count, cells/mm ³	515 (346–730)	543 (388–792)	0.5430
CD4 increase, cells/mm ³	386 (194–574)	339 (207–553)	0.6044
Current CD8 count cells/mm ³	834 (582–1181)	874 (617–1146)	0.6521
CD8 increase, cells/mm ³	555 (144–799)	370 (122–594)	0.1180
Nadir CD4 cell count, cells/mm ³	83 (23–206)	182 (54–314)	0.0091
Nadir CD4 cell count <100 cells/mm ³ , n (%)	24 (55.8)	45 (35.1)	0.0201
Nadir CD4 cell count <200 cells/mm ³ , n (%)	32 (74.4)	71 (55.5)	0.0314
Viral load, log ₁₀ , copies/ml	1.28 (1.28–1.28)	1.28 (1.28–1.55)	0.8307
Undetectable viral load, n (%)	33 (76.7)	91 (70.5)	0.5564
Maximum viral load, log ₁₀ , copies/ml	5.46 (4.70–5.79)	5.19 (4.29–5.58)	0.1479
Maximum viral load ≥5 log ₁₀ , copies/ml	31 (68.9)	75 (55.6)	0.0643
Viral load decrease, log ₁₀ , copies/ml	3.78 (2.84–4.28)	3.58 (2.53–4.16)	0.3725

Values are expressed as median and interquartile range, unless indicated.

*includes 2 patients with post-transfusion HIV and 1 with unknown risk, MsM = men who have sex with men, HTSX = heterosexuals, IDU = intravenous drug users, AIDS = acquired immune deficiency syndrome, HCV = hepatitis C virus, HBV = hepatitis B virus, ml = milliliters.

doi:10.1371/journal.pone.0057347.t001

there is no pathognomic test to etiologically ascribe the etiology of peripheral neuropathy or pancreatitis. Notwithstanding that, none of these well-known causes was present when patients in the study were diagnosed as having peripheral neuropathy or pancreatitis, except for alcohol abuse which in fact was more frequent among controls. Unfortunately, when the study was performed, intracellular d4T-TP concentrations in PBMCs were not available. Third, d4T is a well-known cause of peripheral neuropathy and pancreatitis, especially when combined with didanosine (ddI) or even more when combined with ddI and hydroxyurea [21]. Among our cases, 23 (53.5%) were also exposed to ddI, which may have contributed to the development of pancreatitis and/or peripheral neuropathy. Although cumulated exposure to ddI was not different in cases and controls, taking d4T plus ddI at event was statistically more frequent in cases. However, taking both drugs in combination was not an independent predictor for the development of peripheral neuropathy and/or acute pancreatitis. Fourth, there are a number of gene markers which have been associated with the risk of developing both toxic peripheral neuropathy and pancreatitis in HIV-infected patients on HAART, such as cystic fibrosis trans-membrane conductance regulator (CTR) and serine protease inhibitor kazal-1 (SPINK-1) mutations for pancreatitis and mitochondrial haplotype T for peripheral neuropathy [22–24]. We cannot exclude the possibility that some

of these untested genes may have contributed to acute pancreatitis and/or peripheral neuropathy in our patients.

Acute pancreatitis in the setting of HIV infection and antiretroviral therapy has wide incidence rates ranging from 1.27 to 22.6 events/1000 PY [25–28]. These wide incidence rates may be explained by different diagnostic criteria (clinical vs. laboratory-based), and by the fact that some of the studies were performed in early calendar years, when more toxic drugs were used. Its appearance has been linked to the classical risk factors; i.e. alcohol abuse, hypertriglyceridemia, as well as to the use of antiretroviral drugs such as d4T and ddI [25–27]. The incidence rates for peripheral neuropathy have also been very wide, ranging from 0.7 to 39.7/1000 PY [29]. Known risk factors for developing peripheral neuropathy in HAART-treated patients include alcohol abuse, treatment with other neuropathic drugs (isoniazid, methotrexate,...), factors which were ruled out in our patients [30]. However, sometimes it is difficult to distinguish toxic peripheral neuropathy from exacerbation of HIV-associated neuropathy in HAART-treated patients, which may also contribute to these varying incidence rates. Notwithstanding that, none of our patients had clinical features consistent with peripheral neuropathy prior to taking d4T. The incidence of both acute pancreatitis and peripheral neuropathy in our work fall within these wide ranges, most probably because our study period spans throughout a long

Table 2. Antiretroviral drug exposure in the population studied.

Parameter	Cases (n = 43)	Controls (n = 129)	P value
ART composition at event			0.0154
PI-based, n (%)	33 (76.7)	69 (53.5)	
NNRTI-based, n (%)	10 (23.2)	51 (39.5)	
3 NRTIs, n (%)	0 (0.0)	9 (7.0)	
NRTI backbone at event			
d4T+3TC, n (%)	15 (34.9)	69 (53.5)	0.3335
d4T+ddl, n (%)	23 (53.5)	44 (34.1)	
d4T+ABC, n (%)	2 (4.6)	6 (4.6)	
d4T alone, n (%)	2 (4.6)	6 (4.6)	
d4T+3TC+ABC, n (%)	1 (2.3)	3 (2.3)	
d4T+3TC+ddl, n (%)	0 (0.0)	1 (0.8)	
Current d4T use, n (%)	0(0)	29 (22.5)	0.0015
Current AZT use, n (%)	13 (30.2)	4 (3.1)	<0.0001
ART duration (m)	110.0 (93.0–138.7)	115.0 (94.7–143.2)	0.4783
Individual drug exposure			
AZT exposure, m	43.0 (2.0–70.7)	11.0 (0.0–41.0)	0.0013
AZT exposure, g	559.6 (21.2–998.2)	158.8 (0.0–583.8)	0.0026
d4T exposure, m	26.0 (9.0–47.7)	57.0 (38.7–76.0)	<0.0001
d4T exposure, g	62.9 (21.8–109.8)	135.5 (86.7–177.2)	<0.0001
d4T exposure, mg/kg	1.15 (1.04–1.27)	1.06 (0.92–1.18)	0.0126
3TC/FTC exposure, m	68.0 (38.0–91.7)	70.0 (36.0–103.2)	0.6790
ddl exposure, m	11.0 (3.2–33.0)	17.0 (0.0–60.2)	0.2688
ddC exposure, m	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.7167
ABC exposure, m	2.0 (0.0–46.2)	0.0 (0.0–26.7)	0.0406
TDF exposure, m	0.0 (0.0–45.0)	14.0 (0.0–43.0)	0.3340
EFV exposure, m	0.0 (0.0–0.0)	2.0 (0.0–63.0)	0.0947
NVP exposure, m	6.0 (0.0–46.0)	2.0 (0.0–50.2)	0.6660
ETV exposure, m	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.3068
PI exposure, m	109.0 (28.2–168.7)	46.0 (22.7–97.5)	0.0123
NRTI exposure, m	196.0 (157.2–247.5)	224.0 (172.0–267.2)	0.0819

All parameters expressed as median and (interquartile range) unless indicated. HALS = HIV-1/HAART-associated lipodystrophy syndrome, ART = antiretroviral therapy, PI = protease inhibitor, NNRTI = non-nucleoside reverse transcriptase inhibitor, NRTI = nucleoside reverse transcriptase inhibitor, d4T = stavudine, 3TC = lamivudine, FTC = emtricitabine, TDF = tenofovir, ddl = didanosine, AZT = zidovudine, ddC = zalcitabine, ABC = abacavir, EFV = efavirenz, NVP = nevirapine, ETV = etravirine, m = months, g = grams, kg = kilograms. doi:10.1371/journal.pone.0057347.t002

time period when mitochondrially-toxic drugs were still widely used.

Acute pancreatitis has been associated with mutations in CTR and SPINK-1 genes both in the general population and in HIV-infected patients [22,23]. Among HIV-infected patients with acute clinical pancreatitis in a small study, 40% were carriers of CTR or SPINK-1 mutations [31]. Similarly, a number of gene polymorphisms have been described, associated with an increased risk of developing peripheral neuropathy associated with d4T and ddI use, including mitochondrial haplotype T [32], the mitochondrial polymorphism MTND2/HON4917G, specific to haplogroup G [32], and the 282 C>Y hemochromatosis gene mutation [33].

Table 3. Anthropometric, metabolic and fat data in the population studied.

	Cases (n = 43)	Controls (n = 129)	P value
Weight, kg	64.7 (60.0–73.7)	65.5 (58.0–73.0)	0.7542
BMI, kg/m²	23.5 (20.5–25.8)	23.8 (21.3–25.7)	0.5769
Waist circumference, cm	89.0 (81.2–93.7)	89.0 (82.7–95.0)	0.8006
WHR	0.96 (0.90–1.02)	0.95 (0.90–1.02)	0.8987
LSGS score, units	4.5 (2.0–10.0)	7.0 (3.4–11.1)	0.0386
Facial score, units	1.0 (0.0–2.0)	2.0 (0.7–2.0)	0.0222
Systolic BP, mm Hg	120 (120–138)	120 (110–130)	0.2025
Diastolic BP, mm Hg	75 (70–80)	75 (70–80)	0.6513
Metabolic syndrome, n (%)	21 (46.7)	48 (35.5)	0.2498
Total cholesterol, mmol/l	5.07 (4.18–6.03)	5.08 (4.44–5.92)	0.8966
Triglycerides, mmol/l	1.90 (1.17–3.22)	1.63 (1.12–2.54)	0.2752
HDL cholesterol, mmol/l	1.11 (0.95–1.32)	1.23 (0.97–1.53)	0.1690
LDL cholesterol, mmol/l	2.89 (2.28–3.65)	2.89 (2.46–3.50)	0.6693
VLDL cholesterol, mmol/l	0.88 (0.54–1.19)	0.77 (0.52–1.11)	0.3723
Fasting glucose, mmol/l	5.30 (5.02–5.87)	5.30 (4.90–6.00)	0.8317
HALS, n (%)	18 (41.9)	75 (58.1)	0.0932

All values expressed as median and (interquartile range) unless specified. HALS = HIV-1/HAART-associated lipodystrophy syndrome, BMI = body mass index, WHR = waist-hip ratio, LSGS = lipodystrophy severity grade score, mmol/l = millimoles per liter, HDL = high density lipoprotein, LDL = low density lipoprotein, VLDL = very low density lipoprotein, pmol/l = picomoles per liter, g = grams. doi:10.1371/journal.pone.0057347.t003

Among the factors independently associated with the development of neuropathy or pancreatitis, we found a low CD4 cell count nadir, a common marker for most toxicities and comorbidities [34,35]. Most probably it is a signal of increased inflammation and/or immune activation, which in turn contribute to overexpression of nucleoside transporters potentially leading to increased intracellular d4T-TP levels [36]. The association of TS and MTHFR polymorphisms with mitochondrial toxicity has biological plausibility, since genotypes which have a negative impact on TS activity are associated with higher d4T-TP intracellular levels and with fat redistribution, a well-known toxicity of mitochondrial origin [7,8]. Not surprisingly, we found that low expression TS genotypes are also associated with the appearance of d4T-induced mitochondrial toxicity in the form of peripheral neuropathy whereas the association for acute pancreatitis showed only a trend to statistical significance, most likely because of the small number of cases. A potentially contributing factor could have been that cases had a higher per weight exposure to d4T than controls, but again this was not an independent predictor of development of neuropathy or acute pancreatitis.

MTHFR gene is also polymorphic. The best known polymorphism consists of a 677 C→T transition in exon 4, which results in an alanine to valine substitution in the predicted catalytic domain of MTHFR. This substitution renders the enzyme thermolabile and homozygotes and heterozygotes have about 70% and 35% reduced enzyme activity, respectively [37]. A second common polymorphism in the MTHFR gene is a 1298 A→C transition in exon 7 which results in a glutamate to alanine substitution within a presumed regulatory domain of the protein [38]. The 1298C allele leads to a decreased enzyme activity, and individuals who are compound heterozygous for the 677T and

Table 4. Gene polymorphisms associated with the presence of pancreatitis and/or neuropathy in d4T-exposed patients.

	Cases (n = 43)	Controls (n = 129)	P value
TS genotype			
2R/2R, n (%)	9 (20.9)	19 (13.9)	0.0618
2R/3C, n (%)	15 (34.9)	30 (23.2)	
2R/3G, n (%)	7 (16.3)	35 (27.1)	
3C/3C, n (%)	7 (16.3)	11 (8.5)	
3C/3G, n (%)	5 (11.6)	25 (19.4)	
3G/3G, n (%)	0 (0.0)	10 (7.7)	
Low expression, n (%)	31 (72.1)	60 (46.5)	0.0046
High expression, n (%)	12 (27.9)	69 (53.5)	
MTHFR 677C→T genotype			
C/C, n (%)	21 (48.8)	42 (32.5)	0.1119
C/T, n (%)	18 (41.9)	63 (48.8)	
T/T, n (%)	4 (9.3)	24 (18.6)	
MTHFR 1298A→C genotype			
A/A, n (%)	22 (51.2)	64 (49.6)	0.0708
A/C, n (%)	14 (32.5)	58 (44.9)	
C/C, n (%)	7 (16.3)	7 (5.4)	
MTHFR genotypes and enzymatic activity			
Genotypes with increased activity*, n (%)	34 (79.1)	73 (56.6)	0.0066
Genotypes with decreased activity†, n (%)	9 (20.9)	56 (20.9)	
Combined TS and MTHFR genotypes			
Low expression and increased activity (B+A), n (%)	25 (58.1)	34 (26.3)	0.0013
Low expression and decreased activity (B+B), n (%)	6 (13.9)	26 (20.1)	
High expression and increased activity (A+A), n (%)	9 (20.9)	39 (30.2)	
High expression and decreased activity (A+B), n (%)	3 (6.9)	30 (23.2)	

TS = thymidylate synthase, MTHFR = methylene-tetrahydrofolate reductase,

*Heterozygous and wild-type patients, †Homozygous 677T, homozygous 1298C and compound heterozygous patients.

doi:10.1371/journal.pone.0057347.t004

1298C have a 40–50% reduced MTHFR activity [19]. Therefore, the non-mutated forms of MTHFR exhibit high enzymatic activity and should lead to high levels of intracellular 5,10-methylene tetrahydrofolate that will favor the formation and stability of the inhibitory ternary complex involving TS, 5,10-methylene tetrahydrofolate and deoxyuridine monophosphate [19]. Consequently, patients with non-mutated alleles should have a decreased TS activity, which in turn may lead to increased d4T-TP intracellular levels and to clinical toxicity including pancreatitis and peripheral neuropathy.

Stavudine-associated side effects may seem of only limited relevance today, because its use has greatly decreased in developed countries mainly due to its association with fat distribution abnormalities [39]. However, its use in developing countries as part of the starting antiretroviral regimes, with its associated toll of toxicities [1], is still common despite the WHO recommendation to substitute for less toxic NRTI when feasible [40].

References

- Mateo MG, Gutierrez MM, Vidal F, Domingo P (2012) Drug safety evaluation profile of stavudine plus lamivudine for HIV-1/AIDS infection. *Expert Opin Drug Saf* 11: 473–485.
- Koczor CA, Lewis W (2010) Nucleoside reverse transcriptase inhibitor toxicity and mitochondrial DNA. *Expert Opin Drug Metab Toxicol* 6: 1493–1504.
- Brinkman K, ter Hofstede HJ, Burger DM, Smeitink JA, Koopmans PP (1998) Adverse effects of reverse transcriptase inhibitors: mitochondrial toxicity as common pathway. *AIDS* 12: 1735–1744.
- Brinkman K, Smeitink JA, Romijn JA, Reiss P (1999) Mitochondrial toxicity induced by nucleoside-analogue reverse-transcriptase inhibitors is a key factor in

In summary, our study suggests that d4T-associated acute pancreatitis and/or peripheral neuropathy are associated with low-degree TS expression and MTHFR genotype associated with an increased enzymatic activity. There is a plausible pathogenic mechanism for such an association since it is well-known that, in d4T-treated patients, the presence of low-degree TS expression genotype is associated with increased d4T-TP intracellular concentrations and MTHFR genotype associated with an increased enzymatic activity contributes to a decreased TS functionality. This may be of value in tailoring d4T therapy when needed.

Author Contributions

Recruited the patients and controls: MdCC MdMG MGM EM IF ER. Conceived and designed the experiments: PD F. Vidal MB. Performed the experiments: MdCC MdMG MGM EM IF ER. Analyzed the data: PD F. Vidal MB FT. Contributed reagents/materials/analysis tools: JS MB JCD F. Villarroya. Wrote the paper: PD F. Villarroya F. Vidal FT MB.

- the pathogenesis of antiretroviral-therapy-related lipodystrophy. *Lancet* 354: 1112–1115.
5. Giralt M, Domingo P, Villarroya F (2011) Adipose tissue biology and HIV-infection. *Best Pract Res Clin Endocrinol Metab* 25: 487–499.
 6. Vela JE, Miller MD, Rhodes GR, Ray AS (2008) Effect of nucleoside and nucleotide reverse transcriptase inhibitors of HIV on endogenous nucleotide pools. *Antivir Ther* 13: 789–797.
 7. Domingo P, Cabeza MC, Pruvost A, Torres F, Salazar J, et al. (2010) Relationship between HIV/Highly active antiretroviral therapy (HAART)-associated lipodystrophy syndrome and stavudine-triphosphate intracellular levels in patients with stavudine-based antiretroviral regimens. *Clin Infect Dis* 50: 1033–1040.
 8. Domingo P, Cabeza MC, Pruvost A, Torres F, Salazar J, et al. (2011) Association of thymidylate synthase gene polymorphisms with stavudine triphosphate intracellular levels and lipodystrophy. *Antimicrob Agents Chemother* 55: 1428–1435.
 9. Moore RD, Keruly JC, Chaisson RE (2001) Incidence of pancreatitis in HIV-infected patients receiving nucleoside reverse transcriptase inhibitor drugs. *AIDS* 15: 617–620.
 10. Smith CJ, Olsen CH, Mocroft A, Viard JC, Staszewski S, et al. (2008) The role of antiretroviral therapy in the incidence of pancreatitis in HIV-positive individuals in the EuroSIDA study. *AIDS* 22: 47–56.
 11. Moore RD, Wong WM, Keruly JC, McArthur JC (2000) Incidence of neuropathy in HIV-infected patients on monotherapy versus those on combination therapy with didanosine, stavudine and hydroxyurea. *AIDS* 14: 273–278.
 12. Ellis RJ, Rosario D, Clifford DB, McArthur JC, Simpson D, et al. (2010) Continued high prevalence and adverse clinical impact of human immunodeficiency virus-associated sensory neuropathy in the era of combination antiretroviral therapy: the CHARTER Study. *Arch Neurol* 67: 552–558.
 13. Centers for Disease Control (1993) 1993 revised classification system for HIV infection and expanded surveillance for case definition for AIDS among adolescents and adults. *MMWR* 41 (RR-17): 1–13.
 14. (2001) Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 285: 2486–2497.
 15. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, et al. (2005) Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 112: 2735–2752.
 16. World Health Organization (2000) International Guide for monitoring alcohol consumption and related harm. World Health Organization.
 17. Miller SA, Dykes DD, Polesky HF (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16: 1215.
 18. Kawakami K, Watanabe G (2003) Identification and functional analysis of single nucleotide polymorphism in the tandem repeat sequence of thymidylate synthase gene. *Cancer Res* 63: 6004–6007.
 19. Marcuello E, Altés A, Menoyo A, Rio ED, Baiget M (2006) Methylenetetrahydrofolate reductase gene polymorphisms: genomic predictors of clinical response to fluoropyrimidine-based chemotherapy? *Cancer Chemother Pharmacol* 57: 835–40.
 20. Whitcomb DC (2006) Acute pancreatitis. *N Engl J Med* 354: 2142–2150.
 21. Rutschmann OT, Vernazza PL, Bucher HC, Opravil M, Ledergerber B, et al. (2000) Long-term hydroxyurea in combination with didanosine and stavudine for the treatment of HIV-1 infection. Swiss HIV Cohort Study. *AIDS* 14: 2145–2151.
 22. Cohn JA, Friedman KJ (1998) Relation between mutations of the cystic fibrosis gene and idiopathic pancreatitis. *N Engl J Med* 339: 653–658.
 23. Pfützner RH, Barmada MM (2000) SPINK1/PSTI polymorphisms act as disease modifiers in familial and idiopathic chronic pancreatitis. *Gastroenterology* 119: 615–623.
 24. Hulgan T, Haas DW (2005) Mitochondrial haplogroups and peripheral neuropathy during antiretroviral therapy: an adult AIDS Clinical Trial Group study. *AIDS* 19: 1341–1349.
 25. Riedel DJ, Gebo KA, Moore RD, Lucas GM (2008) A ten-year analysis of the incidence and risk factors for acute pancreatitis requiring hospitalization in an urban HIV clinical cohort. *AIDS Patient Care STDS* 22: 113–121.
 26. Reisler RB, Murphy RL, Redfield RR, Parker RA (2005) Incidence of pancreatitis in HIV-1-infected individuals enrolled in 20 adult AIDS clinical trials group studies: lessons learned. *J Acquir Immune Defic Syndr* 39: 159–166.
 27. Moore RD, Keruly JC, Chaisson RE (2001) Incidence of pancreatitis in HIV-infected patients receiving nucleoside reverse transcriptase inhibitor drugs. *AIDS* 15: 617–620.
 28. Smith CJ, Olsen CH, Mocroft A, EuroSIDA Study Group (2008) The role of antiretroviral therapy in the incidence of pancreatitis in HIV-positive individuals in the EuroSIDA study. *AIDS* 22: 47–56.
 29. Ghosh S, Chandran A, Jansen JP (2012) Epidemiology of HIV-Related Neuropathy: A Systematic Literature Review. *AIDS Res Hum Retroviruses* 28: 36–48.
 30. Evans SR, Ellis RJ, Chen H, Yeh T, Lee AJ, et al. (2011) Peripheral neuropathy in HIV: prevalence and risk factors. *AIDS* 25: 919–928.
 31. Felley C, Morris M (2004) The role of CFTR and SPINK-1 mutations in pancreatitis disorders in HIV-positive patients: a case-control study. *AIDS* 18: 1521–7.
 32. Canter J, Haas D (2008) The mitochondrial pharmacogenomics of haplogroup T: MTND2 (*) LHON4 917G and antiretroviral therapy-associated peripheral neuropathy. *Pharmacogenomics J* 8: 71–77.
 33. Kallianpur A, Hulgan T (2006) Hemochromatosis (HFE) gene mutations and peripheral neuropathy during antiretroviral therapy. *AIDS* 20: 1503–1513.
 34. Guaraldi G, Orlando G, Zona S, Menozzi M, Carli F, et al. (2011) Premature age-related comorbidities among HIV-infected persons compared with the general population. *Clin Infect Dis* 53: 1120–1126.
 35. Lichtenstein KA, Armon C, Buchacz K, Chmiel J, Moorman AC, et al. (2008) Initiation of Antiretroviral Therapy at CD4 Cell Counts >= 350 Cells/mm³ Does Not Increase Incidence or Risk of Peripheral Neuropathy, Anemia, or Renal Insufficiency. *J Acquir Immune Defic Syndr* 47: 27–35.
 36. Guallar JP, Cano-Soldado P, Aymerich I, Domingo JC, Alegre M, et al. (2007) Altered expression of nucleoside transporter genes (SLC28 and SLC29) in adipose tissue from HIV-1-infected patients. *Antivir Ther* 12: 853–863.
 37. Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, et al. (1995) A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet* 10: 111–113.
 38. van der Put NM, Gabreëls F, Stevens EM, Smeitink JA, Trijbels FJ, et al. (1998) A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? *Am J Hum Genet* 62: 1044–1051.
 39. Domingo P, Estrada V, Lopez Aldeguer J, Villarroya F, Martinez E (2012) Fat redistribution syndromes associated with HIV-1 infection and combination antiretroviral therapy: systematic review of current knowledge *AIDS Rev* 14: 112–123.
 40. World Health Organization (2006) Addendum to 2006 HIV Treatment Guidelines. Available: <http://www.who.int/hiv/art/ARTadultsaddendum.pdf>. Accessed 2012 May 23.