

Article

Topsensterols A–C, Cytotoxic Polyhydroxylated Sterol Derivatives from a Marine Sponge *Topsentia* sp.

Min Chen ^{1,2,†}, Xu-Dong Wu ^{1,†}, Qing Zhao ¹ and Chang-Yun Wang ^{1,*}

¹ Key Laboratory of Marine Drugs, The Ministry of Education of China, School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, China; dieying0719@163.com (M.C.); mayrush007@sina.com (X.-D.W.); ouc_zhaoqing@163.com (Q.Z.)

² Marine Science & Technology Institute, College of Environmental Science & Engineering, Yangzhou University, 196#, Huayang West Street, Yangzhou 225127, China

* Correspondence: changyun@ouc.edu.cn; Tel.: +86-532-8203-1536

† These authors contributed equally to this work.

Academic Editor: Kirsten Benkendorff

Received: 14 June 2016; Accepted: 26 July 2016; Published: 1 August 2016

Abstract: Three new polyhydroxylated sterol derivatives topsensterols A–C (1–3) have been isolated from a marine sponge *Topsentia* sp. collected from the South China Sea. Their structures were elucidated by detailed analysis of the spectroscopic data, especially the NOESY spectra. Topsensterols A–C (1–3) possess novel 2 β ,3 α ,4 β ,6 α -tetrahydroxy-14 α -methyl $\Delta^{9(11)}$ steroidal nuclei with unusual side chains. Compound 2 exhibited cytotoxicity against human gastric carcinoma cell line SGC-7901 with an IC₅₀ value of 8.0 μ M. Compound 3 displayed cytotoxicity against human erythroleukemia cell line K562 with an IC₅₀ value of 6.0 μ M.

Keywords: marine sponge; *Topsentia* sp.; polyhydroxylated sterol; topsensterol; cytotoxicity

1. Introduction

Most marine invertebrates are soft bodied and have a sedentary life style, resulting in the formation of distinctive chemical protection strategy [1,2]. The chemical means of defense necessitate the production of toxic compounds that can deter predators and paralyze their prey [2]. Undoubtedly, marine invertebrates represent a rich source of structurally novel and mechanistically unique secondary metabolites for the discovery of new drug leads [3]. Especially, sponges (Porifera) including the genus of *Topsentia* have been recognized as outstanding producers of sterols possessing novel side chains with unique alkylation and dealkylation patterns and displaying a variety of biological activities [4–7].

In our continuing efforts to discover new bioactive substances from marine invertebrates, a chemical investigation of a sponge of *Topsentia* sp. collected from the South China Sea was carried out. As a result, three new polyhydroxylated steroids, topsensterols A–C (1–3) (Figure 1), were isolated from its *n*-butanol extract. Compounds 2 and 3 showed significant cytotoxicity against human gastric carcinoma cell line SGC-7901 and human erythroleukemia cell line K562, respectively. Herein, we report the isolation, structural elucidation, plausible biogenetic pathway and biological evaluation of these steroids.




Figure 1. Chemical structures of compounds 1–3 (relative configuration).

2. Results and Discussion

The frozen sample of sponge *Topsentia* sp. (5.1 kg, wet weight) was exhaustively extracted with 95% EtOH (8 L \times 5 times) at room temperature, and the EtOH solution was evaporated under reduced pressure to give the crude extract. This extract was further partitioned between H₂O and EtOAc. After evaporation of the organic solvents, the remaining aqueous suspension was extracted with *n*-butanol. Then the organic layer was concentrated to offer the *n*-butanol extract (27.1 g). This extract was subjected to a silica gel column chromatograph (CC) and the fractions were purified by octadecylsilyl silica gel CC and semi-preparative HPLC to obtain compounds 1–3.

Topsensterol A (1) was obtained as a white amorphous powder. Its molecular formula was determined as C₃₂H₅₀O₈ on the basis of HR-ESI-MS (Supplementary Materials, Figure S9), indicating 8 degrees of unsaturation. The ¹H NMR spectrum (Table 1, Supplementary Materials, Figure S1) showed five methyl singlets including two methoxy signals (δ_{H} 3.70, 3.66, 1.34, 0.82, and 0.70) and two methyl doublets (δ_{H} 1.12 and 0.91) together with four oxygenated methine signals (δ_{H} 4.25, 4.11, 3.98, and 3.91). In addition, two olefinic signals (δ_{H} 5.86 (s), 5.32 (d, $J = 5.2$ Hz)) were observed. The ¹³C NMR spectrum (Table 2, Supplementary Materials, Figure S2) exhibited two ester carbonyls, two trisubstituted double bonds, seven methyls, seven methylenes, nine methines including four oxygenated methines, and three quaternary sp³ carbons. Therefore, a steroid nature was suggested for this molecule. The HMBC correlations (Figure 2, Supplementary Materials, Figure S5) observed from Me-30 to C-8, C-14, and C-15 demonstrated that a methyl group anchored at C-14 of the steroid nucleus. Interpretation of the COSY correlations (Figure 2, Supplementary Materials, Figure S4) from H-1 to H-8 indicated a contiguous sequence including four oxygenated methines, assigning four hydroxy groups were at C-2, C-3, C-4, and C-6, respectively. The COSY crosspeaks of H-11/H-12 and HMBC correlations from H-11 to C-8, C-10, and C-13 inferred the $\Delta^{9(11)}$ -unsaturated structure in steroid nucleus. The above spectroscopic data indicated that 1 belongs to 14 α -methyl $\Delta^{9(11)}$ -unsaturated steroids previously isolated from marine sponge, *Topsentia* sp. [8]. In addition, HMBC correlations from 26-OMe to C-26 and from 28-OMe to C-28 showed 26-OMe and 28-OMe were connected to C-26 and C-28, respectively. HMBC correlations from H-27 to C-24, C-25, C-26, and C-28, and from H-24 to C-23, C-25, and C-26 revealed the double bond was between C-25 and C-27. Consequently, a 2-substituted-dimethyl maleate unit was suggested in the side chain. Therefore the planar structure of 1 was established.

The stereochemistry of the steroid nucleus was established on the basis of coupling constants, COSY and NOESY data. Small coupling constants between H-2 and both H₂-1 suggested that H-2 adopted an equatorial orientation. A large coupling constant between H-5 and H-6 ($J = 10.8$ Hz) and a small coupling constant between H-5 and H-4 ($J = 2.4$ Hz), implied that both H-5 and H-6 were axial, and H-4 was equatorial. In the COSY spectrum, a conspicuous *W*-type long-range crosspeaks between H-2 and H-4 further confirmed both H-2 and H-4 were equatorial. While *W*-type long-range crosspeaks between H-1 β and H-3 indicated that H-3 occupied an equatorial orientation. Therefore, the hydroxyl groups 2-OH, 3-OH, 4-OH, and 6-OH occupied the axial, axial, axial, and equatorial orientation, respectively. In the NOESY spectrum (Figure 3, Supplementary Materials, Figures S6 and S7), the absence of a crosspeak between Me-19 and H-5 suggested a *trans*-fused A/B ring junction. The NOESY crosspeaks of Me-19/H-8, Me-18/H-8, Me-18/H-12 β , and H-6/H-8 indicated these

protons were β -orientation, while crosspeaks of Me-30/H-17, and Me-30/H-12 α revealed these protons were α -orientation. Furthermore, NOESY crosspeaks observed between Me-29 and H-27 suggested the double bond occupied *Z*-configuration, confirming the presence of a 2-substituted-dimethyl maleate unit in **1**.

Table 1. ^1H NMR spectroscopic data (400 MHz, CD_3OD) for **1**–**3**.

H	1	2	3
1 α	1.96 (dd, 13.0, 4.8)	2.00 (m)	1.98 (m)
1 β	2.40 (m)	2.46 (br d, 13.0)	2.46 (br d, 14.0)
2	4.25 (br s)	4.28 (br s)	4.28 (br s)
3	3.91 (t, 2.4)	3.98 (t, 2.8)	3.97 (t, 3.2)
4	3.98 (br d, 2.4)	4.04 (br d, 2.8)	4.03 (br d, 3.2)
5	1.41 (dd, 10.8, 2.0)	1.46 (dd, 10.8, 2.0)	1.45 (dd, 10.8, 2.0)
6	4.11 (td, 10.8, 4.4)	4.16 (td, 10.8, 4.4)	4.15 (td, 10.8, 4.4)
7 α	1.51 (m)	1.53 (m)	1.53 (m)
7 β	1.91 (m)	1.91 (m)	1.91 (m)
8	1.95 (m)	1.95 (m)	1.95 (m)
11	5.32 (d, 5.2)	5.36 (d, 5.2)	5.36 (br d, 5.2)
12 α	2.12 (d, 17.5)	2.15 (br d, 17.5)	2.15 (br d, 17.5)
12 β	1.93 (m)	1.93 (m)	1.93 (m)
15a	1.38 (m)	1.38 (m)	1.38 (m)
15b	1.44 (m)	1.44 (m)	1.44 (m)
16a	1.37 (m)	1.37 (m)	1.37 (m)
16b	1.92 (m)	1.92 (m)	1.92 (m)
17	1.66 (q, 9.4)	1.69 (q, 9.4)	1.68 (q, 9.4)
18	0.70 (s)	0.73 (s)	0.73 (s)
19	1.34 (s)	1.38 (s)	1.38 (s)
20	1.42 (m)	1.42 (m)	1.42 (m)
21	0.91 (d, 6.4)	0.95 (d, 6.4)	0.95 (d, 6.4)
22a	1.06 (m)	1.11 (m)	1.11 (m)
22b	1.43 (m)	1.48 (m)	1.48 (m)
23a	1.42 (m)	1.47 (m)	1.47 (m)
23b	1.54 (m)	1.55 (m)	1.55 (m)
24	2.42 (m)	2.52 (q, 6.4)	2.53 (q, 6.4)
27	5.86 (s)	7.32 (br s)	6.93 (br s)
28		4.87 (br s)	5.87 (br s)
29	1.12 (d, 6.8)	1.21 (d, 6.8)	1.20 (d, 6.8)
30	0.82 (s)	0.85 (s)	0.85 (s)
26-OCH ₃	3.70 (s)		
28-OCH ₃	3.66 (s)		3.58 (s)





Figure 2. ^1H - ^1H COSY and key HMBC correlations of compounds **1**–**3**.

Table 2. ^{13}C NMR spectroscopic data (100 MHz, CD_3OD) for 1–3.

C	1	2	3
1	39.9, CH ₂	39.7, CH ₂	39.8, CH ₂
2	71.5, CH	71.2, CH	71.3, CH
3	72.5, CH	72.0, CH	72.2, CH
4	72.8, CH	72.4, CH	72.6, CH
5	48.1, CH	48.8, CH	49.0, CH
6	66.7, CH	66.4, CH	66.5, CH
7	37.9, CH ₂	37.5, CH ₂	37.6, CH ₂
8	41.4, CH	41.0, CH	41.1, CH
9	147.9, C	147.4, C	147.5, C
10	39.8, C	39.5, C	39.5, C
11	117.0, CH	116.9, CH	116.9, CH
12	38.4, CH ₂	38.1, CH ₂	38.1, CH ₂
13	45.7, C	45.4, C	45.4, C
14	48.1, C	47.8, C	47.9, C
15	34.9, CH ₂	34.7, CH ₂	34.7, CH ₂
16	29.0, CH ₂	28.8, CH ₂	28.8, CH ₂
17	52.3, CH	51.9, CH	51.9, CH
18	15.1, CH ₃	15.1, CH ₃	15.1, CH ₃
19	27.1, CH ₃	27.0, CH ₃	27.0, CH ₃
20	37.5, CH	37.0, CH	37.1, CH
21	18.9, CH ₃	18.9, CH ₃	18.9, CH ₃
22	34.7, CH ₂	34.4, CH ₂	34.5, CH ₂
23	32.7, CH ₂	32.5, CH ₂	32.4, CH ₂
24	40.5, CH	31.8, CH	31.9, CH
25	157.6, C	139.6, C	144.2, C
26	170.7, C	176.3, C	172.8, C
27	119.4, CH	146.1, CH	143.1, CH
28	167.1, C	71.7, CH ₂	104.0, CH
29	19.3, CH ₃	18.8, CH ₃	18.8, CH ₃
30	19.0, CH ₃	18.9, CH ₃	18.9, CH ₃
26-OCH ₃	52.3, CH ₃		
28-OCH ₃	52.7, CH ₃		57.0, CH ₃


**Figure 3.** Key NOESY correlations for the sterol nucleus of compound 1.

Careful inspection of the NMR spectra of 1–3 (Tables 1 and 2) showed that the common and highly conserved signals of the steroidal nuclei for 1–3 are strikingly similar, including the signals of three angular methyl groups, a C-9(11)-double bond, four oxygenated methines, and four hydroxyl groups. Combined with their NOESY data and comparison with the configurations of the known similar metabolites such as topsentiasterol sulfates A–F, chlorotopsentiasterol sulfate D, and iodotopsentiasterol sulfate D also isolated from the marine sponge *Topsentia* sp. [7,8], compounds 1–3 were suggested to possess the same $2\beta,3\alpha,4\beta,6\alpha$ -tetrahydroxy-14 α -methyl $\Delta^{9(11)}$ sterol framework. The differences between these compounds were restricted to the respective aliphatic side chains.

Topsensterol B (2) was isolated as a white amorphous powder with a molecular formula of $C_{30}H_{46}O_6$ determined on the basis of HR-ESI-MS (Supplementary Materials, Figure S16). The COSY spectrum (Supplementary Materials, Figure S13) revealed the connectivities of $H_3-21/H-20/H_2-22/H_2-23/H-24/H_3-29$ and the terminal γ -lactone in the side chain was deduced by interpretation of HMBC data (Supplementary Materials, Figure S14) together with NMR signals (Tables 1 and 2, Supplementary Materials, Figures S10 and S11) for an oxymethylene (δ_H 4.87, δ_C 71.7), a trisubstituted double bond (δ_H 7.32, δ_C 139.6, 146.1) and an ester carbonyl (δ_C 176.3). These NMR signals also corresponded well with those reported for 2-alkyl butenolide [8].

Topsensterol C (3), with molecular formula of $C_{31}H_{48}O_7$, was also isolated as a white amorphous powder. The 1H and ^{13}C NMR signals (Tables 1 and 2, Supplementary Materials, Figures S17 and S18) of the side chain were very similar to those of 2. The only significant difference was that a methoxy (δ_H 3.58, δ_C 57.0) and an oxygenated methine (δ_H 5.87, δ_C 104.0) in 3 replaced the oxymethylene (δ_H 4.87, δ_C 71.7) in 2. Interpretation of HMBC data (Supplementary Materials, Figure S21) of 3 indicated that a 4-methoxy-2-alkyl butenolide moiety connected at the terminal of the side chain for 3.

Polyhydroxylated sterols with various novel side chains are common secondary metabolites from marine sponges. In present study, topsensterol A (1) possesses a unique side chain terminated with a 2-substituted-dimethyl maleate unit. To the best of our knowledge, it is the first report of polyhydroxylated sterol with a 2-substituted-dimethyl maleate side chain. The plausible biosynthesis mechanism to form the side chains of 1–3 was proposed in Scheme 1. According to the literature [6,7], the side chains present in 1–3 are likely formed via methylations by *S*-adenosylmethionine (SAM) to the side chain present in parkeol, along with oxidation and methoxylation. Scheme 1 illustrates that: (i) the addition of a methyl from SAM to C-24 and a loss of a proton to generate the 25,27-double bond; (ii) SAM adds a methyl to C-27 to form the C-25 cation; (iii) the C-25 cation followed by oxidation and tautomerization to form topsensterol B (2) (pathway a); and (iv) the C-25 cation followed by oxidation and methoxylation to form topsensterol C (3) and topsensterol A (1) (pathway b).



Scheme 1. Plausible biosynthesis mechanism to form the side chains of 1–3.

Marine sponges of the genus *Topsentia* were reported to produce various structurally unique steroids including polyhydroxylated sterols [9] and polysulfated steroids [10–12]. In the present study, all of the isolated polar steroids were polyhydroxylated sterols, as the desulfated derivatives of similar sterol sulfates [8]. It seems that our isolated compounds maybe a group of desulfated artifacts. However, the desulfation of polysulfated sterols should be not so easy to occur during the ordinary extraction and separation processes. To date many polysulfated sterols have been reported to be obtained from marine sponges via ordinary separation processes, indicating that polysulfated sterols are stable enough. More importantly, on the basis of biogenetic considerations, the 2 β ,3 α ,4 β ,6 α -tetrahydroxy-14 α -methyl $\Delta^9(11)$ steroidal nuclei pattern could be biosynthesized directly from parkeol in marine sponges [6,13,14]. Given the above, compounds 1–3 were more likely to be produced by ecological conditions, but the probability of desulfated artifacts could not be ruled out.

Compounds 1–3 were assessed for their cytotoxic activity against human gastric carcinoma SGC-7901 and human erythroleukemia K562 cell lines by MTT method [15]. Compound 2 exhibited cytotoxicity against SGC-7901 and K562 cell lines with IC₅₀ values of 8.0 and 20 μ M, respectively. Compound 3 exhibited cytotoxicity against SGC-7901 and K562 cell lines with IC₅₀ values of 28 and 6.0 μ M, respectively. However, compound 1 showed no cytotoxicity against the two cell lines. The above results revealed that the terminal butenolide moiety in the side chain may play a key role in the cytotoxicity.

Compounds 1–3 were also tested for their antimicrobial activity against human pathogenic bacteria including *Staphylococcus aureus* (ATCC 51650), Methicillin-resistant *Staphylococcus aureus* (ATCC 9551), and *Candida albicans* (ATCC 10231) using the method developed by Fromtling et al. [16]. However, no compound was found to be active against these bacteria.

3. Experimental Section

3.1. General Experimental Procedures

Optical rotation values were measured on a Jasco-P-1020 digital polarimeter (Jasco Corp., Tokyo, Japan). NMR spectra were recorded on a Bruker AV-400 NMR spectrometer (Bruker Corp., Fallanden, Switzerland), and chemical shifts were recorded as δ values (400 MHz for ¹H and 100 MHz for ¹³C). ESIMS spectra were obtained from a Micromass Q-TOF spectrometer (Waters Corp., Milford, MA, USA). Semipreparative HPLC was performed on a Waters 1525 system using a semipreparative C₁₈ column (5 μ m, 10 \times 250 mm, Kromasil, Sweden) coupled with a Waters 2996 photodiode array detector (Waters Corp., Milford, MA, USA). Silica gel (200–300 mesh, Qingdao Haiyang Chemical Factory, Qingdao, China), and octadecylsilyl (ODS) silica gel (45–60 mm; Merck KGaA, Darmstadt, Germany) were used for column chromatography. Thin-layer chromatography (TLC) was performed on precoated silica gel 60 GF254 plates (Yantai Zifu Chemical Group Co., Yantai, China).

3.2. Animal Material

The marine sponge *Topsentia* sp. was collected from Xuwen, Guangdong Province, China, in April 2006, and was identified by Nicole J. de Voogd, National Museum of Natural History, The Netherlands. A voucher specimen (GD-XW-20060007) was deposited at the Key Laboratory of Marine Drugs, Ministry of Education, Ocean University of China, Qingdao, China.

3.3. Extraction and Isolation

The *n*-butanol extract (27.1 g) was subjected to a silica gel column chromatograph (CC) and eluted with a gradient of CHCl₃/MeOH (50:1–0:1, *v:v*) to collect 3 fractions (Fr.A–Fr.C). Fr.A displayed cytotoxicity and the ¹H NMR features of diverse sterol analogues. This fraction was fractionated upon a octadecylsilyl silica gel CC and eluted with a gradient of H₂O/MeOH (50:1–0:1, *v:v*) to obtain three subfractions (Fr.A1–Fr.A3). Fr.A1 was subjected to semi-preparative HPLC eluted with MeOH/H₂O

(80/20, v/v) to obtain compounds **2** (4.2 mg) and **3** (6.4 mg). Fr.A2 was subjected to semi-preparative HPLC eluted with MeOH/H₂O (85/15, v/v) to yield compound **1** (2.8 mg).

Topsensterol A (**1**): white amorphous powder; $[\alpha]_D^{20} +11$ (c 0.1, MeOH); for ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 563.3600 [M + H]⁺ (calcd. for C₃₂H₅₁O₈⁺, 563.3584).

Topsensterol B (**2**): white amorphous powder; $[\alpha]_D^{20} +14$ (c 0.1, MeOH); for ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 520.3621 [M + NH₄]⁺ (calcd. for C₃₀H₅₀NO₆⁺, 520.3633).

Topsensterol C (**3**): white amorphous powder; $[\alpha]_D^{20} +17$ (c 0.1, MeOH); for ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 550.3732 [M + NH₄]⁺ (calcd. for C₃₁H₅₂NO₇⁺, 550.3738).

4. Conclusions

In summary, three new polyhydroxylated steroid derivatives topsensterols A–C (**1–3**) possessing unusual side chains were successfully isolated from the marine sponge *Topsentia* sp. The plausible biosynthesis mechanism to form the uncommon side chains of **1–3** was proposed. Compounds **2** and **3** exhibited significant cytotoxicities, demonstrating that the terminal butenolide moiety in the side chain may act an important part in the cytotoxicity.

Supplementary Materials: The following are available online at www.mdpi.com/1660-3397/14/8/146/s1, HRESIMS, 1D and 2D NMR data of compounds **1–3**.

Acknowledgments: We thank Chang-Lun Shao, School of Medicine and Pharmacy, Ocean University of China, for collection of the sponge samples. This work was supported by the National Natural Science Foundation of China (No. 41130858; 41322037; U1406402-1), and the Taishan Scholars Program, China.

Author Contributions: Min Chen and Xu-Dong Wu contributed to extraction, isolation, NMR analysis, structure elucidation and manuscript preparation; Qing Zhao contributed to bioactivities test; and Chang-Yun Wang was the project leader, organized and guided the experiments and wrote the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Jiang, C.S.; Muller, W.E.; Schroder, H.C.; Guo, Y.W. Disulfide- and polysulfide-containing metabolites from marine organisms. *Chem. Rev.* **2012**, *112*, 2179–2207. [[CrossRef](#)] [[PubMed](#)]
2. Haefner, B. Drugs from the deep: Marine natural products as drug candidates. *Drug Discov. Today* **2003**, *8*, 536–544. [[CrossRef](#)]
3. Gomes, N.G.; Dasari, R.; Chandra, S.; Kiss, R.; Kornienko, A. Marine invertebrate metabolites with anticancer activities: Solutions to the “supply problem”. *Mar. Drugs* **2016**, *14*. [[CrossRef](#)] [[PubMed](#)]
4. Aiello, A.; Fattorusso, E.; Menna, M. Steroids from sponges: Recent reports. *Steroids* **1999**, *64*, 687–714. [[CrossRef](#)]
5. Dai, J.; Sorribas, A.; Yoshida, W.Y.; Kelly, M.; Williams, P.G. Topsentinols, 24-isopropyl steroids from the marine sponge *Topsentia* sp. *J. Nat. Prod.* **2010**, *73*, 1597–1600. [[CrossRef](#)] [[PubMed](#)]
6. Whitson, E.L.; Bugni, T.S.; Chockalingam, P.S.; Concepcion, G.P.; Harper, M.K.; He, M.; Hooper, J.N.A.; Mangalindan, G.C.; Ritacco, F.; Ireland, C.M. Spheciosterol sulfates, PKC ζ inhibitors from a Philippine sponge *Spheciospongia* sp. *J. Nat. Prod.* **2008**, *71*, 1213–1217. [[CrossRef](#)] [[PubMed](#)]
7. Guzii, A.G.; Makarieva, T.N.; Denisenko, V.A.; Dmitrenok, P.S.; Burtseva, Y.V.; Krasokhin, V.B.; Stonik, V.A. Topsentiasterol sulfates with novel iodinated and chlorinated side chains from the marine sponge *Topsentia* sp. *Tetrahedron Lett.* **2008**, *49*, 7191–7193. [[CrossRef](#)]
8. Fusetani, N.; Takahashi, M.; Matsunaga, S. Topsentiasterol sulfates, antimicrobial sterol sulfates possessing novel side chains, from a marine sponge, *Topsentia* sp. *Tetrahedron* **1994**, *50*, 7765–7770. [[CrossRef](#)]
9. Luo, X.; Li, F.; Shinde, P.B.; Hong, J.; Lee, C.O.; Im, K.S.; Jung, J.H. 26,27-Cyclosterols and other polyoxygenated sterols from a marine sponge *Topsentia* sp. *J. Nat. Prod.* **2006**, *69*, 1760–1768. [[CrossRef](#)] [[PubMed](#)]
10. McKee, T.C.; Cardellina, J.H.; Tischler, M.; Snader, K.M.; Boyd, M.R. Ibisterol sulfate, a novel HIV-inhibitory sulfated sterol from the deep water sponge *Topsentia* sp. *Tetrahedron Lett.* **1993**, *34*, 389–392. [[CrossRef](#)]

11. Sun, H.H.; Gross, S.S.; Gunasekera, M.; Koehn, F.E. Weinbersterol disulfates A and B, antiviral steroid sulfates from the sponge *Petrosia weinbergi*. *Tetrahedron* **1991**, *47*, 1185–1190. [[CrossRef](#)]
12. Fusetani, N.; Matsunaga, S.; Konosu, S. Bioactive marine metabolites II. Halistanol sulfate, an antimicrobial novel steroid sulfate from the marine sponge *halichondria* cf. *moorei* bergquist. *Tetrahedron Lett.* **1981**, *22*, 1985–1988. [[CrossRef](#)]
13. Sarma, N.S.; Krishna, M.S.R.; Rao, S.R. Sterol ring system oxidation pattern in marine sponges. *Mar. Drugs* **2005**, *3*, 84–111. [[CrossRef](#)]
14. Massey, I.J.; Djerassi, C. Mass spectrometry in structural and stereochemical problems. 252. Structural and stereochemical applications of mass spectrometry in the marine sterol field. Synthesis and electron impact induced mass spectral fragmentation of Δ^{24} - and $\Delta^{24(28)}$ - 3β -hydroxy- Δ^5 -sterols. *J. Org. Chem.* **1979**, *44*, 2448–2456.
15. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55–63. [[CrossRef](#)]
16. Fromtling, R.A.; Galgiani, J.N.; Pfaller, M.A.; Espinel-Ingroff, A.; Bartizal, K.F.; Bartlett, M.S.; Body, B.A.; Frey, C.; Hall, G.; Roberts, G.D. Multicenter evaluation of a broth macrodilution antifungal susceptibility test for yeasts. *Antimicrob. Agents Chemother.* **1993**, *37*, 39–45. [[CrossRef](#)] [[PubMed](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).