


Corrigendum: *In silico* serotyping of *E. coli* from short read data identifies limited novel O-loci but extensive diversity of O:H serotype combinations within and between pathogenic lineages

Danielle J. Ingle,^{1,2,3,*} Mary Valcanis,⁴ Alex Kuzevski,⁴ Marija Tauschek,¹ Michael Inouye,^{2,5,6} Tim Stinear,¹ Myron M. Levine,⁶ Roy M. Robins-Browne^{1,7} and Kathryn E. Holt^{2,3}

Microbial Genomics 2016;2, doi: 10.1099/mgen.0.000064

There was an error in Fig. 1 in the published article. In the key, under Antimicrobial Resistance, the orange colour indicates the presence of the OXA-1 class D β -lactamase (not Carbapenemases as originally labelled).

The authors apologise for any inconvenience caused.



Received 15 February 2017; Accepted 15 February 2017

Author affiliations: ¹Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Parkville, Victoria 3010, Australia; ²Centre for Systems Genomics, University of Melbourne, Parkville, Victoria 3010, Australia; ³Department of Biochemistry and Molecular Biology, Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria 3010, Australia; ⁴Microbiological Diagnostic Unit Public Health Laboratory, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Victoria 3010, Australia; ⁵School of BioSciences, University of Melbourne, Victoria 3010, Australia; ⁶Department of Medicine, University of Maryland School of Medicine, Baltimore, MD 21201, USA; ⁷Murdoch Childrens Research Institute, Royal Children's Hospital, Victoria 3010, Australia.

*Correspondence: Danielle J. Ingle, ingled@student.unimelb.edu.au