

## VIEWPOINTS

### Vini Vidi Stenti

Christopher J. Cooper , MD; Joseph I. Shapiro , MD; Lance D. Dworkin , MD

In a recent issue of the *Journal of the American Heart Association (JAHA)*, Reinhard and colleagues presented an interesting cohort series of people enrolled at 2 Danish centers undergoing renal artery stenting.<sup>1</sup> As described by the authors, they established their registry after the reporting of the CORAL (Cardiovascular Outcomes in Renal Atherosclerotic Lesions), ASTRAL (Angioplasty and Stenting for Renal Artery Lesions), and STAR (Stenting in Renal Dysfunction Caused by Atherosclerotic Renal Artery Stenosis) randomized trials and enrolled people with  $\geq 70\%$  renal artery stenosis in predominantly 3 groups: those with hypertension not yet controlled; those with declining renal function; and, finally, a subgroup with heart failure. Of the 102 patients, 5 were not treated because of an occluded renal artery, and there were 12 procedure-related complications, including 4 significant renal artery injuries, 2 renal artery emboli, 2 respiratory events requiring intensive care, and 4 access site pseudoaneurysms. Over a median of 2 years, the serious adverse event rate was high. Ten died and 5 initiated renal replacement therapy, including 1 who started dialysis 2 days after stenting, and another 5 underwent contralateral nephrectomy for blood pressure control. Furthermore, of the 5 patients who had unsuccessful attempts at stenting, 4 died within 6 months, and the other required renal replacement therapy.

---

**See Viewpoint by Narita et al.**

---

The authors carefully measured blood pressure using 24-hour ambulatory monitoring and observed a 20-mmHg decline in systolic pressure at 3 months, which was sustained to the 24-month follow-up period. Also observed was a reduction in the number of antihypertensive medications. The use and type of antihypertensive medications was not specified before and after stenting, and there was a relatively low use of renin-angiotensin system antihypertensives.

The current study should be viewed in the context of a long history of investigation in the treatment of people with atherosclerotic renal artery stenosis. Some 90 years ago, Goldblatt<sup>2</sup> observed that partial ligation of a renal artery resulted in increasing blood pressure. Later, it was determined that activation of the renin-angiotensin system was important in the pathogenesis of hypertension, and this was subsequently augmented by work demonstrating the importance of other mechanisms including sympathetic activations.<sup>3-5</sup>

Importantly, there is a mature body of work evaluating surgical renal artery bypass,<sup>6</sup> renal artery angioplasty,<sup>7</sup> and nephrectomy,<sup>8</sup> with limitations of these therapies including high surgical mortality,<sup>9</sup> restenosis,<sup>10</sup> and loss of kidney function for each of these treatments, respectively. With the advent of the Palmaz balloon-expandable stent, there emerged a percutaneous treatment that was highly effective in opening an obstructed renal artery and had lower rates of restenosis. In 1998, Gerald Dorros and colleagues published their center's experience with 163 people with atherosclerotic renal artery stenosis treated with the

**Key Words:** ischemic nephropathy ■ nephrectomy ■ renal artery stenosis ■ renovascular hypertension ■ stent

---

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

Correspondence to: Christopher J. Cooper, MD, College of Medicine and Life Sciences, University of Toledo, 3000 Arlington Ave, MS 1018, Toledo, OH 43614. Email: [christopher.cooper@utoledo.edu](mailto:christopher.cooper@utoledo.edu)

For Disclosures, see page 3.

© 2022 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: [www.ahajournals.org/journal/jaha](http://www.ahajournals.org/journal/jaha)

Palmaz stent.<sup>11</sup> Dorros's result was a heady affirmation of what we believed: find renal artery stenosis, stent renal artery stenosis, improve blood pressure and kidney function, improve clinical outcomes.

At the center of this belief was that (1) the improvements in blood pressure after stenting were caused by stent treatment; (2) medical therapies, while important, were insufficient to achieve this outcome; and (3) a reduction in blood pressure or improvement in estimated glomerular filtration rate would necessarily translate into improved clinical outcomes. In many of these observational studies, we noted that preprocedural rapid declines in renal function were associated with stabilization or even improved renal function in follow-up, and elevated blood pressures also improved. What we did not contemplate included phenomena such as regression to the mean where blood pressure and kidney function vary over time and high values tend to regress toward normal values simply because of variation and not because of treatment, and that concomitant attention to effective medical treatment also resulted in improved blood pressure and stabilized kidney function.

Importantly, there have been a number of high-quality cohort studies that followed Dorros's seminal publication. These are generally quite similar: average age 70; half being women; visual estimates of "severe" renal artery stenosis averaging  $\approx 80\%$ ; modest numbers of people with stage 4+ chronic kidney disease; a subset of people with heart failure, with some having pulmonary edema. With the large number of observational studies demonstrating impressive reductions in blood pressure and "stabilization" of kidney function, it seems quite obvious that if we come to the interventional suite and see "severe" renal artery stenosis, we should treat it; *vini vidi stenti*, or I came, I saw, I stented. However, the best evidence, the published randomized trials, fails to support this approach.

To Reinhard's credit, in the current cohort series, they have enriched the proportion of several high-risk subgroups including people on dialysis at enrollment, those with relatively high blood pressures at baseline, and those with baseline heart failure or pulmonary edema. Reinhard and colleagues contend that the people in their cohort study were different from those in randomized trials, which is partly true. Specifically, people hospitalized with acute heart failure and those on hemodialysis were excluded from the randomized studies. The organizers of the randomized trials did so because of the challenges of disentangling clinical events that were actively occurring that were also study end points, that is, adjudicating the occurrence of kidney disease requiring renal replacement therapy in a person being treated with dialysis and adjudicating heart failure when the person is in heart failure at the time of treatment. Consequently, these small subsets of patients are not represented in the randomized studies, and clinicians need better data to sort out

when to treat these critically ill patients since the risks and rewards are significant.

The critical question that the current study is unable to address is whether these patients would have had similar or even better outcomes had the study group simply treated them with aggressive and optimal medical therapy. What we know is that this high-risk group had some clinical improvements, such that 2 of 4 on dialysis no longer required dialysis. What we also know is that the other 2 on dialysis at the time of enrollment did not recover renal function; an additional 5 patients ended up on dialysis, including 1 who initiated treatment 2 days after stenting; 4 of 5 with failed revascularization required dialysis; and 10 died within 2 years. In the era of surgical revascularization, Hallett<sup>12</sup> observed that  $\approx 1$  in 4 people with azotemia preoperatively improved kidney function, half were stable, and 1 in 4 worsened after surgery. As a clinician in a conversation with a very ill patient, is an attempt at revascularization warranted, knowing that the rewards are potentially significant and the risks are also high?

To underscore this point, CORAL enrolled 364 patients with stenoses  $>80\%$ , and there was no difference in clinical outcome between medical therapy alone and medical therapy with stent revascularization.<sup>13</sup> Similarly, although systolic blood pressure was predictive of clinical outcomes, stent treatment did not modify the effect, including in those with systolic blood pressures  $>180$  mmHg at study entry.<sup>14</sup> Similarly, the presence of global renal ischemia, defined as bilateral stenosis or a stenosis in a solitary functioning kidney, which was documented in 140 CORAL patients, was not associated with a difference in clinical outcomes. Of note, these subgroups are larger than many of the observational studies including the current Reinhard cohort.

Stenting to improve blood pressure has a limited role when clinicians are committed to assuring that their patients receive optimal medical therapy. Effective medical therapy is critically important since renal artery stenosis occurs in the setting of systemic risk factors and systemic atherosclerosis, which is oftentimes widespread. Optimal medical therapy includes aggressive medical therapy to define blood pressure thresholds using long-acting renin-angiotensin blockade and calcium antagonists with excellent trough-to-peak blood pressure ratios, intensive lipid lowering, smoking cessation, and antiplatelet therapy. In CORAL, stent treatment did lead to statistically significantly better blood pressures than medical therapy alone; however, the difference was modest (2 mmHg better) and was not associated with fewer adverse cardiovascular or renal events.<sup>13</sup>

There are some people who will experience an increase in serum creatinine with effective blood pressure control (with or without renin-angiotensin inhibition); however, this should not be mistaken for "kidney failure" and most often does not necessitate cessation of treatment. Reinhard reported that 41% of people enrolled had

renin-angiotensin inhibition treatment discontinued because of a  $\geq 30\%$  increase in plasma creatinine. People treated with medical therapy alone who have an acute rise in creatinine often improve renal function absent any intervention, and importantly, their clinical outcomes are not impacted.<sup>15</sup> This phenomenon is so common that many studies of progression of chronic kidney disease exclude changes observed in the first few months of therapy and calculate the real effect of therapy on glomerular filtration rate beginning at  $\geq 3$  months after blood pressure is controlled.<sup>16</sup> Consequently, if a patient's creatinine rises from 1.0 to 1.3 mg/dL, or from 2.0 to 2.6 mg/dL with effective blood pressure control, the majority will do quite well without changing medical treatment and will improve or stabilize kidney function without revascularization.

Reinhard and colleagues suggest in their concluding remarks about the potential value of a randomized trial and highlight the difficulties of conducting such a study. As clinicians who are invested in providing the best possible care for our patients, additional cohort studies are unlikely to improve the situation. The core challenge is that high-risk patients have the potential for high-yield outcomes, such as the 2 patients who no longer required renal replacement therapy after stenting, and are at high risk of serious or fatal outcomes, such as seen with the current study.

What should busy clinicians remember? The vast majority of people with atherosclerotic renal artery stenosis need to be treated with aggressive and optimal medical therapy alone and do not benefit from stenting: *vini vidi optimus medicamentum*. There remains a modest group of people who were not represented in the randomized trials such as those with stage 5 or end-stage kidney disease and acute heart failure, for which difficult conversations must be had about the potential benefits and risks of revascularization. Finally, blood pressure control is rarely a reason to implant stents in renal arteries when we commit to ensuring optimal medical therapy, understand that an increase in creatinine during optimization of blood pressure control does occur, and in most people does not necessitate revascularization. Exceptional cases in which medical therapy fails to achieve blood pressure control should be evaluated by clinicians experienced with managing patients with complicated renal artery stenosis. When revascularization is performed, invariably these patients require ongoing medical therapy to achieve and maintain optimal blood pressure control.

## ARTICLE INFORMATION

### Affiliations

University of Toledo, OH (C.J.C., L.D.D.); and Marshall University, Huntington, WV (J.I.S.).

## Disclosures

None.

## REFERENCES

- Reinhard M, Schousboe K, Andersens U, Buus NH, Rantanen JM, Bech JN, Mafi HM, Langfeldt S, Bharadwaz A, Hørlyck A, et al. Renal artery stenting in consecutive high-risk patients with atherosclerotic renovascular disease: a prospective 2-center cohort study. *J Am Heart Assoc*. 2022;11:e024421. doi: [10.1161/JAHA.121.024421](https://doi.org/10.1161/JAHA.121.024421)
- Goldblatt H, Lynch J, Hanzal RF, Summerville WW. Studies on experimental hypertension; production of persistent elevation of systolic blood pressure by means of renal ischemia. *J Exp Med*. 1934;59:347–379. doi: [10.1084/jem.59.3.347](https://doi.org/10.1084/jem.59.3.347)
- Johansson M, Elam M, Rundqvist B, Eisenhofer G, Herlitz H, Friberg P. Increased sympathetic nerve activity in renovascular hypertension. *Circulation*. 1999;99:2537–2542. doi: [10.1161/01.cir.99.19.2537](https://doi.org/10.1161/01.cir.99.19.2537)
- Miyajima E, Yamada Y, Yoshida Y, Matsukawa T, Shionoiri H, Tochikubo O, Ishii M. Muscle sympathetic nerve activity in renovascular hypertension and primary aldosteronism. *Hypertension*. 1991;17:1057–1062. doi: [10.1161/01.hyp.17.6.1057](https://doi.org/10.1161/01.hyp.17.6.1057)
- Mathias CJ, Kooner JS, Peart S. Neurogenic components of hypertension in human renal artery stenosis. *Clin Exp Hypertens A*. 1987;9(suppl 1):293–306. doi: [10.3109/10641968709160180](https://doi.org/10.3109/10641968709160180)
- Hunt JC, Sheps SG, Harrison EG, Strong CG, Bernatz PE. Renal and renovascular hypertension. A reasoned approach to diagnosis and management. *Arch Intern Med*. 1974;133:988–999.
- van Jaarsveld BC, Krijnen P, Pieterman H, Derckx FHM, Deinum J, Postma CT, Dees A, Woittiez AJJ, Bartelink AKM, Velv AJM, et al. The effect of balloon angioplasty on hypertension in atherosclerotic renal-artery stenosis. *N Engl J Med*. 2000;342:1007–1014. doi: [10.1056/NEJM200004063421403](https://doi.org/10.1056/NEJM200004063421403)
- Oskin TC, Hansen KJ, Deitch JS, Craen TE, Dean RH. Chronic renal artery occlusion: nephrectomy versus revascularization. *J Vasc Surg*. 1999;29:140–149. doi: [10.1016/s0741-5214\(99\)70355-1](https://doi.org/10.1016/s0741-5214(99)70355-1)
- Modrall JG, Rosero E, Smith ST, Arko FR, Valentine RJ, Clagett GP, Timaran CH. Operative mortality for renal artery bypass in the United States: results from the National Inpatient Sample. *J Vasc Surg*. 2008;48:317–322. doi: [10.1016/j.jvs.2008.03.014](https://doi.org/10.1016/j.jvs.2008.03.014)
- van de Ven PJ, Kaatee R, Beutler JJ, Beek FJ, Woittiez AJ, Buskens E, Koomans HA, Mali WP. Arterial stenting and balloon angioplasty in ostial atherosclerotic renovascular disease: a randomised trial. *Lancet*. 1999;353:282–286. doi: [10.1016/S0140-6736\(98\)04432-8](https://doi.org/10.1016/S0140-6736(98)04432-8)
- Dorros G, Jaff M, Mathiak L, Dorros II, Lowe A, Murphy K, He T. Four-year follow-up of Palmaz-Schatz stent revascularization as treatment for atherosclerotic renal artery stenosis. *Circulation*. 1998;98:642–647. doi: [10.1161/01.cir.98.7.642](https://doi.org/10.1161/01.cir.98.7.642)
- Hallett JW, Textor SC, Kos PB, Nicpon G, Bower TC, Cherry KJ, Gloviczki P, Pairolero PC. Advanced renovascular hypertension and renal insufficiency: trends in medical comorbidity and surgical approach from 1970 to 1993. *J Vasc Surg*. 1995;21:750–759. doi: [10.1016/s0741-5214\(05\)80006-0](https://doi.org/10.1016/s0741-5214(05)80006-0)
- Cooper CJ, Murphy TP, Cutlip DE, Jamerson K, Henrich W, Reid DM, Cohen DJ, Matsumoto AH, Steffes M, Jaff MR, et al. Stenting and medical therapy for atherosclerotic renal-artery stenosis. *N Engl J Med*. 2014;370:13–22. doi: [10.1056/NEJMoa1310753](https://doi.org/10.1056/NEJMoa1310753)
- Murphy T, Cooper C, Matsumoto A, Cutlip DE, Pencina KM, Jamerson K, Tuttle KR, Shapiro JI, D'Agostino R, Massaro J, et al. Renal artery stent outcomes. *J Am Coll Cardiol*. 2015;66:2487–2494. doi: [10.1016/j.jacc.2015.09.073](https://doi.org/10.1016/j.jacc.2015.09.073)
- Cooper EL, Xie Y, Nguyen H, Brewster PS, Sholl H, Sharrett M, Ren K, Chen T, Tuttle KR, Haller ST, et al. Early rapid decline in kidney function in medically managed patients with atherosclerotic renal artery stenosis. *J Am Heart Assoc*. 2019;8:e012366. doi: [10.1161/JAHA.119.012366](https://doi.org/10.1161/JAHA.119.012366)
- Levey AS, Inker LA, Matsushita K, Greene T, Willis K, Lewis E, de Zeeuw D, Cheung AK, Coresh J. GFR decline as an end point for clinical trials in CKD: a scientific workshop sponsored by the National Kidney Foundation and the US Food and Drug Administration. *Am J Kidney Dis*. 2014;64:821–835. doi: [10.1053/j.ajkd.2014.07.030](https://doi.org/10.1053/j.ajkd.2014.07.030)