

Case report—right atrial mass: a very rare presentation of endometrial cancer metastasis

Emre Erturk ^{1*}, Onur Soyler ², Fatma Seher Pehlivan ³, and Cagatay Arslan ⁴

¹Department of Cardiology, Izmir University of Economics Medicalpoint Hospital, Imbatli Mah., 1825 sokak, No:12, Izmir 35575, Turkey; ²Department of Cardiovascular Surgery, Izmir University of Economics Medicalpoint Hospital, Izmir, Turkey; ³Mikro Private Pathology Laboratory, Izmir, Turkey; and ⁴Department of Medical Oncology, Izmir University of Economics Medicalpoint Hospital, Izmir, Turkey

Received 15 August 2022; revised 29 August 2023; accepted 11 September 2023; online publish-ahead-of-print 14 September 2023

Background

We report a case of a 47-year-old woman with right atrial metastasis of endometrioid adenocarcinoma, which is an uncommon clinical presentation for patients with endometrial cancer (EC). The principal aim of this case is to demonstrate the possibility of distant metastasis, something rarely encountered among this group of patients.

Case summary

Our patient, diagnosed with EC and receiving chemotherapy and radiotherapy after surgery, was found to have enhanced 18-fluorodeoxyglucose uptake inside the right atrium on the repeat positron emission tomography–computed tomography scan at the ninth month after initial diagnosis. Following trans-oesophageal echocardiography, cardiac magnetic resonance imaging showed a hyper-vascular mass with right atrial lateral wall involvement likely to be malignant in nature. A right atrial tumour was successfully removed by cardiovascular surgeons, and a pericardial patch was placed at the site of the excised atrium. The pathological examination showed EC metastasis. Following surgery, systemic treatment was planned for recurrent EC. The patient had an uneventful recovery after the surgery.

Discussion

Endometrial cancer is the most common gynaecologic malignancy and the fourth most common cancer in women. The lymphatic pathway is the main metastatic behaviour of EC; however, haematogenous metastases are not uncommon, especially in patients with higher stages of the disease. Our patient did not show any signs and symptoms of cardiac involvement. Nevertheless, clinicians should be alert for symptoms of cardiac involvement like new-onset murmur, embolism, or dyspnoea. Having known the behavioural pattern of the primary tumour, timely utilization of diagnostic imaging methods in accordance with clinical suspicions in patients with rapidly growing tumours can be lifesaving.

Keywords

Cardiac tumours • Cardiac MRI (magnetic resonance imaging) • Endometrial carcinoma • Case report

ESC curriculum

2.1 Imaging modalities • 2.2 Echocardiography • 2.3 Cardiac magnetic resonance • 2.5 Nuclear techniques • 6.8 Cardiac tumours

* Corresponding author. Tel: +90 505 696 1545, Email: emerturk@gmail.com; emre.erturk@ieu.edu.tr

Handling Editor: Luca Arcari

Peer-reviewers: Maria Mattioli; Melissa Yen Moey

Compliance Editor: Polyvios Demetriades

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Learning points

- This case is educative proof of the fact that malignancies that are quite susceptible to distant metastasis, as we have witnessed, should be evaluated thoroughly.
- Susceptibility can be surmised on observing rapid growth rate, when considered alongside a locally advanced stage at the initial diagnosis. Although routine computed tomography scans might not succeed in detecting metastatic foci, a positron emission tomography–computed tomography scan could prove particularly beneficial for such diagnostic approaches.
- Distant organ malignancies might metastasize to the heart; thus, cardiac involvement should not be overlooked during the management of these patients, even when the patients are asymptomatic.
- Multi-modality imaging may help with detection and visualization of rare foci and therefore should be implemented during routine follow-up of oncology patients with certain malignancies, especially those shown to have frequent distant organ metastasis. Detection of rare metastasis sites can guide the treatment of patients.
- A multi-disciplinary approach for oncology patients is shown to be quite beneficial, and treatment and follow-up decisions should be taken by a designated team. When necessary, this group of patients should be referred to tertiary care centres where highly organized oncology groups are present.

Introduction

Endometrial cancer (EC) is the most common gynaecologic malignancy. It is the fourth most common cancer among women in the USA after breast, lung, and colorectal cancers. Risk factors are related to excessive unopposed exposure of the endometrium to oestrogen.¹ The most frequently occurring histological sub-type is endometrioid adenocarcinoma.² Higher-stage endometrioid carcinoma, non-endometrioid histology, molecular factors like p53 mutation, and stage of the disease are predictors of recurrence, and adjuvant treatments are based on these prognostic variables. Lymph node metastasis is the main metastatic pathway of EC. The most common metastasis sites of EC are pelvic lymph nodes, such as common iliac, internal iliac, and external iliac lymph nodes, followed by retroperitoneal lymph nodes.³ Studies have found that ~5% of patients suffered at least one site of distant metastases. The most common metastasis sites are the lung followed by distant lymph nodes, liver, bone, and brain.⁴ In literature, the heart is one of the rare distant sites found to have metastasis from EC^{5,6} (Table 1). Although primary cardiac tumours are extremely uncommon, the heart can have metastases due to any malignant neoplasm which is able to spread to distant sites. Cardiac metastases incidence seems to range from 2.3% to 18.3%.¹⁴ Any type of tumour can affect the heart. The probability of cardiac involvement is multi-factorial and depends on anatomic considerations, stage of disease, individual tumour, and host biology. Primary lung cancer represents 36–39% of cardiac metastases, followed by breast cancer (10–12%) and haematologic malignancies (10–21%). The pericardium is the most frequently involved site of cardiac metastasis followed by epicardium, myocardium, endocardium, and intracavitary metastasis, respectively.¹⁵ Here, we report a case of intracavitary metastasis of EC, which is an unusual presentation.

Summary figure

Day 0	Initial presentation covering symptoms and endometrial carcinoma diagnosis
During the first month following initial diagnosis	CT scan showing enlarged para-aortic lymph nodes followed by PET-CT scan for confirmation of metastasis. PET-CT scan showed para-aortic metastasis
1st month following initial diagnosis	Surgical intervention following PET-CT scan and staging of the disease. Initiation of chemotherapy
9th month following initial diagnosis	Repeat CT scan showing conglomerated para-aortic lymph nodes followed by PET-CT scan for confirmation of metastasis. PET-CT scan showed para-aortic metastasis along with right atrial involvement.
9th month following initial diagnosis	TTE, TOE, and MRI for diagnosing and probing the right atrial mass further before surgery
10th month following initial diagnosis	Surgical removal of right atrial mass. Pathological diagnostic confirmation

Case presentation

A 46-year-old-female who has been on medication for hypertension and Type 2 diabetes was admitted with ongoing abdominal pain. The patient was diagnosed with endometrial adenocarcinoma Stage III following surgical procedure and pathological assessment. Besides endometrium, left adnexal and uterine tube and para-aortic lymph node involvements were detected. Six cycles of adjuvant paclitaxel and carboplatin were administered followed by adjuvant radiotherapy (RT). After 3 months from the end of adjuvant RT, a control thoracic and abdominal computerized tomography (CT) was done. Para-aortic lymph node metastasis was detected as a result of the abdominal CT scan. Regarding surgical removal planning of the para-aortic lymph nodes, an 18-fluorodeoxyglucose (FDG) positron emission tomography CT scan was done to re-stage the disease for any other metastases on the 9th month after initial diagnosis. It revealed a mass and enhanced glucose uptake in the right atrium of the heart and also metastases at the para-aortic lymph nodes (Figure 1).

On initial examination, there were no distinctive cardiac exam findings. The electrocardiogram was sinus rhythm, with normal rate, PR (140 ms), QT intervals (460 ms), and QRS (100 ms) width. Trans-thoracic echocardiography showed an irregular mass inside the right atrium, which was hyper-echoic, irregular, and attached to the lateral wall. The mobility of the mass was limited, and there was no tricuspid valve involvement with trivial tricuspid regurgitation.

Trans-oesophageal echocardiography (TOE) was planned to obtain more precise images and to establish any vascular or valvular connections with the tumour but came back with no particular involvement. The best images were obtained in bi-caval view which showed the tumour was attached to the lateral wall with a wide base. The longest diameters were 3.3 × 2.3 cm wide. Neither superior nor inferior caval veins were affected by the tumoural body. Following TOE, cardiac magnetic

Table 1 Endometrial carcinoma cases with cardiac metastasis

Endometrial carcinoma cases in the literature with metastasis to the heart				
No	Case	Author	Year	Reference no
1	Right ventricular metastasis of undifferentiated endometrial cancer	Liu T. <i>et al.</i>	2014	6
2	Endometrial adenocarcinoma with pericardium metastasis	Liu Guang <i>et al.</i>	2019	7
3	Right ventricular metastasis of poorly differentiated mixed endometrial and clear cell carcinoma	Glenn E. Bigsby <i>et al.</i>	2004	8
4	Endometrial adenocarcinoma metastasis to the right ventricle	Mario Castillo-Sang <i>et al.</i>	2007	9
5	Right atrial metastasis of endometrial adenocarcinoma	Khalil Kanjwal <i>et al.</i>	2010	10
6	Endometrial adenocarcinoma with right ventricular metastasis	Marcos Danillo P. Oliveira <i>et al.</i>	2019	5
7	Endometrial carcinoma metastasis to the right atrium and right ventricle	Masi Javeed <i>et al.</i>	2022	11
8	Right atrial metastasis of endometrial adenocarcinoma	Cipullo I <i>et al.</i>	2021	12
9	Endometrial carcinoma metastasis to the right ventricle, mimicking acute pulmonary embolism	Rajesh Kunadharaju <i>et al.</i>	2021	13




Figure 1 Positron emission tomography–computed tomography images: upper row-1st month of diagnosis showing para-aortic lymph node involvement and no cardiac uptake, lower row-9th month of diagnosis – enhanced glucose uptake in right atrial zone (arrows). PET–CT, positron emission tomography–computed tomography.

resonance imaging (MRI) was planned to detect the nature, vascularization, and local invasion of the lesion and to guide surgery (Figure 2).

Cardiac MRI showed a hyper-vascular mass with right atrial lateral wall involvement which was more likely to be malignant in nature (Figure 3).

A three-stage treatment was planned for the patient. Firstly, surgical removal of the cardiac metastasis was conducted. Systemic treatment

(chemotherapy) was planned after the surgery, and a second surgery was planned for the para-aortic lymph node metastasis after systemic treatment. The tumour inside the right atrium was successfully removed by cardiovascular surgeons, and a pericardial patch was placed at the site of the excised atrium (Figure 4). The patient had no complications and had an uneventful recovery and was discharged home on the seventh day following the surgery.




Figure 2 Trans-oesophageal echocardiography showing inter-atrial septum and the mass in the right atrium. This is a modified bi-caval view showing the tumour has no relationship with inter-atrial septum and inferior and superior caval veins. It is an irregularly shaped, dense tumour which is 3.3 × 2.3 cm in size.




Figure 3 Cardiac magnetic resonance imaging showing right atrial mass (arrows). Contrast uptake on the second image is an indicator of hyper-vascularity of tumour tissue. RV, right ventricle; LV, left ventricle; PA, pulmonary artery.



Figure 4 Tumour tissue removed from the right atrium.




Figure 5 (A) Adenocarcinoma metastasis to endocardium (HE [hematoxylin and eosin] $\times 40$), (B) adenocarcinoma infiltration to endocardium, predominantly lymphoid cells (HE $\times 40$), (C) poorly differentiated adenocarcinoma containing lymphoid cell-rich stroma (HE $\times 100$), (D) pancytokeratin positive tumour cells (DAB $\times 40$), (E) Pax8 nuclear positive tumour cells (DAB $\times 100$), and (F) focal and weak ER positive tumour cells (DAB $\times 100$).

The pathological examination showed EC metastasis. Diaminobenzidine (3,3'-DAB)-staining revealed pancytokeratin and Pax8 nuclear positive cells with focal and weak oestrogen receptor positivity (Figure 5). Following surgery, the patient was put on paclitaxel and carboplatin plus pembrolizumab treatment, while tumour tissue was positive for programmed cell death ligand 1 (PDL-1) staining. The patient will be planned for surgical resection of para-aortic lymph nodes if they are found to be resectable following the fourth cycle of systemic treatment.

Discussion

Endometrial cancer is the most common gynaecologic malignancy and the fourth most common cancer in women. Studies have shown that cardiac metastasis from EC is a very rare condition. Metastatic endometrial carcinoma cases in the literature, which include the heart, are shown in Table 1. Right ventricular metastasis has been the most common cardiac metastasis site. We reported a case of intracavitary metastasis of EC, which is an unusual presentation and should be taken into consideration for patients with malignancies displaying rapid growth rate and higher metastatic potential.

Our patient had an intracavitary cardiac metastasis, which is, by any means, an extremely rare spreading pattern of EC to a distant organ. If the 18-FDG positron emission tomography-computed tomography (PET-CT) had not been done, cardiac metastasis would have been missed, and an inadequate treatment would have been carried out.

Recent cardio-oncology guideline by European Society of Cardiology (ESC) underscored the importance of serial evaluation of oncology patients on certain treatment protocols. The primary cardiovascular concern with patients undergoing chemotherapy protocols is myocardial dysfunction which can be induced by both agents given to our patient (paclitaxel and carboplatin). Cisplatin was also found to be correlated with coronary artery disease, hypertension, and thrombosis. However, carboplatin has less cardiovascular toxicity. Cardiovascular risk factor monitoring is recommended during and after therapy for such patients.¹⁶ However, even in the most recent documents such as this guideline, there is no definitive recommendation regarding the optimal timing for routine metastasis checks for patient sub-groups.

This case is a crucial demonstration of timely utilization of diagnostic imaging methods in accordance with clinical suspicion in patients with rapidly growing tumours. The management of this patient might raise the question of the necessity of having every single oncology patient meticulously checked for distant metastasis. Although the literature and experience do not provide a precise answer to the issue of meticulously checking every single oncology patient for distant metastasis, certain characteristics of individual patients and malignancies should be used for decision-making. Haematogenous spreading to unexpected organs and the possibility of rare metastatic foci should always be taken into account when assessing and staging these malignancies and giving treatment decisions. In our case, findings on the initial PET CT scan indicated the rapid spreading rate and high possibility of metastasis which necessitated more frequent follow-up and subsequently another PET-CT scan at 9 months, the diagnosis being made upon these complementary imaging techniques. Considering the growth rate of the tumour tissue, our patient would likely have been symptomatic due to the gradual cessation of blood flow through the right atrium or embolic events unless it was diagnosed. The multi-disciplinary approach along with multimodality imaging methods has been quite beneficial in the management of this rare case. Incorporating such patients to the literature may invoke awareness on the intensity of investigations

that should be implemented on the care of patients with certain disease characteristics.

Lead author biography



Emre Erturk, MD, who is affiliated with IEU Medicalpoint Hospital and specializes in the Department of Cardiology, completed his medical school education at Dokuz Eylul University Faculty of Medicine and pursued his residency at Cerrahpasa Medical School in the Department of Cardiology. His primary areas of expertise include interventional cardiology, heart failure, and cardiac imaging. Dr. Erturk is an active member of several professional organizations, including the European Society of Cardiology, European Heart Failure Association, EAPCI, and Turkish Society of Cardiology.

Acknowledgements

MRI acquisition, Mohammed Abusharekh, MD; surgical intervention, Gokhan Albayrak MD and Koray Aykut MD.

Consent: The authors confirm that written consent for the submission and publication of this case, including images, has been obtained from the patient and his parents in line with COPE guidance.

Conflict of interest: None declared.

Funding: None declared.

Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

References

- Braun MM, Overbeek-Wager EA, Grumbo RJ. Diagnosis and management of endometrial cancer. *Am Fam Physician* 2016;**93**:468–474.
- Morice P, Leary A, Creutzberg C, Abu-Rustum N, Darai E. Endometrial cancer. *Lancet* 2016;**387**:1094–1108.
- Mariani A, Webb MJ, Keeney GL, Podratz KC. Routes of lymphatic spread: a study of 112 consecutive patients with endometrial cancer. *Gynecol Oncol* 2001;**81**:100–104.
- Li J, Sun L, Zhang Y, Cai S. Patterns of distant metastases in patients with endometrial carcinoma: a SEER population-based analysis. *J Clin Oncol* 2019;**37**:e17109.
- Oliveira MDP, Florenzano MT, Santos LFC, Navarro PLB, Filho RCS. Cardiac metastasis secondary to endometrial cancer: an extremely rare presentation. *Arq Bras Cardiol Imagem cardiovasc* 2019;**32**:138–140.
- Liu T, Khan S, Behr S, Aparici CM. Diagnosis of cardiac metastasis from endometrial cancer by F-18 FDG-PET/CT. *Nucl Med Mol Imaging* 2014;**48**:237–240.
- Liu G, Zhang Q, Li Z, Chen X, Zhang N, Zhang J. Endometrial carcinoma complicated by malignant pericardial effusion: a case report on the therapeutic regimen. *Medicine (Baltimore)* 2019;**98**:e17584.
- Biggsby GE IV, Holloway RW, Weppelman B, Reynolds RB, Williams B. Endometrioid adenocarcinoma of the uterus with cardiac metastasis. A case report and six-year follow-up. *Gynecol Oncol* 2005;**97**:256–259.
- Castillo-Sang M, Slam K, Gociman B, Durham SJ, Booth R. Endometrial adenocarcinoma metastatic to the right ventricle: a case report and review of the literature. *Cardiovasc Pathol* 2009;**18**:178–182.
- Kanjwal K, Colyer W Jr. Transthoracic echocardiography-guided biopsy of a metastatic endometrial adenocarcinoma in the right atrium: a review of diagnosis and treatment of cardiac masses. *Am J Ther* 2010;**17**:e118–e125.
- Javeed M, Ravuri R, Javeed Z, Taylor S, Akel R. A rare case of intracavitary cardiac metastasis of endometrial carcinosarcoma. *Cureus* 2022;**14**:e25583.
- Cipullo I, Campisi P, Centofanti P, Musumeci G, Cirillo S, Sanjinez Pesua JOS, et al. 1089 Cardiac metastasis of endometrial cancer: a multidisciplinary approach. *Int J Gynecol Cancer* 2021;**31**:A137–A138.

13. Kunadharaju R, Bhura S, Mengiste H, Natalzia P, Al Ghoul Y, Sethi S. Metastatic endometrial carcinoma mimicking acute pulmonary embolism with clot in transit. *Chest* 2021; **160**:A2196–A2197.
14. Bussani R, De-Giorgio F, Abbate A, Silvestri F. Cardiac metastases. *J Clin Pathol* 2007;**60**:27–34.
15. Goldberg AD, Blankstein R, Padera RF. Tumors metastatic to the heart. *Circulation* 2013; **128**:1790–1794.
16. Lyon AR, López-Fernández T, Couch LS, Asteggiano R, Aznar MC, Bergler-Klein J, et al. 2022 ESC guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS): developed by the task force on cardio-oncology of the European Society of Cardiology (ESC). *Eur Heart J* 2022;**43**:4229–4361.