



Intervention Programs Targeting Burnout in Health Professionals: A Systematic Review

**Diana Araújo*¹, *Ana Bárto*^{1,2,3,4}, *Claúdia Fernandes*⁵, *Anabela Pereira*^{6,7,8},
Sara Monteiro^{1,4,9}

1. Department of Education and Psychology, University of Aveiro, Aveiro, Portugal
2. CINTESIS@RISE, CINTESIS.UPT, Portucalense University, Porto, Portugal
3. Research in Education and Community Intervention, Piaget Institute – ISEIT/Viseu, Viseu, Portugal
4. Center for Global Studies, Open University, Lisboa, Portugal
5. CATIM -Technological Center for the Metal Working Industry, Porto, Portugal
6. Department of Psychology, Colégio Pedro da Fonseca, University of Évora, Évora, Portugal
7. Center for Research in Education and Psychology (CIEP), University of Évora, Évora, Portugal
8. William James Center for Research – WJCR/UA, Aveiro, Portugal
9. Department of Social Sciences and Management, Universidade Aberta, Lisboa, Portugal

*Corresponding Author: Email: diana.araujo@ua.pt

(Received 15 Jun 2023; accepted 10 Sep 2023)

Abstract

Background: Burnout is an occupational syndrome, with a higher prevalence in professionals whose close involvement with other people is significant. There is a great diversity of professionals at risk of burnout, and therefore the implementation of intervention programs is relevant, as helping people to maintain their mental and emotional health enables them to become more meaningfully involved in their communities and become more effective and active global citizens. We aimed to review systematically the characteristics of interventions targeting burnout in health professionals.

Methods: The search was conducted in three databases: Scopus, Web of Science, and PubMed, following the PRISMA model, and 16 eligible articles were identified between 2012 and 2023.

Results: Mindfulness showed great efficacy in preventing burnout, by reducing stress levels and promoting empathy. In addition, relaxation and breathing techniques, yoga, and music therapy showed to improve burnout and occupational stress levels. Balint groups have also helped slow the progression of burnout. Implications about the need for future research to foster the promotion of well-being and mental health of health professionals are mentioned.

Conclusion: Mindfulness, relaxation and breathing techniques, yoga, music therapy and balint groups proved to be effective in preventing burnout.

Keywords: Burnout; Prevention; Intervention program; Health professionals

Introduction

Burnout is a syndrome that results from chronic stress at work, with several consequences for the

well-being and health of workers (1) and for the productivity of organizations. It is an occupational



Copyright © 2024 Araújo et al. Published by Tehran University of Medical Sciences.
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license.
(<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited

phenomenon (2) that can lead to several psychological problems, including anxiety and depression, which, can lead to the worker leaving the profession (3) or to high rates of absenteeism.

Although this issue is one of the most addressed mental health problems in society, burnout is not yet officially recognized as a mental disorder (4), and there are no differential diagnosis criteria for this issue, which is sometimes confused with depression (5). Burnout is not included in the DSM-5 (6) however, it appears as an occupational phenomenon in the ICD-11 (7). According to the WHO (2), and the ICD-11 definition; burnout results from chronic stress in the workplace characterized by feelings of exhaustion, increased feelings of negativity towards work and reduced professional effectiveness.

Due to the implications that burnout can have on health professionals' lives and the negative impact it has on patient care and other professional duties, e.g., decreased enthusiasm, motivation, and a sense of hopelessness in one's career, ineffective communication, less safe and effective clinical practices (8), as well as errors due to fatigue, lack of attention, and impaired decision-making abilities (9), it is essential to act not only at a remedial level but at a preventive level. Although burnout and stress prevention programs are scarce (10), they can be found in most of the above-mentioned professions, namely for physicians, nurses, and teachers. However, the auxiliary population is not studied as much, and minimal intervention programs are carried out with these professionals. Healthcare professionals have been developing increasingly higher rates of illness, absenteeism, burnout, and distress when compared to other professional sectors (11), which corroborates the urgent need for prevention in this area.

Some professions are more predisposed to the development of burnout syndrome, as is the case of police workers (4); health professionals such as nurses (5) and physicians, from surgeons (6) to several health sector trainees (7); emergency service professionals such as firefighters (2); teachers (8); military personnel (9); pilots (10); air traffic controllers and marine navigators (11). There is

also a high risk of burnout in formal caregivers of older people, namely nurses and aides (12).

Burnout prevention can help people maintain their mental and emotional health, which enables them to engage more meaningfully in their communities and become more effective and active global citizens (8). By investing in burnout prevention in health professionals, we are encouraging the empowerment of people to effectively manage stress, maintain a healthy work-life balance and develop communication and conflict resolution skills (13). By making professionals more functional in the various spheres of their lives, we are contributing to the promotion of global citizenship behaviors, as professionals will be able to maintain their energy and motivation to act as engaged and responsible citizens (14).

There has been increasing investment in the development of intervention programs aimed at reducing burnout among specific professional groups (e.g. physicians and nurses) (15,16). Despite this, review studies are needed to systematize valid and comprehensive results from different intervention approaches. We aimed to systematically review the characteristics of interventions targeting burnout in health professionals. This review is expected to provide a reference to guide researchers and mental health providers to select appropriate interventions and identify gaps in need of further exploration.

Materials and Methods

Overview

This systematic review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) 2020. The protocol for this systematic review was registered on PROSPERO (CRD42022352460).

Literature Search Strategy and Study Selection

To carry out this review, a search was conducted in the following online databases: Scopus, Web of Science, and PubMed. The keywords used were "burnout" AND "prevention" AND "intervention programs". Specific filters related to restrictions

on publication date and language were used. Articles in English, published between 2012 and 2023, were searched. Additional sources were identified by citation searching. The first search was performed in each database in Apr 2021, and then it was re-run in Oct 2023 to identify possible further studies.

Eligibility Criteria

Studies were eligible for final inclusion in the systematic review if they met the following criteria: (i) published in a peer-reviewed journal between the

years of 2012 and 2023; (ii) written in English; (iii) employing a quantitative design; and (v) developing and evaluating interventions for managing burnout in health professionals (e.g. nurses, physicians).

Systematic reviews and meta-analyses, incomplete articles or those that only presented abstracts, studies in languages other than English, studies with populations not related to the health sector, as well as comparative studies, literature reviews, validations, or administrations of scales were excluded (Table 1).

Table 1: Inclusion and exclusion criteria

<i>Criteria</i>	
Inclusion	Exclusion
Written in English	Written in other language than English
Year of publication 2012 to 2023	Publication out of the range 2012-2023
Study population healthcare workers	Population not related to healthcare workers
Intervention programs for managing burnout	Systematic reviews/meta-analyses

Data Extraction Process

The data extraction process was conducted according to the PRISMA model.

First, in the initial search, articles that met the inclusion criteria, which are potentially eligible articles, were selected. Next, all duplicate articles were removed. After this, the titles and abstracts were analysed. Taking into consideration the inclusion criteria (Table 1), relevant articles were extracted and those that did not fit were excluded (17) (Fig. 1).

The following information was summarized: name of the author(s) and year of publication, sample, randomized study type of intervention, duration of sessions, duration of follow-up, instruments used, and study results (Table 2).

Study quality assessment

The quality assessment of eligible studies was appraised using the Joanna Briggs Institute (JBI) methodology (18). Each item on the JBI checklist was appraised as “yes”, “no”, “unclear”, or “not

applicable”. The assessment of the methodological quality of the studies was carried out based on the information available in the articles. Two authors conducted the appraisal. Disagreements were solved by consulting with a third author.

Results

Study selection

After initial searches, 510 potentially eligible studies were identified: 115 in Scopus, 157 in Web of Science, and 238 in PubMed. In addition to these, articles were identified through citation searching (n=4). Using the JBI tool, we found that the overall quality of the studies was moderate to high. After the removal of duplicates (n=137), 373 possibly eligible studies remained. At the end of the selection of titles and abstracts, 280 articles were excluded, leaving 93 studies. Of these, 79 were excluded. Four articles were selected by citation research, leaving 16 studies that met all the selection criteria, according to the flowchart (Fig. 1).




Fig. 1: Flow chart of the study selection process

Study characteristics and participants

The majority of participants are women, except for one study in which males predominate (19). There are 7 studies that do not refer to the gender of the participants encompassing 578 study subjects (10, 20-25). In one study, only the gender of participants in the experimental group is mentioned (26). In the total of 1295 subjects, there are 474 women, 234 men, and 587 undefined/unidentified.

Most of the analyzed articles do not refer to the average age of the participants. This fact only happens in 6 studies (27-32). The average age is 37.7 yr old, with the minimum average being 22.8 yr and the maximum being 51.2 yr.

Intervention characteristics and effectiveness

Sixteen articles were included in the present review, whose data were systematized in Table 2.

Burnout prevention programs can have different levels of intervention – they can be applied in institutions, in individuals, or even at societal or sectorial levels. Generally, programs targeting individuals adopt a cognitive-behavioral approach, aiming at reducing burnout through the development of skills, coping strategies, and social support (N=8) (19).

From the total number of articles (N=16), it is possible to show that, even though all of them are with people in the healthcare sector, only one concerns a mental health intervention program, namely nurses (19). The others occur with medical (26) and nursing students (31), primary care professionals (10), operating theatre staff (29), oncology staff (21), and physicians from various specialties, including internal medicine (20,24), nurses (32), general surgery, anaesthesiology, and physical medicine/rehabilitation (20), obstetrics/gynaecol-

ogy, family medicine and paediatrics (24) and psychiatry (20,24). In one of the included studies, the study population consists of health professionals who are war veterans (30).

Some studies have been carried out on this subject, however, there are some gaps, e.g., most of these studies are carried out with physicians and nurses, leaving aside other non-medical health professionals, such as assistants.

The years of publications of the selected studies occur from 2014 to 2023, with no article published dated 2016 and 2022. The year with the highest number of published articles ($n=6$) is 2019 [Min=0; year 2016, 2022; Max=6; year 2019; SD= 1.62]. Of the studies included in this review, only two do not use assessment instruments (19,24). The instruments administered are heterogeneous, with the most common in most studies being the Maslach Burnout Inventory (MBI), used in nine studies (10, 19, 22, 23, 27, 28, 29, 33). The other used instruments were: Health Promoting Lifestyle Profile (HPLP), Freiburg Mindfulness Inventory (FMI) (24); Professional Quality of Life (ProQol) (20, 32), Revised Impact of Event Scale (IES-R) (21); Brief Psychiatric Rating Scale (BPRS), Struening and Cohen's Opinion about Mental Illness questionnaire (OMI), Font-Roja Job Satisfaction Questionnaire (FR) (22); Beck Anxiety Inventory (BAI), Authentic Happiness Inventory (AHI), Fordyce Happiness Scale (FHS) (26); Minnesota Satisfaction Questionnaire (MSQ) (28); Perceived Stress Scale (PSS) (29, 31); SOC Questionnaire (SCQ) (30); Copenhagen Burnout Inventory (CBI) (25); Five Facet Mindfulness Questionnaire (FFMQ), Perceived Stress Questionnaire (PSQ), Jefferson Scale of Physician Empathy (JSPE), Goldberg Anxiety and Depression Scale (GADS) (10); Mindful Self-Care Scale (MSCS) (31); Self-Compassion Scale (SCS) (10, 28); Quantitative Workload Inventory (QWI), Job Insecurity Scale (JIS), Work Design Questionnaire (WSQ), Work Acceptance and Action Questionnaire (WAAQ), Utrecht Work Engagement Scale (UWES) (33), Posttraumatic Growth Inventory (PTGI), Brief Resilience Scale (BRS), Self-Reflection and Insight Scale (SRIS), Self-Compassion

Scale-Short Form (SCS-SF), Psychological Empowerment Instrument (PEI), General Self-Efficacy Scale (GSE) and Brief Index of Affective Job Satisfaction (BIAJS) (32).

Follow-up only occurs in seven studies (10, 19, 21, 31, 32, 33), most commonly three months after the intervention (10, 19, 21, 30), however, studies with follow-up of six months (19, 21), ten months (22), one month (19) and four weeks (31) were found. Three of the selected articles present follow-up at various time points: 1, 3, and 6 months (19, 32) and 3 and 6 months (21).

There is a clear preference for face-to-face studies, only online intervention occurred in one of the studies included in the review (27).

The selected articles present diversity regarding the country in which the intervention program was implemented: Saudi Arabia (19), Spain (10, 22), China (28), Tunisia (29), United States of America (24-26, 31, 32) and the Netherlands (33). Five of the articles do not identify in which country the intervention program was implemented.

A minority of randomized studies is clear, only five studies follow this type of method (10, 19, 27, 28, 32).

Regarding the duration of sessions, the least long-lasting intervention is about 2 days (19, 21), followed by 4 wk (31), 6 wk (29), 8 wk (27), 9 wk (22), 10 wk (33), 2 months (32), 3 months (20, 30), 4 months (24), 5 months (23), 6 months (28), 10 months (33) and 1 year (25). There is one study that conducts the intervention at two time points, 4 and 8 wk (10).

It is possible to see that most of the analyzed articles compare groups (10, 19, 22-26, 28, 29, 32, 33). Of eleven studies, there are nine which use experimental and control groups. Of these, there are five which refer to the size of the effect (10, 19, 22, 27, 29, 32).

There are several types of intervention in the selected articles, all of which aim to intervene in the promotion of mental health, either by directly preventing burnout or indirectly by reducing occupational stress, as well as promoting job satisfaction. The intervention programs found are mostly psychoeducational. There are programs based on psychotherapeutic interventions (10, 22, 26, 31, 32),

personal and social development (19-21, 23, 25, 27, 29, 30) and personal and professional well-being (19-21, 23, 25, 27, 29, 30). The studies present a significant diversity of interventions, including wellness programs, yoga, focus groups, music

therapy, mindfulness, and coaching. This variety suggests the need for approaches tailored to the needs of health professionals, recognizing that different individuals may respond better to different strategies.

Table 2: Summary of the studies

Reference No.	Sample Size	Participants	Randomized	Comparison Groups / Effect Size	Intervention Format	Duration/ Number of Sessions	Follow-up	Instruments	Results
(20)	n=188	Residents	No	No / No	Wellness Intervention	12 wk	-	-	Residents chose to continue the weekly sessions voluntarily after the initial 12-week curriculum
(19)	n=296	Mental Health Nurses	Yes (For 1 stage)	Yes / Yes	Burnout Prevention Programme	2 days	1-, 3-, and 6-months intervals	MBI ¹	The intervention was effective in managing burnout
(27)	n=40	Nurses	Yes	Yes / Yes	Yoga Intervention	8 wk	-	HPLP II ² ; FMI ³ ; MBI	Participants reported higher self-care, lower emotional exhaustion and depersonalization
(21)	n=15	Oncology Staff	No	No / No	Compassion Fatigue Program	4 sessions	3 and 6 months	ProQol IV ⁴ ; IES-R ⁵	Compassion fatigue resilience programs have been shown to be effective, feasible, and scalable
(22)	n=38	General Practitioners	No	Yes / Yes	Multimodal Training Programme (MTP) Integrated Brief Systemic Therapy (IBST) approach	9 weekly sessions	10 months	BPRS ⁶ ; OMI ⁷ ; FR ⁸ ; MBI-HSS ⁹	MTP showed significant improvements, better scores on overall psychopathological status, and better evolution of job satisfaction
(31)	n=67	Nursing Students	No	No/ No	Mindfulness sessions	4 wk	4 wk	PSS ¹⁰ ; MSCS ¹¹	Mindfulness education increased self-care in nursing students who practiced mindfulness outside class

¹ MBI - Maslach Burnout Inventory

² HPLP - Health Promoting Lifestyle Profile

³ FMI - Freiburg Mindfulness Inventory

⁴ ProQol - Professional Quality of Life

⁵ IES-R Revised Impact of Event Scale

⁶ BPRS - Brief Psychiatric Rating Scale

⁷ OMI - Struening and Cohen's Opinion about Mental Illness

⁸ FR - Font-Roja Job Satisfaction Questionnaire

⁹ MBI-HSS - MBI - Human Service Survey

¹⁰ PSS - Perceived Stress Scale

¹¹ MSCS - Mindful Self-Care Scale

Table 2: Continued...

(23)	n=46	Emergency Medicine Residents	No	Yes / No	The Happiness Practice (THP) – Corporate-based well-ness intervention	5 months; 6 monthly sessions	-	MBI	The program shows a decreased burnout, while increase individual happiness, resilience, innovation and sustainability
(26)	n=93	Medical Students	No	Yes / No	The Joy Initiative	10 wk	-	BAI ¹² ; AHI ¹³ ; FHS ¹⁴	Students reported improvements in life satisfaction, as well as increased ability to cope with stressors
(28)	n=36	Resident Physicians	Yes	Yes / No	Balint Group Intervention	10 sessions/6 months	-	MBI-HSS; MSQ ¹⁵	The results indicated that Balint groups helped to slow the progression of burnout
(24)	n=120	Core Faculty and Residents	No	Yes / No	Focus Groups	4 months/14 focus groups	-	-	The results emphasize the importance of interventions targeted at the social environments in which residents work
(29)	n=34	Operating Room Staff	No	Yes / Yes	Music Therapy Program	6 wk	-	PSS-10 ¹⁶ ; MBI	Music therapy improved the stress levels of the operating room staff
(30)	n=39	Health Care Workers within the Veteran Affairs	No	No / No	Mantram Repetition Program (MRP)	3 months	3 months	MBI-GS ¹⁷ ; SCQ ¹⁸	An MRP intervention may reduce burnout, particularly in those individuals who are naïve to practicing meditation
(25)	n=39	Residents	No	No / No	Implemented Wellness Program	1 year	-	CBI ¹⁹	The results show a clinically significant overall reduction in mean burnout
(10)	n=132	Primary Care Professionals	Yes	Yes / Yes	MINDUDD – Intervention based on Mindfulness-Based Stress Reduction	8 wk and 4 wk	3 months	FFMQ ²⁰ ; SCS ²¹ ; PSQ ²² ; MBI-GS; JSPE ²³ ; GADS ²⁴	Results show that there is a correlation between mindfulness and self-compassion and increased resilience and psychological well-being

¹² BAI - Beck Anxiety Inventory¹³ AHI - Authentic Happiness Inventory¹⁴ FHS - Fordyce Happiness Scale¹⁵ MSQ - Minnesota Satisfaction Questionnaire¹⁶ PSS-10 - Perceived Stress Scale Version 10¹⁷ MBI-GS - MBI-General Survey¹⁸ SCQ - SOC Questionnaire¹⁹ CBI - Copenhagen Burnout Inventory²⁰ FFMQ – Five Facet Mindfulness Questionnaire²¹ SCS - Self-Compassion Scale²² PSQ - Perceived Stress Questionnaire²³ JSPE - Jefferson Scale of Physician Empathy²⁴ GADS - Goldberg Anxiety and Depression Scale

Table 2: Continued...

(32)	n=77	Nurses	Yes	Yes/Yes	Psychoeducational Group Program	2 months	1, 3 and 6 months	PTGI ²⁵ ; ProQoL; BRS ²⁶ ; SRIS ²⁷ ; SCS-SF ²⁸ ; PEI ²⁹ ; GSE ³⁰ ; PSS; BIAJS ³¹	Psychoeducational group program can be an effective intervention for improving mental wellbeing
(33)	n=114	Residents and Specialists	No	Yes / No	Coaching Intervention	10 months / 6 sessions	-	Some items: QWI ³² ; JIS ³³ ; WAFCS ³⁴ ; SCS; WDQ ³⁵ ; MBI; UWES ³⁶	Results show that the coaching group reported a reduction in burnout symptoms

Discussion

Over the last decade, an increasing number of studies on the issue of burnout have emerged. The present review seeks to analyze 16 articles, published between 2014 and 2023, which aimed at sharing burnout intervention programs with professionals from the healthcare sector.

There are advantages in the implementation of burnout intervention programs in individuals' mental health.

There is a predominance of intervention programs focused on the promotion of well-being, as a way of preventing occupational pathologies, such as burnout. By enhancing self-awareness, the individual improves self-care, which improves psychological well-being (10), which, in turn, helps prevent burnout. Furthermore, programs that work to prevent compassion fatigue indirectly prevent burnout, as compassion fatigue can progress to burnout (22). Strategies that proved to be effective in

preventing/reducing burnout were relaxation exercises, breathing training, music therapy, and yoga.

Concerning psychotherapeutic interventions, mindfulness shows benefits in reducing stress and anxiety levels (22, 31), as well as promoting the development of empathy in professionals (10), so it may be a path to be followed in burnout prevention. Several intervention programs are based on Mindfulness-Based Stress Reduction (MBSR) (34). Mindfulness-based interventions have consistently shown positive effects in reducing burnout, as increasing self-awareness and the ability to cope with stress seems to be a promising avenue for improving the well-being of healthcare professionals. The practice of mindfulness has not only proved effective in managing stress, but also in promoting empathy among health professionals.

Studies incorporating practices such as yoga and music therapy have also shown promising results, as improved self-care, reduced perceived stress

²⁵ PTGI - Posttraumatic Growth Inventory

²⁶ BRS - Brief Resilience Scale

²⁷ SRIS - Self-Reflection and Insight Scale

²⁸ SCS-SF - Self-Compassion Scale-Short Form

²⁹ PEI - Psychological Empowerment Instrument

³⁰ GSE - General Self-Efficacy Scale

³¹ BIAJS - Brief Index of Affective Job Satisfaction

³² QWI - Quantitative Workload Inventory

³³ JIS - Job Insecurity Scale

³⁴ WDQ - Work Design Questionnaire

³⁵ WAAQ - Work Acceptance and Action Questionnaire

³⁶ UWES - Utrecht Work Engagement Scale

and emotional exhaustion highlight the importance of holistic approaches that integrate body and mind.

Balint groups have been shown to be effective in promoting mental well-being since social interaction and sharing experiences seem to play a significant role in preventing burnout (28, 35-37).

WHO has emphasized the need for healthcare professionals to have access to training and education programs that promote well-being and resiliency (2). Education and training are instrumental in enhancing professional knowledge and skills, and transversal competencies, alongside promoting global citizenship. In patient care environments, special training can be introduced that teaches professionals about different practices, and ways to empower teams and individuals, promoting, for example, transversal competencies (e.g., coping strategies, meditation, and yoga).

There is a minority of studies with follow-up draws our attention to the need to evaluate better the effectiveness of intervention programs. Interventions should take into consideration their effectiveness and without follow-up, it is difficult to understand the benefits of interventions in the medium and long term and to design and manage intervention programs based on key performance indicators.

Organizations that adopt responsible social practices have a healthier and more balanced work environment, which can help prevent burnout among their employees (38). However, organizations that neglect social and environmental issues and do not value the well-being of their employees have a more stressful work environment, which can lead to an increase in burnout (39).

Limitations were found in this systematic review. Firstly, the characteristics of the studies vary in several aspects, from the number of sessions, type/format of intervention, and target audience. Second, there is little variability regarding burnout instruments, with most studies using the MBI.

Finally, restricting the search, may have limited the discovery of relevant articles.

Some flaws need to be addressed in future research. One suggestion for future studies is to have more interventions in organizations as there

are more long-term benefits (40). Organizations should invest in promoting employee health through burnout promotion programs as it can help create a healthier working environment, which can lead to a stronger organizational culture that is more likely to support global citizenship initiatives (14).

Other forms of burnout prevention were not addressed due to the study criteria and the nature of the investigation, such as volunteering, as getting involved in volunteer activities can promote positive emotions, provide a sense of accomplishment, and promote connections, which can have a positive impact on preventing burnout (41). Volunteer opportunities create a sense of shared experience and purpose for the work done, help to build mutual interests, and promote empathy. Besides that, digital connectivity can help to reduce burnout among healthcare workers by providing access to social support, information and resources (42). Internet access and digital tools provide individuals with access to information, allowing them to connect with others and begin to engage in the world's conversation and best practices, reducing isolation and promoting, for example, belonging to digital communities.

It is important to consider a greater diversity of target audiences for burnout intervention programs, such as medical assistants or direct care nurses, whose studies are still scarce, with a clear prevalence of intervention programs in physicians and nurses. Early intervention may also help to prevent the onset of burnout, being, therefore, a suggestion for future research.

Conclusion

This systematic review provides a comprehensive overview of burnout intervention programs in the healthcare sector, emphasizing the importance of a multifaceted approach that encompasses individual well-being, organizational practices, and collaborative initiatives within the healthcare community.

These results have implications for promoting well-being. Implementing effective strategies can

not only benefit workers but also contribute to a healthier and more sustainable working environment. However, it is essential to recognize the complexity of burnout and the need for approaches that consider both individual and organizational factors. The path to preventing burnout requires an ongoing commitment to mental health and well-being, both on the part of professionals and institutions.

The significant influence of the work environment on burnout experiences underscores the responsibility of organizations in promoting mental health. Socially responsible organizational practices not only create more balanced environments but also contribute to the prevention of burnout.

The identified gaps provide a roadmap for future research, guiding efforts toward more inclusive and sustainable strategies for preventing burnout in healthcare professionals.

Journalism Ethics considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors declare that there is no conflict of interests.

References

1. Salvagioni DAJ, Melanda FN, Mesas AE, et al. (2017). Physical, psychological and occupational consequences of job burnout: A systematic review of prospective studies. *PLoS One*, 12 (10): e0185781.
2. World Health Organization. Burn-out an "occupational phenomenon": International Classification of Diseases. 2019.
3. De la Fuente Solana EI, Pradas-Hernández L, González-Fernández CT, et al. (2021). Burn-out syndrome in paediatric nurses: A multi-centre study. *Int J Environ Res Public Health*, 18 (3): 1324.
4. Heinemann LV, Heinemann T (2017). Burnout research: Emergence and scientific investigation of a contested diagnosis. *SAGE Open*, 7 (1): 215824401769715.
5. Kaschka WP, Korczak D, Broich K (2011). Burn-out: A fashionable diagnosis. *Dtsch Arztebl Int*, 108 (46): 781–787.
6. American Psychiatric Association (2013). *Diagnostic and statistical manual of mental disorders*. 5th ed. American Psychiatric Publishing, England, pp.: 8-40.
7. World Health Organization. 2022. International Classification of Diseases 11th Revision. <https://icd.who.int/en>
8. Maslach C, Leiter MP (2016). Understanding the burnout experience: Recent research and its implications for psychiatry. *World Psychiatry*, 15 (2): 103-111.
9. Shanafelt TD, Hasan O, Dyrbye LN, et al. (2015). Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clin Proc*, 90(12):1600-13.
10. Pérula-De Torres LA, Atalaya JC, García-Campayo J, et al. (2019). Controlled clinical trial comparing the effectiveness of a mindfulness and self-compassion 4-session programme versus an 8-session programme to reduce work stress and burnout in family and community medicine physicians and nurses: MINDUDD study protocol. *BMC Fam Pract*, 20 (1) :24.
11. Brand SL, Coon JT, Fleming LE, et al. (2017). Whole-system approaches to improving the health and wellbeing of healthcare workers: A systematic review. *PLoS One*, 12 (12): e0188418.
12. Potard C, Landais C (2021). Relationships between frustration intolerance beliefs, cognitive emotion regulation strategies and burnout among geriatric nurses and care assistants. *Geriatr Nurs*, 42 (3): 700–707

13. Oliveira SM, Alcantara Sousa LV, Gadelha MDS, et al. (2019). Prevention Actions of Burnout Syndrome in Nurses: An Integrating Literature Review. *Clin Pract Epidemiol Ment Health*, 15: 64-73.
14. Van Rompay-Bartels I, Tuninga RSJ (2023). Toward a model of global citizenship in business education. *Journal of Transnational Management*, 28 (1–2): 5–34.
15. Kalani SD, Azadfallah P, Oreyzi H, et al. (2018). Interventions for Physician Burnout: A Systematic Review of Systematic Reviews. *Int J Prev Med*, 9:81.
16. Zhang XJ, Song Y, Jiang T, et al. (2020). Interventions to reduce burnout of physicians and nurses: An overview of systematic reviews and meta-analyses. *Medicine (Baltimore)*, 99 (26): e20992.
17. Page MJ, McKenzie JE, Bossuyt PM, et al. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 372:n71.
18. Aromataris E, Fernandez R, Godfrey C, et al. (2015). Summarizing systematic reviews: methodological development, conduct and reporting of an umbrella review approach. *Int J Evid Based Healthc*, 13 (3): 132-140.
19. Alenezi A, McAndrew, S, Fallon P (2019). Burning out physical and emotional fatigue: Evaluating the effects of a programme aimed at reducing burnout among mental health nurses. *Int J Ment Health Nurs*, 28 (5): 1045–1055.
20. Aggarwal R, Deutsch JK, Medina J, et al. (2017). Resident Wellness: An Intervention to Decrease Burnout and Increase Resiliency and Happiness. *MedEdPORTAL*, 13:10651.
21. Back AL, Deignan PF, Potter PA (2014). Compassion, Compassion Fatigue, and Burnout: Key Insights for Oncology Professionals. *Am Soc Clin Oncol Educ Book*, 2014:e454-9.
22. Barcons C, García B, Sarri C, et al. (2019). Effectiveness of a multimodal training programme to improve general practitioners' burnout, job satisfaction and psychological well-being. *BMC Fam Pract*, 20 (1): 155.
23. Hart D, Paetow G, Zarzar R (2019). Does implementation of a corporate wellness initiative improve burnout? *West J Emerg Med*, 20 (1): 138–144.
24. Ironside K, Becker D, Chen I, et al. (2019). Resident and Faculty Perspectives on Prevention of Resident Burnout: A Focus Group Study. *Perm J*, 23:18-185.
25. Mari S, Meyen R, Kim B (2019). Resident-led organizational initiatives to reduce burnout and improve wellness. *BMC Med Educ*, 19 (1): 437.
26. Hlubocky FJ, Rose M, Epstein RM (2017). Mastering Resilience in Oncology: Learn to Thrive in the Face of Burnout. *Am Soc Clin Oncol Educ Book*, 37 (1): 771–781.
27. Alexander GK, Rollins K, Walker D, et al. (2015). Yoga for Self-Care and Burnout Prevention among Nurses. *Workplace Health Saf*, 63 (10): 462–470.
28. Huang L, Harsh J, Cui H, et al. (2020). A Randomized Controlled Trial of Balint Groups to Prevent Burnout among Residents in China. *Front Psychiatry* 10:957.
29. Kacem I, Kahloul M, El Arem S, et al. (2020). Effects of music therapy on occupational stress and burn-out risk of operating room staff. *Libyan J Med*, 15 (1): 1768024.
30. Leary S, Weingart K, Topp R, Bormann J (2018). The Effect of Mantram Repetition on Burnout and Stress Among VA Staff. *Workplace Health Saf*, 66 (3): 120–128.
31. Burner LR, Spadaro KC (2023). Self-care skills to prevent burn-out: a pilot study embedding mindfulness in an undergraduate nursing course. *J Holist Nurs*, 41 (3): 265-274.
32. Sawyer AT, Tao H, Bailey AK (2023). The Impact of a Psychoeducational Group Program on the Mental Well-Being of Unit-Based Nurse Leaders: A Randomized Controlled Trial. *Int J Environ Res Public Health*, 20 (11): 6035.
33. Solms L, Van Vianen A, Koen J, et al. (2021). Turning the tide: A quasi-experimental study on a coaching intervention to reduce burn-out symptoms and foster personal resources among medical residents and specialists in the Netherlands. *BMJ Open*, 11 (1): e041708.
34. Klein A, Taieb O, Xavier S, et al. (2020). The benefits of mindfulness-based interventions on burnout among health professionals: A systematic review. *Explore (NY)*, 16 (1): 35–43.
35. Hanson P, Clarke A, Villarreal M, et al. (2020). Burnout, resilience, and perception of mindfulness programmes among GP trainees: A mixed-methods study. *BJGP Open*, 4 (3): bjpopen20X101058.

36. Van Roy K, Vanheule S, Inslegers R (2015). Research on Balint groups: A literature review. *Patient Educ Couns*, 98 (6): 685–694.
37. Huang H, Zhang H, Xie Y, et al. (2020). Effect of Balint group training on burnout and quality of work life among intensive care nurses: A randomized controlled trial. *Neurol Psychiatry Brain Res*, 35:16–21.
38. Ahmad N, Ullah Z, Ryu HB, et al. (2023). From Corporate Social Responsibility to Employee Well-Being: Navigating the Pathway to Sustainable Healthcare. *Psychol Res Behav Manag*, 16:1079-1095.
39. Schabracq MJ, Winnubst JAM, Cooper CL (2014). A critical review of the Burnout concept: thirty-five years of research and theorizing. In: *The Handbook of Work and Health Psychology*. Eds, Schaufeli and Taris. 3rd ed, John Wiley & Sons, UK, pp. 383-425.
40. Awa WL, Plaumann M, Walter, U (2010). Burnout prevention: A review of intervention programs. *Patient Educ Couns*, 78 (2): 184–190.
41. Nichol B, Wilson R, Rodrigues A, Haighton C (2023). Exploring the Effects of Volunteering on the Social, Mental, and Physical Health and Well-being of Volunteers: An Umbrella Review. *Voluntas*, 1-32.
42. Esmacilzadeh P, Mirzaei T (2021). Using Electronic Health Records to Mitigate Workplace Burnout among Clinicians during the COVID-19 Pandemic: Field Study in Iran. *JMIR Med Inform*, 9(6): e28497.