



Case report

Bilateral coronary ostial disease following mediastinal irradiation: a case report

Salman Waqar*, Rajwinder Jutley, Richard Mount and Pradip Sarkar

Address: Department of Cardiothoracic Surgery, Northern General Hospital, Sheffield Teaching Hospitals NHS Trust, Sheffield, S5 7AU, UK

Email: SW* - salmanwqr@gmail.com; RJ - jutley@btinternet.com; RM - md00ram@doctors.org.uk; PS - pradip.sarkar@sth.nhs.uk

* Corresponding author

Received: 1 March 2009 Accepted: 29 July 2009 Published: 25 August 2009

Cases Journal 2009, 2:7792 doi: 10.4076/1757-1626-2-7792

This article is available from: <http://casesjournal.com/casesjournal/article/view/7792>

© 2009 Waqar et al.; licensee Cases Network Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Introduction: Ostial coronary artery disease is rare with a reported incidence of 0.07 to 0.25% in all patients undergoing angiography. It has a strong association with previous mediastinal irradiation, which induces specific histological changes distinct from atherosclerotic lesions. The radiation also affects the myocardium and surrounding structures, which can alter the surgical approach.

Case presentation: We present a case of a 62-year-old female who developed bilateral ostial coronary artery stenosis 32 years following therapeutic radiotherapy for Hodgkin's disease. She underwent successful coronary artery bypass surgery using a combination of arterial and venous conduits. Postoperatively she developed a clinical picture of diastolic impairment not detected pre-operatively. She was managed appropriately and made a successful recovery.

Conclusions: This case highlights the cardiac pathology associated with mediastinal irradiation, which should be suspected during surgical assessment, especially in long-term survivors. It heightens the surgeon's awareness so a more thorough evaluation of coronary anatomy, ventricular function and potential conduits is made prior to surgery.

Introduction

Ostial coronary artery disease is rare with a reported incidence of 0.07 to 0.25% in all patients undergoing angiography [1]. It has a strong association with previous mediastinal irradiation, which induces specific histological changes distinct from atherosclerotic lesions. The radiation also affects the myocardium and surrounding structures, which can alter the surgical approach.

Case presentation

A 62-year-old British white Caucasian female presented with a 2-year history of Canadian Cardiovascular Society

(CCS) Class II exertional angina and mild dyspnoea. Her medical history consisted of hypertension, hypercholesterolaemia and Hodgkin's disease 32 years previously for which she underwent radiotherapy to the chest and pelvis. The total radiotherapy dose delivered was 5330 rads (cGy). She underwent coronary angiography following a strongly positive exercise test. This showed severe (>90%) stenosis at the origin of left main stem associated with a significant pressure drop across the lesion. There was also significant stenosis in the proximal circumflex artery, intermediate artery and proximal left anterior descending (LAD) artery (Figure 1). Injection of the right coronary




Figure 1. Injection of the left coronary system showed ostial coronary disease associated with a pressure drop across the lesion.

system demonstrated severe ostial stenosis (>90%) again associated with a pressure drop across the lesion (Figure 2). Her left ventricular function was preserved on ventriculogram. The patient underwent urgent CABG using saphenous vein to bypass the right coronary artery (RCA), a radial artery graft to the obtuse marginal (OM) and left internal mammary artery (LIMA) to the mid-LAD. She was weaned from cardiopulmonary bypass (CPB) without inotropic support. Her initial recovery was




Figure 2. The right coronary system also had similar features on injection of the contrast.

unremarkable, and she had a fast-tracked extubation. Within a few hours, however, she developed signs of low cardiac output syndrome with oliguria. Fluid challenges were administered resulting in rapidly raised central venous pressures (CVP) and dyspnoea secondary to pulmonary oedema. Invasive monitoring with a pulmonary artery wedge catheter (PAWC) demonstrated high filling pressures (pulmonary artery wedge pressure of 20 mmHg) with a low cardiac index of 1.8. The features were suggestive of severe diastolic dysfunction. Inotropic support with adrenaline was commenced with good clinical response. She improved with circulatory support and continuous positive airway pressure (CPAP) over the following 3 days and was discharged home 8 days later.

Discussion

Radiotherapy is an established treatment modality for a variety of tumours including Hodgkin's disease. Radiotherapy is also used as adjuvant treatment in breast cancer. Both breast cancer and Hodgkin's disease account for the most important causes of radiation-induced cardiac damage with the latter resulting in a higher relative risk estimate for fatal cardiovascular events (2.2 to 7.2) [2]. The risk correlates with younger age at irradiation, length of follow-up and dose volume used [3].

Cardiac damage can manifest in a variety of ways especially if >65% of the heart has been irradiated [4]. Clinical presentation ranges from most commonly pericarditis to valvular dysfunction, conduction abnormalities and myocardial infarction [4,5]. A rare complication of mediastinal irradiation is the development of coronary artery disease [6,7]. The pattern of disease is unusual in that it tends to affect the coronary ostia, presumably due to their relatively central location within the radiation field. Histologically, radiation-induced coronary lesions vary from atherosclerotic disease although it has been suggested that both pathologies may act in synergy whereby atherosclerosis is accelerated if risk factors such as hypercholesterolaemia are present [8]. The intimal plaque is similar in both radiation-induced disease and atherosclerosis. Virani *et al* demonstrated that characteristically with radiation damage there is medial thinning and adventitial fibrosis [9]. Other histological features include intimal foam cell collections with calcification and necrosis within the central core of the plaque. Irradiation damage to the heart may extend beyond the coronary arteries. A surgical autopsy study by Vienot and Edwards at the Mayo Clinic on 27 patients with previous mediastinal radiation showed that 71% of cases had radiation injury to the valves with the mitral valve the most affected. 63% of cases had radiation-induced fibrosis most severe in patients who received a radiation dose greater than 3000 rad (cGy) radiation dose [10]. It is likely the diastolic dysfunction observed in our patient was secondary to

radiation induced subendocardial fibrosis as she received a cumulative-dose of 5330 rad (cGy). Although theoretically less frequent with modern techniques of radiotherapy that employ lower doses and cardiac shielding, fibrosis would have prevented myocardial stretch with hindrance of Starling's forces. This dysfunction was not apparent on reviewing the pre-operative ventriculogram, which tends to evaluate systolic function.

The Internal Mammary Arteries (IMA) may also be included within the radiation field and damaged thereby precluding its use for coronary bypass grafting [11]. However, in a comprehensive review of forty-nine patients with previous mediastinal irradiation who underwent elective coronary bypass surgery using at least one IMA, Nasso *et al* found no incidence of graft failure [12]. Intra operative mammary flow rates assessed by Doppler showed no difference between irradiated and non-irradiated patients. The group did, however, suggest the use of skeletonisation during conduit harvesting due to the abnormally high incidence of fibrous tissue around the vessel to achieve maximum length.

Our report highlights the complexities in dealing with patients with previous mediastinal irradiation, more so when historically higher doses with lack of cardiac shielding were employed. In support of the observations made by Fuzellier *et al*, we recommend a multi-disciplinary approach when managing such patients, especially if the patient has previously received a radiation dose of more than 3000 rad (cGy) [7]. It is important that patients are carefully assessed for ostial coronary lesions, preferably to also assess the internal mammary arteries and echocardiography performed routinely to evaluate ventricular function. As long-term survivors following mediastinal irradiation (especially for Hodgkin's disease) are at higher risk of cardiovascular fatality, particular attention must be paid to this subset of patients. Surgery is the treatment of choice and the use of the IMA over saphenous vein grafts should be encouraged as it provides a prognostic benefit [13].

Abbreviations

CABG, coronary artery bypass graft; CCS, canadian cardiovascular society; CPAP, continuous positive airway pressure; CPB, cardio-pulmonary bypass; CVP, central venous pressure; LAD, left anterior descending; LIMA, left internal mammary artery; OM, obtuse marginal; RCA, right coronary artery.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review from the journal's Editor-in-Chief.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SW, RJ, RM and PKS managed the patient, analyzed and interpreted the data including the coronary angiograms. All authors contributed to writing the manuscript including the discussion and have read and approved the final manuscript.

References

1. Grigorov V, Goldberg L, Mekel J: **Isolated bilateral ostial coronary stenosis with proximal right coronary artery occlusion.** *Int J Cardiovasc Intervent* 2000, **3**:47-49.
2. Adams MJ, Hardenbergh PH, Constine LS, Lipshultz SE: **Radiation-associated cardiovascular disease.** *Crit Rev Oncol Hematol* 2003, **45**:55-75.
3. Adams MJ, Lipshultz SE, Schwartz C, Fajardo LF, Coen V, Constine LS: **Radiation associated cardiovascular disease: manifestations and management.** *Semin Radiat Oncol* 2003, **13**:346-356.
4. Gaya AM, Ashford RF: **Cardiac complications of radiation therapy.** *Clin Oncol* 2005, **17**:153-159.
5. Kaplan BM, Miller AJ, Bharati S, Lev M, Martin Grais I: **Complete AV block following mediastinal radiation therapy: electrocardiographic and pathologic correlation and review of the world literature.** *J Interv Card Electrophysiol* 1997, **1**:175-188.
6. Sachithanandan A, Ahmed A, O'Kane H: **Bilateral isolated coronary ostial stenosis following mediastinal irradiation.** *Asian Cardiovasc Thorac Ann* 2004, **12**:78-88.
7. Fuzellier JF, Mauran P, Metz D: **Radiation-induced Bilateral Coronary Ostial Stenosis in a 17-year-Old Patient.** *J Card Surg* 2006, **21**:600-602.
8. Stewart JR, Fajardo LF: **Radiation-induced heart disease: an update.** *Prog Cardiovasc Dis* 1984, **27**:173-194.
9. Virmani R, Farb A, Carter AJ, Jones RM: **Pathology of radiation-induced coronary artery disease in human and pig.** *Cardiovasc Radiat Med* 1999, **1**:98-101.
10. Veinot JP, Edwards WD: **Pathology of radiation-induced heart disease: a surgical and autopsy study of 27 cases.** *Hum Pathol* 1996, **27**:766-773.
11. Khan MH, Ettinger SM: **Post mediastinal radiation coronary artery disease and its effects on arterial conduits.** *Catheter Cardiovasc Inter* 2001, **52**:242-248.
12. Nasso G, Canosa C, De Filippo CM, Modugno P, Anselmi A, Gaudino M, Alessandrini F: **Thoracic radiation therapy and suitability of internal thoracic arteries for myocardial revascularization.** *Chest* 2005, **128**:1587-1592.
13. Loop FD, Lytle BW, Cosgrove DM, Stewart RW, Goormastic M, Williams GW, Golding LA, Gill CC, Taylor PC, Sheldon WC et al.: **Influence of the internal-mammary-artery graft on 10-year survival and other cardiac events.** *N Engl J Med* 1986, **314**:1-6.