

# Regulatory Mechanisms and Physiological Impacts of Quorum Sensing in Gram-Negative Bacteria

Xiaobing Chu<sup>1,2</sup>, Qiwen Yang<sup>1,3</sup>

<sup>1</sup>Department of Clinical Laboratory, State Key Laboratory of Complex Severe and Rare Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, 100730, People's Republic of China; <sup>2</sup>Graduate School, Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, People's Republic of China; <sup>3</sup>Key Laboratory of Pathogen Infection Prevention and Control, Peking Union Medical College, Ministry of Education, Beijing, People's Republic of China

Correspondence: Qiwen Yang, Email yangqiwen81@vip.163.com

**Abstract:** The Quorum sensing (QS) system is a widely existing communication mechanism, which regulates bacterial community behaviors and the expression of specific genes. The most common pathogenic bacteria in clinical infections are gram-negative bacteria, and QS plays an important regulatory role in the production of virulence factors and development of antibiotic resistance. This article reviews the QS systems of gram-negative bacteria and provides an overview of how they regulate their physiological functions.

**Keywords:** quorum sensing, gram-negative bacteria, physiological impacts


Currently, Gram-negative bacteria are the predominant pathogens responsible for hospital-acquired infections, exceeding the incidence of infections caused by Gram-positive pathogens by 20%.<sup>1</sup> The 2023 China Antimicrobial Surveillance Network data indicate that clinical infections continue to be predominantly caused by gram-negative bacteria, accounting for 70.1% of the cases.<sup>2</sup> The most common pathogens in descending order are *Escherichia coli* (18.02%), *Klebsiella pneumoniae* (14.06%), *Acinetobacter baumannii* (8.76%), and *Pseudomonas aeruginosa* (7.46%).<sup>2</sup> These Gram-negative strains are the most common among carbapenem-resistant gram-negative bacilli and represent drug-resistant bacteria of the greatest concern in clinical settings. In 2019, approximately 4.95 million people died from diseases related to bacterial resistance, with approximately 1.27 million of these deaths directly attributed to bacterial resistance.<sup>3</sup> There are multiple reasons for the development of antibiotic resistance in bacteria, such as biofilm formation, genetic material mutations, latent gene activation, and the transfer of genetic material (for example, horizontal gene transfer mediated by phage transduction or plasmid conjugation).

Quorum sensing (QS) is a mechanism for intra- or intercellular communication and the regulation of gene expression, which is achieved through the production, release, receptor binding, and induction of downstream signals by signaling molecules.<sup>4</sup> Many physiological functions of Gram-negative bacteria are regulated by QS. In terms of virulence regulation, upon invading the human body, pathogens utilize QS systems to sense the population density and develop offensive and defensive strategies. Under low-density conditions, bacteria downregulate the expression of virulence factors to evade the host immune attack. When the population density increases to a level that the human immune system cannot cope with, bacteria express a large number of virulence factors in a short period, thereby delivering a devastating blow to the host's immune system.<sup>5</sup> QS also plays a crucial role in bacterial antibiotic resistance and it has been confirmed that the signaling molecule system regulates bacterial efflux pumps.<sup>6</sup> Under the pressure of antibiotics and host defenses, QS systems can regulate the formation of biofilms, persister cells, and even small colony variants in pathogens.<sup>7</sup> Surface-active agents regulated by QS can enhance the hydrophobicity of cells, thereby facilitating their adhesion to solid substrates.<sup>8</sup> QS is one of the most extensively studied targets in antivirulence therapy research. Since this process enables the coordinated regulation of multiple virulence factors without being essential for bacterial growth, targeting QS can control bacterial pathogenesis while limiting selective pressure for survival and the development of

antibiotic resistance. The link between QS and pathogen virulence and resistance has led to increasing interest in using QS as a new approach to treat bacterial infections. Therefore, regulation of bacterial physiological functions by QS should be considered. Research on the QS systems of bacteria can aid in gaining a comprehensive understanding of bacterial communities.

## An Overview of QS Systems in Gram-Negative Bacterial The AHL-Based QS System

In Gram-negative bacteria, QS is commonly mediated by N-acyl homoserine lactones (AHLs), also known as auto-inducer-1 (AI-1) (Figure 1A). AHLs were first discovered in the bioluminescence system of the marine bacterium *Vibrio*



**Figure 1** Schematic of three different natural QS systems. **(A)** *V. fischeri*'s LuxI/LuxR QS system. AHL is one of the QS signal molecules. LuxI is an AHL synthetase, and LuxR binds to AHL to form the LuxR-AHL complex, which initiates transcription of the *lux* cluster by binding to the plus promoter. The *lux CDABE* in the *Lux* cluster is responsible for bioluminescence. **(B)** AI-2 QS system. AI-2 is one of the QS signaling molecules synthesized by *luxS*. AI-2 is transported into and out of the cell by membrane materials, and *lsrK* can phosphorylate AI-2. In the absence of phosphorylated AI-2, *LsrR* inhibits transcription of *P<sub>luxR</sub>* and *P<sub>luxS</sub>* by binding to these promoters. When phosphorylated AI-2 appears and binds to *LsrR*, transcription is inhibited by the release of *LsrR* promoter binding. **(C)** The AI - 3 / epinephrine, norepinephrine signaling pathway. AI - 3 and epinephrine/norepinephrine appear to be recognized by the same receptor and interact with sensor kinases located in the periplasm. The adrenergic receptor, QseC, makes up part of the QseB/C two-component system that regulates the flagellumegulon and thus motility.

**Abbreviations:** IM, inner membrane; OM, outer membrane.

*fischeri*, and this phenomenon has been positively correlated with the bacterial population density.<sup>9</sup> The AHL-based QS system comprises three components: (1) AHL signaling molecules, (2) synthase proteins that produce AHL signals, and (3) receptor proteins that perceive and respond to AHLs. AHLs consist of a series of fatty acid side chains of varying lengths that are connected to a homoserine lactone (HSL) ring. Most known AHLs have fatty acid side chains ranging from 4 to 20 carbon atoms in length. Their specificity is primarily determined by the length of the fatty acid side chain, modifications of the carbon atoms (mainly at the C3 position, with some AHLs showing biological activity only after modification), and degree of saturation of the fatty acid side chain.<sup>10,11</sup> The AHL-based QS system exhibits four fundamental characteristics: 1. Autoinducers are typically synthesized from S-adenosylmethionine (SAM) and can traverse the cell membrane, 2. Autoinducers can bind to receptors located in the membrane or cytoplasm, and 3. The AHL-QS pathway activates numerous genes that regulate various cellular processes, 4. Regulation of AHL-QS involves the generation of autoinducers through a positive feedback loop. In gram-negative bacterial species, multiple homologs of AHL-mediated QS system signal-receptor pairs have been identified, including LuxI/R, LasI/R, RhII/R, Afel/R, BtaI/R, EsaI/R, and TofI/R.<sup>12</sup> Some organisms, such as *E. coli*, *K. pneumoniae*, *Salmonella* spp., and *Shigella* spp., lack AHL synthase enzymes, and thus do not produce AHLs. However, they possess a LuxR homolog known as SdiA, which can bind to AHLs produced by other microorganisms and influence gene expression.<sup>13,14</sup>

In Gram-negative bacteria that utilize AHL as QS signals, signal transduction pathways are diverse. Among these, *P. aeruginosa*, which is the most extensively researched species, primarily comprises four QS systems: *lasR/lasI*, *rhlR/rhII*, *pqs*, and *iqs*. The first system is the *lasR/lasI* system, which is composed of the transcriptional activator LasR and acyl-homoserine lactone synthase LasI. *LasI* guides the synthesis of N-3-oxo-dodecanoyl-homoserine lactone (3-oxo-C12-HSL) and secretes it into the extracellular environment during active transport. At a certain threshold concentration, it binds to LasR, activating the transcription of virulence factors including alkaline protease, exotoxin A, and elastase, thereby enhancing the expression of virulence genes in *P. aeruginosa*.<sup>15</sup> The second system is the *rhlR/rhII* system, in which RhlR acts as a transcriptional regulatory factor, and *RhII* encodes an AHL synthase. This system produces homoserine lactones with a C4-HSL structure that can freely traverse the cell membrane and regulate the expression of numerous genes such as chitinase, cyanide, and pyocyanin.<sup>16</sup> In addition to typical AHLs, *P. aeruginosa* produces 2-heptyl-3-hydroxy-4(1H)-quinolone [Pseudomonas quinolone signal (PQS)] as a QS signal. PQS belongs to the 2-alkyl-4-quinolone (AQs) family<sup>17</sup> and represents the third recently discovered QS system specific to *P. aeruginosa*. The signaling molecules of the quinolone system possess antimicrobial activity and are insoluble in water.<sup>18</sup> The *pqs* system primarily functions through two signalling molecules, PQS and 2-heptyl-4-hydroxyquinoline (HHQ), both of which can bind to the LysR-type transcriptional regulator PqsR to form a PqsR-PQS (or PqsR-HHQ) complex, activating the expression of various virulence genes.<sup>19,20</sup> In addition to its role in QS regulation through binding to PqsR, PQS can independently regulate gene expression in a manner that does not rely on PqsR.<sup>21</sup> The PQS can link two systems: *lasR/lasI* and *rhlR/rhII*. On the one hand, *lasR/lasI* and *rhlR/rhII* control PQS production. In contrast, PQS influenced *lasR/lasI* and *rhlR/rhII*. The *iqs* system utilizes IQS as a QS signal molecule, and its structure has been identified as that of 2-(2-hydroxyphenyl)-thiazole-4-ethanal. The genes involved in *iqs* synthesis belong to the non-ribosomal peptide synthetase gene cluster, *ambBCDE*. Under low-phosphate conditions, mutations in *ambBCDE* lead to reduced production of PQS and C4-HSL signals, as well as decreased expression of virulence factors such as pyocyanin, rhamnolipids, and elastase.<sup>22</sup> In addition to the aforementioned four QS systems, another *P. aeruginosa* QS auxiliary system, namely the GacS/GacA system, plays a significant role in enhancing bacterial motility, releasing rhamnolipids, and promoting biofilm formation, among other functions<sup>23</sup>

## AI-2 QS System

### Synthesis of AI-2

The AI-2 QS system utilizes furanone borate diester compounds as signaling molecules, and this QS system is present in both Gram-negative and Gram-positive bacteria. The classical biosynthetic pathway of AI-2 is a part of the activated methyl cycle, where AI-2 is generated from methionine. It goes through four enzymatic reactions to produce AI-2. Methionine is converted into S-adenosylmethionine (SAM) by S-adenosylmethionine synthase (MetK). SAM served as a methyl donor to generate S-adenosylhomocysteine (SAH). The synthesis of AI-2 involves the conversion of SAH to

homocysteine, which can be accomplished through a one-step reaction using SAH hydrolase (SahH), or a two-step reaction that requires SAH nucleosidase (Pfs) and LuxS. LuxS catalyzes the cleavage of the sulfur-ether bond in SRH, yielding homocysteine and 4,5-dihydroxy-2,3-pentanedione (DPD). The former generates methionine to enter the methyl cycle, whereas the latter can be rearranged to produce two different conformations, (2S,4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran (S-THMF) and (2R,4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran (R-THMF), which can interconvert and serve as ligands for *Vibrio Harveyi* LuxP-type receptors and *Salmonella enterica* LsrB-type receptors.<sup>24</sup>

## The Perception of AI-2

The AI-2 mediated QS system exists in over 60 bacterial species, including *V. fischeri*, *E. coli*, *K. pneumoniae*, *Salmonella* Typhi, *Bacillus subtilis*, *Staphylococcus aureus*, and *Yersinia* Pestis.<sup>25</sup> To date, four types of AI-2 receptors have been discovered: periplasmic binding proteins homologous to LuxP and LsrB,<sup>26</sup> CahR-type receptors containing the dCache\_1 domain<sup>27</sup> and YeaJ-type AI-2 receptors containing the GAPES1 domain.<sup>28</sup>

LuxP and LsrB are two homologs of periplasmic binding proteins (PBPs), with only 11% nucleotide similarity. These two receptors bind to different chemical properties of AI-2 precursor molecules derived from DPD.<sup>29</sup> The LuxP receptor is only found in *Vibrio* species and belongs to the high-affinity substrate-binding protein family. It regulates the activity of the membrane-bound sensor histidine kinases (HKs) protein using AI-2, effectively modulating the transmission of phosphorylation signals and impacting downstream gene expression. When AI-2 is present at low concentrations, LuxQ undergoes phosphorylation and transmits a phosphorylation signal through LuxU to LuxO. Phosphorylated LuxO cooperates with the alternative sigma factor  $\sigma^{54}$  to activate sRNA transcription. sRNA forms a complex with the partner protein Hfq, leading to the disruption of LuxR, a transcriptional regulator protein that controls the transcription of the *lux* operon. This inhibition suppressed transcription of the *lux* operon. However, when AI-2 is present at high concentrations, it binds to the LuxP receptor protein and activates its phosphatase activity. This leads to the dephosphorylation of the phosphotransferase LuxU and the transcriptional regulator LuxO. Dephosphorylated LuxO relieves its inhibitory effect on the transcription activator LuxR, initiating the transcription of the bioluminescence operon and resulting in bioluminescence phenomenon.<sup>25</sup>

Unlike the LuxP receptor, which is found only in *S. enterica*, the LsrB receptor was initially identified in *Vibrio* species. It has also been found in members of other *Enterobacteriaceae*, *Rhizobiaceae*, *Bacillaceae*, and *Clostridia*.<sup>30,31</sup> The LsrB receptor, a high-affinity substrate-binding periplasmic protein and component of the ATP-binding cassette (ABC) transport system, can internalize AI-2.<sup>32,33</sup> When the extracellular accumulation of AI-2 reaches a threshold, AI-2 is recognized, binds to the LsrB receptor, and enters the cytoplasm through a heterodimeric transmembrane channel composed of LsrC and LsrD, a process energized by ATP hydrolysis by LsrA. Once in the cytoplasm, AI-2 is phosphorylated by the kinase LsrK, and is further processed AI-2 by LsrF, which can bind to the inhibitory protein LsrR of the *lsrACDBFGE* operon, releasing it from the *Plsr* promoter and enhancing the expression of the *lsr* operon, facilitating the rapid transport of extracellular AI-2<sup>34</sup>(Figure 1B).

Studies have shown that many bacteria lacking LuxP or LsrB receptors respond to AI-2, indicating the presence of unidentified AI-2 receptors. In recent years, two new types of AI-2 receptors, the CahR-type<sup>27,35</sup> and YeaJ-type,<sup>28</sup> have been discovered.

The CahR-type receptor is a newly discovered sensor of extracellular signal transduction proteins, in which AI-2 recognizes the dCache\_1 domain of the CahR-type receptor, thus activating downstream gene expression. Studies have shown that CahR-type AI-2 receptors are widely present in prokaryotes and regulate chemotaxis and biofilm formation in *P. aeruginosa* PAO1.<sup>27</sup>

The YeaJ-type AI-2 receptor is characterized by the presence of a GAPES1 domain and GGDEF domain responsible for the synthesis of cyclic di-GMP (c-di-GMP). The interaction of YeaJ and YedQ with AI-2 and host-derived bile components taurocholate and taurodeoxycholate leads to an increase in intracellular concentrations of c-di-GMP. This elevation in c-di-GMP levels subsequently enhances the abundance of the T3SS chaperone SicA. The binding of SicA to InvF, SipB, and SipC was diminished, leading to a decrease in the transcription levels of T3SS-1 genes, including *sopB*, *sopE2*, *sicA*, *sipB*, and *sipC*. Specifically, in *S. Typhi*, binding of AI-2 to the GAPES1 domain of the YeaJ-type receptor



induces c-di-GMP synthesis, thereby inhibiting the expression of T3SS-1 genes. Studies indicate that homologs of YeaJ are present in many species of the *Enterobacteriaceae* family.<sup>28</sup>

### AI-3 QS System

In recent years, research has shown that in *E. coli* mutants lacking the LuxS protein, which is incapable of synthesizing the AI-2 signaling molecule, a novel signaling molecule distinct from AI-2, termed AI-3, is produced to maintain inter-species communication (Figure 1C). However, the structure of AI-3 remains elusive. However, epinephrine (Epi) and norepinephrine (NE) can promote toxin secretion by enterohemorrhagic *E. coli*, indicating that the AI-3 type QS system regulates bacterial virulence through AI-3, epinephrine, and norepinephrine.<sup>36</sup> In 2020, AI-3 was identified as 3,6-dimethylpyrazin-2-ol (3,6-DPO), a small molecule derived from threonine metabolism. This discovery underscores the complexity of the QS system in bacteria and highlights the sophisticated mechanisms by which bacteria interact with their hosts and regulate virulence factors.<sup>37</sup>

## Other Signaling Molecules in Gram-Negative Bacteria

### Diffusible Signal Factors

The diffusible signal factor (DSF) family is an intriguing group of QS signals found in various Gram-negative bacteria. The first identified DSF was cis-11-methyl-dodecenoic acid (XcDSF), discovered in the plant pathogen *Xanthomonas campestris* pv. *campestris* (*Xcc*). Subsequently, other DSFs were discovered. DSF-mediated interspecies communication pathways typically fall into three distinct groups, represented by *Xcc*, *B. cenocepacia*, and *P. aeruginosa*.<sup>38</sup> In addition, DSF can mediate interspecies communication. DSF plays a crucial role in regulating biofilm formation, virulence, motility, toxin production, exopolysaccharide synthesis, and extracellular enzyme activity.

### $\alpha$ -Hydroxyketones

In recent years, a novel class of signaling molecules known as AHKs has been identified in human pathogens such as *Legionella pneumophila* and *Vibrio cholerae*. *V. cholerae* synthesizes cholera autoinducer (CAI)-1 (3-hydroxytridecan-4-one) through the *cqs* gene, while *L. pneumophila* produces autoinducer (LAI)-1 (3-hydroxypentadecan-4-one) via the *lqs* gene cluster. Both CAI-1 and LAI-1 are diffusible and volatile autoinducers that play crucial roles in regulating biofilm formation, motility, and pathogenicity.<sup>39</sup>

### Indole


Various bacteria and some plants produce significant amounts of indole, leading to its widespread presence in both prokaryotic and eukaryotic communities. Recently, indole has been demonstrated to function as an intercellular, interspecies, and interkingdom signaling molecule. Indole is stable in indole-producing bacteria, but many non-indole-producing bacteria and eukaryotes can modify or degrade indole through various oxygenases. Recently, indole has been shown to regulate antibiotic resistance and persistence formation in *E. coli*.<sup>40</sup> Indole stimulates biofilm formation in *P. aeruginosa* and alters gene expression in a manner opposite to that of AHL, downregulating virulence and QS-regulated phenotypes in *P. aeruginosa*. Numerous studies have reported that indole controls various bacterial phenotypes and exhibits anti-virulence properties against non-indole-producing pathogens. Possible anti-virulence mechanisms include the inhibition of bacterial QS, biofilm formation, toxin production, and various bacterial adhesion factors.<sup>41</sup>

## An Overview of Physiological Functions Mediated by QS Systems

The fact that QS can be perceived by a wide range of receptors from different species demonstrates its extensive involvement in the regulation of various physiological functions of bacteria, such as bioluminescence, motility, aggregation, biofilm formation, expression of virulence factors, bacterial escape, competition, antibiotic resistance, plasmid conjugation, spore formation, and colonization.(Figure 2 and Table 1)

## Regulation of Bacterial Motility, Chemotaxis, and Autoaggregation by QS Systems

In nature, bacteria exist in two primary life forms: a planktonic state and an attached state that forms biofilms on surfaces. Biofilm formation aids bacteria in resisting adverse external factors, and supports their growth and proliferation. *E. coli*



**Figure 2** Physiological Functions Mediated by QS System.

utilizes AI-2 as its sole QS signal, which mediates aggregation of *E. coli* through the induction of chemotactic responses. This process can be divided into three stages. Stage One does not rely on the chemotactic ability of the cells. *E. coli* expressing Ag43 initially form “seeding” aggregates through random collisions. Ag43 belongs to the autotransporter protein family, in which adjacent cells Ag43 can interact to form dimers. Since the physiological density of the cells is not high at this time, relying solely on random collisions for further growth is inefficient. Stage Two: The “seeding” aggregates formed in the initial stage secrete sufficient concentrations of AI-2, thereby gradient-mediated chemotaxis of individual cells towards the aggregate. The increase in the growth rate of the aggregates may be due to the increase in local cell density mediated by chemotaxis, which slows as the aggregates and motile cells reach equilibrium. Stage Three: As the concentration of AI-2 further increased, the high levels of AI-2 in the external environment gradually disrupted the chemotaxis of *E. coli* towards the aggregates, resulting in more separated cells within the aggregates than those connecting them, leading to the dissolution of the aggregates. Curli fibers are a major protein component of the *E. coli* biofilm matrix. Research indicates that AI-2 induced chemotaxis promotes early stage autoaggregation of *E. coli* mediated by curli fibers during low-temperature growth. This aggregation enhances the AI-2-mediated signal transduction, biofilm formation, and resistance to adverse conditions.<sup>105</sup>

## Regulation of Bacterial Environmental Stress Response by QS Systems

The ability of bacteria to resist stress is a crucial determinant of their survival in natural environments and represents a significant marker of their adaptability. Extensive research has shown that the QS system plays a vital role in regulating bacterial stress resistance. Studies have found that after the deletion of the *luxS* gene in *Porphyromonas gingivalis*, its survival rate significantly increased under conditions of 50°C, pH 9.0, and exposure to 0.35 mM hydrogen peroxide.<sup>106</sup>

**Table I** Informative Table of Different Gram-Negative Bacterial QS Systems and Function

Bacterium	QS systems	Virulence-related phenotypes	Resistance phenotypes	Reference
<i>Pseudomonas aeruginosa</i>	(1) LasI (S)-LasR (R)-HSL (s)	Expression of several virulence factors including: LasA protease, LasB elastase, Apr alkaline protease	Regulating biofilm formation influences resistance to most antibiotics	[42–44]
	(2) RhlI(S)-RhlR(R)-HSL(s)	Expression of several virulence factors including: LasB elastase, RhlAB rhamnolipids, Phz pyocyanin, Hen hydrogen cyanid	Regulating biofilm formation influences resistance to most antibiotics	[45,46]
	(3)PqsABCD(S)-PqsR (R)-Alkyl quinolones (PQS)	Expression of several virulence factors including: Phz pyocyanin, Hian hydrogen cyanidel, LecA lectin PQS may exert cytotoxicity on cells by targeting the inhibition of the cellular respiratory chain	Regulating biofilm formation influences resistance to most antibiotics	[47]
	(4)AmbBCDE(S)-IqsR (R)- integrated quorum sensing system(IQS)	Expression of several virulence factors including: pyocyanin, rhamnolipids and elastase	Regulating biofilm formation influences resistance to most antibiotics	[15]
Enterohemorrhagic <i>Escherichia coli</i>	(1) Unknown (S)-SdiA (R)-3OC8HSL (s)	Growth and division, Colonization, Adhesion and subversion of epithelial intestinal cells	Regulating biofilm formation influences resistance to most antibiotics Regulating the arcAB pathway of the efflux pump through SdiA affects Resistance to nalidixic acid, Chloramphenicol, Quinolones (ciprofloxacin, Norfloxacin)	[48–50]
	(2) LuxS (S)-LsrB (R)-AI-2 (s)	Motility, Adhesion and subversion of epithelial intestinal cells, Exotoxin, Persistent infection, Stress survival	Regulating biofilm formation influences resistance to most antibiotics	[51–54]
	(3) Unknown (S)-QseC (R)-AI-3 (s)	Motility, Adhesion and subversion of epithelial intestinal cells		[55,56]
	(4) Indole	Adhesion and subversion of epithelial intestinal cells, Persistent infection	Regulating biofilm formation influences resistance to most antibiotics	[57–59]
<i>Klebsiella pneumoniae</i>	(1) Unknown (S)-SdiA (R)- HSL (s)	Controlling fimbriae expression, biofilm formation		[60]
	(2) LuxS (S)-LsrB (R)-AI-2 (s)	Regulating the synthesis of LPS and PNAG and the structure of biofilms		[61]
	(3) QseB(S)-QseC (R)-AI-3 (s)	Biofilm formation, serum resistance		[62]
<i>Salmonella.Typhi</i>	(1) Unknown (S)-SdiA (R)-3OC8HSL (s)	Competitive advantage, Motility, Adhesion, Invasion, Resistance to complement, Persistent infection	Regulating biofilm formation influences resistance to most antibiotics Influencing resistance to Nisin through PhoPQ and PmrAB	[63–66]
	(2) LuxS (S)-LsrB (R)-AI-2 (s)	Stress survival (oxidative stress), Motility, Adhesion, Invasion		[67–69]
	(3) Unknown (S)-QseC (R)-AI-3 (s)	Motility, Invasion, Intracellular survival, Persistent infection	Regulating biofilm formation influences resistance to most antibiotics	[70,71]
	(4) Indole	Motility, Invasion	Influencing resistance to multi-antibiotics through the Phage-shock response and oxidative stress pathway mediated by OxyR	[72,73]
	(5) DSF	Invasion		[74,75]
<i>Burkholderia cepacea</i>	(1) CepI (S)-CepR (R)-C8HSL, C6HSL (s)	Expression of swarm motility genes and several virulence factors including: proteases, siderophores, toxins, antifungal agents and biofilm formation		[76]
	(2) CciI (S)-CciR (R)-C8HSL, C6HSL (s)	Expression of swarm motility genes and several virulence factors including: proteases, siderophores, toxins, antifungal agents and biofilm formation		[77]
<i>Vibrio Harveyi</i>	(1) LuxM (S)-LuxN (R)- 3OHC4HSL (s)	Expression of bioluminescence genes		[78]
	(2) LuxS (S)-LuxP (R)-AI-2 (s)	Expression of several virulence factors including: extracellular protease; swimming and swarming abilities		[79]
	(3) CqsA (S)-CqsS (R)-CAI-I (s)	Expression of bioluminescence genes, metalloprotease production, and type III secretion		[80]

(Continued)

**Table I** (Continued).

Bacterium	QS systems	Virulence-related phenotypes	Resistance phenotypes	Reference
<i>Vibrio Cholerae</i>	(1) LuxS (S)-LuxP (R)-AI-2 (s)	Expression of several virulence factors including: Biofilm formation and EPS production, host colonization	Aminoglycosides	[81,82]
	(2) CqsA(S)-CqsS (R)-CAI-1 (s)	Expression of several virulence factors including: Biofilm formation and EPS production, host colonization		[83]
<i>Acinetobacter</i> spp	(1) Abal (S)-AbaR (R)-3OHC12HSL (s)	Expression of virulence factors including biofilm formation	Influencing resistance to most antibiotics through Biofilm formation and VBNC formation	[84]
<i>Haemophilus influenzae</i>	(1) LuxS (S)-RbsB (R)-AI-2 (s)	Invasion, lipooligosaccharide composition, Persistent infection		[85–87]
<i>Campylobacter jejuni</i>	(1) AHL QS	Transmission ability		[88]
<i>Yersinia Pestis</i>	(2) LuxS (S)-unknown (R)-AI-2 (s)	Motility, Adhesion and invasion, Stress survival, Transmission ability, Exotoxin		[89–92]
	(1) Yspl(S)- YspR(R)-3OC6HSL, 3OC8HSL	Survival ability in fleas, Stress survival (Starvation stress), Growth and division Invasion		[93–95]
<i>Yersinia Enterocolitica</i>	(2) Ypel(S)-YpelR(R)-unknown	Survival ability during host infection, Survival ability in fleas, Adhesion, Invasion, Membrane modeling	[96,97]	
	(1) Yen(S)- YenR(R)-HSL(s)			Swarming motility and cell aggregation, Host cell attachment, Virulence Plasmid Maintenance (pYVe)
<i>Yersinia Pseudotuberculosis</i>	(1) YpsI(S)-YpsR(R)-HSL(s)	Swarming motility and cell aggregation, Survival ability, Damage host cells	[100,101]	
	(2) YtbI(S)-YtbR(R)-HSL(s)			
<i>Brucella melitensis</i>	(1) Unknown (S)-vjbR (R)- C12-HSL (2) Unknown (S)-BlxR(R)-	Motility, Survival ability in <i>Brucella</i> containing vacuoles (BCV), Adhesion, Intracellular replication, Immune evasion	Influencing rifampin resistance through the virB operon (virB7–11)	[102–104]

## Regulation of Bacterial Biofilms by QS Systems

The formation of biofilms is a result of cell density-dependent gene expression and provides an appropriate microenvironment for intercellular communication. The main events involved in biofilm formation include the initial attachment of planktonic cells to a surface, followed by cell differentiation, EPS secretion, maturation, and dispersal.<sup>107</sup> QS plays a significant regulatory role in coordinating biofilm formation across multiple species, although it may not be the primary regulatory factor, with varying effects from initial attachment to maturation. QS-induced alterations in cell surface characteristics contribute to biofilm phenotypic variation, cell aggregation, and the synthesis of extracellular polysaccharides or adhesins. As a component of biofilms, EPS offers microorganisms a stable environment to withstand environmental stresses such as high temperatures, elevated alcohol levels, and acidity. Additionally, the polysaccharides and proteins in EPS serve as nutritional sources for microbial growth and metabolism. Firstly, QS affects the physico-chemical properties of cells and enhances biofilm formation in the biofilm lifestyle. QS-regulated surfactants can increase cell hydrophobicity, promoting their adhesion to solid substrates.<sup>8</sup> Further findings indicate a direct correlation between QS and EPS production. In wastewater treatment, the exogenous addition of 5 nmol/L C6-HSL significantly boosted EPS secretion.<sup>108</sup> Other AHL molecules, such as 3-O-C6-HSL, C8-HSL, and C12-HSL, are also significantly related to EPS formation.<sup>109</sup> Therefore, microbial communities can be regulated by QS through EPS and biofilm formation.

The majority of bacterial infections in the human body are associated with biofilms, and the formation of biofilms is a critical factor contributing to the difficulty in treating and preventing the recurrence of clinical bacterial infections; AI-2 plays a crucial regulatory role during biofilm formation.<sup>6</sup> When the *luxS* gene in *Haemophilus influenzae*, responsible for the QS system, is lacking, its ability to form biofilms is significantly enhanced.<sup>85,110</sup> In *K. pneumoniae*, *luxS* mutants can form mature biofilms but with a reduced capacity for microcolony development, primarily during the early stages of



biofilm formation.<sup>61</sup> The regulation of biofilms by QS varies considerably among bacterial strains and potentially exhibits strain-specific characteristics.

## The Regulation of Bacterial Antibiotic Resistance by QS Systems

### QS Systems Collaborate with Biofilm to Mediate Antibiotic Resistance

The collective QS in conjunction with biofilms mediates the dissemination of antibiotic resistance. Studies have demonstrated that obstructing the QS system within biofilms further elucidates the interplay between QS resistance and QS.<sup>111</sup> Protective polymers on the surfaces of biofilms attenuate bacterial growth and respiration rates within biofilms, rendering natural and artificial antibiotics insufficient to completely eradicate infectious biofilm communities.<sup>112</sup> When mature bacterial biofilms are formed, the bacteria within the biofilm exhibit significantly increased resistance, with cells showing antibiotic resistance levels 10–1000 times higher than those of planktonic bacteria of the same species at similar growth stages.<sup>113</sup> Pathogenic bacteria within biofilms use QS mechanisms to activate virulence factors and to develop antibiotic resistance. Additionally, antibiotic-degrading enzymes produced within the bacterial biofilms and certain gene products contribute to bacterial resistance. Antibiotic-degrading enzymes acquire antibiotic resistance via QS as a stress response. Studies have shown that the  $\beta$ -lactamases produced by bacteria are critical factors in conferring resistance to  $\beta$ -lactam antibiotics in biofilms. The reduced permeability of antibiotics through biofilms is attributed to the potential binding of antibiotics to the structural components within the biofilm matrix, rather than a decrease in antibiotic diffusion within the biofilm matrix.<sup>114</sup>

### QS Systems Regulate the Active Efflux of Antibiotics

In addition to preventing drugs from entering the cell, bacteria can also actively pump drugs. Efflux pumps are transmembrane proteins that transport various toxic compounds, including antibiotics, in an energy-dependent manner across the bacterial membranes. Although all bacteria contain multiple efflux pumps, they are particularly important as mechanisms of antimicrobial resistance (AMR) in gram-negative bacteria. Efflux transport proteins are classified into six families, including members of the resistance-nodulation-division (RND) family, which exhibit the most clinically relevant levels of resistance in gram-negative bacteria. In Gram-negative bacteria, the signaling molecule AHL actively translocates through the cell membrane via the MexAB-OprM efflux pump. Some researchers have found that auto-inducer molecules can upregulate the expression of the multidrug resistance pump MexAB-OprM, leading to multidrug resistance in *P. aeruginosa*.<sup>115</sup> Overexpression of the QS regulator SdiA leads to increased expression of the AcrAB efflux pump, which also plays a role in the multidrug resistance efflux pump system in *E. coli*.<sup>116</sup> However, the expression levels of efflux pumps can also affect the signaling molecules. Some researchers have found that overexpression of the MexCD-OprJ multidrug resistance efflux pump can reduce the signaling molecule response in *P. aeruginosa*.<sup>117</sup> Furthermore, a previous study demonstrated that chemical induction of the MexEF-oprN efflux pump can inhibit QS in *P. aeruginosa*.<sup>118</sup> The notable contribution of RND efflux pumps to multidrug resistance, especially in clinically relevant gram-negative pathogens, offers a distinct opportunity to restore drug sensitivity by targeting the structure, function, and regulation of these transport proteins.

### QS Systems Regulate the Formation of Persistent Bacteria

Persistent bacteria represent a specific subset of bacterial cells that display transient antibiotic tolerance, are often characterized by reduced growth or growth arrest, and can resume growth after exposure to lethal stress. The presence of these persistent cells significantly contributes to the persistent and recurrent nature of stubborn bacterial infections and is closely linked to an increased risk of developing antibiotic resistance during the course of treatment. In *E. coli*, the *hipB* gene, which is associated with the toxin-antitoxin system, is involved in the formation of persisters related to exposure to nanoaluminum, triggering antibiotic persistence in *E. coli* through the quorum-sensing regulators LrsF and QseB.<sup>119</sup> Research has also found that substances secreted by *E. coli*, which can serve as QS signal molecules, such as indole, can lead to an increase in the persistence of *S. Typhi*.<sup>120</sup>

## The Regulation of Antibiotic Resistance Genes (ARGs) Expression by QS Systems

A study found that in *P. aeruginosa*, cis-2-unsaturated fatty acid signaling molecules are involved in inter-species communication, which can lead to biofilm formation and alterations in susceptibility to various antibiotics.<sup>121</sup> The LasI/R circuit in *P. aeruginosa* regulates the expression of ampC  $\beta$ -lactamase. After three days of exposure to imipenem, the expression of *ampC* in the *AlasI* strain was reduced by one-fourth compared to that in the wild-type strain.<sup>122</sup> Study also discovered that under antibiotic induction, the activity of AHLs signaling molecules can promote the expression of resistance genes, potentially leading to the gradual development of multidrug resistance in *A. baumannii*, causing bacteria to evolve into multidrug-resistant strains<sup>123</sup> Further comprehensive investigation is warranted to elucidate the influence of QS on the expression of ARGs.

## The Regulation of Horizontal Gene Transfer by QS Systems

In the process of bacterial biofilm formation involving QS, bacteria in biofilms can also transfer and spread antibiotic resistance genes through the conjugative transfer of plasmids.<sup>124</sup> The bonding efficiency of plasmids in the biofilm was 700 times that of suspended cells.<sup>125</sup> When Licht et al<sup>126</sup> studied the conjugational transfer of plasmids, they found that the conjugational transfer frequency between suspended cells was much lower than that in the biofilm system (4–50 times), and the conjugational transfer frequency of plasmids in the biofilm system peaked in a short time. In addition, the frequency of plasmid conjugative transfer was high when bacteria containing anti-plasmids were introduced into bacteria that formed biofilms. A study also demonstrated that bacterial QS facilitates conjugative transfer of the multidrug-resistant plasmid RP4 within bacterial biofilms, significantly affecting the dissemination and spread of antibiotic resistance genes<sup>127</sup>.

Furthermore, in a study by Wang Ting et al<sup>128</sup>, it was found that AI-2 acts as a self-inducer and enhances conjugation frequency by regulating the expression levels of conjugative transfer-related genes, thereby promoting conjugative transfer of tetracycline resistance genes in *Vibrio Parahaemolyticus*. Exogenous AI-2 can upregulate the expression of TEM-type enzymes in a *LsrR*-dependent manner, increasing antibiotic resistance in *E. coli* strains<sup>129</sup>.

### 2.5 The Regulation of Bacterial Virulence by QS Systems.

## QS Systems Regulate Virulence Through the Secretion Systems

The bacterial secretion system is a macromolecular complex that exists in the bacterial cell membrane. It is a complex transmembrane molecular machine that can provide a secretion pathway for a variety of bacterial effectors. Nine bacterial secretory systems, T1SS–T9SS, have been found to play important roles in bacterial survival and pathogenicity. Transcriptomic analysis of *P. aeruginosa* reveals that the Type 1 secretion System (T1SS) is positively regulated by the QS system, as the expression of its effector alkaline protease AprA depends on QS. The Type 2 secretion System (T2SS) is responsible for secreting folded proteins from the periplasm of Gram-negative bacteria<sup>130</sup>. The Xcp system in *P. aeruginosa* secretes QS-regulated virulence factors including elastase and exotoxins. Interestingly, the Xcp system itself is positively regulated by QS<sup>131</sup>. The QS system is directly associated with Type IV Secretion System (T4SS) in *Brucella*. In *Brucella*, the LuxR-type QS system is responsible for the virulence characteristics regulated by the *virB* operon, which encodes the T4SS controlled by *VjbR*<sup>132</sup>.

## QS Systems Regulate Virulence Phenotypes

A study revealed that deletion of the *luxS* gene in *Actinobacillus pleuropneumoniae* significantly reduced the transcription level of the virulence gene *apxIIA*. Mouse experiments confirmed a 96-fold decrease in LD50 for the *luxS* deletion strain, and bacterial loads in the peritoneal lavage fluid, spleen, and lung tissues of mice were also significantly reduced. These findings indicate that QS positively regulates the virulence of *A. pleuropneumoniae*<sup>133</sup>. However, the opposite experimental results were found in *H. influenzae*, where the *luxS* deletion strain exhibited enhanced invasiveness in NCI-H292 cells. Infection experiments in chinchillas demonstrated that the *luxS* deletion strain of *H. influenzae* displayed increased pathogenicity in chinchillas, with significantly higher levels isolated from the middle ear than the wild-type strain. Additionally, histopathological results showed stronger inflammation and increased goblet cell hyperplasia in mice infected with the *luxS* deletion strain of *H. influenzae* 10 d post-infection<sup>86</sup>. This indicated that the QS system regulates virulence differently in different bacterial strains.

## Discussion and Prospects

In summary, in recent decades, there has been a significant advancement in our understanding of QS. The regulation of physiological functions by QS in Gram-negative bacteria has been studied extensively. This review provides a brief overview of QS systems in gram-negative bacteria and summarizes the regulation of chemotaxis, biofilm formation, resistance, virulence, and other QS aspects. This helps us to understand the role of the QS system in the physiological regulation of gram-negative bacteria.

Currently, research on QS systems of gram-negative bacteria has mainly focused on AHL signaling molecules, and *P. aeruginosa* has become a model strain. In contrast, relatively little research has been conducted on the AI-2 QS system in Gram-negative bacteria, and many aspects of AI-2 regulation remain unclear, posing numerous questions worthy of further investigation: 1) There is limited research on the interaction of AI-2 with the host, and the role of AI-2 signaling molecules in the microbial-host interaction process remains unclear. 2) Existing studies on the interaction of AI-2 with the host are mainly confined to in vitro cell culture experiments, and there is an urgent need for animal experiments to understand whether AI-2 can exert harmful biological effects by coordinating microbial behavior and regulating microbial networks or individual bacteria. 3) As an inter-species signaling molecule, there is currently limited research on the role of AI-2 in bacterial–bacterial interactions. 4) With the development of advanced technologies, AI-2-related research should focus on the combined use of metagenomics, transcriptomics, proteomics, metabolomics, and other multiomics technologies to elucidate the regulatory role of AI-2. Addressing these issues can help researchers gain a better understanding of the microbial world and provide a theoretical basis for the prevention and control of related pathogenic bacteria as well as the development of antimicrobial drugs.

This article focuses on the regulatory role of QS in Gram-negative bacteria, but QS, as a ubiquitous form of communication in microbial communities, also exerts significant effects on Gram-positive bacteria and fungi. Gram-positive bacteria, for instance, regulate the production of toxins and extracellular enzymes through QS. *S. aureus*, for example, utilizes a QS system known as the Agr system to regulate the expression of toxins (such as  $\alpha$ -hemolysin) and other virulence factors. This enables the bacteria to coordinate the expression of virulence in response to population density during host infection, thereby enhancing their infectivity. In fungi, such as *Candida albicans*, virulence often depends on QS signals. When the population density increases, QS activates virulence-related genes, promoting the transition of the fungus from a yeast form to a filamentous form, thus enhancing its invasiveness. Furthermore, QS also plays a regulatory role in biofilm formation and antimicrobial resistance in both Gram-positive bacteria and fungi. Through QS, microorganisms can more efficiently regulate gene expression, cooperate in attacking the host, and improve their competitive survival. In the context of antibiotic and antifungal treatments, QS mechanisms may serve as new therapeutic targets. Therefore, a deeper understanding of the mechanisms of QS could help develop new strategies for combating bacterial and fungal infections, especially in the face of growing antimicrobial resistance.

## Author Contributions

All authors have made significant contributions to the work reported. Xiaobing Chu completed the conception of the article, data acquisition, initial drafting, and figure drawing, while Qiwen Yang completed the conception of the article, as well as the review and revision work. All authors have no objections to the current version of the manuscript, agree to submit the article to this journal, and take full responsibility for all aspects of the work.

## Funding

This work was funded by the National High Level Hospital Clinical Research Funding (2022-PUMCH-B-028 and PUMCH-C-060), National Key Research and Development Program of China (2021YFC2301002), and National Natural Science Foundation of China (82272380 and 82072318).

## Disclosure

The authors declare that they have no competing interests.

## References

1. Changxin Z, Changyong Z. Drug resistance analysis of gram-negative bacteria in our hospital in 2014. *Chin Licensed Pharmacist*. 2015;2015:1.
2. Chinet. Available from: <http://www.chinets.com/Data/AntibioticDrugFast>. Accessed 16, Feb 2024.
3. Ranjbar R, Alam M, Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Evid Based Nurs*. 2022;27:16.
4. Waters CM, Bassler BL. Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol*. 2005;21:319–346. doi:10.1146/annurev.cellbio.21.012704.131001
5. Kariminik A, Baseri-Salehi M, Kheirkhah B. Pseudomonas aeruginosa quorum sensing modulates immune responses: an updated review article. *Immunol Lett*. 2017;190:1–6. doi:10.1016/j.imlet.2017.07.002
6. Zhao X, Yu Z, Ding T. Quorum-sensing regulation of antimicrobial resistance in bacteria. *Microorganisms*. 2020;8:425. doi:10.3390/microorganisms8030425
7. Häffner N, Bär J, Dengler Haunreiter V, et al. Intracellular environment and agr system affect colony size heterogeneity of Staphylococcus aureus. *Front Microbiol*. 2020;11:1415. doi:10.3389/fmicb.2020.01415
8. Aleti G, Lehner S, Bacher M, et al. Surfactin variants mediate species-specific biofilm formation and root colonization in bacillus. *Environ Microbiol*. 2016;18:2634–2645. doi:10.1111/1462-2920.13405
9. Chong G, Kimyon O, Manefield M. Quorum sensing signal synthesis may represent a selective advantage independent of its role in regulation of bioluminescence in vibrio fischeri. *PLoS One*. 2013;8:e67443.
10. Raffa RB, Iannuzzo JR, Levine DR, et al. Bacterial communication (“quorum sensing”) via ligands and receptors: a novel pharmacologic target for the design of antibiotic drugs. *J Pharmacol Exp Ther*. 2005;312:417–423. doi:10.1124/jpet.104.075150
11. Vannini A, Volpari C, Gargioli C, et al. The crystal structure of the quorum sensing protein TraR bound to its autoinducer and target DNA. *EMBO J*. 2002;21:4393–4401. doi:10.1093/emboj/cdf459
12. Liu L, Zeng X, Zheng J, Zou Y, Qiu S, Dai Y. AHL-mediated quorum sensing to regulate bacterial substance and energy metabolism: a review. *Microbiol Res*. 2022;262:127102. doi:10.1016/j.micres.2022.127102
13. Sabag-Daigle A, Dyszel JL, Gonzalez JF, Ali MM, Ahmer BMM. Identification of sdiA-regulated genes in a mouse commensal strain of Enterobacter cloacae. *Front Cell Infect Microbiol*. 2015;5:47. doi:10.3389/fcimb.2015.00047
14. Case RJ, Labbate M, Kjelleberg S. AHL-driven quorum-sensing circuits: their frequency and function among the Proteobacteria. *ISME J*. 2008;2:345–349. doi:10.1038/ismej.2008.13
15. Lee J, Wu J, Deng Y. A cell-cell communication signal integrates quorum sensing and stress response | nature Chemical Biology. *Nat Chem Biol*. 2013.
16. Nadal Jimenez P, Koch G, Thompson JA, Xavier KB, Cool RH, Quax WJ. The multiple signaling systems regulating virulence in pseudomonas aeruginosa. *Microbiol Mol Biol Rev*. 2012;76:46–65. doi:10.1128/MMBR.05007-11
17. Dubern J-F, Diggle SP. Quorum sensing by 2-alkyl-4-quinolones in Pseudomonas aeruginosa and other bacterial species - molecular biosystems (RSC Publishing). *Mol Biosyst*. 2008;4(9):882. doi:10.1039/b803796p
18. Farrow JM, Pesci EC. Two distinct pathways supply anthranilate as a precursor of the Pseudomonas quinolone signal. *J Bacteriol*. 2007;189:3425–3433. doi:10.1128/JB.00209-07
19. Lin J, Cheng J, Wang Y, Shen X. The Pseudomonas quinolone signal (PQS): not just for quorum sensing anymore. *Front Cell Infect Microbiol*. 2018;8:230. doi:10.3389/fcimb.2018.00230
20. Kostylev M, Kim DY, Smalley NE, Salukhe I, Greenberg EP, Dandekar AA. Evolution of the Pseudomonas aeruginosa quorum-sensing hierarchy. *Proc Natl Acad Sci U S A*. 2019;116:7027–7032. doi:10.1073/pnas.1819796116
21. Rampioni G, Falcone M, Heeb S, et al. Unravelling the genome-wide contributions of specific 2-Alkyl-4-quinolones and PqsE to quorum sensing in Pseudomonas aeruginosa. *PLoS Pathog*. 2016;12:e1006029.
22. Lee J, Zhang L. The hierarchy quorum sensing network in Pseudomonas aeruginosa. *Protein Cell*. 2015;6:26–41. doi:10.1007/s13238-014-0100-x
23. Ventre I, Goodman AL, Vallet-Gely I, et al. Multiple sensors control reciprocal expression of Pseudomonas aeruginosa regulatory RNA and virulence genes. *Proc Natl Acad Sci U S A*. 2006;103:171–176. doi:10.1073/pnas.0507407103
24. Wang Y, Liu B, Grenier D, Yi L. Regulatory mechanisms of the LuxS/AI-2 system and bacterial resistance. *Antimicrob Agents Chemother*. 2019;63:e01186–19. doi:10.1128/AAC.01186-19
25. Ge C, Sheng H, Chen X, et al. Quorum sensing system used as a tool in metabolic engineering. *Biotechnol J*. 2020;15:e1900360.
26. Papenfort K, Bassler BL. Quorum sensing signal-response systems in gram-negative bacteria. *Nat Rev Microbiol*. 2016;14:576–588. doi:10.1038/nrmicro.2016.89
27. Zhang L, Li S, Liu X, et al. Sensing of autoinducer-2 by functionally distinct receptors in prokaryotes. *Nat Commun*. 2020;11:5371. doi:10.1038/s41467-020-19243-5
28. Li S, Sun H, Li J, et al. Autoinducer-2 and bile salts induce c-di-GMP synthesis to repress the T3SS via a T3SS chaperone. *Nat Commun*. 2022;13:6684. doi:10.1038/s41467-022-34607-9
29. Miller ST, Xavier KB, Campagna SR, et al. Salmonella typhimurium recognizes a chemically distinct form of the bacterial quorum-sensing signal AI-2. *Molecular Cell*. 2004;15:677–687. doi:10.1016/j.molcel.2004.07.020
30. Pereira CS, de Regt AK, Brito PH, Miller ST, Xavier KB. Identification of functional LsrB-Like autoinducer-2 receptors. *J Bacteriol*. 2009;191:6975–6987. doi:10.1128/JB.00976-09
31. Torcato IM, Kasal MR, Brito PH, Miller ST, Xavier KB. Identification of novel autoinducer-2 receptors in clostridia reveals plasticity in the binding site of the LsrB receptor family. *J Biol Chem*. 2019;294:4450–4463. doi:10.1074/jbc.RA118.006938
32. Bassler BL, Wright M, Silverman MR. Multiple signalling systems controlling expression of luminescence in Vibrio harveyi: sequence and function of genes encoding a second sensory pathway. *Mol Microbiol*. 1994;13:273–286. doi:10.1111/j.1365-2958.1994.tb00422.x
33. Jung SA, Chapman CA, Ng W-L. Quadruple quorum-sensing inputs control vibrio cholerae virulence and maintain system robustness. *PLoS Pathog*. 2015;11:e1004837.



34. Thijs IM, Zhao H, De Weerd A. The AI-2-dependent regulator LsrR has a limited regulon in salmonella typhimurium. *Cell Res.* 2010;20(8):966–969. doi:10.1038/cr.2010.104
35. Liu X, Liu Q, Sun S, et al. Exploring AI-2-mediated interspecies communications within rumen microbial communities. *Microbiome.* 2022;10:167. doi:10.1186/s40168-022-01367-z
36. Kim CS, Gatsios A, Cuesta S, et al. Characterization of autoinducer-3 structure and biosynthesis in *E. coli*. *ACS Cent Sci.* 2020;6:197–206. doi:10.1021/acscentsci.9b01076
37. Lallement C, Goldring WP, Jelsbak L. Global transcriptomic response of the AI-3 isomers 3,5-DPO and 3,6-DPO in salmonella typhimurium. *Arch Microbiol.* 2023;205(4):. doi:10.1007/s00203-023-03450-x
38. He Y-W, Deng Y, Miao Y, et al. DSF-family quorum sensing signal-mediated intraspecies, interspecies, and inter-kingdom communication. *Trends Microbiol.* 2023;31:36–50. doi:10.1016/j.tim.2022.07.006
39. Tiaden A, Spirig T, Hilbi H. Bacterial gene regulation by alpha-hydroxyketone signaling. *Trends Microbiol.* 2010;18:288–297. doi:10.1016/j.tim.2010.03.004
40. Lee HH, Molla MN, Cantor CR, Collins JJ. Bacterial charity work leads to population-wide resistance. *Nature.* 2010;467:82–85. doi:10.1038/nature09354
41. Lee J, Attila C, Cirillo SLG, Cirillo JD, Wood TK. Indole and 7-hydroxyindole diminish *Pseudomonas aeruginosa* virulence. *Microb Biotechnol.* 2009;2:75–90. doi:10.1111/j.1751-7915.2008.00061.x
42. Gambello MJ, Iglewski BH. Cloning and characterization of the *Pseudomonas aeruginosa* lasR gene, a transcriptional activator of elastase expression. *J Bacteriol.* 1991;173:3000–3009. doi:10.1128/jb.173.9.3000-3009.1991
43. Gambello MJ, Kaye S, Iglewski BH. LasR of *Pseudomonas aeruginosa* is a transcriptional activator of the alkaline protease gene (apr) and an enhancer of exotoxin A expression. *Infect Immun.* 1993;61:1180–1184. doi:10.1128/iai.61.4.1180-1184.1993
44. Passador L, Cook JM, Gambello MJ, Rust L, Iglewski BH. Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication. *Science.* 1993;260:1127–1130. doi:10.1126/science.8493556
45. Jensen V, Löns D, Zaoui C, et al. RhlR expression in *Pseudomonas aeruginosa* is modulated by the *Pseudomonas* quinolone signal via PhoB-dependent and -independent pathways. *J Bacteriol.* 2006;188:8601–8606. doi:10.1128/JB.01378-06
46. Jackowski JT, Szeplalusi Z, Wanner DA. Effects of *P. aeruginosa*-derived bacterial products on tracheal ciliary function: role of O2 radicals. *Am J Physiol Lung Cell Mol Physiol.* 1991;260(2):L61–L67. doi:10.1152/ajplung.1991.260.2.L61
47. Rieger B, Thierbach S, Ommer M, Dienhart FSV, Fetzner S, Busch KB. *Pseudomonas* Quinolone Signal molecule PQS behaves like a B Class inhibitor at the IQ site of mitochondrial complex I. *FASEB Bioadv.* 2020;2:188–202. doi:10.1096/fba.2019-00084
48. Dyszel JL, Soares JA, Swearingen MC, Lindsay A, Smith JN, Ahmer BMM. *E. coli* K-12 and EHEC genes regulated by SdiA. *PLoS One.* 2010;5(e8946):e8946. doi:10.1371/journal.pone.0008946
49. Kanamaru K, Kanamaru K, Tatsuno I, Tobe T, Sasakawa C. SdiA, an *Escherichia coli* homologue of quorum-sensing regulators, controls the expression of virulence factors in enterohaemorrhagic *Escherichia coli* O157:H7. *Mol Microbiol.* 2000;38:805–816. doi:10.1046/j.1365-2958.2000.02171.x
50. Sperandio V. SdiA sensing of acyl-homoserine lactones by enterohemorrhagic *E. coli* (EHEC) serotype O157:H7 in the bovine rumen. *Gut Microbes.* 2010;1:432–435. doi:10.4161/gmic.1.6.14177
51. Sperandio V, Mellies JL, Nguyen W, Shin S, Kaper JB. Quorum sensing controls expression of the type III secretion gene transcription and protein secretion in enterohaemorrhagic and enteropathogenic *Escherichia coli*. *Proc Natl Acad Sci U S A.* 1999;96:15196–15201. doi:10.1073/pnas.96.26.15196
52. Sperandio V, Torres AG, Girón JA, Kaper JB. Quorum sensing is a global regulatory mechanism in enterohaemorrhagic *Escherichia coli* O157:H7. *J Bacteriol.* 2001;183:5187–5197. doi:10.1128/JB.183.17.5187-5197.2001
53. Park H, Lee K, Yeo S, Shin H, Holzappel WH. Autoinducer-2 quorum sensing influences viability of *Escherichia coli* O157:H7 under osmotic and in vitro gastrointestinal stress conditions. *Front Microbiol.* 2017;8:1077. doi:10.3389/fmicb.2017.01077
54. González Barrios AF, Zuo R, Hashimoto Y, Yang L, Bentley WE, Wood TK. Autoinducer 2 controls biofilm formation in *Escherichia coli* through a novel motility quorum-sensing regulator (MqsR, B3022). *J Bacteriol.* 2006;188:305–316. doi:10.1128/JB.188.1.305-316.2006
55. Moreira CG, Sperandio V. The epinephrine/norepinephrine/autoinducer-3 interkingdom signaling system in *Escherichia coli* O157:H7. *Adv Exp Med Biol.* 2016
56. Sharma VK, Casey TA. *Escherichia coli* O157:H7 lacking the qseBC-encoded quorum-sensing system outcompetes the parental strain in colonization of cattle intestines. *Appl Environ Microbiol.* 2014;80:1882–1892. doi:10.1128/AEM.03198-13
57. Hirakawa H, Kodama T, Takumi-Kobayashi A, Honda T, Yamaguchi A. Secreted indole serves as a signal for expression of type III secretion system translocators in enterohaemorrhagic *Escherichia coli* O157:H7. *Microbiology.* 2009;155(Pt 2):541–550. doi:10.1099/mic.0.020420-0
58. Vega NM, Allison KR, Khalil AS, Collins JJ. Signaling-mediated bacterial persister formation. *Nat Chem Biol.* 2012;8:431–433. doi:10.1038/nchembio.915
59. Hirakawa H, Inazumi Y, Masaki T, Hirata T, Yamaguchi A. Indole induces the expression of multidrug exporter genes in *Escherichia coli*. *Mol Microbiol.* 2005;55(4):1113–1126. doi:10.1111/j.1365-2958.2004.04449.x
60. Pacheco T, Gomes AÉI, Siqueira NMG, et al. SdiA, a quorum-sensing regulator, suppresses fimbriae expression, biofilm formation, and quorum-sensing signaling molecules production in *Klebsiella pneumoniae*. *Front Microbiol.* 2021;12:597735. doi:10.3389/fmicb.2021.597735
61. Chen L, Wilksch JJ, Liu H, et al. Investigation of LuxS-mediated quorum sensing in *Klebsiella pneumoniae*. *J Med Microbiol.* 2020;69:402–413. doi:10.1099/jmm.0.001148
62. Lv J, Zhu J, Wang T, et al. The role of the two-component QseBC signaling system in biofilm formation and virulence of hypervirulent *Klebsiella pneumoniae* ATCC43816. *Front Microbiol.* 2022;13:817494. doi:10.3389/fmicb.2022.817494
63. Dyszel JL, Smith JN, Lucas DE, et al. *Salmonella enterica* serovar typhimurium can detect acyl homoserine lactone production by *Yersinia enterocolitica* in mice. *J Bacteriol.* 2010;192:29–37. doi:10.1128/JB.01139-09
64. Luiz de Freitas L, Aparecida Dos Santos CI, Carneiro DG, Dantas Vanetti MC. Nisin and acid resistance in *Salmonella* is enhanced by *N*-dodecanoyl-homoserine lactone. *Microb Pathogenesis.* 2020;147:104320. doi:10.1016/j.micpath.2020.104320
65. Abed N, Grépinet O, Canepa S, et al. Direct regulation of the pefl-srgC operon encoding the Rck invasin by the quorum-sensing regulator SdiA in *Salmonella Typhimurium*. *Mol Microbiol.* 2014;94:254–271. doi:10.1111/mmi.12738



66. Liu Z, Que F, Liao L, et al. Study on the promotion of bacterial biofilm formation by a Salmonella conjugative plasmid and the underlying mechanism. *PLoS One*. 2014;9:e109808.
67. Choi J, Park J, Ryu S. Repressed Quorum sensing by overexpressing LsrR Hampers Salmonella evasion from oxidative killing within macrophages. *J Microbiol Biotechnol*. 2010;20(12):1624–1629.
68. Soni KA, Jesudhasan PR, Cepeda M. Autoinducer AI-2 is involved in regulating a variety of cellular processes in salmonella typhimurium. *Foodborne Pathogens Dis*. 2008;5(2):147–153. doi:10.1089/fpd.2007.0050
69. Choi J, Shin D, Kim M, Park J, Lim S, Ryu S. LsrR-mediated quorum sensing controls invasiveness of Salmonella typhimurium by regulating SPI-1 and flagella genes. *PLoS One*. 2012;7:e37059.
70. Moreira CG, Weinshenker D, Sperandio V. QseC mediates salmonella enterica serovar typhimurium virulence in vitro and in vivo. *Infect Immun*. 2010;78:914–926. doi:10.1128/IAI.01038-09
71. Bustamante VH, Martínez LC, Santana FJ, Knodler LA, Steele-Mortimer O, Puente JL. HilD-mediated transcriptional cross-talk between SPI-1 and SPI-2. *Proc Natl Acad Sci U S A*. 2008;105:14591–14596. doi:10.1073/pnas.0801205105
72. Kohli N, Crisp Z, Riordan R, Li M, Alaniz RC, Jayaraman A. The microbiota metabolite indole inhibits Salmonella virulence: involvement of the PhoPQ two-component system. *PLoS One*. 2018;13:e0190613.
73. Vega NM, Allison KR, Samuels AN, Klempner MS, Collins JJ. Salmonella typhimurium intercepts Escherichia coli signaling to enhance antibiotic tolerance. *Proc Natl Acad Sci U S A*. 2013;110:14420–14425. doi:10.1073/pnas.1308085110
74. Bosire EM, Eade CR, Schiltz CJ, et al. Diffusible signal factors act through AraC-type transcriptional regulators as chemical cues to repress virulence of enteric pathogens. *Infect Immun*. 2020;88:e00226–20. doi:10.1128/IAI.00226-20
75. Chowdhury R, Pavinski Bitar PD, Keresztes I, Condo AM, Altier C. A diffusible signal factor of the intestine dictates Salmonella invasion through its direct control of the virulence activator HilD. *PLoS Pathog*. 2021;17:e1009357.
76. Huber B, Riedel K, Hentzer M. The cep quorum-sensing system of Burkholderia cepacia H111 controls biofilm formation and swarming motility. *Microbiology*. 2001;147(9):2517–2528. doi:10.1099/00221287-147-9-2517
77. Malott RJ, Baldwin A, Mahenthalingam E, Sokol PA. Characterization of the cciIR quorum-sensing system in Burkholderia cenocepacia. *Infect Immun*. 2005;73(8):4982–4992. doi:10.1128/IAI.73.8.4982-4992.2005
78. Bassler BL, Wright M, Showalter RE, Silverman MR. Intercellular signalling in Vibrio harveyi: sequence and function of genes regulating expression of luminescence. *Mol Microbiol*. 1993;9:773–786. doi:10.1111/j.1365-2958.1993.tb01737.x
79. Zhang Y-Q, Deng Y-Q, Feng J, et al. LuxS modulates motility and secretion of extracellular protease in fish pathogen Vibrio harveyi. *Can J Microbiol*. 2022;68:215–226. doi:10.1139/cjm-2021-0311
80. Henke JM, Bassler BL. Three parallel quorum-sensing systems regulate gene expression in vibrio harveyi. *J Bacteriol*. 2004;186:6902–6914. doi:10.1128/JB.186.20.6902-6914.2004
81. Carvalho A, Krin E, Korlowski C, Mazel D, Baharoglu Z. Interplay between sublethal aminoglycosides and quorum sensing: consequences on survival in V. cholerae. *Cells*. 2021;10:3227. doi:10.3390/cells10113227
82. Hsiao A, Ahmed AMS, Subramanian S, et al. Members of the human gut microbiota involved in recovery from Vibrio cholerae infection. *Nature*. 2014;515:423–426. doi:10.1038/nature13738
83. Zhu J, Mekalanos JJ. Quorum sensing-dependent biofilms enhance colonization in Vibrio cholerae. *Dev Cell*. 2003;5:647–656. doi:10.1016/S1534-5807(03)00295-8
84. Anbazhagan D, Mansor M, Yan GOS, Md Yusof MY, Hassan H, Sekaran SD. Detection of quorum sensing signal molecules and identification of an autoinducer synthase gene among biofilm forming clinical isolates of Acinetobacter spp. *PLoS One*. 2012;7(e36696):e36696. doi:10.1371/journal.pone.0036696
85. Armbruster CE, Pang B, Murrah K, et al. RbsB (NTHI\_0632) mediates quorum signal uptake in nontypeable Haemophilus influenzae strain 86-028NP. *Mol Microbiol*. 2011;82:836–850. doi:10.1111/j.1365-2958.2011.07831.x
86. Daines DA, Bothwell M, Furrer J, et al. Haemophilus influenzae luxS mutants form a biofilm and have increased virulence. *Microb Pathog*. 2005;39:87–96. doi:10.1016/j.micpath.2005.06.003
87. Armbruster CE, Hong W, Pang B, et al. LuxS promotes biofilm maturation and persistence of nontypeable Haemophilus influenzae in vivo via modulation of lipooligosaccharides on the bacterial surface. *Infect Immun*. 2009;77:4081–4091. doi:10.1128/IAI.00320-09
88. Moorhead SM, Griffiths MW. Expression and characterization of cell-signalling molecules in Campylobacter jejuni. *J Appl Microbiol*. 2011;110:786–800. doi:10.1111/j.1365-2672.2010.04934.x
89. Elvers KT, Park SF. Quorum sensing in Campylobacter jejuni: detection of a luxS encoded signalling molecule. *Microbiology*. 2002;148(Pt 5):1475–1481. doi:10.1099/00221287-148-5-1475
90. Reeser RJ, Medler RT, Billington SJ, Jost BH, Joens LA. Characterization of Campylobacter jejuni biofilms under defined growth conditions. *Appl Environ Microbiol*. 2007;73:1908–1913. doi:10.1128/AEM.00740-06
91. He Y, Frye JG, Strobaugh TP, Chen C-Y. Analysis of AI-2/LuxS-dependent transcription in Campylobacter jejuni strain 81-176. *Foodborne Pathog Dis*. 2008;5:399–415. doi:10.1089/fpd.2008.0106
92. Plummer PJ. LuxS and quorum-sensing in Campylobacter. *Front Cell Infect Microbiol*. 2012;2:22. doi:10.3389/fcimb.2012.00022
93. Kirwan JP, Gould TA, Schweizer HP, Bearden SW, Murphy RC, Churchill MEA. Quorum-sensing signal synthesis by the Yersinia pestis acyl-homoserine lactone synthase YspI. *J Bacteriol*. 2006;188:784–788. doi:10.1128/JB.188.2.784-788.2006
94. LaRock CN, Yu J, Horswill AR, Parsek MR, Minion FC. Transcriptome analysis of acyl-homoserine lactone-based quorum sensing regulation in Yersinia pestis [corrected]. *PLoS One*. 2013;8:e62337.
95. Ritzert JT, Minasov G, Embry R, Schipma MJ, Satchell KJF. The cyclic AMP receptor protein regulates quorum sensing and global gene expression in Yersinia pestis during planktonic growth and growth in biofilms. *mBio*. 2019;10:e02613–19. doi:10.1128/mBio.02613-19
96. Yu J, Madsen ML, Caruthers MD, et al. Analysis of autoinducer-2 quorum sensing in Yersinia pestis. *Infect Immun*. 2013;81:4053–4062. doi:10.1128/IAI.00880-13
97. Fitts EC, Andersson JA, Kirtley ML, et al. New insights into autoinducer-2 signaling as a virulence regulator in a mouse model of pneumonic plague. *mSphere*. 2016;1:e00342–16. doi:10.1128/mSphere.00342-16
98. Atkinson S, Sockett RE, Cámara M, Williams P. Quorum sensing and the lifestyle of Yersinia. *Curr Issues Mol Biol*. 2006;8:1–10.

99. Atkinson S, Chang C-Y, Sockett RE, Cámara M, Williams P. Quorum sensing in *Yersinia enterocolitica* controls swimming and swarming motility. *J Bacteriol.* 2006;188:1451–1461. doi:10.1128/JB.188.4.1451-1461.2006
100. Atkinson S, Chang C-Y, Patrick HL, et al. Functional interplay between the *Yersinia pseudotuberculosis* YpsRI and YtbRI quorum sensing systems modulates swimming motility by controlling expression of flhDC and fliA. *Mol Microbiol.* 2008;69:137–151. doi:10.1111/j.1365-2958.2008.06268.x
101. Atkinson S, Goldstone RJ, Joshua GWP, et al. Biofilm development on *Caenorhabditis elegans* by *Yersinia* is facilitated by quorum sensing-dependent repression of type III secretion. *PLoS Pathog.* 2011;7:e1001250.
102. Rambow-Larsen AA, Rajashekara G, Petersen E, Splitter G. Putative quorum-sensing regulator blxR of *Brucella melitensis* regulates virulence factors including the Type IV secretion system and flagella. *J Bacteriol.* 2008;190:3274–3282. doi:10.1128/JB.01915-07
103. Kleinman CL, Sycz G, Bonomi HR, Rodríguez RM, Zorreguieta A, Sieira R. ChIP-seq analysis of the LuxR-type regulator VjbR reveals novel insights into the *Brucella* virulence gene expression network. *Nucleic Acids Res.* 2017;45:5757–5769. doi:10.1093/nar/gkx165
104. Yang X, Wu T, Liu W, et al. Cell membrane components of *Brucella melitensis* play important roles in the resistance of low-level rifampicin. *PLoS Negl Trop Dis.* 2020;14:e0008888.
105. Laganenka L, Colin R, Sourjik V. Chemotaxis towards autoinducer 2 mediates autoaggregation in *Escherichia coli*. *Nat Commun.* 2016;7:12984. doi:10.1038/ncomms12984
106. Yuan L, Hillman JD, Progulsk-Fox A. Microarray analysis of quorum-sensing-regulated genes in *Porphyromonas gingivalis*. *Infect Immun.* 2005;73:4146–4154. doi:10.1128/IAI.73.7.4146-4154.2005
107. Mangwani N, Shukla SK, Rao TS, Das S. Calcium-mediated modulation of *Pseudomonas mendocina* NR802 biofilm influences the phenanthrene degradation. *Colloids Surf B Biointerfaces.* 2014;114:301–309. doi:10.1016/j.colsurfb.2013.10.003
108. Shuai J, Hu X, Wang B, et al. Response of aerobic sludge to AHL-mediated QS: granulation, simultaneous nitrogen and phosphorus removal performance. *Chin Chem Lett.* 2021;32:3402–3409. doi:10.1016/j.ccllet.2021.04.061
109. Ji Y, Yu H, Cao R, Xu X, Zhu L. Promoting the granulation process of aerobic sludge via a sustainable strategy of effluent reflux in view of AHL-mediated quorum sensing. *J Environ Manage.* 2022;303:114091. doi:10.1016/j.jenvman.2021.114091
110. Pang B, Armbruster CE, Foster G, Learman BS, Gandhi U, Swords WE. Autoinducer 2 (AI-2) production by nontypeable *Haemophilus influenzae* 86-028NP promotes expression of a predicted glycosyltransferase that is a determinant of biofilm maturation, prevention of dispersal, and persistence in vivo. *Infect Immun.* 2018;86:e00506–18. doi:10.1128/IAI.00506-18
111. Quorum-sensing regulation of antimicrobial resistance in bacteria - PMC. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7143945/>. Accessed 4, Jun 2024.
112. Pople IK, Bayston R, Hayward RD. Infection of cerebrospinal fluid shunts in infants: a study of etiological factors. *J Neurosurg.* 1992;77:29–36. doi:10.3171/jns.1992.77.1.0029
113. Davies D. Understanding biofilm resistance to antibacterial agents. *Nat Rev Drug Discov.* 2003;2:114–122. doi:10.1038/nrd1008
114. Wu H, Moser C, Wang H-Z, Hoiby N, Song Z-J. Strategies for combating bacterial biofilm infections. *Int J Oral Sci.* 2015;7:1–7. doi:10.1038/ijos.2014.65
115. Maseda H, Sawada I, Saito K, Uchiyama H, Nakae T, Nomura N. Enhancement of the mexAB-oprM efflux pump expression by a quorum-sensing autoinducer and its cancellation by a regulator, MexT, of the mexEF-oprN Efflux pump operon in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother.* 2004;48:1320–1328. doi:10.1128/AAC.48.4.1320-1328.2004
116. Yang S, Lopez CR, Zechiedrich EL. Quorum sensing and multidrug transporters in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 2006;103:2386–2391. doi:10.1073/pnas.0502890102
117. Alcalde-Rico M, Olivares-Pacheco J, Alvarez-Ortega C, Cámara M, Martínez JL. Role of the multidrug resistance efflux pump MexCD-OprJ in the *Pseudomonas aeruginosa* quorum sensing response. *Front Microbiol.* 2018;9:2752. doi:10.3389/fmicb.2018.02752
118. Kristensen R, Andersen JB, Rybtke M, et al. Inhibition of *Pseudomonas aeruginosa* quorum sensing by chemical induction of the MexEF-oprN efflux pump. *Antimicrob Agents Chemother.* 2024;68(2):e01387–23. doi:10.1128/aac.01387-23
119. Wang S, Zhao C, Xue B, et al. Nanoalumina triggers the antibiotic persistence of *Escherichia coli* through quorum sensing regulators *lrsF* and *qseB*. *J Hazard Mater.* 2022;436:129198. doi:10.1016/j.jhazmat.2022.129198
120. Van den Bergh B, Fauvart M, Michiels J. Formation, physiology, ecology, evolution and clinical importance of bacterial persisters. *FEMS Microbiol Rev.* 2017;41:219–251. doi:10.1093/femsre/fux001
121. An S, Murtagh J, Twomey KB, et al. Modulation of antibiotic sensitivity and biofilm formation in *Pseudomonas aeruginosa* by interspecies signal analogues. *Nat Commun.* 2019;10:2334. doi:10.1038/s41467-019-10271-4
122. The role of quorum sensing system in antimicrobial induced ampC expression in *Pseudomonas aeruginosa* biofilm - Zhao - 2015 - journal of basic microbiology - Wiley online library. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/jobm.201300987>. Accessed 4, Jun 2024.
123. Dou Y, Song F, Guo F, et al. *Acinetobacter baumannii* quorum-sensing signalling molecule induces the expression of drug-resistance genes. *Mol Med Rep.* 2017;15:4061–4068. doi:10.3892/mmr.2017.6528
124. Biswas R, Panja AS, Bandopadhyay R. Molecular mechanism of antibiotic resistance: the untouched area of future hope. *Indian J Microbiol.* 2019;59:254–259. doi:10.1007/s12088-019-00781-6
125. Król JE, Wojtowicz AJ, Rogers LM, et al. Invasion of *E. coli* biofilms by antibiotic resistance plasmids. *Plasmid.* 2013;70:110–119. doi:10.1016/j.plasmid.2013.03.003
126. Licht TR, Christensen BB, Krogfelt KA, Molin S. Plasmid transfer in the animal intestine and other dynamic bacterial populations: the role of community structure and environment. *Microbiology.* 1999;145(Pt 9):2615–2622. doi:10.1099/00221287-145-9-2615
127. Zhu L, Chen T, Xu L, et al. Effect and mechanism of quorum sensing on horizontal transfer of multidrug plasmid RP4 in BAC biofilm. *Sci Total Envi.* 2020;698:134236. doi:10.1016/j.scitotenv.2019.134236
128. Wang T, Y LC, Yao L, et al. Effects of signaling molecule AI-2 on conjugation transfer of tetracycline resistance genes in *Vibrio parahaemolyticus*. *J Food Saf Qual.* 2022;13(2):505-510.
129. Xue T, Yu L, Shang F, et al. Short communication: the role of autoinducer 2 (AI-2) on antibiotic resistance regulation in an *Escherichia coli* strain isolated from a dairy cow with mastitis. *J Dairy Sci.* 2016;99:4693–4698. doi:10.3168/jds.2015-10543
130. Hentzer M, Wu H, Andersen JB, et al. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J.* 2003;22:3803–3815. doi:10.1093/emboj/cdg366

131. Maura D, Hazan R, Kitao T, Ballok AE, Rahme LG. Evidence for direct control of virulence and defense gene circuits by the *Pseudomonas aeruginosa* quorum sensing regulator, MvfR. *Sci Rep*. 2016;6:34083. doi:10.1038/srep34083
132. Li P, Tian M, Bao Y, et al. Brucella rough mutant induce macrophage death via activating IRE1 $\alpha$  pathway of endoplasmic reticulum stress by enhanced T4SS secretion. *Front Cell Infect Microbiol*. 2017;7:422. doi:10.3389/fcimb.2017.00422
133. Li L, Zhou R, Li T, et al. Enhanced biofilm formation and reduced virulence of *actinobacillus pleuropneumoniae* luxS mutant. *Microb Pathog*. 2008;45:192–200. doi:10.1016/j.micpath.2008.05.008

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>