



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.


Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

Contents lists available at ScienceDirect

Brain, Behavior, and Immunity

journal homepage: www.elsevier.com/locate/ybrbi

Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: A community-based cohort study of 387,109 adults in UK



Mark Hamer^{a,*}, Mika Kivimäki^b, Catharine R. Gale^{c,d}, G. David Batty^b

^a Division of Surgery and Interventional Sciences, Faculty Medical Sciences, University College London, London, UK

^b Department of Epidemiology and Public Health, University College London, UK

^c MRC Lifecourse Epidemiology Unit, University of Southampton, UK

^d Lothian Birth Cohorts, Department of Psychology, University of Edinburgh, UK

ARTICLE INFO

Keywords:

Physical activity

Smoking

Obesity

Infection

Coronavirus

C-reactive protein

Population cohort

ABSTRACT

We conducted the first large-scale general population study on lifestyle risk factors (smoking, physical inactivity, obesity, and excessive alcohol intake) for COVID-19 using prospective cohort data with national registry linkage to hospitalisation. Participants were 387,109 men and women (56.4 ± 8.8 yr; 55.1% women) residing in England from UK Biobank study. Physical activity, smoking, and alcohol intake, were assessed by questionnaire at baseline (2006–2010). Body mass index, from measured height and weight, was used as an indicator of overall obesity. Outcome was cases of COVID-19 serious enough to warrant a hospital admission from 16-March-2020 to 26-April-2020. There were 760 COVID-19 cases. After adjustment for age, sex and mutually for each lifestyle factor, physical inactivity (Relative risk, 1.32, 95% confidence interval, 1.10, 1.58), smoking (1.42; 1.12, 1.79) and obesity (2.05; 1.68, 2.49) but not heavy alcohol consumption (1.12; 0.93, 1.35) were all related to COVID-19. We also found a dose-dependent increase in risk of COVID-19 with less favourable lifestyle scores, such that participants in the most adverse category had 4-fold higher risk (4.41; 2.52–7.71) compared to people with the most optimal lifestyle. C-reactive protein levels were associated with elevated risk of COVID-19 in a dose-dependent manner, and partly (10–16%) explained associations between adverse lifestyle and COVID-19. Based on UK risk factor prevalence estimates, unhealthy behaviours in combination accounted for up to 51% of the population attributable fraction of severe COVID-19. Our findings suggest that an unhealthy lifestyle synonymous with an elevated risk of non-communicable disease is also a risk factor for COVID-19 hospital admission, which might be partly explained by low grade inflammation. Adopting simple lifestyle changes could lower the risk of severe infection.

1. Introduction

For non-communicable disease outcomes, lifestyle risk factors have been consistently associated with morbidity, mortality and loss of disease-free years of life (Colpani et al., 2018; Schlesinger et al., 2020; Nyberg et al., 2020; Li et al., 2020). There are also population cohort data on possible adverse effects of poor lifestyle on serious respiratory infections. For example, physical inactivity and smoking appear to be independently associated with higher risk of community-acquired pneumonia and pneumonia mortality (Wang et al., 2014; Baik et al., 2000; Inoue et al., 2007; Hamer et al., 2019; Paulsen et al., 2017). Evidence for alcohol intake and diet on risk of respiratory infection are less clear (Hamer et al., 2019; Paulsen et al., 2017).

A better understanding of the links between lifestyle risk factors and

COVID-19 has obvious implications for prevention of severe outcomes and also in identifying characteristics of those people most at risk. We are, however, unaware of any existing data on the relation of lifestyle risk factors with COVID-19. Accordingly, we examined the association of lifestyle risk factors with new cases of COVID-19-hospitalisations in a general population-based cohort study.

2. Methods

2.1. Study population

We used data from UK Biobank, a prospective cohort study, the sampling and procedures of which have been well described (Sudlow et al., 2015). Baseline data collection took place between 2006 and

* Corresponding author at: Division of Surgery and Interventional Sciences, Faculty Medical Sciences, University College London, 43-45 Foley Street, London W1W 7TS, UK.

E-mail address: m.hamer@ucl.ac.uk (M. Hamer).

<https://doi.org/10.1016/j.bbi.2020.05.059>

Received 18 May 2020; Received in revised form 21 May 2020; Accepted 21 May 2020

Available online 23 May 2020

0889-1591/ © 2020 Elsevier Inc. All rights reserved.

2010 across twenty-two research assessment centres in the UK giving rise to a sample of 502,655 people aged 40 to 69 years (response rate 5.5%) (Sudlow et al., 2015). Ethical approval was received from the North-West Multi-centre Research Ethics Committee. The research was carried out in accordance with the Declaration of Helsinki of the World Medical Association, and participants gave informed consent. No specific ethical approval was required for the present analyses of anonymised data.

2.2. Lifestyle measures

Physical activity, smoking, and alcohol consumption were assessed by questionnaire at baseline. These characteristics have demonstrated face validity in the UK Biobank sample through their associations with mortality and cardiovascular disease (Said et al., 2018). Participants were categorised into never, previous, and current smokers. From information on the weekly intake of beer and cider (1 pint = 2 units), wines (1 standard glass = 2 units) and spirits (1 shot = 1 unit), we aggregated units of alcohol intake per week. Heavy alcohol intake was defined as ≥ 14 units in women and ≥ 21 units in men (Nyberg et al., 2020). Physical activity was assessed using the International Physical Activity Questionnaire (IPAQ) short form (Craig et al., 2003) that measures duration and frequency of moderate-to-vigorous physical activity (MVPA) from all domains in the last week. Meeting activity guidelines was defined as ≥ 150 min/week MVPA or ≥ 75 min/week vigorous PA (Nyberg et al., 2020). Body weight was measured using Tanita BC418MA scales and standing height using a Seca height measure. Body mass index (BMI) was calculated [weight (kilograms)/height (meters²squared] and categorised into standard groups: healthy weight < 25 ; overweight $25 - < 30$; obese ≥ 30 kg/m (Nyberg et al., 2020).

2.3. Covariates

During the clinic visit, data were collected via self-report for ethnicity (White, South Asian, Black, Chinese, other), educational attainment (college/degree; A-level; O-level; CSEs or equivalent; National vocational Qualifications/ Higher National Diploma or equivalent; other professional qualification; none), and self-reported physician diagnosed cardiovascular diseases (heart attack, angina, stroke) and diabetes. Hypertension was defined as elevated measured blood pressure ($\geq 140/90$ mmHg) and /or use of anti-hypertensive medication.

2.4. Ascertainment of hospitalisation for COVID-19

Provided by Public Health England, data on COVID-19 status covered the period from 16th March 2020, after which testing was restricted to those with symptoms in hospital (<http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=40100>). For the present analyses COVID-19 testing results up to 26th April 2020 were included. These data can be regarded as a proxy for hospitalisations for severe cases of the disease for England only; study members from Scotland and Wales were therefore omitted from our analytical sample. COVID-19 disease tests were performed on samples from combined nose/throat swabs, using real time polymerase chain reaction (RT-PCR) in accredited laboratories (NHS England and NHS Improvement).

2.5. Statistical analyses

Analyses were performed using SPSS Version 26. We assigned points to different levels of each lifestyle behaviour: smoking history (0 = never; 1 = past; 2 = current), physical activity (0 = meeting guidelines; 1 = active but below guideline; 2 = inactive), alcohol (0 = moderate intake within guidelines; 1 = never or very occasional; 2 = heavy intake exceeding guidelines), obesity (0 = healthy weight; 1 = overweight; 2 = obese). Thus, scores ranged from 0 (optimal) to 8 (worst). We fitted regression models to estimate relative risk (RR) and

95% confidence intervals for associations between lifestyle scores and COVID-19. Relative risks were first adjusted for age and sex, followed by education, ethnicity, diabetes, hypertension, and cardiovascular diseases. We calculated Population Attributable Fraction (PAF) using our mutually-adjusted effect estimates on lifestyle factors and COVID-19, and prevalence of adverse lifestyle from Health Survey for England (NHS Digital, 2018; NHS Digital, 2018) (a representative, population-based survey of adults aged 16 and over living in private households in England) to evaluate the proportion of severe COVID-19 cases that could be avoided if high-risk people adopted a healthier lifestyle:

$$PAF \text{ in group } j = p_j (RR_j - 1) / \left[1 + \sum_{i=1}^k P(RR_i - 1) \right]$$

$$PAF = \sum_{j=1}^K PAF_j$$

where p_i = proportion of the population in group i ; RR_i = rate ratio in group i ; k = number of non-reference risk groups.

3. Results

The analytical sample comprised 387,109 participants (56.2 ± 8.0 years; 55.1% women) who were alive up to 5th March 2020, and had available data on lifestyle exposures and covariates. Participants were largely white British (94.5%). Of the lifestyle factors, 33.5% exceeded alcohol intake guidelines, 23.5% were obese, 9.7% smokers, 17.8% physically inactive, and 4.9% had a diabetes diagnosis, 56.1% hypertension, and 5.2% cardiovascular disease. Around 0.2% ($N = 760$) of the sample were hospitalized with a COVID-19 infection during the follow-up period, and their risk profile was characterized as being male, older age, smokers, physically inactive, less highly educated, non-white ethnicity, and higher prevalence of cardiometabolic comorbidity (Table 1).

3.1. Lifestyle and COVID-19

There was a dose-dependent association between the risk of COVID-19 with worsening lifestyle scores, such that participants in the most unfavourable category had 4-fold higher risk ($RR = 4.41$; 95% CI, 2.52, 7.71) (Table 2). These associations were little attenuated after adjustment for covariates. Risk ratios adjusted for age, sex and mutually for each lifestyle factor were raised for physical inactivity (1.32; 1.10, 1.58), smoking (1.42; 1.12, 1.79), obesity (2.05; 1.68, 2.49) but not for

Table 1
Baseline characteristics of sample in relation to COVID-19.

	COVID-19 hospitalisation	
	No	Yes
Age (yrs)	56.4 \pm 8.0	57.1 \pm 9.0
Sex (% men)	44.8	55.3
Smokers	9.8	11.9
Physical inactivity	17.8	25.0
Moderate alcohol intake	36.2	28.6
Degree educated	32.8	26.7
White ethnicity	94.5	86.7
Diabetes	4.8	9.5
Hypertension	56.1	63.9
Cardiovascular disease	5.2	9.4
Body mass index (kg/m ²)	27.3 \pm 4.7	29.0 \pm 5.4
Waist-Hip ratio	0.87 \pm 0.1	0.91 \pm 0.1
Total cholesterol (mmol/l)	5.7 \pm 1.1	5.4 \pm 1.2
HDL cholesterol (mmol/l)	1.5 \pm 0.4	1.3 \pm 0.3
Glycated haemoglobin (mmol/mol)	35.9 \pm 6.5	38.0 \pm 8.8
C-reactive protein (log units)	0.98 \pm 0.64	1.12 \pm 0.68

Results are expressed as percentage or mean \pm SD.

Table 2
Combined and individual lifestyle behavioral risk factors in relation to COVID-19 hospitalisation (N = 387,109).

Total lifestyle score	CASES/N	Relative Risk (95% confidence interval)	
		Model 1	Model 2
0 (optimal)	13/19,776	1.0 (ref)	1.0 (ref)
1	55/52,053	1.58 (0.86, 2.59)	1.48 (0.81, 2.71)
2	142/77,861	2.73 (1.55, 4.81)	2.43 (1.38, 4.29)
3	163/87,998	2.76 (1.57, 4.85)	2.41 (1.37, 4.25)
4	160/75,123	3.12 (1.77, 5.49)	2.70 (1.53, 4.75)
≥ 5 (worst)	227/74,298	4.41 (2.52, 7.71)	3.73 (2.12, 6.54)
p-trend		<0.001	<0.001
Individual behaviours			
<i>Smoking</i>			
Never	354/214,828	(ref)	1.0 (ref)
Past	313/134,855	1.34 (1.15, 1.56)	1.36 (1.15, 1.59)
Current	93/37,426	1.45 (1.16, 1.83)	1.36 (1.08, 1.71)
<i>Physical activity</i>			
Sufficient	382/209,489	(ref)	1.0 (ref)
Insufficient	192/108,707	0.98 (0.83, 1.17)	0.99 (0.84, 1.18)
None	186/68,913	1.51 (1.27, 1.81)	1.38 (1.15, 1.64)
<i>Alcohol consumption</i>			
Below guideline	216/140,908	(ref)	1.0 (ref)
Rarely/never	304/116,389	1.88 (1.55, 2.24)	1.57 (1.31, 1.88)
Above guideline	240/129,812	1.23 (1.00, 1.45)	1.24 (1.03, 1.50)
<i>Body mass index</i>			
Healthy weight	166/131,162	(ref)	1.0 (ref)
Overweight	317/165,052	1.41 (1.16, 1.70)	1.32 (1.09, 1.60)
Obesity	277/90,895	2.28 (1.88, 2.77)	1.97 (1.61, 2.42)

Model 1 adjusted for age and sex.

Model 2 adjusted for age, sex, education, ethnicity, diabetes, hypertension, cardiovascular disease (heart attack, angina, or stroke).

heavy alcohol consumption (1.12; 0.93, 1.35) in relation to COVID-19 compared to optimal reference categories.

3.2. Population attributable fraction

Using the Health Survey for England prevalence estimates (17% for current smoking, 25% for ex-smoking, 27% for physical inactivity, 35% for overweight and 28% for obesity), the PAF for the three unhealthy lifestyle factors in combination was 51.4% (13.3% for smoking, 8.6% for physical inactivity, and 29.5% for overweight and obesity).

3.3. Inflammatory mechanisms

We further explored potential mechanisms, specifically if low grade inflammation might partly explain associations between adverse lifestyle and risk of COVID-19. Data on high sensitivity C-reactive protein (hsCRP), measured at baseline at least 10 years before possible infection, (Fry et al.) were available in a sub-sample of participants (n = 363,263). We observed an association between adverse lifestyle score and higher hsCRP levels (B = 0.10, 95% CI, 0.09, 0.11) after adjustment for age, sex, education, ethnicity, diabetes, hypertension, cardiovascular disease. hsCRP levels were associated with elevated risk of COVID-19 in a dose-dependent manner (Table 3). When the association between lifestyle score and COVID-19 was adjusted for hsCRP, the effect estimates were attenuated by 10–16 % suggesting a possible mediating effect (Table 3).

4. Discussion

The present study demonstrates associations between adverse lifestyle and higher risk of COVID-19 in a large community-dwelling cohort. The associations were not explained by taking into account covariates such as education, ethnicity and self-reported cardiometabolic diseases, although further adjustment for hsCRP did partially attenuate the association. Based on UK risk factor prevalence estimates,

Table 3
Lifestyle risk factors, C-reactive protein, and Hospital Admission for COVID-19 in A Sub-sample with Available Biomarkers (N = 363,263).

Lifestyle score	Relative Risk (95% confidence interval)	
	Model 1 *	Model 2 †
0 (optimal)	1.0 (ref)	1.0 (ref)
1	1.46 (0.78, 2.74)	1.41 (0.75, 2.65)
2	2.44 (1.35, 4.40)	2.30 (1.27, 4.16)
3	2.44 (1.39, 4.39)	2.26 (1.25, 4.08)
4	2.77 (1.54, 5.00)	2.52 (1.39, 4.55)
≥ 5 (worst)	3.74 (2.09, 6.72)	3.30 (1.83, 5.95)
<i>C-reactive protein quintile</i>		
≤ 0.55 mg/L	–	1.0 (Ref)
0.56 – 1.02 mg/L	–	1.18 (0.90, 1.54)
1.03 – 1.75 mg/L	–	1.32 (1.01, 1.71)
1.76 – 3.33 mg/L	–	1.48 (1.15, 1.92)
> 3.33 mg/L	–	1.47 (1.13, 1.91)

* Adjusted for age, sex, education, ethnicity, diabetes, hypertension, cardiovascular diseases.

† Additionally adjusted for high-sensitivity C-reactive protein.

unhealthy behaviors in combination accounted for up to 51% of the population attributable fraction of severe COVID-19.

Physical activity has been previously shown to protect against serious community acquired infections in population cohort studies (Wang et al., 2014; Baik et al., 2000; Inoue et al., 2007; Hamer et al., 2019; Paulsen et al., 2017). Other studies (Schwellnus et al., 2016; Spence et al., 2007; Matthews et al., 2002; Nieman et al., 2011) in athletic populations have described a “J” shaped association between exercise volume and infection with optimal protection at moderate levels of activity. In the present study, protective associations of physical activity on COVID-19 were observed even at relatively low levels of activity below the current guidelines (i.e., < 150 min moderate to vigorous activity) and no dose-response effect was observed for higher levels. There are plausible biological mechanisms explaining the immunological benefits of exercise (Schwellnus et al., 2016), for example, anti-inflammatory effects and beneficial effects on adaptive immune responses (Pascoe et al., 2014).

The existing evidence on obesity and infection have been mixed. Some data suggested BMI above 25 kg.m⁻² was protective against pneumonia mortality (Inoue et al., 2007; Hamer et al., 2019) whilst others have suggested that overweight and obesity was associated with higher risk of respiratory and skin infections whilst protective against viral and fungal infections (Harpsoe et al., 2016). In a large Norwegian cohort, overweight and obesity were associated with higher 30 day mortality risk after detection of blood borne bacterial infection (Paulsen et al., 2017). Our results suggested both overweight and obesity were risk factors for severe COVID-19 infection, consistent with emerging data in small clinical studies (Sattar et al., 2020). The potential mechanisms have been linked to immune hyper-reactivity, impaired metabolic responses, and the adverse effects of obesity on lung function, diminishing forced expiratory volume and forced vital capacity (Sattar et al., 2020).

We found only weak evidence for a link between excessive alcohol intake and COVID-19, which was attenuated to the null in models mutually adjusted for other behavioral risk factors. This is largely consistent with our previous work on alcohol and infectious disease mortality (Hamer et al., 2019). Interestingly, ‘none drinkers’ were at greater risk of COVID-19, which is likely non-causal as this group have often stopped drinking due to prescribed medication and underlying health conditions.

The role of low-grade inflammation in susceptibility to severe COVID-19 infection remains poorly understood. Our data suggests low grade inflammation was a risk factor for severe COVID-19, and partially explained links between lifestyle behaviors and infection. C-reactive protein is known to play an important role in immune function (Del

Giudice and Gangestad, 2018) thus the findings are plausible.

There are several caveats to our work. Some cases of COVID-19 could have been captured in patients originally hospitalized for reasons other than the infection. We did not capture COVID-19 infections treated outside hospital settings; rather, our outcome was people with the infection of sufficient severity to warrant in-patient care. The response rate to the original baseline survey in UK Biobank was 5.5%. As such, this is a select group: relative to the general population, the study sample is healthier and better educated. While this means that estimates of the occurrence of disease, including COVID-19, have little utility, because exposures range is wide and the study sample is large, risk factors associations are not affected (Batty et al., 2020). PAF reflects the prevalence of the risk factor in the population and the strength of its association with the outcome being considered; the core assumption is that the risk factor has a causal link to the outcome. As our results are based on observational data rather than an intervention, the present PAF-findings may overestimate of the proportion of COVID-19 hospitalisation that could be prevented by lifestyle change.

In conclusion, these data suggest that adopting simple lifestyle changes could lower the risk of severe COVID-19 infection.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

Funding: GDB is supported by the UK Medical Research Council (MR/P023444/1) and the US National Institute on Aging (1R56AG052519-01; 1R01AG052519-01A1); MK by the UK Medical Research Council (MR/R024227), US National Institute on Aging (NIH), US (R01AG056477), NordForsk (75021), and Academy of Finland (311492); There was no direct financial or material support for the work reported in the manuscript.

Acknowledgement: We thank the UK Biobank study member for their generosity in participating.

Access to data: This research has been conducted using the UK Biobank Resource under Application 10279 (<http://www.ukbiobank.ac.uk/>).

Contributions: MH and GDB generated the idea for the present paper, and with MK formulated an analytical plan; CRG prepared the data set; MH carried out all the data analyses and wrote the manuscript; All authors commented on an earlier version of the manuscript. MH will act as guarantors for this work. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Role of the funding source: The funders of the studies had no role in study design, data collection, data analysis, data interpretation, or report preparation. CRG/MH had full access to UK Biobank data. MH takes responsibility for the decision to submit the manuscript for publication.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bbi.2020.05.059>.

References

Colpani, V., Baena, C.P., Jaspers, L., van Dijk, G.M., Farajzadegan, Z., Dhana, K., Tielmans, M.J., Voortman, T., Freak-Poli, R., Veloso, G.G.V., Chowdhury, R., Kavousi, M., Muka, T., Franco, O.H., 2018. Lifestyle factors, cardiovascular disease

- and all-cause mortality in middle-aged and elderly women: a systematic review and meta-analysis. *Eur. J. Epidemiol.* 33 (9), 831–845.
- Schlesinger, S., Neuenschwander, M., Ballon, A., Nöthlings, U., Barbaresko, J., 2020. Adherence to healthy lifestyles and incidence of diabetes and mortality among individuals with diabetes: a systematic review and meta-analysis of prospective studies. *J. Epidemiol. Community Health* 74 (5), 481–487.
- Nyberg, S.T., Singh-Manoux, A., Pentti, J., Madsen, I.E.H., Sabia, S., Alfredsson, L., Björner, J.B., Borritz, M., Burr, H., Goldberg, M., Heikkilä, K., Jokela, M., Knutsson, A., Lallukka, T., Lindbohm, J.V., Nielsen, M.L., Nordin, M., Oksanen, T., Pejtersen, J.H., Rahkonen, O., Rugulies, R., Shipley, M.J., Sipilä, P.N., Stenholm, S., Suominen, S., Vahtera, J., Virtanen, M., Westerlund, H., Zins, M., Hamer, M., Batty, G.D., Kivimäki, M., 2020. Association of healthy lifestyle with years lived without major chronic diseases. *JAMA Int. Med.* <https://doi.org/10.1001/jamainternmed.2020.0618>. [Epub ahead of print].
- Li, Y., Schoufour, J., Wang, D.D., Dhana, K., Pan, A., Liu, X., Song, M., Liu, G., Shin, H.J., Sun, Q., Al-Shaar, L., Wang, M., Rimm, E.B., Hertzmark, E., Stampfer, M.J., Willett, W.C., Franco, O.H., Hu, F.B., 2020 Jan. Healthy lifestyle and life expectancy free of cancer, cardiovascular disease, and type 2 diabetes: prospective cohort study. *BMJ* 8 (368), 16669. <https://doi.org/10.1136/bmj.16669>.
- Wang, H.E., Baddley, J., Griffin, R.L., et al., 2014. Physical inactivity and long-term rates of community-acquired sepsis. *Prev. Med.* 65, 58–64.
- Baik, I., Curhan, G.C., Rimm, E.B., Bendich, A., Willett, W.C., Fawzy, W.W., 2000. A prospective study of age and lifestyle factors in relation to community-acquired pneumonia in US men and women. *Arch. Intern. Med.* 160 (20), 3082–3088.
- Inoue, Y., Koizumi, A., Wada, Y., et al., 2007. Risk and protective factors related to mortality from pneumonia among middle aged and elderly community residents: the JACC Study. *J. Epidemiol.* 17 (6), 194–202.
- Hamer, M., O'Donovan, G., Stamatakis, E., 2019. Lifestyle risk factors, obesity and infectious disease mortality in the general population: linkage study of 97,844 adults from England and Scotland. *Prev. Med.* 123, 65–70.
- Paulsen, J., Askim, A., Mohus, R.M., et al., 2017. Associations of obesity and lifestyle with the risk and mortality of bloodstream infection in a general population: a 15-year follow-up of 64 027 individuals in the HUNT Study. *Int. J. Epidemiol.* 46 (5), 1573–1581.
- Sudlow, C., Gallacher, J., Allen, N., et al., 2015. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* 12 (3), e1001779.
- Said, M.A., Verweij, N., van der Harst, P., 2018. Associations of combined genetic and lifestyle risks with incident cardiovascular disease and diabetes in the UK biobank study. *JAMA Cardiol.* 3 (8), 693–702.
- Craig, C.L., Marshall, A.L., Sjoström, M., Bauman, A.E., Booth, M.L., Ainsworth, B.E., Pratt, M., Ekelund, U.L., Yngve, A., Sallis, J.F., Oja, P., 2003. International physical activity questionnaire: 12-country reliability and validity. *Med. Sci. Sport Exerc.* 35 (8), 1381–1395.
- NHS England and NHS Improvement. COVID-19 virus testing in NHS laboratories. Accessed 5/3/20 <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/03/guidance-and-sop-covid-19-virus-testing-in-nhs-laboratories-v1.pdf>.
- NHS Digital. Health Survey for England 2018 Overweight and obesity in adults and children. Accessed 5/5/20 <https://files.digital.nhs.uk/52/FD7E18/HSE18-Adult-Child-Obesity-rep.pdf>.
- NHS Digital. Health Survey for England 2018 Adult's health-related behaviours. Accessed 5/5/20 <https://files.digital.nhs.uk/B5/771AC5/HSE18-Adult-Health-Related-Behaviours-rep-v3.pdf>.
- Fry, D., Almond, R., Moffat, S., Gordon, M., Singh, P. UK Biobank Biomarker Project: Companion document to accompany serum biomarker data Accessed 5/5/20 https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/serum_biochemistry.pdf.
- Schwelunus, M., Soligard, T., Alonso, J.M., et al., 2016. How much is too much? (Part 2) International Olympic Committee consensus statement on load in sport and risk of illness. *Br. J. Sports Med.* 50 (17), 1043–1052.
- Spence, L., Brown, W.J., Pyne, D.B., et al., 2007. Incidence, etiology, and symptomatology of upper respiratory illness in elite athletes. *Med. Sci. Sports Exerc.* 39 (4), 577–586.
- Matthews, C.E., Ockene, I.S., Freedson, P.S., Rosal, M.C., Merriam, P.A., Hebert, J.R., 2002. Moderate to vigorous physical activity and risk of upper-respiratory tract infection. *Med. Sci. Sports Exerc.* 34 (8), 1242–1248.
- Nieman, D.C., Henson, D.A., Austin, M.D., Sha, W., 2011. Upper respiratory tract infection is reduced in physically fit and active adults. *Br. J. Sports Med.* 45 (12), 987–992.
- Pascoe, A.R., Fiatarone Singh, M.A., Edwards, K.M., 2014. The effects of exercise on vaccination responses: a review of chronic and acute exercise interventions in humans. *Brain Behav. Immun.* 39, 33–41.
- Harpoe, M.C., Nielsen, N.M., Friis-Møller, N., et al., 2016. Body mass index and risk of infections among women in the Danish national birth cohort. *Am. J. Epidemiol.* 183 (11), 1008–1017.
- Sattar, N., McInnes, I.B., McMurray, J.J.V., 2020. Obesity a risk factor for severe COVID-19 infection: multiple potential mechanisms. *Circulation.* <https://doi.org/10.1161/CIRCULATIONAHA.120.047659>. [Epub ahead of print].
- Del Giudice, M., Gangestad, S.W., 2018. Rethinking IL-6 and CRP: why they are more than inflammatory biomarkers, and why it matters. *Brain Behav. Immun.* 70, 61–75.
- Batty, G.D., Gale, C.R., Kivimäki, M., Deary, I.J., Bell, S., 2020. Comparison of risk factor associations in UK Biobank against representative, general population based studies with conventional response rates: prospective cohort study and individual participant meta-analysis. *BMJ* 368, m131. <https://doi.org/10.1136/bmj.m131>.