

What healthcare leaders need to do to protect the psychological well-being of frontline staff in the COVID-19 pandemic

Neil Greenberg ,¹ Derek Tracy^{1,2}

Delivery of high-quality care is dependent on staff with the right skills, experience and motivation. National Health Service (NHS) staff are performing in extraordinary ways while being scrutinised by a watchful media that currently describes them as heroes and a presently enthusiastic public that regularly claps them.

Nonetheless, it is inevitable that exposure to significant stressors will lead to some experiencing mental health problems. Like the rest of the population, staff are likely to be anxious about the future while contending the challenges of societal shut-down, which makes it difficult to destress by having a drink with a friend, playing sport or being part of social gatherings. Some will no doubt worry about their exposure to COVID-19 that could directly endanger them and their families. News of unexplained higher mortality in BAME groups and inadequate hospital supply chains for personal protective equipment (PPE) may also cause disquiet.

At work, staff wearing burdensome PPE will find that patients die despite their best efforts. Distress resulting from feeling that a life could have been saved if only you knew more, had better equipment, experience or training, or if there were not so many patients to deal with may result in 'moral injury'. This describes the suffering that occurs when circumstances clash with one's moral or ethical code.¹ Our understanding of moral injury remains somewhat immature, but many people with moral injuries develop mental illnesses such as depression or post-traumatic stress disorder (PTSD).²

Leaders will know that protecting staff mental health forms part of the legal duty of care.^{3 4} However, 'presenteeism'⁵—people being at work but operating suboptimally because of health complaints—is a much bigger problem than absenteeism. In safety critical healthcare, making poor, slow or incorrect decisions can have disastrous consequences.

Preventive medicine provides a useful model for healthcare leaders. First, prevent ill-health onset (primary prevention); next, prevent progression by identifying and intervening against early indicators of impending ill-health (secondary prevention); and last, when illness occurs, provide early treatment to optimise rapid recovery (tertiary prevention).

With the above model in mind, leaders must provide staff with frank information about what lies ahead while simultaneously positively emphasising how important their roles are and the organisation's honest commitment to support them. This helps individuals make informed choices about their role suitability and to prepare cognitively, emotionally and practically. Additionally, provision of coping skills training ('psychological PPE') may help foster resilience, although evidence that it prevents longer term mental ill-health is lacking.

Stronger evidence shows that reinforcing social bonds, between colleagues and supervisors, is highly protective of mental health.⁶ Leaders should 'buddy up' shift staff and ensure end-of-shift reviews are conducted. Psychological debriefing techniques should not be used as they have been conclusively found to be unhelpful and may cause harm⁷; unsurprisingly, the National Institute for Health and Care Excellence (NICE) warn against their use. Staff require basic needs met: shift patterns, rest areas and suitable safety equipment, and leaders should ensure that up-to-date and accurate information on local and national supportive services are well advertised, using media such as posters, emails and handouts. While many 'wellness' approaches are in vogue, such as mindfulness or virtual yoga—and unlikely to cause harm—evidence of benefit is often slim.⁸

Commonly, people developing mental health difficulties fail to seek help. Secondary preventive measures require supervisors, and trained peers, to be alert for early signs of distress. Leaders should ensure that supervisors can have psychologically savvy supportive conversations as evidence shows this leads to teams performing better and taking less

sick leave.^{5 9} Yet equally, many feel more comfortable sharing concerns with their peers; indeed, such concerns may relate to their managers. Peer-supporters, properly trained and supervised, can help maintain staff resilience; one example, is the 'TRiM' Trauma Risk Management programme developed by the UK military and now used within the NHS.¹⁰ While not 'penicillin for trauma', it is evidenced to support traumatised staff, reduce sickness and facilitate access to professional care. Leaders should thus ensure that structured peer support is available for staff while noting that organisational mental health screening programmes are not effective.¹¹ There are many reasons for this, including concerns about being labelled as weak, having a negative impact on one's career and perceiving support as a tick-box exercise.

However, despite well-constructed primary and secondary prevention strategies, some staff will need professional assessment and care. The current crisis is likely to cause an increase in anxiety and depressive disorders; some also may develop PTSD. Leaders should ensure there is rapid access to 'frontline' mental health professionals (MHPs), who focus on helping staff to return to duty and thus conserve staffing levels. These MHPs should use the four PIES principles (proximity, immediacy, expectancy and simplicity), which 'de-medicalise' normal (even uncomfortable) responses and reinforce practical measures that can often be implemented by supervisors. Most staff experiencing early trauma symptoms will find that managed proactively, these settle-down spontaneously, and the use of PIES has been shown protect staff's mental health in the longer term.¹² However, NICE recommends active monitoring of trauma-exposed staff and a minority will undoubtedly need formal, rather than PIES, treatment.¹³ The earlier such treatment commences, the greater the likelihood that long term disability will be avoided.

We strongly encourage healthcare leaders to follow the preventive medicine evidence; ensure your staff are properly prepared for their role practically and psychologically, provide basic equipment and training, empower teams to support each other—with a particular focus on helping supervisors feeling confident to speak to team members about their mental health—and adopt a 'nip it in the bud' approach; have MHPs in a supportive and supervisory role with them providing early 'return to duty' focused care for those who need it; and lastly, ensure that

¹King's College London, London, UK

²Oxleas NHS Foundation Trust, Dartford, UK

Correspondence to

Professor Neil Greenberg, King's College London, London WC2R 2LS, UK; neil.greenberg@kcl.ac.uk

all trauma-exposed staff are actively monitored as the country begins its return to the new normal. An effective staff mental health plan fulfils legal requirements under duty of care and maximises the numbers of functional staff available to carry out life saving care while protecting the organisation's quality of care, staff morale and reputation.

Twitter Neil Greenberg @profngreenberg and Derek Tracy @derektracy1

Acknowledgements We would like to acknowledge all of the mental health and welfare team at the London Nightingale Hospital.

Contributors Both authors contributed to the development and editing of the manuscript and are content with the author order.

Funding The study was funded by Public Health Research Programme, grant number (NIHR/KCL/2020).

Disclaimer The views expressed are those of the authors and not necessarily those of the NHS, NIHR, Department of Health and Social Care, or Public Health England.

Competing interests Both NG and DT have provided input to the London Nightingale hospital mental health plan. NG runs a psychological health consultancy that provides resilience training for a wide range of organisations, including a few National Health Service (NHS) teams. The work was supported by the National Institute for Health Research (NIHR) Health Protection Research Unit in Emergency Preparedness and Response at King's College London, in partnership with Public Health England and in collaboration with the University of East Anglia and Newcastle University.

Patient consent for publication Not required.

Provenance and peer review Commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



To cite: Greenberg N, Tracy D. *BMJ Leader* Published Online First: [please include Day Month Year]. doi:10.1136/leader-2020-000273

Received 27 April 2020
Accepted 1 May 2020

BMJ Leader 2020;**0**:1–2.
doi:10.1136/leader-2020-000273

ORCID iD

Neil Greenberg <http://orcid.org/0000-0003-4550-2971>

REFERENCES

- 1 Greenberg N, Docherty M, Gnanapragasam S, et al. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ* 2020;368:m1211.
- 2 Williamson V, Stevelink SAM, Greenberg N. Occupational moral injury and mental health: systematic review and meta-analysis. *Br J Psychiatry* 2018;212:339–46.
- 3 Brooks SK, Rubin GJ, Greenberg N. Traumatic stress within disaster-exposed occupations: overview of

the literature and suggestions for the management of traumatic stress in the workplace. *Br Med Bull* 2019;129:25–34.

- 4 UKPT Society. Traumatic stress management guidance; 2014. <http://www.ukpts.co.uk/site/assets/UKPTS-Guidance-Docment-120614.pdf>
- 5 Biron C, Brun Jean-Pierre, Ivers H, et al. At work but ill: psychosocial work environment and well-being determinants of presenteeism propensity. *J Public Ment Health* 2006;5:26–37.
- 6 Tam CWC, Pang EPF, Lam LCW, et al. Severe acute respiratory syndrome (SARS) in Hong Kong in 2003: stress and psychological impact among frontline healthcare workers. *Psychol Med* 2004;34:1197–204.
- 7 Rose S, Bisson J, Wessely S. A systematic review of single-session psychological interventions ('debriefing') following trauma. *Psychother Psychosom* 2003;72:176–84.
- 8 Kearns MC, Ressler KJ, Zatzick D, et al. Early interventions for PTSD: a review. *Depress Anxiety* 2012;29:833–42.
- 9 Brooks SK, Dunn R, Sage CAM, et al. Risk and resilience factors affecting the psychological wellbeing of individuals deployed in humanitarian relief roles after a disaster. *J Ment Health* 2015;24:385–413.
- 10 Greenberg N, Langston V, Jones N. Trauma risk management (TRIM) in the UK armed forces. *J R Army Med Corps* 2008;154:124–7.
- 11 Dunn R, Brooks SK, Greenberg N. Psychological impact of traumatic events: guidance for trauma exposed organisations. *Occupational Health at Work* 2015;12:17–21.
- 12 Solomon Z, Shklar R, Mikulincer M. Frontline treatment of combat stress reaction: a 20-year longitudinal evaluation study. *Am J Psychiatry* 2005;162:2309–14.
- 13 National Institute for Health and Care Excellence (NICE). National Institute for Health and Care Excellence. Post-traumatic stress disorder. NG116, 2018. Available: <https://www.nice.org.uk/guidance/ng116>