



DATA NOTE

The genome sequence of the Arran brown, *Erebia ligea* (Linnaeus, 1758) [version 1; peer review: 2 approved]

Konrad Lohse ¹, Alex Hayward², Dominik R. Laetsch ¹, Roger Vila ³, Kay Lucek ⁴, Wellcome Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective, Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, UK

²College of Life and Environmental Sciences, Department of Biosciences, University of Exeter, Exeter, UK

³Institut de Biologia Evolutiva, CSIC - Universitat Pompeu Fabra, Barcelona, Spain, Spain

⁴Department of Environmental Sciences, University of Basel, Basel, Switzerland

V1 First published: 13 Oct 2022, 7:259
<https://doi.org/10.12688/wellcomeopenres.18115.1>
 Latest published: 13 Oct 2022, 7:259
<https://doi.org/10.12688/wellcomeopenres.18115.1>

Abstract

We present a genome assembly from an individual male *Erebia ligea* (Arran brown; Arthropoda; Insecta; Lepidoptera; Nymphalidae). The genome sequence is 506 megabases in span. The majority (99.92%) of the assembly is scaffolded into 29 chromosomal pseudomolecules, with the Z sex chromosome assembled. The complete mitochondrial genome was also assembled and is 15.2 kilobases in length.

Keywords

Erebia ligea, Arran brown, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life](#) gateway.

Open Peer Review

Approval Status

	1	2
version 1		
13 Oct 2022	view	view

1. **Markus Moest**, University of Innsbruck, Innsbruck, Austria

2. **Manuela Lopez Villavicencio** , Museum National d'Histoire Naturelle Réseau des bibliothèques du museum, Paris, France

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: **Lohse K:** Investigation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Hayward A:** Investigation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Laetsch DR:** Investigation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Vila R:** Investigation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Lucek K:** Writing – Original Draft Preparation;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (206194) and the Darwin Tree of Life Discretionary Award (218328). KL and DL are supported by an ERC grant (ModelGenomLand 757648). KL was also supported by a NERC fellowship (NE/L011522/1). AH is supported by a Biotechnology and Biological Sciences Research Council (BBSRC) David Phillips Fellowship (BB/N020146/1). RV was supported by the Spanish government through grant PID2019-107078GB-I00/MCIN/AEI/ 10.13039/501100011033.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2022 Lohse K *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Lohse K, Hayward A, Laetsch DR *et al.* **The genome sequence of the Arran brown, *Erebia ligea* (Linnaeus, 1758) [version 1; peer review: 2 approved]** Wellcome Open Research 2022, 7:259 <https://doi.org/10.12688/wellcomeopenres.18115.1>

First published: 13 Oct 2022, 7:259 <https://doi.org/10.12688/wellcomeopenres.18115.1>

Species taxonomy

Eukaryota; Metazoa; Ecdysozoa; Arthropoda; Hexapoda; Insecta; Pterygota; Neoptera; Endopterygota; Lepidoptera; Glossata; Ditrysia; Papilionoidea; Nymphalidae; Satyrinae; Satyrini; Erebiina; *Erebia*; *Erebia ligea* (Linnaeus, 1758) (NCBI:txid111903).

Background

The Arran brown, *Erebia ligea*, is one of the most widespread species of the genus *Erebia*, occurring from the Russian Kamchatka Peninsula and Japan in eastern Asia (Dubatolov *et al.*, 1998) to central and northern Europe (Kudrna *et al.*, 2015). Although the species takes its common name from the Isle of Arran in Scotland, where it was first recorded in 1803, the current and historic presence of this butterfly in the British Isles remains disputed (Salmon, 1995). The intraspecific phenotypic diversity present throughout the distribution of *E. ligea* has triggered the description of several subspecies (Dubatolov *et al.*, 1998; Warren, 1937; Zakharova & Tatarinov, 2016), however, a formal biogeographic assessment remains lacking.

E. ligea is characterised as a woodland species associated with clearings and meadows, and occurs at relatively low altitudes compared to most other *Erebia* butterflies (Kleckova *et al.*, 2014). Recorded host plants include a variety of grasses (Poaceae) and sedges (*Carex*, Cyperaceae). It is univoltine and in some northern localities it is recorded only every second year (Tolman & Lewington, 2008). Although *E. ligea* is considered a species of Least Concern according to the IUCN Red List (Europe) (van Swaay *et al.*, 2010), the species can be locally endangered (Fichefet *et al.*, 2008).

While the first karyotypic analysis suggested that male *Erebia ligea* from Finland have 29 chromosomes (Federley, 1938), Japanese individuals from Hokkaido were found to have only 28 chromosomes (Saitoh & Abe, 1997). These values are close to the most common and putatively ancestral chromosomal state for Lepidoptera ($n=31$; Robinson, 1971), although *Erebia* is one of the most karyologically diverse known genera of butterflies (Robinson, 1971; de Vos *et al.*, 2020).

Genome sequence report

The genome was sequenced from a single male *E. ligea* (Figure 1) collected from Borzont, Joseni, Harghita, Romania (latitude 46.664, longitude 25.317). A total of 34-fold coverage of Pacific Biosciences single-molecule circular consensus (HiFi) long reads and 63-fold coverage of 10X Genomics read clouds were generated. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data. Manual assembly curation corrected 47 missing/misjoins and removed 10 haplotypic duplications, reducing the assembly length by 3.59% and the scaffold number by 39.39%, and increased the scaffold N50 by 4.29%.

The final assembly has a total length of 506 Mb in 40 sequence scaffolds, with a scaffold N50 of 19.1 Mb (Table 1). The majority, 99.92%, of assembly sequence was assigned to 40 chromosomal-level scaffolds, representing 28 autosomes



Figure 1. Forewings and hindwings of the male *Erebia ligea* specimen from which the genome was sequenced. Dorsal (left) and ventral (right) surface view of wings from specimen RO_EE_997 (ilEreLige1) from Borzont, Joseni, Harghita, Romania, used to generate Pacific Biosciences, 10X genomics and Hi-C data.

Table 1. Genome data for *Erebia ligea*, ilEreLige1.2.

Project accession data	
Assembly identifier	ilEreLige1.2
Species	<i>Erebia ligea</i>
Specimen	ilEreLige1 (genome assembly, Hi-C)
NCBI taxonomy ID	NCBI:txid111903
BioProject	PRJEB42125
BioSample ID	SAMEA7523313
Isolate information	Male, whole organism
Raw data accessions	
PacificBiosciences SEQUEL II	ERR7141799
10X Genomics Illumina	ERR6745725-ERR6745728
Hi-C Illumina	ERR6745729-ERR6745732
Genome assembly	
Assembly accession	GCA_917051295.2
Span (Mb)	506
Number of contigs	78
Contig N50 length (Mb)	14.9
Number of scaffolds	40
Scaffold N50 length (Mb)	19.1
Longest scaffold (Mb)	22.7
BUSCO* genome score	C:97.9%[S:97.4%,D:0.5%,F:0.2%,M:1.9%,n:5,286

*BUSCO scores based on the lepidoptera_odb10 BUSCO set using v5.2.2. C= complete [S= single copy, D=duplicated], F=fragmented, M=missing, n=number of orthologues in comparison. A full set of BUSCO scores is available at <https://blobtoolkit.genomehubs.org/view/ilEreLige1.2/dataset/CAKAVA02/busco>.

(numbered by sequence length), and the Z sex chromosome (Figure 2–Figure 5; Table 2). The assembly has a BUSCO v5.2.2 (Manni *et al.*, 2021) completeness of 97.9% (single 97.4%, duplicated 0.5%) using the lepidoptera_odb10 reference set (n=5,286). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited.

Methods

Sample acquisition and nucleic acid extraction

A single male *E. ligea* specimen (ilEreLige1, genome assembly, HiC) was collected from Borzont, Joseni, Harghita, Romania (latitude 46.664, longitude 25.317) using a handnet by Konrad Lohse, Dominik Laetsch (both University of Edinburgh) and Alex Hayward (University of Exeter). The sample was identified

by Roger Vila (Institut de Biologia Evolutiva, Barcelona) and snap-frozen from live in a dry shipper.

DNA was extracted at the Scientific Operations Core, Wellcome Sanger Institute. The ilEreLige1 sample was weighed and dissected on dry ice with tissue set aside for Hi-C sequencing. Whole organism tissue was disrupted by manual grinding with a disposable pestle. Fragment size analysis of 0.01–0.5 ng of DNA was then performed using an Agilent FemtoPulse. High molecular weight (HMW) DNA was extracted using the Qiagen MagAttract HMW DNA extraction kit. Low molecular weight DNA was removed from a 200-ng aliquot of extracted DNA using 0.8X AMPure XP purification kit prior to 10X Chromium sequencing; a minimum of 50 ng DNA was submitted for 10X sequencing. HMW DNA was




Figure 2. Genome assembly of *Erebia ligea*, ilEreLige1.2: metrics. The BlobToolKit Snailplot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 506,397,422 bp assembly. The distribution of chromosome lengths is shown in dark grey with the plot radius scaled to the longest chromosome present in the assembly (22,722,498 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 chromosome lengths (19,149,538 and 12,368,103 bp), respectively. The pale grey spiral shows the cumulative chromosome count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the lepidoptera_odb10 set is shown in the top right. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/ilEreLige1.2/dataset/CAKAVA02/snail>.




Figure 3. Genome assembly of *Erebia ligea*, ilEreLige1.2: GC coverage. BlobToolKit GC-coverage plot. Scaffolds are coloured by phylum. Circles are sized in proportion to scaffold length. Histograms show the distribution of scaffold length sum along each axis. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/ilEreLige1.2/dataset/CAKAVA02/blob>.

sheared into an average fragment size between 12–20 kb in a Megaruptor 3 system with speed setting 30. Sheared DNA was purified by solid-phase reversible immobilisation using AMPure PB beads with a 1.8X ratio of beads to sample to remove the shorter fragments and concentrate the DNA sample. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

Sequencing

Pacific Biosciences HiFi circular consensus and 10X Genomics read cloud DNA sequencing libraries were constructed according to the manufacturers' instructions. Sequencing was performed by the Scientific Operations core at the WSI on Pacific Biosciences SEQUEL II (HiFi) and Illumina HiSeq X (10X) instruments. Hi-C data were also generated from remaining whole organism tissue of ilEreLige1 using the Arima v1 Hi-C kit and sequenced on an Illumina HiSeq X (10X) instrument.




Figure 4. Genome assembly of *Erebia ligea*, iEreLige1.2: cumulative sequence. BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/iEreLige1.2/dataset/CAKAVA02/cumulative>.

Genome assembly

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021); haplotypic duplication was identified and removed with purge_dups (Guan *et al.*, 2020). One round of polishing was performed by aligning 10X Genomics read data to the assembly with longranger align, calling variants with freebayes (Garrison & Marth, 2012). The assembly was then scaffolded with Hi-C

data (Rao *et al.*, 2014) using SALSA2 (Ghurye *et al.*, 2019). The assembly was checked for contamination and corrected using the gEVAL system (Chow *et al.*, 2016) as described previously (Howe *et al.*, 2021). Manual curation (Howe *et al.*, 2021) was performed using gEVAL, HiGlass (Kerpedjiev *et al.*, 2018) and Pretext. The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2021), which




Figure 5. Genome assembly of *Erebia ligea*, ilEreLige1.2: Hi-C contact map. Hi-C contact map of the ilEreLige1.2 assembly, visualised in HiGlass. Chromosomes are shown in size order from left to right and top to bottom. The interactive Hi-C map can be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/l/?d=L3267sjSyakmh-bayAgPg>.

Table 2. Chromosomal pseudomolecules in the genome assembly of *Erebia ligea*, ilEreLige1.2.

INSDC accession	Chromosome	Size (Mb)	GC%
OU785219.1	1	22.72	37.2
OU785220.1	2	22.11	37.3
OU785221.1	3	22.01	37.4
OU785223.1	4	21.42	37.3
OU785224.1	5	20.97	37.3
OU785225.1	6	20.77	37.1
OU785226.1	7	20.52	37.1
OU785227.1	8	20.11	37.4
OU785228.1	9	20.06	37.3
OU785229.1	10	19.31	37.4
OU785230.1	11	19.22	37.3
OU785231.1	12	19.15	37.2
OU785232.1	13	18.89	37.3
OU785233.1	14	18.5	37.4
OU785234.1	15	18.36	37.3

INSDC accession	Chromosome	Size (Mb)	GC%
OU785235.1	16	17.54	37.4
OU785236.1	17	17.22	37.7
OU785237.1	18	16.82	37.2
OU785238.1	19	16.82	37.4
OU785239.1	20	16.64	37.5
OU785240.1	21	15.51	37.3
OU785241.1	22	15.26	37.6
OU785242.1	23	12.47	37.7
OU785243.1	24	12.37	38.2
OU785244.1	25	11.94	37.5
OU785245.1	26	10.28	37.7
OU785246.1	27	9.22	37.8
OU785247.1	28	8.14	38.1
OU785222.1	Z	21.64	37.3
OU785248.1	MT	0.02	20.2
-	Unplaced	0.39	41.4

performed annotation using MitoFinder (Allio *et al.*, 2020). The genome was analysed and BUSCO scores generated within the BlobToolKit environment (Challis *et al.*, 2020). Table 3 contains a list of all software tool versions used, where appropriate.

Table 3. Software tools used.

Software tool	Version	Source
Hifiasm	0.12	Cheng <i>et al.</i>, 2021
purge_dups	1.2.3	Guan <i>et al.</i>, 2020
SALSA2	2.2	Ghurye <i>et al.</i>, 2019
longranger align	2.2.2	https://support.10xgenomics.com/genome-exome/software/pipelines/latest/advanced/other-pipelines
freebayes	1.3.1-17-gaa2ace8	Garrison & Marth, 2012
MitoHiFi	1	Uliano-Silva <i>et al.</i>, 2021
HiGlass	1.11.6	Kerpedjiev <i>et al.</i>, 2018
PretextView	0.1.x	https://github.com/wtsi-hpag/PretextView
BlobToolKit	3.0.5	Challis <i>et al.</i>, 2020

Data availability

European Nucleotide Archive: *Erebia ligea* (Arran brown). Accession number [PRJEB42125](#); <https://identifiers.org/ena.embl/PRJEB42125>.

The genome sequence is released openly for reuse. The *E. ligea* genome sequencing initiative is part of the Darwin Tree of Life (DTOL) project. All raw sequence data and the assembly have been deposited in INSDC databases. Raw data and assembly accession identifiers are reported in Table 1.

Author information

Members of the Wellcome Sanger Institute Tree of Life programme are listed here: <https://doi.org/10.5281/zenodo.6866293>.

Members of Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective are listed here: <https://doi.org/10.5281/zenodo.5746904>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.6125046>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.6418363>.

References

- Allio R, Somaker-Bastos A, Romiguier J, *et al.*: **MitoFinder: Efficient Automated Large-Scale Extraction of Mitogenomic Data in Target Enrichment Phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit - interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, *et al.*: **Haplotype-Resolved *de Novo* Assembly Using Phased Assembly Graphs with Hifiasm.** *Nat Methods.* 2021; **18**(2): 170–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chow W, Brugger K, Caccamo M, *et al.*: **gEVAL — a Web-Based Browser for Evaluating Genome Assemblies.** *Bioinformatics.* 2016; **32**(16): 2508–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- de Vos JM, Augustijn H, Båtscher L, *et al.*: **Speciation through Chromosomal Fusion and Fission in Lepidoptera.** *Philos Trans R Soc Lond B Biol Sci.* 2020; **375**(1806): 20190539.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dubatolov VV, Korshunov YP, Gorbunov PY, *et al.*: **A Review of the *Erebia ligea*-Complex (Lepidoptera, Satyridae) from Eastern Asia.** *Lepidoptera Science.* 1998; **49**(3): 177–93.
[Publisher Full Text](#)
- Federley H: **Chromosomenzahlen Finnländischer Lepidopteren: I. Rhopalocera.** *Hereditas.* 1938; **24**(4): 397–464.
- Fichet V, Barbier Y, Baugnée JY, *et al.*: **Papillons de Jour de Wallonie: (1985–2007).** 2008.
[Reference Source](#)
- Garrison E, Marth G: **Haplotype-Based Variant Detection from Short-Read Sequencing.** 2012; arXiv: 1207.3907.
[Publisher Full Text](#)
- Ghurye J, Rhie A, Walenz BP, *et al.*: **Integrating Hi-C Links with Assembly Graphs for Chromosome-Scale Assembly.** *PLoS Comput Biol.* 2019; **15**(8): e1007273.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and Removing Haplotypic Duplication in Primary Genome Assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Howe K, Chow W, Collins J, *et al.*: **Significantly Improving the Quality of Genome Assemblies through Curation.** *GigaScience.* 2021; **10**(1): g1aa153.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: Web-Based Visual Exploration and Analysis of Genome Interaction Maps.** *Genome Biol.* 2018; **19**(1): 125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kleckova I, Konvicka M, Klecka J: **Thermoregulation and Microhabitat Use in Mountain Butterflies of the Genus *Erebia*: Importance of Fine-Scale Habitat Heterogeneity.** *J Therm Biol.* 2014; **41**: 50–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kudrna O, Pennerstorfer J, Lux K: **Distribution Atlas of European Butterflies and Skippers.** Schwanfeld: Wiss. Verl. Peks. 2015.
[Reference Source](#)
- Manni M, Berkeley MR, Seppely M, *et al.*: **BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rao SS, Huntley MH, Durand NC, *et al.*: **A 3D Map of the Human Genome at Kilobase Resolution Reveals Principles of Chromatin Looping.** *Cell.* 2014; **159**(7): 1665–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Robinson R: **Lepidoptera Genetics**. Oxford: Pergamon Press, 1971.
[Reference Source](#)

Saitoh K, Abe A: **The Chromosomes of *Erebia ligea* Rishirizana**. *Nota Lepidopterol.* 1997; **20**(3/4): 326–29.
[Reference Source](#)

Salmon MA: **Further Observations of the *Erebia ligea* (Linnaeus) and Other Controversies**. *The Entomologist's Record and Journal of Variation.* 1995; **107**: 117.

Tolman T, Lewington R: **Collins Butterfly Guide: The Most Complete Guide to the Butterflies of Britain and Europe**. HarperCollins Publishers Ltd. 2008.
[Reference Source](#)

Uliano-Silva M, Nunes JGF, Krashenninnikova K, *et al.*: **marcelauliano/MitoHiFi:**

mitohifi_v2.0. 2021.

[Publisher Full Text](#)

van Swaay CAM, Cuttelod A, Collins S, *et al.*: **European Red List of Butterflies**. 2010; 47.

[Publisher Full Text](#)

Warren BCS: **On the Evolution of Subspecies, as Demonstrated by the Alternation of Variability Existing in the Subspecies of the Genus *Erebia* (Lepidoptera)**. *Zool J Linn Soc.* 1937; **40**(271): 305–23.

[Publisher Full Text](#)

Zakharova EY, Tatarinov AG: **Chrono-geographical Approach to Analysis of Variability of Bicyclic *Erebia ligea* (L.) (Lepidoptera: Satyridae) Species in the Urals**. *Contemp Probl Ecol.* 2016; **9**(3): 272–81.

[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 19 June 2023

<https://doi.org/10.21956/wellcomeopenres.20086.r60428>

© 2023 Lopez Villavicencio M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Manuela Lopez Villavicencio 

Museum National d'Histoire Naturelle Réseau des bibliothèques du musée, Paris, Île-de-France, France

This article reports the genome assembly of a male individual of the very widespread butterfly species *Erebia ligea*. The assembly included the autosomes, the Z sexual chromosome and the mitochondrial genome.

The article is clearly written and pleasant to read. It shows convincing evidences for a high-quality assembly based on BUSCO scores. The methods for genome assembly, quality test and Hi-C scaffolding are relevant and up-to-date. There are sufficient details of methods and materials provided to allow replication by others.

It is important to highlight that the assembly sequence was assigned to 40 chromosomal-level scaffolds, representing 28 autosomes plus the sexual Z chromosome and this is relevant because previous karyotypic analysis have found different chromosomal numbers for this species.

As I am interested in butterfly genomes, I would have loved to have other interesting information as genome-wide level of heterozygosity (estimated with jellyfish and genomescope).

Overall, I think the release of this well-assembled and annotated genome is a very useful contribution and I recommend the indexing of this article.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.**Reviewer Expertise:** Evolutionary biology, Genomics**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 14 December 2022

<https://doi.org/10.21956/wellcomeopenres.20086.r53486>

© 2022 Moest M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Markus Moest**

Department of Ecology, University of Innsbruck, Innsbruck, Austria

Lohse et al. present a high-quality reference genome for a common Eurasian butterfly species. The quality of the assembly is impressive and represents a very valuable resource for future studies on biogeography and population genomics of the species as well as speciation research in the genus *Erebia*.

I only have few minor comments/questions:

Background:

"relatively low altitudes"... I would suggest to provide an approximate altitudinal range or range of altitudinal zones that are inhabited by this species

"in some northern localities it is recorded only every second year"...is the entire population really only present every other year or should this just mean that the completion of development can take up to 2 years?

Genome Sequence Report:

"sequence was assigned to 40 chromosomal-level scaffolds, representing 28 autosomes (numbered by sequence length), and the Z sex chromosome"... this section and the fact that in the legend in Figure2 41 scaffolds are indicated confuses me. My interpretation of "chromosomal-level scaffolds" is 1 scaffold = 1 chromosome, but that does not seem to be the case here (40 scaffolds = 28 autosomes + 1 Z) - could well be that my interpretation is the problem here. The 40 scaffolds in the text and the 41 scaffolds in Figure2 are maybe due to including/excluding the mitochondrial scaffold but I think this is not explicitly stated.

Methods:

Some more details on the specific tissues that were used for extraction would be helpful - e.g. whether only thorax or muscle tissues were used, whether the abdomen and guts were excluded, etc. as this information is relevant in order to assess the risk of potential contamination. Moreover, the description of the DNA extraction procedure is a bit unclear. For example, was the DNA used for PacBio sequencing first size selected in the same way as for the 10X sequencing, how much DNA was used for PacBio and how was the tissue set aside for Hi-C processed?

"A **single male** *E. ligea* specimen (ilEreLige1, genome assembly, HiC) was collected from Borzont, Joseni, Harghita, Romania (latitude 46.664, longitude 25.317) using a handnet by **Konrad Lohse, Dominik Laetsch** (both University of Edinburgh) and **Alex Hayward** (University of Exeter)."...obviously an extraordinarily strong specimen...

General comment to Methods in data notes (not a comment to the authors!):

I think it would be useful to provide a link to a general but more detailed description of the - I assume standardised - sequencing methods used in the DToL project, if that's available. This would provide interested readers with some more details and background and the method sections in the notes can remain short, concise and focussed on deviations from the general protocol.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Evolutionary Ecology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.