

SHORT REPORT

Open Access

Ca²⁺-activated mitochondrial biogenesis and functions improve stem cell fate in Rg3-treated human mesenchymal stem cells



Taeui Hong^{1,2,3†}, Moon Young Kim^{1,3,4,5†}, Dat Da Ly^{1,2†}, Su Jung Park^{1,3}, Young Woo Eom^{4,5}, Kyu-Sang Park^{1,2*} and Soon Koo Baik^{1,3,4,5*}

Abstract

Although mitochondrial functions are essential for cell survival, their critical roles in stem cell fate, including proliferation, differentiation, and senescence, remain elusive. Ginsenoside Rg3 exhibits various biological activities and reportedly increases mitochondrial biogenesis and respiration. Herein, we observed that Rg3 increased proliferation and suppressed senescence of human bone marrow-derived mesenchymal stem cells. Osteogenic, but not adipogenic, differentiation was facilitated by Rg3 treatment. Rg3 suppressed reactive oxygen species production and upregulated mitochondrial biogenesis and antioxidant enzymes, including superoxide dismutase. Consistently, Rg3 strongly augmented basal and ATP synthesis-linked respiration with high spare respiratory capacity. Rg3 treatment elevated cytosolic Ca²⁺ concentration contributing to mitochondrial activation. Reduction of intracellular or extracellular Ca²⁺ levels strongly inhibited Rg3-induced activation of mitochondrial respiration and biogenesis. Taken together, Rg3 enhances capabilities of mitochondrial and antioxidant functions mainly through a Ca²⁺-dependent pathway, which improves the proliferation and differentiation potentials and prevents the senescence of human mesenchymal stem cells.

Keywords: Mesenchymal stem cells, Ginsenoside Rg3, Cellular senescence, Oxidative stress, Mitochondria

Introduction

Mitochondria are essential for energy metabolism, calcium homeostasis, cell survival, and apoptosis; however, their role in stem cell proliferation, differentiation, and senescence has not been fully elucidated. In a stem cell niche, quiescent cells have a low mitochondrial mass in a low oxygen environment [1]. These cells have less bioenergetic reliance on oxidative phosphorylation and

more glycolytic activity. Furthermore, superoxide production from mitochondrial activities inhibits stemness, negatively affecting stem cell fate [2]. Nevertheless, counteracting evidences demonstrate mitochondrial metabolism exhibits an essential role in stem cell fate and defense against senescence [3]. Active mitochondrial respiration is required to maintain quiescence and the differentiation ability of stem cells [4]. After mitochondria transfer, recipient bone marrow-derived mesenchymal stem cells (BMSCs) displayed increased oxidative phosphorylation, which enhanced proliferation and osteogenic differentiation [5]. Conversely, mitochondrial DNA mutations, lower mitochondrial content, and impaired

* Correspondence: qsang@yonsei.ac.kr; baiksk@yonsei.ac.kr

[†]Taeui Hong, Moon Young Kim and Dat Da Ly contributed equally to this work.

¹Mitohormesis Research Center, Yonsei University Wonju College of Medicine, 20 Ilsan-ro, Wonju, Gangwon-Do 26426, Republic of Korea
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

mitochondrial metabolism lead to stem cell senescence [6].

Rg3 isolated from *Panax ginseng* has various pharmacological actions, including antioxidant [7], anti-inflammatory [8], anti-cancer [9], neuroprotective [10], and anti-aging activities [11]. Rg3 increases insulin sensitivity and inhibits adipogenic differentiation associated with AMP-activated protein kinase (AMPK) activation and peroxisome proliferator-activated receptor γ (PPAR γ) inhibition, respectively [12, 13]. Rg3 inhibits oxidative stress by elevating catalase and superoxide dismutase (SOD) activities and decreasing xanthine oxidase activity [7]. Additionally, Rg3-related increases in mitochondrial biogenesis and respiration have been reported [14]. However, most of investigation about Rg3 actions had been performed in cancer cells, and little is known about its role in stem cells. In this study, we aimed to resolve the effect of Rg3-induced mitochondrial activation on the fate of human BMSCs.

Methods

Cell culture and reagents

Human BMSCs were purchased from American Type Culture Collection. Mononuclear cells were cultured with low-glucose Dulbecco's modified Eagle's medium (Gibco) containing 10% fetal bovine serum (Gibco) and 1% penicillin/streptomycin (Gibco) at 37 °C in a 5% CO₂. Cells with passages 3 or 4 were used for the experiments treated with ginsenoside Rg3 (Cayman Chemical) or 0.25% ethanol as a vehicle. The detailed information is described in Supplemental Methods.

Quantitative polymerase chain reaction (qPCR)

Total RNA was isolated from BMSCs and used to synthesize cDNA. Transcript levels were measured by QuantStudio™ 6 Real-Time PCR System (Applied Biosystems) using sequence-specific primers listed in supplementary figure 1. The cycle threshold (Ct) values of the target genes were normalized to control (peptidylprolyl isomerase A), and the relative amount was calculated by using the equation, $2^{-\Delta\Delta Ct}$.

Western blot analysis

The cells were lysed using RIPA lysis buffer, and lysates were separated by 10–15% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred to polyvinylidene fluoride membranes. The membranes were incubated overnight with primary antibodies, the detailed information of which is described in Supplemental Methods. After washing, the membranes were incubated for 1 h in horseradish peroxidase-conjugated secondary antibodies. The membranes were incubated in a developing solution (GE Healthcare), and signal was detected using Chemi-Doc System (Bio-rad).

Senescence-associated β -galactosidase (SA- β -gal) staining

SA- β -gal activity was assessed with a senescence β -gal staining kit (Cell Signaling Technologies) according to the manufacturer's instructions.

Adipogenic and osteogenic differentiation

After incubation with vehicle or Rg3 for 5 days, BMSCs were maintained for 2 or 3 weeks in either osteogenic or adipogenic medium. The osteogenic medium contains 10 mM β -glycerophosphate (Sigma), 0.2 mM ascorbic acid-2-phosphate (Sigma), and 100 nM dexamethasone (Sigma). After 14 days, Alizarin Red staining was performed and the absorbance at 540 nm was detected by spectrophotometer. The adipogenic medium contains 1 mM dexamethasone, 0.5 mM 1-methyl-3-isobutyl-xanthine (Sigma), 100 μ M indomethacin (Sigma), and 10 μ g/mL insulin (Sigma). After 21 days, Oil Red O staining was performed and the absorbance at 520 nm was detected.

Oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) measurement

OCR and ECAR were measured in BMSCs using the Seahorse XFe96 Analyzer (Agilent Technologies) in response to XF Cell Mito Stress Test Kit or Glycolysis Stress Test Kit (Agilent Technologies). Supplemental Experimental Procedures provides detailed information about OCR and ECAR measurement from BMSCs.

Cytosolic ROS measurement

BMSCs were loaded with 5 μ M DCFH₂DA (Invitrogen) for 20 min at 37 °C, and fluorescence intensity reflecting ROS production was measured at the excitation and emission wavelengths of 485 and 538 nm using a fluorescence-activated cell sorting (FACS Aria III, BD Biosciences, USA).

Live-cell Ca²⁺ imaging

BMSCs seeded on 12-mm coverslip were loaded with 5 μ M Fura-2/AM (Thermo Fisher Scientific) for 30 min at 37 °C. Then, cells were transferred to a perfusion chamber on an inverted microscope (IX73, Olympus). Fluorescence images with an illuminator (pe-340Fura; CoolLED) were captured at 510 nm with a CCD camera (Prime-BSI; Teledyne Photometrics), and the ratio of fluorescence intensities (F_{340}/F_{380}) reflecting intracellular Ca²⁺ was analyzed by MetaFluor 6.1 (Molecular Devices).

Statistical analysis

Student's *t* test was used to compare means of two data groups. One-way ANOVA was used to compare means of two or more data groups. All graphs and statistical analysis were performed using software (Prism version

6.0, GraphPad Software). Data are presented as means ± SEMs, and $P < 0.05$ was considered significant.

Results

Rg3 augments proliferation and reduces senescence in human BMSCs


Human BMSCs exposed to Rg3 for 5 days showed increased proliferation in a dose-dependent manner, with a shortened population doubling time (Fig. 1a). The expression of stemness genes, including *OCT4*, *NANOG*, and *SOX2*, tends to be increased by Rg3 treatment (Fig. 1b). As an indicator of senescence, β-galactosidase staining showed less positive (senescent) cells in the Rg3-treated cells than in the control cells (Fig. 1c). Rg3 treatment downregulated the transcriptional levels of *p21* and *p53*, which are important molecules involved in cell cycle arrest and senescence (Fig. 1d). To investigate differentiation potency, BMSCs were treated with vehicle or Rg3 for 5 days and then cultured in osteogenic and adipogenic differentiation media for 14 and 21 days, respectively. Rg3 increased Ca^{2+} deposition, as measured by Alizarin Red S staining (Fig. 1e). Osteogenic genes such as RUNX family transcription factor 2 (*RUNX2*) and alkaline phosphatase (*ALP*) were upregulated by Rg3 (Fig. 1f). Conversely, adipogenic differentiation, detected via Oil Red O staining, was attenuated by Rg3 (Fig. 1g). The expression of the adipogenic genes, *PPARγ* and CCAAT-enhancer-binding protein α (*C/EBPα*), was lower in Rg3-treated cells (Fig. 1h).


Rg3 reduces oxidative stress and increases mitochondrial biogenesis and metabolism

Reactive oxygen species (ROS) production, known to cause stem cell senescence, was reduced by 5 days of Rg3 treatment, both under basal and increased oxidative stress conditions (Fig. 2a). The expression of antioxidant enzymes, including mitochondrial SOD (*SOD2*) and catalase, was increased by Rg3 (Fig. 2b and Fig. S1A). As upstream regulators of mitochondrial biogenesis, PPARγ coactivator 1α (*PGC1α*) and mitochondrial transcription factor A (*Tfam*) were also significantly upregulated upon Rg3 exposure. The p70S6 kinase (*S6K*), a downstream mammalian target of rapamycin (mTOR), was activated by Rg3. Components of the respiratory chain, including succinate dehydrogenase B (*SDHB*) and ubiquinol-cytochrome c reductase complex 2 (*UQCRC2*), were upregulated by Rg3 (Fig. 2c). Mitochondrial oxygen consumption by BMSCs was markedly increased following 5 days of Rg3 treatment, as was basal and maximal respiration, ATP-linked consumption, spare capacity, and proton leak (Fig. 2d). Glycolytic activity was also augmented, and the glycolysis reserve was diminished by Rg3 (Fig. 2e). Activation of mitochondrial respiration and glycolysis by Rg3 shifted the metabolic phenotype of BMSCs from a quiescent to an energetic state (Fig. 2f).

Rg3 induces mitochondrial biogenesis and activation mediated by Ca^{2+} increase

To identify the molecular mechanism underlying mitochondrial biogenesis and functional activation, we






performed a real-time measurement of intracellular Ca²⁺ concentration ([Ca²⁺]_i) in BMSCs loaded with Fura-2 (Fig. 3a). Acute exposure to Rg3 increased [Ca²⁺]_i under either 1.5 mM extracellular Ca²⁺ (Fig. 3b) or Ca²⁺ free (Fig. S2A) condition. Mitochondrial respiration was dose-dependently increased by Rg3 over 1 h treatment (Fig. 3c). Rg3 activated AMPK and cAMP response element-binding protein (CREB), which are involved in Ca²⁺ activated signaling (Fig. S1B and S1C). Mitochondrial biogenesis and activation by acute (1 h) and chronic (5 days) Rg3 incubation were abrogated by intracellular Ca²⁺ chelation (Fig. S2B) or by reducing extracellular Ca²⁺ with 0.4 mM EGTA (Fig. 3d), respectively. Upregulation of mitochondrial proteins, including Tfam and SDHB, by Rg3 was abolished by extracellular EGTA addition (Fig. 3e, f).

Discussion

Mitochondria are important bioenergetic organelles, but functional consequences of mitochondrial activation in stem cells have been underestimated. Recent advances have shown that adequate mitochondrial function in stem cells is essential to maintain proliferation and differentiation abilities [2, 3]. Inhibition of mitochondrial fatty acid oxidation

or impaired clearance of dysfunctional mitochondria deteriorates stem cell function and fate [15, 16]. Here, we demonstrated that upregulated mitochondrial biogenesis and respiratory capacity by ginsenosides Rg3 are related to improved stem cell fates, including increased proliferation, reduced senescence, and facilitated osteogenic differentiation of human BMSCs. Mitochondrial activation via increased [Ca²⁺]_i by Rg3 leads to energetic and healthy mitochondria with reduced oxidative stress. Overall, [Ca²⁺]_i-mediated activation of mitochondrial metabolism and the antioxidant system improves stem cell fate in human BMSCs.

Previous reports on mitochondrial actions of Rg3 or ginseng extracts in myoblast cells are consistent with our findings in stem cells [14, 17]. Furthermore, Rg3 treatment protected lipopolysaccharide-induced mitochondrial dysfunction in hepatocytes by increasing mitochondrial biogenesis and functions via AMPK signaling [18]. We observed that Rg3 increased [Ca²⁺]_i by intracellular Ca²⁺ release and extracellular Ca²⁺ influx. This [Ca²⁺]_i rise was responsible for early and late increases in basal, maximal, and ATP synthesis-linked respiration. Rg3 activated AMPK and CREB at different time points, both of which are involved in Ca²⁺-activated upregulation of PGC1α and mitochondrial



biogenesis. Additionally, mitochondrial Ca^{2+} uptake increases activities of matrix dehydrogenases, oxygen consumption, and ATP synthesis [19]. Therefore, it is conceivable that increased mitochondrial respiration could be attributed to cytosolic and mitochondrial Ca^{2+} rises and their consequences on mitochondrial biogenesis.

Accumulating evidence suggests that oxidative stress in stem cells accelerates senescence, inhibits proliferation, and enhances adipogenic but reduces osteogenic differentiation [20, 21]. We demonstrated that Rg3 suppressed cytosolic ROS levels under both basal and H_2O_2 -treated conditions. Antioxidant defense mechanisms were reinforced by Rg3 incubation, as observed by increased SOD2 and catalase, which is compatible with the result of ROS measurement. While Rg3-induced increases in SOD activity have been previously reported, the molecular mechanisms underlying this response have not been fully elucidated [22]. It has been shown that PGC1 α activated antioxidant genes, including SOD and catalase [23, 24]. We suggest that Rg3 increases antioxidant enzyme expression with mitochondrial biogenesis, thereby attenuating senescence by suppressing oxidative stress.

In conclusion, Rg3 increases mitochondrial biogenesis and activities along with antioxidant effects, which are associated with improved stem cell functions in BMSCs. Of note, Rg3 induced intracellular Ca^{2+} elevation, mediating Rg3-induced mitochondrial upregulation. We observed that Rg3 increases mitochondrial fitness, as estimated by a bioenergetic health index (Fig. S2C), which indicates efficient ATP synthesis and ample reserve capacity [25]. While further investigations are necessary to demonstrate the molecular mechanisms underlying the causal relationship between mitochondrial metabolism and stemness maintenance in BMSCs, we nonetheless suggest that mitochondrial activation via appropriate Ca^{2+} elevation could be an effective strategy for stem cell maintenance and differentiation, aimed at increasing the number of healthy mitochondria and inducing a stronger defense against oxidative stress. The therapeutic preservation of well-functioning mitochondria and a robust antioxidant defense could benefit stem cell transplantation, as well as slow the aging process of endogenous stem cells.

Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1186/s13287-020-01974-3>.

Additional file 1: Supplementary Fig. S1. Rg3 activates cell signaling and mitochondrial biogenesis. (A) Transcriptional regulations of main regulator of mitochondrial biogenesis (PGC1 α) and antioxidant enzymes (SOD2 and catalase) by Rg3 were measured by quantitative PCR. (B, C) Effects of acute (1 h; B) and chronic (5 days; C) exposure of Rg3 on AMP-activated protein kinase (AMPK) and cAMP response element-binding protein (CREB) activation. Data are presented as means \pm SEMs, * $P < 0.05$ and ** $P < 0.01$. **Supplementary Fig. S2.** Ca²⁺ increase by Rg3 activates mitochondrial respiration. (A) Increased cytosolic Ca²⁺ concentration ([Ca²⁺]_i) under extracellular Ca²⁺ free condition. [Ca²⁺]_i measurement using fluorescence imaging system after Fura-2-AM dye loading. Rg3-induced [Ca²⁺]_i elevations under Ca²⁺ free and normal Ca²⁺ conditions were compared by the area under the curve (AUC) above the baseline. (B) Role of intracellular Ca²⁺ increase on acute Rg3-induced mitochondrial activation. BAPTA-AM (10 or 20 μ M) were pretreated for chelating intracellular Ca²⁺. Effects of intracellular Ca²⁺ reduction on mitochondrial oxygen consumption rate (OCR) changes by 1 h exposure of Rg3 were evaluated. (C) Rg3-induced augmentation of biological health index. Based on oxygen consumption rate (OCR) data in Fig. 2D, enhanced biological health index by Rg3 was calculated by the equation as follow; BHI = log₁₀[(ATP synthesis-linked) \times (Spare Respiratory capacity) / (Proton leak) \times (Non-mitochondrial respiration)]. Data are presented as means \pm SEMs, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ and **** $P < 0.0001$. **Supplementary Fig. S3.** Role of sustained Ca²⁺ elevation by Rg3 on stem cell fates. (DOCX 3166 kb)

Abbreviations

ALP: Alkaline phosphatase; AMPK: AMP-activated protein kinase; BMSCs: Bone marrow-derived mesenchymal stem cells; [Ca²⁺]_i: Intracellular Ca²⁺ concentration; C/EBP α : CCAAT-enhancer-binding protein α ; CREB: cAMP response element-binding protein; PGC1 α : PPAR γ coactivator 1 α ; PPAR γ : Peroxisome proliferator-activated receptor γ ; ROS: Reactive oxygen species; RUNX2: RUNX family transcription factor 2; SDHB: Succinate dehydrogenase B; SOD: Superoxide dismutase; S6K: p70S6 kinase; Tfam: Mitochondrial transcription factor A; UQCRC2: Ubiquinol-cytochrome c reductase complex 2

Acknowledgements

This work was supported by the Medical Research Center Program (2017R1A5A2015369) and Bio & Medical Technology Development Program (NRF-2020M3A9D8039920) from the Ministry of Science, ICT, and the Yonsei University Wonju Campus Future-Leading Research Initiative of 2018 (2018-62-0055).

Authors' contributions

Conceptualization by T.H., M.Y.K., K.P., and S.K.B.; performed experiments by T.H., D.D.L., and S.J.P.; data analysis by T.H., M.Y.K., D.D.L., S.J.P., Y.W.E., and K.P.; writing by T.H., M.Y.K., D.D.L., K.P., and S.K.B.; funding acquisition by K.P. and M.Y.K. All authors read and approved the final manuscript.

Funding

The research grants of Medical Research Center Program (2017R1A5A2015369) and Bio & Medical Technology Development Program (NRF-2020M3A9D8039920) from the Ministry of Science, ICT, Republic of Korea and the Yonsei University Wonju Campus Future-Leading Research Initiative of 2018 (2018-62-0055) were used in performing experiments, data analysis, and writing the manuscript.

Availability of data and materials

The authors agree to share data and materials related to this manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare that there are no conflicts of interest related to this work.

Author details

¹Mitohormesis Research Center, Yonsei University Wonju College of Medicine, 20 Il-san-ro, Wonju, Gangwon-Do 26426, Republic of Korea. ²Department of Physiology, Yonsei University Wonju College of Medicine, 20 Il-san-ro, Wonju, Gangwon-Do 26426, Republic of Korea. ³Department of Internal Medicine, Yonsei University Wonju College of Medicine, 20 Il-san-ro, Wonju, Gangwon-Do 26426, Republic of Korea. ⁴Cell Therapy and Tissue Engineering Center, Yonsei University Wonju College of Medicine, Il-san-ro 20, Wonju 26426, Gangwon-Do, Republic of Korea. ⁵Regeneration Medicine Research Center, Yonsei University Wonju College of Medicine, 20 Il-san-ro, Wonju 26426, Gangwon-Do, Republic of Korea.

Received: 12 August 2020 Accepted: 13 October 2020

Published online: 04 November 2020

References

- Takubo K, Nagamatsu G, Kobayashi CI, Nakamura-Ishizu A, Kobayashi H, Ikeda E, Goda N, Rahimi Y, Johnson RS, Soga T, et al. Regulation of glycolysis by Pdk functions as a metabolic checkpoint for cell cycle quiescence in hematopoietic stem cells. *Cell Stem Cell*. 2013;12(1):49–61.
- Papa L, Djedaini M, Hoffman R. Mitochondrial role in stemness and differentiation of hematopoietic stem cells. *Stem Cells Int*. 2019;4067162.
- Zhang H, Menzies KJ, Auwerx J. The role of mitochondria in stem cell fate and aging. *Development*. 2018;145(8):dev143420.
- Ansó E, Weinberg SE, Diebold LP, Thompson BJ, Malinge S, Schumacker PT, Liu X, Zhang Y, Shao Z, Steadman M, et al. The mitochondrial respiratory chain is essential for haematopoietic stem cell function. *Nat Cell Biol*. 2017; 19(6):614–25.
- Guo Y, Chi X, Wang Y, Heng BC, Wei Y, Zhang X, Zhao H, Yin Y, Deng X. Mitochondria transfer enhances proliferation, migration, and osteogenic differentiation of bone marrow mesenchymal stem cell and promotes bone defect healing. *Stem Cell Res Ther*. 2020;11(1):245.
- Stoll EA, Cheung W, Mikheev AM, Sweet IR, Bielas JH, Zhang J, Rostomily RC, Horner PJ. Aging neural progenitor cells have decreased mitochondrial content and lower oxidative metabolism. *J Biol Chem*. 2011;286(44):38592–601.
- Wei X, Su F, Su X, Hu T, Hu S. Stereospecific antioxidant effects of ginsenoside Rg3 on oxidative stress induced by cyclophosphamide in mice. *Fitoterapia*. 2012;83(4):636–42.
- Yoon S-J, Park J-Y, Choi S, Lee J-B, Jung H, Kim T-D, Yoon SR, Choi I, Shim S, Park Y-J. Ginsenoside Rg3 regulates S-nitrosylation of the NLRP3 inflammasome via suppression of iNOS. *Biochem Biophys Res Commun*. 2015;463(4):1184–9.
- Kim D-G, Jung KH, Lee D-G, Yoon J-H, Choi KS, Kwon SW, Shen H-M, Morgan MJ, Hong S-S, Kim Y-S. 20 (S)-Ginsenoside Rg3 is a novel inhibitor of autophagy and sensitizes hepatocellular carcinoma to doxorubicin. *Oncotarget*. 2014;5(12):4438.
- Tian J, Fu F, Geng M, Jiang Y, Yang J, Jiang W, Wang C, Liu K. Neuroprotective effect of 20 (S)-ginsenoside Rg3 on cerebral ischemia in rats. *Neurosci Lett*. 2005;374(2):92–7.
- Lee H, Hong Y, Tran Q, Cho H, Kim M, Kim C, Kwon SH, Park S, Park J, Park J. A new role for the ginsenoside RG3 in antiaging via mitochondria function in ultraviolet-irradiated human dermal fibroblasts. *J Ginseng Res*. 2019;43(3): 431–41.
- Hwang JT, Lee MS, Kim HJ, Sung MJ, Kim HY, Kim MS, Kwon DY. Antiobesity effect of ginsenoside Rg3 involves the AMPK and PPAR- γ signal pathways. *Phytother Res*. 2009;23(2):262–6.
- Ginsberg HN, Maccallum PR. The obesity, metabolic syndrome, and type 2 diabetes mellitus pandemic: II. Therapeutic management of atherogenic dyslipidemia. *J Clin Hypertens*. 2009;11(9):520–7.
- Kim MJ, Koo YD, Kim M, Lim S, Park YJ, Chung SS, Jang HC, Park KS. Rg3 improves mitochondrial function and the expression of key genes involved in mitochondrial biogenesis in C2C12 myotubes. *Diabetes Metab J*. 2016; 40(5):406–13.
- Ito K, Carracedo A, Weiss D, Arai F, Ala U, Avigan DE, Schafer ZT, Evans RM, Suda T, Lee C-H, et al. A PML-PPAR- δ pathway for fatty acid oxidation regulates hematopoietic stem cell maintenance. *Nat Med*. 2012;18(9):1350–8.

16. Vannini N, Campos V, Girotra M, Trachsel V, Rojas-Sutterlin S, Tratwal J, Ragusa S, Stefanidis E, Ryu D, Rainer PY, et al. The NAD-booster nicotinamide riboside potently stimulates hematopoiesis through increased mitochondrial clearance. *Cell Stem Cell*. 2019;24(3):405–18.
17. Huang Y, Kwan KKL, Leung KW, Yao P, Wang H, Dong TT, Tsim KWK. Ginseng extracts modulate mitochondrial bioenergetics of live cardiomyoblasts: a functional comparison of different extraction solvents. *J Ginseng Res*. 2019;43(4):517–26.
18. Xing W, Yang L, Peng Y, Wang Q, Gao M, Yang M, Xiao X. Ginsenoside Rg3 attenuates sepsis-induced injury and mitochondrial dysfunction in liver via AMPK-mediated autophagy flux. *Biosci Rep*. 2017;37(4):BSR20170934.
19. Quan X, Nguyen TT, Choi S-K, Xu S, Das R, Cha S-K, Kim N, Han J, Wiederkehr A, Wollheim CB, et al. Essential role of mitochondrial Ca²⁺ uniporter in the generation of mitochondrial pH gradient and metabolism-secretion coupling in insulin-releasing cells. *J Biol Chem*. 2015;290(7):4086–96.
20. Denu RA, Hematti P. Effects of oxidative stress on mesenchymal stem cell biology. *Oxidative Med Cell Longev*. 2016;2016:2989076.
21. García-Prat L, Martínez-Vicente M, Perdiguero E, Ortet L, Rodríguez-Ubreva J, Rebollo E, Ruiz-Bonilla V, Gutarra S, Ballestar E, Serrano AL, et al. Autophagy maintains stemness by preventing senescence. *Nature*. 2016;529(7584):37–42.
22. Sun M, Huang C, Wang C, Zheng J, Zhang P, Xu Y, Chen H, Shen W. Ginsenoside Rg3 improves cardiac mitochondrial population quality: mimetic exercise training. *Biochem Biophys Res Commun*. 2013;441(1):169–74.
23. St-Pierre J, Drori S, Uldry M, Silvaggi JM, Rhee J, Jäger S, Handschin C, Zheng K, Lin J, Yang W, et al. Suppression of reactive oxygen species and neurodegeneration by the PGC-1 transcriptional coactivators. *Cell*. 2006; 127(2):397–408.
24. Iacovelli J, Rowe GC, Khadka A, Diaz-Aguilar D, Spencer C, Arany Z, Saint-Geniez M. PGC-1 α induces human RPE oxidative metabolism and antioxidant capacity. *Invest Ophthalmol Vis Sci*. 2016;57(3):1038–51.
25. Chacko BK, Kramer PA, Ravi S, Benavides GA, Mitchell T, Dranka BP, Ferrick D, Singal AK, Ballinger SW, Bailey SM, et al. The Bioenergetic Health Index: a new concept in mitochondrial translational research. *Clin Sci (Lond)*. 2014; 127(6):367–73.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

