



Abubidentin A, New Oleanane-type Triterpene Ester from *Abutilon bidentatum* and its antioxidant, cholinesterase and antimicrobial activities

Gadah A. Al-Hamoud¹ Nawal M. Al-Musayeib¹ Musarat Amina¹
Sabrin R.M. Ibrahim²

¹Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia

²Pharmacognosy, Faculty of Pharmacy, Assiut University, Assiut, Egypt

ABSTRACT

Background. This work describes the phytochemical and biological investigation of aerial parts of *Abutilon bidentatum* Hochst. Of Saudi origin.

Methodology. Petroleum ether fraction of ethanolic extract *A. bidentatum* was fractionated on a silica gel column and further purified with different chromatographic procedures for the isolation of chemical compounds. The chemical structures of all the pure isolated compounds were elucidated by the interpretation of their spectral data using IR, UV, ¹H, ¹³C NMR, and MS spectroscopy and chemical methods (alkaline hydrolysis) as well as comparison with data reported in the literature. The extract and isolated compounds were evaluated for antioxidant, cholinesterase inhibitory, and antimicrobial activities.

Results. A new oleanane-type triterpene ester, namely abubidentin A (3) (α , 3β , 30-trihydroxy-29-carboxy-olean-9(11), 12-diene-3-dotriacontanoate), along with two known compounds: 2-hydroxydocosanoic acid (1) and stigmasta-22-ene-3- β -ol (2) were isolated from the aerial parts of *Abutilon bidentatum* Hochst. (Malvaceae). Concerning the biological potential, the abubidentinA displayed antioxidant, cholinesterase inhibitory and antimicrobial activities. AbubidentinA possessed strong antioxidant activity against DPPH and ABTS⁺ radical scavenging assays. This new triterpene exhibited high inhibition against acetylcholinesterase (IC₅₀ 38.13 \pm 0.07 μ g mL⁻¹) and butyrylcholinesterase (IC₅₀ 32.68 \pm 0.37 μ g mL⁻¹). Abubidentin A displayed promising antimicrobial activity against *Escherichia coli*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* (125–150 μ g mL⁻¹).

Conclusion. These findings suggest *A. bidentatum* can contribute as a source of new biologically active compounds, especially antioxidants and antimicrobial agents.

Subjects Biochemistry, Biomolecules, Bioorganic Chemistry, Natural Products

Keywords *Abutilon bidentatum*, Malvaceae, Abubidentin A, Oleanane-type triterpene, Antioxidant, Cholinesterase, Antimicrobial

Submitted 25 October 2021
Accepted 9 February 2022
Published 8 March 2022

Corresponding authors

Gadah A. Al-Hamoud,
galhamoud@ksu.edu.sa
Nawal M. Al-Musayeib,
nalmusayeib@ksu.edu.sa

Academic editor

Carlos Fernández Marcos

Additional Information and
Declarations can be found on
page 13

DOI 10.7717/peerj.13040

© Copyright

2022 A. Al-Hamoud et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

INTRODUCTION

Abutilon Miller is the extensive genus of family Malvaceae that contains around 150 species of annual or perennial herbs, shrubs or undershrubs, and small trees. This plant species is native to temperate, subtropical, and tropical areas of Africa, Asia, America and Australia (Arbat, 2012). The word Abutilon is the Greek ancient name of the mulberry tree and has been assigned to genus due to its close similarity to the morphology of leaves. The genus Abutilon showed the existence of precious insoluble fibers isolated from different species of this genus and has noteworthy importance (Gomaa et al., 2016). The species of Abutilon genus have been used to cure specific health issues including; rheumatoid arthritis, diuretic and demulcent due to the occurrence of high mucilage content (Ali et al., 2014; Baquar, 1989). Numerous important pharmacological attributes such as hepatoprotective, antioxidant, analgesic, antipyretic, anti-inflammatory, antimicrobial, anticancer, diuretic, anti-hyperglycemic, CNS activity, immunostimulant, anti-hyperlipidemias, anti-hypertensive, antidiarrheal, anti-urolithiatic, and wound healing activities have been assigned to the plant species of this genus (Khadabadi & Bhajipale, 2010; Agrawal, 2017). Phytochemical studies claimed that the genus contain different secondary metabolites including phenolic acids, flavonoids, triterpenes, sterols, coumarins, quinones, alkaloids, anthocyanins, iridoids, saponins, megastigmanes, and fatty acids (Gomaa et al., 2018). In Saudi Arabia, it is represented by five species, namely *A. bidentatum*, *A. figarianum*, *A. fruticosum*, *A. hirtum*, and *A. pannosum* (Migahid, 1978).

Abutilon bidentatum Hochst. locally known as Ren-Umbro is an erect shrub, 1–2 m in height with stellate pubescent branches mixed with long simple hairs. Leaves broadly ovate, long-petiolate blade up to c. 12(–17) × 10(–13) cm, deeply cordate at base, sparse, stellate tomentose, serrate, acuminate at apex, velvety on both sides with serrate-dentate margins. Flowers are present in the leaf axils or on a shoot axillary. Seeds are papillose, 2.5 mm long stellately spreading, and black. Different parts of *A. bidentatum* have been utilized to treat several ailments in ethno-medicine, especially its leaves are used to cure infections. Root powder is used in folk-traditional medicine to treat dysentery, colitis, and diarrhea (Al-Shanwani, 1996). Literature scrutiny revealed that this plant remained unexplored for phytochemical and biological properties. Only three studies could be traced concerning the chemical constituents and pharmacological activities till date. These studies reported the antibacterial, antioxidant (Survase, Jamdhade & Chavan, 2012; Shahwar et al., 2010) and hepatoprotective potential of aerial parts of *A. bidentatum* (Yasmin, Akram Kashmiri & Anwar, 2011). However, a single study with respect to its chemical constituents, reporting the isolation of cholestane derivative from the *A. bidentatum* has been reported in the literature (Jain, Jain & Arora, 1996). The present study reports the isolation and structural characterization of a new oleanane-type triterpene ester: abubidentin A (3, Fig. 1) and two known compounds (1 and 2) from the aerial parts of *A. bidentatum*.




Figure 1 Structures of isolated compounds 1–3.

Full-size DOI: 10.7717/peerj.13040/fig-1

MATERIALS & METHODS

Chemicals and reagents

All the chemicals and reagents used for this study including; ethanol (96%), methanol (99.8%), petroleum ether (40–60 °C), dichloromethane ($\geq 99.8\%$), chloroform (99.9%), n-butanol (99.8%), dimethyl sulfoxide (DMSO), hydrochloric acid (HCl), potassium hydroxide (KOH, $\geq 85\%$), 1,1-diphenyl-2-picrylhydrazyl (DPPH, 95%), 2, 20-azino-bis[3-ethyl benzo thiazoline-6-sulphonic acid] (ABTS), potassium persulfate ($K_2S_2O_8$), acetylthiocholine iodide ($\geq 99.0\%$), S-butyrylthiochoilne iodide ($\geq 98.0\%$), sodium phosphate (Na_3PO_4 , 96%), ascorbic acid, resazurin, 2-nitrobenzoic acid and were obtained from Sigma Aldrich (Hamburg, Germany). Muller Hinton agar and Muller Hinton broth were procured from HiMedia (Himedia Laboratories Pvt Ltd., Mumbai, India). Donepezil, galantamine and chloramphenicol were purchased from the local drug store.

Instrumentation

Optical rotation: Perkin-Elmer Model 341 LC polarimeter (Perkin-Elmer, Waltham, MA, USA). UV spectra: in MeOH using a Perkin-Elmer Lambda 25 UV/VIS spectrophotometer (Perkin-Elmer, Waltham, MA, USA). IR spectra: Shimadzu Infrared-400 spectrophotometer (Shimadzu, Kyoto, Japan). ESIMS spectra: Agilent 6320 Ion trap mass spectrometer (Agilent Technologies, Santa Clara, CA, USA) equipped with an electrospray ionization interface. NMR spectra: Bruker DRX 700 spectrometer (Bruker, Rheinstetten, Germany). Column chromatographic separations: silica gel 60 (0.04–0.063

mm, Merck, Darmstadt, Germany). TLC analyses: pre-coated silica gel F254 aluminum sheets (Merck, Darmstadt, Germany).

Enzymes and pathogenic strains

The acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) were obtained from the mouse brain and human blood at the Pharmacology Department of King Saud University, Saudi Arabia. *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), and *Staphylococcus aureus* were provided by Microbiology Department, King Khaled University Hospital (KKUH), Saudi Arabia.

Plant material

The fresh aerial parts of *Abutilon bidentatum* Hochst. were collected from Jazan city, Aseer region, Saudi Arabia in March 2009. The plant was kindly identified by Dr. Mohamed Yusuf at the Pharmacognosy Department, College of Pharmacy, King Saud University. A voucher specimen (#16022) was deposited in the herbarium of the Pharmacognosy Department.

Extraction and isolation

Air dried-powdered aerial parts of *A. bidentatum* (900 g) were exhaustively extracted by maceration with 96% EtOH (3L × 7) with shaking. The combined EtOH extract was dried under reduced pressure at 45 °C to give a dry total extract weighing 89 g (EE). A part of the dried residue (87 g) was suspended in 450 mL distilled water and fractionated by shaking with petroleum ether (500 mL × 6), dichloromethane (500 mL × 4), ethyl acetate (500 mL × 5), and n-BuOH (500 mL × 6), respectively. Each combined extract was individually evaporated under reduced pressure to give the petroleum ether (PEF, 13 g), dichloromethane (DCF, 4 g), ethyl acetate (EAF, 6 g), n-BuOH (BF, 9 g), and aqueous soluble (AF, 55 g) fractions. Therefore, part of PEF (11.5 g) was chromatographed on SiO₂ column (330 g, 5 × 90 cm). Elution was started with petroleum ether and the polarity was gradually increased with CH₂Cl₂, followed by CH₂Cl₂: MeOH mixtures up to 100% MeOH to afford seven subfractions: AB-1 to AB-7. TLC pattern of subfractions AB-2 and AB-5 showed the presence of prominent compounds and were selected for further column chromatography. Subfraction AB-2 (33 mg) was chromatographed over silica gel using petroleum ether:CH₂Cl₂ gradient to give **1** (6.7 mg, white powder). Subfraction AB-5 (311 mg) was applied on SiO₂ column (15 g, 70 × 1 cm), using petroleum ether:CH₂Cl₂ (95:5 to 80:20) to give two subfractions AB-5.1 and AB-5.2 containing two major spots on TLC. Subfraction AB-5.1 (43 mg) was purified on SiO₂ column, eluted with petroleum ether:CH₂Cl₂ gradient to yield **2** (10 mg, white amorphous powder). Sub fraction AB-5.2 was similarly treated as subfraction AB-5.1 to give **3** (12 mg, white amorphous powder).

Abubidentin A (1 α ,3 β ,30-Trihydroxy-29-carboxy-olean-9(11), 12-diene-3-dotriacontanoate; **3**)

White amorphous powder. $[\alpha]_D^{+45}$ ($c = 0.1$, MeOH). UV (MeOH) λ_{\max} ($\log \epsilon$): 212 (3.10), 275 (2.21) nm. IR (KBr): 3,420, 2,947, 1,732, 1,712, 1,636, 1,245 cm⁻¹. NMR data (CDCl₃, 700 and 176 MHz) see Table 1. (+) ESI-MS m/z : 971.4 [M+Na]⁺; (-) ESI-MS m/z : 947.8 [M-H]⁻.

Table 1 NMR spectral data of compound 3 (500 and 125 MHz).

Position	δ_H [mult., J (Hz)]	δ_C (mult.)	HMBC
1	4.17 (<i>d</i> , $J = 2.8$)	71.6 CH	2, 3, 5, 10, 25
2	1.65–1.66 (m) 1.42–1.43 m	30.4 CH ₂	3, 4, 10
3	5.09 (<i>dd</i> , $J = 12.0, 5.0$)	74.3 CH	1', 1, 4, 23, 24
4	–	39.6 C	–
5	1.48–1.50 (m)	44.0 CH	3, 4, 10, 24, 25
6	1.61–1.63 (m) 1.45–1.47 (m)	16.2 CH ₂	–
7	2.05–2.07 (m) 1.81–1.82 (m)	30.9 CH ₂	–
8	–	37.1 C	–
9	–	149.5 C	–
10	–	43.4 C	–
11	5.73 (<i>d</i> , $J = 6.0$)	116.0 CH	8, 9, 10, 13
12	5.58 (<i>d</i> , $J = 6.0$)	119.6 CH	9, 14, 18
13	–	146.0 C	–
14	–	39.9 C	–
15	1.68–1.70 (m) 1.25–1.28 (m)	24.1 CH ₂	13
16	1.88–1.91 (m) 0.98–1.00 (m)	24.6 CH ₂	14
17	–	32.8 C	–
18	2.18 (<i>dd</i> , $J = 12.8, 5.6$)	41.4 CH	12, 13, 19, 22, 29, 30
19	1.63–1.65 (m) 1.05–1.08 (m)	45.1 CH ₂	17, 21, 29, 30
20	–	30.2 C	–
21	1.15–1.17 (m) 1.00–1.02 (m)	33.5 CH ₂	29, 30
22	1.48–1.50 (m) 1.32–1.34 (m)	33.8 CH ₂	–
23	0.97 (s)	24.6 CH ₃	3, 4, 5, 24
24	0.91 (s)	15.8 CH ₃	3, 4, 5, 23
25	1.26 (s)	22.5 CH ₃	1, 5, 9
26	1.17 (s)	19.4 CH ₃	9, 14
27	1.07 (s)	19.2 CH ₃	8, 13
28	0.92 (s)	27.2 CH ₃	17, 18, 22
29	–	173.0 C	–
30	4.06 (<i>d</i> , $J = 10.2$) 3.81 (<i>d</i> , $J = 10.2$)	69.6 CH ₂	19, 20, 21, 29
1'	–	172.3 C	–
2'	2.33 (<i>t</i> , $J = 6.8$)	32.0 CH ₂	1'
3'	1.62–1.65 (m)	24.1 CH ₂	–
(CH ₂) ₂₆	1.31–1.27 (m)	28.7–28.1 CH ₂	–

(continued on next page)

Table 1 (continued)

Position	δ_H [mult., J (Hz)]	δ_C (mult.)	HMBC
30	1.44–1.46 (m)	31.9 CH ₂	32'
31'	1.12–1.14 (m)	21.7 CH ₂	32'
32'	0.89 (t, J = 6.8)	13.1 CH ₃	31', 30'

Alkaline hydrolysis of compound 3

Compound 3 (4 mg) was dissolved in five mL of 3% KOH/MeOH solution and kept undisturbed for 15 min at ambient temperature. After 15 min, 1 N HCl/MeOH was added to the solution for the neutralization. The solution was then partitioned with CHCl₃, CHCl₃ layer was separated, evaporated and the obtained residue was taken up for column chromatography on SiO₂ (mesh 60–120) using as eluent hexane: EtOAc gradient to provide methyl ester of dotriacontanoic acid, which was verified by GC/MS (Ibrahim, Mohamed & Ross, 2016; El-Shanawany et al., 2015; Al-Musayeib et al., 2013). A 500-Clarus Perkin-Elmer GC/MS (Waltham, MA, USA) was applied for GC/MS analysis. The integrator combined with software (4.5.0.007 version) controller was turbo mass. A 5 MS/GC elite capillary (30 × 0.25 mm × 0.5 μm) column and helium (He) a carrier gas at 2 ml/min flow rate (55.8 cm/s flow initial with 32 p.s.i., split; 1:40) were applied. Temperature conditions including; source temperature, inlet line temperature, emission trap and electron energy were adjusted at 150 °C, 200 °C, 100 °C and 70 eV, respectively. The injector temperature at 220 °C was maintained. Whereas, the temperature of the column was set at 50 °C for 5 min, raised to 220 °C at the 20 C/min rate. MS was scanned from 50 to 650 m/z.

Antioxidant activity

DPPH radical scavenging activity

The ability of the samples to scavenge DPPH radical was determined by Wang, Chen & Hou (2019) method with slight modification (Wang, Chen & Hou, 2019). A total of 20 μL of different sample concentrations (5.25–50 mg mL⁻¹) solutions was reacted with 180 μL of DPPH• (6–5 M) dissolved in methanol (80%) in a 96-well plate. Samples were incubated at ambient temperature under dark conditions for 30 min and the absorbance wavelength was measured at 517 nm against a blank sample. DPPH• solution and ascorbic acid were used as blank samples and positive control, respectively. The inhibitory concentration (IC₅₀) was expressed as the concentration that inhibits the 50% of DPPH and calculated using the following equation

$$\text{Inhibition percentage (\%)} = \frac{(A_b - A_s)}{A_b} \times 100$$

where A_b and A_s are the absorbance values of the blank and test samples, respectively.

ABTS radical scavenging activity

ABTS radical cation decolorization assay was performed to measure the total antioxidant activity of the samples by obeying the previously described method (Re et al., 1999). In brief, ABTS•⁺ radicals were generated by treating 7 mM ABTS⁺ aqueous solution with 2.4 mM potassium persulfate for 12–16 h at room temperature in the dark. Prior to use, this solution was diluted with ethanol (approx. 1:89 v/v) and equilibrated to 0.7,000 ±

0.02 at 300 °C absorbance at 734 nm to obtain the working solution. Afterward, a 1.0 mL of working solution was reacted with 20 μL (1 mg mL^{-1}) samples and incubated for 30 min. After incubation, samples were scanned at 734 nm absorbance wavelength and the percentage of inhibition was determined. Ascorbic acid (AA) was applied reference standard.

Evaluation of cholinesterase inhibitory activities

The cholinesterase inhibitory effect of test sample was investigated on two enzymes (acetylcholinesterase and butyrylcholinesterase) by spectrophotometric method by following Ellman et al. procedure with little modification (Ellman et al., 1961). Crude enzymes, AChE (acetylcholinesterase) and BChE (butyrylcholinesterase) were collected from brain of mice and blood of humans, respectively, by obeying earlier described procedure (Asaduzzaman et al., 2014; Uddin et al., 2015). The AChE and BChE assays were tested by using two chemical substrates acetylthiocholine iodide and S-butyrylthiocholine iodide, respectively. In brief, 10 μL of each enzyme was reacted individually with equal volume (10 μL) of different concentration ($25\text{--}400 \mu\text{g mL}^{-1}$) test sample and reference standard followed by incubation at 37 °C for 15 min for the complete interaction. Afterwards, 2-nitrobenzoic acid (1 mM, 62 μL), sodium phosphate buffer (50 mM, 25 μL , pH 8) provided with bovine albumin serum (0.1%) and acetylcholine iodide (0.5 mM, 13 μL) were added separately into each reaction mixture. Each reaction mixture was individually incubated further for 15 min at 37 °C and absorbance at 405 nm was immediately noted against the blank. Donepezil and galantamine were used as reference compounds for AChE and BChE activity, respectively. All the experiments were performed in triplicates to avoid error and the results were estimated through the two-tailed Student's *t*-test at a $p < 0.05$ significance. The inhibition percentage of cholinesterase activity was calculated using the following equation

$$\text{I\%} = \frac{A_c - A_s}{A_c} \times 100$$

where A_c and A_s is the absorbance of control and sample or reference compound. IC₅₀ values could be determined from the dose response curve obtained by plotting the percent inhibition values against test concentrations of each compound.

Antimicrobial activity

Minimum inhibitory concentration (MIC) assay was performed to evaluate the *in vitro* antimicrobial activities using a broth microdilution method (Nascente, 2009). Three bacterial strains *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), and *Staphylococcus aureus* were applied to examine the antimicrobial activity of samples. Briefly, the microbial strains were transferred to Muller Hinton agar (MHB, HiMedia, India) and 24-h colonies were individually suspended in 10 mL Muller Hinton broth (MHB, HiMedia, India). The suspension of each microbial strain was standardized at 575 nm wavelength using a spectrophotometer (CRAIC Technologies, CA, USA), to match the McFarland scale ($1.5 \times 10^8 \text{ CFU mL}^{-1}$). The standardized suspension was further diluted to obtain a final concentration of $5 \times 10^5 \text{ CFU mL}^{-1}$. The samples prepared in DMSO

at 1 mgmL^{-1} and different concentrations ($30\text{--}500 \text{ }\mu\text{gmL}^{-1}$) were attained after dilution in Mueller Hinton broth. Three inoculated wells containing different concentrations of DMSO (4% to 1% range), one non-inoculated well without an antimicrobial agent and negative controls were included. The inoculated well was used to monitor whether the broth was sufficient for the microbial strain to grow. Chloramphenicol (500 to 30 mgmL^{-1}) was applied as a positive control. The sample treated 96-well microplates were sealed and incubated for 24 h at 37°C . After a 1-day incubation, 30 ml of 0.02% resazurin solution was added into each well to examine the viability of the microbial strain (Palomino *et al.*, 2002). The minimum used concentration of test sample that inhibited the microorganism growth (MIC value) was calculated as the minimum concentration of the test samples required to prevent the color change of the resazurin solution. All the assays were performed in triplicates.

Statistical analysis

The experiments were conducted in triplicates and results were expressed as mean \pm standard deviation. Statistical and graphical analysis were performed on Graph Pad Prism (version 8.0.1) and Microsoft Excel 2010. *T*-test was carried out to determine the statistical significance between the average values and ($p < 0.05$) was considered significant.

RESULTS & DISCUSSION

Compound **3** was isolated as a white amorphous powder. It gave a positive Liebermann-Burchard reaction, indicating its triterpenoidal nature (Al-Musayeb *et al.*, 2013; Ibrahim *et al.*, 2012). The molecular formula of **3** was determined as $\text{C}_{62}\text{H}_{108}\text{O}_6$ on the basis of the ESI-MS pseudo-molecular ion peaks at m/z 971.4 $[\text{M}+\text{Na}]^+$ and 947.8 $[\text{M}-\text{H}]^-$. The IR spectrum showed the presence of hydroxyl ($3,420 \text{ cm}^{-1}$), ester carbonyl ($1,732 \text{ cm}^{-1}$), acid carbonyl ($1,712 \text{ cm}^{-1}$), and double bond ($1,636 \text{ cm}^{-1}$) functionalities. The ^{13}C and DEPT NMR spectra of **3** displayed resonances for 62 carbons: six tertiary methyl, one primary methyl, 39 methylenes, one of them for an oxymethylene $\delta(\text{C})$ 69.6 (C-30), six methines, and ten quaternary carbon signals, including one ester carbonyl at $\delta(\text{C})$ 172.3 (C-1'), acid carbonyl $\delta(\text{C})$ 173.0 (C-29), and two quaternary olefinic carbons at $\delta(\text{C})$ 149.5 (C-9) and 146.0 (C-13). The ^1H NMR spectrum showed six methyl singlets at $\delta(\text{H})$ 0.97 H-C(23), 0.91 H-C(24), 1.26 H-C(25), 1.17 H-C(26), 1.07 H-C(27), and 0.92 H-C(28), characteristic for an oleanane-type triterpenoid (Ibrahim, Mohamed & Ross, 2016; Al Musayeb *et al.*, 2014; Mahato & Kundu, 1994). They correlated to the carbon signals, resonating at $\delta(\text{C})$ 24.6, 15.8, 22.5, 19.4, 19.2, and 27.2, respectively in the HSQC spectrum. Moreover, two coupled olefinic protons at $\delta(\text{H})$ 5.73 (d , $J = 6.0$, H-C(11)) and 5.58 (d , $J = 6.0$, H-C(12)) were observed in the ^1H NMR and 1H-1H COSY spectra, indicating the presence of two tri-substituted olefinic double bonds (Fig. 2). They showed HSQC cross peaks to the carbons at $\delta(\text{C})$ 116.0 C(11) and 119.6 C(12). The placement of the double bonds at C9-C11 and C12-C13 was established based on the HMBC cross peaks of H-C(11) to C(8), C(9), C(10), and C(13), H-C(12) to C(9), C(13), and C(14), H-C(18) to C(12) and C(13), H-C(27) to C(13), and H-C(25) and H-C(26) to C(9), indicating that **3** had an olean-9(11),12-diene skeleton (Caceres-Castillo *et al.*, 2008;

Mahato & Kundu, 1994) (Fig. 2). The ^1H and ^{13}C NMR spectra displayed two oxymethine signals at $\delta(\text{H})$ 4.17 (*d*, $J = 2.8$, H-C(1))/ $\delta(\text{C})$ 71.6 (C-1) and 5.09 (dd, $J = 12.0$, 5.0 Hz, H-C(3))/74.3 (C-3). Their assignment was established based on the observed 1H-1H COSY of H-C(2) to H-C(1) and H-C(3) as well as the HMBC cross peaks of H-C(1), H-C(23), and H-C(24) to C(3) and H-C(3) and H-C(25) to C(1). The coupling constants of H-C(1) ($J_{1,2} = 2.8$ Hz) and H-C(3) ($J_{3,2} = 12.0$, 5.0 Hz) revealed that the hydroxyl groups at C(1) and C(3) were α - and β -configured, respectively, based on comparison with the previously reported triterpenoids [$J_{1,2} = 2.5$ –3.0 Hz and $J_{3,2} = 10.0$ –12.8, 4.2–7.2 Hz] (*Ibrahim et al., 2019; Litaudon et al., 2009; Rogers & Subramony, 1988*). The signals at $\delta(\text{H})$ 4.06 and 3.81 (2H, each *d*, $J = 10.2$, H-C(30))/ $\delta(\text{C})$ 69.6 (C-30) and $\delta(\text{C})$ 173.0 (C-29) revealed the presence of an oxymethylene and carboxyl functionalities in **3**. Their placement at (C-20) was established by the HMBC cross peaks of H-C(19) and H-C(21) to (C-29) and (C-30) and H-C(30) to (C-29). The multiple aliphatic protons at $\delta(\text{H})$ 1.31–1.27 were attributed to the presence of a long chain aliphatic moiety. The ^{13}C NMR signal at $\delta(\text{C})$ 172.3 (C-1') was assigned to an ester carbonyl, which was confirmed by the IR absorption band at $1,732\text{ cm}^{-1}$. While the signals at $\delta(\text{H})$ 0.89 (t, $J = 6.8$)/ $\delta(\text{C})$ 13.1 were assigned to the terminal methyl group of a straight-chain fatty acid (*Al-Musayeib et al., 2013; El-Shanawany et al., 2015; Ibrahim, Mohamed & Ross, 2016*). Upon alkaline hydrolysis of **3**, it gave a methyl ester of dotriacontanoic acid, which was identified by GC-MS molecular ion peak at m/z 494 $[\text{M}]^+$ and confirmed by the ESI-MS fragment ion peak at m/z 486 $[\text{M}+\text{H}-(\text{dotriacontanoyl moiety})]^+$. The attachment of the fatty acyl moiety at (C-3) was established by the HMBC cross peak of H-C(3) to (C-1') and confirmed by the downfield shift of HC-(3) $\delta(\text{H})$ 5.09. The relative stereochemistry at stereocenters was assigned by comparing the J values and ^1H and ^{13}C chemical shifts with those of related triterpenes (*Ibrahim et al., 2019; Litaudon et al., 2009*). Based on these findings, **3** was assigned as 1 α ,3 β ,30-trihydroxy-29-carboxy-olean-9(11),12-diene-3-dotriacontanoate and named abubidentin A.

The known compounds were identified as 2-hydroxydocosanoic acid (**1**) (*Inagaki et al., 2001*) and stigmasta-22-ene-3- β -ol (**2**) (*Connor et al., 2006*) by comparing their NMR spectral and physical data in the literature.

Antioxidant activity assessments

DPPH and ABTS⁺ assays are widely applied to measure the compound's ability to determine its antioxidant potential. Both the spectrophotometric methods used for evaluating antioxidant activity are based on electron transfer reactions and visually rely on the reduction of a colored oxidant. The obtained results from these two assays expressed good correlation. Fig. 3A and Table 2 represents the free radical inhibition of isolated compounds (**1**, **2**, and **3**) and ascorbic acid (AA, standard) at different concentrations. The results showed that the DPPH scavenging potential of compounds **1**, **2**, and **3** incrementally increased with the increase in concentration of compounds. The IC₅₀ values obtained were 10.82 ± 0.24 , 7.60 ± 0.42 , and $4.67 \pm 0.28\ \mu\text{g mL}^{-1}$ for compounds **1**, **2**, and **3** were, respectively, indicating that compound **3** possesses the highest radical scavenging potential




Figure 2 Some Key ^1H - ^1H COSY (–) and HMBC (H → C) correlations of 3.

Full-size DOI: 10.7717/peerj.13040/fig-2




Figure 3 (A) DPPH and (B) ABTS free radical scavenging activity of compound 1, 2, 3 and ascorbic acid. Values were obtained as mean \pm standard deviation and *mean significant difference compared to control ($p < 0.05$).

Full-size DOI: 10.7717/peerj.13040/fig-3

and was 1.55 fold lower than ascorbic acid (IC_{50} $3.12 \pm 0.24 \mu\text{g mL}^{-1}$), followed by compounds 2 and 1 ($p < 0.05$).

The ABTS^+ assay is an additional important procedure for the quantification of radical scavenging potential that can provide parallel results to those obtained in the DPPH assay. The results showed that all the three isolated compounds exerted significant ABTS^+ free radical scavenging activity and had an antioxidant potential proportional to that of ascorbic acid (Fig. 3B and Table 2). Compound 3 was found to be the most active radical scavenger with an IC_{50} value of $6.42 \pm 0.25 \mu\text{g mL}^{-1}$ and 3.12 fold lower than that of ascorbic acid, followed by compound 2 and 1 with IC_{50} values of 8.18 ± 0.13 and 11.45

Table 2 Antioxidant activity and cholinesterase inhibitory of the extract and pure isolated compounds (1, 2 and 3) from of *A bidentatum*.

Sample	Antioxidant activity		Cholinesterase inhibitory	
	DPPH IC ₅₀ ($\mu\text{g mL}^{-1}$)	ABTS ⁺ IC ₅₀ ($\mu\text{g mL}^{-1}$)	AChE IC ₅₀ ($\mu\text{g mL}^{-1}$)	BChE IC ₅₀ ($\mu\text{g mL}^{-1}$)
Ethanol extract	16.34 \pm 0.25e	18.65 \pm 0.67e	132.56 \pm 1.4e	142.35 \pm 0.54e
Compound 1	10.82 \pm 0.24 ^d	11.45 \pm 0.37 ^d	121.97 \pm 1.61 ^d	137.76 \pm 0.67 ^d
Compound 2	7.60 \pm 0.42 ^c	8.18 \pm 0.13 ^c	68.65 \pm 0.56 ^c	49.52 \pm 0.35 ^c
Compound 3	4.67 \pm 0.28 ^b	6.42 \pm 0.25 ^b	38.13 \pm 0.07 ^b	32.68 \pm 0.37 ^b
Ascorbic acid	3.12 \pm 0.24 ^a	4.45 \pm 0.17 ^a	–	–
Donepezil	–	–	9.32 \pm 0.38 ^a	–
Galantamine	–	–	–	10.27 \pm 0.88 ^a

Notes.

Means in each column with different subscript letters (a, b, c, d, e) differ significantly ($P < 0.05$).

$\pm 0.37 \mu\text{g mL}^{-1}$, respectively ($p < 0.05$). Compound 3 displayed both the highest DPPH and ABTS radical scavenging activities. A correlation between DPPH and ABTS methods applied to determine the antioxidant potential of the tested compounds was examined. The DPPH radical activity showed a strong correlation with ABTS radical activity ($R_2 = 96.67\%$).

Cholinesterase inhibitory activity

The soluble fraction of the extract and compounds (1–3) were screened for AChE and BChE inhibition at different concentrations. The percent inhibition of AChE by the test compounds were presented in Fig. 4A. Donepezil used as a reference AChE inhibitor showed an IC₅₀ value of $9.32 \pm 0.38 \mu\text{g mL}^{-1}$. The result revealed that all the investigated compounds (1–3) showed inhibition of the AChE enzyme in a dose-dependent manner. Among the compounds, high activity was displayed by compound 3 with IC₅₀ values of $38.13 \pm 0.07 \mu\text{g mL}^{-1}$, followed by compound 2 (IC₅₀ value = $68.65 \pm 0.56 \mu\text{g mL}^{-1}$). However, compound 1 showed low activity with IC₅₀ of $121.97 \pm 1.61 \mu\text{g mL}^{-1}$ (Table 2). Similarly, in BChE inhibitory assay, compounds 3 and 2 exerted high inhibitory potentials with IC₅₀ of 32.68 ± 0.37 and $49.52 \pm 0.35 \mu\text{g mL}^{-1}$, respectively (Table 2, Fig. 4B). The IC₅₀ value of compound 1 was $137.76 \pm 0.67 \mu\text{g mL}^{-1}$ exhibited very negligible effects. Thus, compound 3 had appreciable activity towards both AChE and BChE enzymes.

Antimicrobial activity

The antimicrobial potential of isolated compounds (1–3) was assessed by calculating MIC values and their ability to inhibit the growth of the tested microbial strain. The assay was performed in 96-well microplates by applying resazurin as a developer. The well plates that exhibited the blue color after the addition of resazurin were considered as the MIC value for each microorganism (Ostrosky et al., 2008). The pure isolated compounds 1–3 were examined against *E. coli*, *P. aeruginosa*, and *S. aureus* bacterial strains for their antimicrobial potential. All the tested compounds exerted antimicrobial effect in the range of $125\text{--}1,000 \mu\text{g mL}^{-1}$ towards the pathogenic strains (Table 3). However, the prominent activity was shown by compound 2 with MIC $\leq 125 \mu\text{g mL}^{-1}$, $\leq 250 \mu\text{g mL}^{-1}$, $\leq 150 \mu\text{g mL}^{-1}$




Figure 4 Cholinesterase inhibitory activities of compound 1–3 isolated from *A. bidentatum* (a) Inhibition of acetylcholinesterase (AChE) by isolated compounds of *A. bidentatum* and standard donepezil, (B) Inhibition of butyrylcholinesterase (BChE) by compounds isolated from *A. bidentatum* and standard galantamine.

Full-size [DOI: 10.7717/peerj.13040/fig-4](https://doi.org/10.7717/peerj.13040/fig-4)

Table 3 Minimum inhibitory concentration (µg mL⁻¹) of extract and pure isolated compounds (1, 2 and 3) against selected pathogens.

Sample	Microbial strain		
	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. aureus</i>
Ethanol extract	≤600	≤1,000	≤1,000
Compound 1	≤500	≤1,000	≤1,000
Compound 2	≤125	≤250	≤150
Compound 3	≤150	≤125	≤125
Chloramphenicol	≤40	≤40	≤40

and compound 3 with ≤150 µg mL⁻¹, ≤125 µg mL⁻¹, ≤125 µg mL⁻¹, against *E. coli*, *P. aeruginosa*, and *S. aureus*, respectively. There is no consistent classification with respect to MIC values (Aligiannis *et al.*, 2001), but the values obtained ≤1,000 µg mL⁻¹ were considered as satisfactory and sensitive (Webster *et al.*, 2008). Thus, the MIC value of compounds 3 and 2 can be considered as promising.

CONCLUSION

A new oleanane-type triterpene ester, namely abubidentin A together with two known 2-hydroxydocosanoic acid and stigmasta-22-ene-3-β-ol were isolated from aerial parts of *A. bidentatum*. The extracts and compounds were investigated for antioxidant, cholinesterase inhibitory, and antimicrobial activities. The outcomes demonstrated that the newly isolated compound possesses a strong antioxidant effect towards DPPH and ABTS+ radical scavenging assays. This new triterpene exhibited high inhibition against acetylcholinesterase and butyrylcholinesterase. In addition, the new compound also showed promising antimicrobial effects against tested microorganisms. These results suggested that *A. bidentatum* is a promising source of useful natural products and the new compound offers opportunities to develop a novel drug.

ACKNOWLEDGEMENTS

This work was supported by Researchers Supporting Project No.(RSP-2021/294), King Saud University, Riyadh, Saudi Arabia.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by Researchers Supporting Project No. RSP-2021/294, King Saud University, Riyadh, Saudi Arabia. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Researchers Supporting: RSP-2021/294.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Gadah A. Al-Hamoud conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Nawal M. Al-Musayeib analyzed the data, prepared figures and/or tables, and approved the final draft.
- Musarat Amina performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Sabrin R.M. Ibrahim conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:
The raw data is available in the [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13040#supplemental-information>.

REFERENCES

- Agrawal T.** 2017. Abutilon a vulnerable genus: a review. *Research and Reviews: Journal of Pharmacognosy and Phytochemistry* 5(1):44–46.
- Al Musayeib NM, Mothana RA, Ibrahim SR, El Gamal AA, Al-Massarani SM.** 2014. Klodorone A and klodorol A: new triterpenes from *Kleinia odora*. *Natural Product Research* 28(15):1142–1146 DOI 10.1080/14786419.2014.915831.

- Al-Musayeib NM, Mohamed GA, Ibrahim SR, Ross SA. 2013.** Lupeol-3-O-decanoate, a new triterpene ester from *Cadaba farinosa* Forssk. growing in Saudi Arabia. *Medicinal Chemistry Research* **22(11)**:5297–5302
DOI [10.1007/s00044-013-0536-1](https://doi.org/10.1007/s00044-013-0536-1).
- Al-Shanwani M. 1996.** *Plants used in the Saudi folk medicine*. Riyadh: King Abdul Aziz City for Science and Technology, 146.
- Ali B, Ibrahim M, Hussain I, Hussain N, Imran M, Nawaz H, Jan S, Khalid M, Ghous T, Akash MSH. 2014.** Pakistamide C, a new sphingolipid from *Abutilon pakistanicum*. *Revista Brasileira de Farmacognosia* **24**:277–281
DOI [10.1016/j.bjp.2014.07.002](https://doi.org/10.1016/j.bjp.2014.07.002).
- Aliagiannis N, Kalpoutzakis E, Mitaku S, Chinou IB. 2001.** Composition and antimicrobial activity of the essential oils of two *Origanum* species. *Journal of Agricultural and Food Chemistry* **49(9)**:4168–4170 DOI [10.1021/jf001494m](https://doi.org/10.1021/jf001494m).
- Arbat AA. 2012.** Pharmacognostic studies of stem of *Abutilon pannosum* (Forst f.). *Bioscience Discovery* **3(3)**:317–320.
- Asaduzzaman M, Uddin MJ, Kader MA, Alam AH, Rahman AA, Rashid M, Kato K, Tanaka T, Takeda M, Sadik G. 2014.** In vitro acetylcholinesterase inhibitory activity and the antioxidant properties of *Aegle marmelos* leaf extract: implications for the treatment of Alzheimer's disease. *Psychogeriatrics* **14(1)**:1–0
DOI [10.1111/psyg.12031](https://doi.org/10.1111/psyg.12031).
- Baquar SR. 1989.** Medicinal and poisonous plants of Pakistan. In: *Medicinal and poisonous plants of Pakistan*. Karachi, Pakistan: Printas.
- Caceres-Castillo D, Mena-Rejon GJ, Cedillo-Rivera R, Quijano L. 2008.** 21 β -Hydroxy-oleanane-type triterpenes from *Hippocratea excelsa*. *Phytochemistry* **69(4)**:1057–1064
DOI [10.1016/j.phytochem.2007.10.016](https://doi.org/10.1016/j.phytochem.2007.10.016).
- Connor WE, Wang Y, Green M, Lin DS. 2006.** Effects of diet and metamorphosis upon the sterol composition of the butterfly *Morpho peleides*. *Journal of Lipid Research* **47(7)**:1444–1448 DOI [10.1194/jlr.M600056-JLR200](https://doi.org/10.1194/jlr.M600056-JLR200).
- El-Shanawany MA, Sayed HM, Ibrahim SR, Fayed MA. 2015.** Stigmasterol tetra-cosanoate, a new stigmasterol ester from the Egyptian *Blepharis ciliaris*. *Drug Research* **65(07)**:347–353 DOI [10.1055/s-0034-1382064](https://doi.org/10.1055/s-0034-1382064).
- Ellman GL, Courtney KD, Andres Jr V, Featherstone RM. 1961.** A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* **7(2)**:88–95 DOI [10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9).
- Gomaa AA, Samy MN, Desoukey SY, Kamel MS. 2016.** Pharmacognostical studies of leaf, stem, root and flower of *Abutilon hirtum* (Lam.) Sweet. *International Journal of Pharmacognosy Phytochemistry Research* **8**:199–216
Available at <http://www.ijppr.com>.
- Gomaa AA, Samy MN, Desoukey SY, Kamel MS. 2018.** Phytochemistry and pharmacological activities of genus *Abutilon*: a review (1972–2015). *Journal of Advanced Biomedical and Pharmaceutical Sciences* **1(2)**:56–74
DOI [10.21608/jabps.2018.3333.1000](https://doi.org/10.21608/jabps.2018.3333.1000).

- Ibrahim SRM, Khedr AIM, Mohamed GA, Zayed MF, El-Kholy AAS, Al Haidari RA. 2019.** Cucumol B, a new triterpene benzoate from *Cucumis melo* seeds with cytotoxic effect toward ovarian and human breast adenocarcinoma. *Journal of Asian Natural Products Research* **21**(11):1112–1118 DOI [10.1080/10286020.2018.1488832](https://doi.org/10.1080/10286020.2018.1488832).
- Ibrahim SR, Mohamed GA, Ross SA. 2016.** Integracides F and G: New tetracyclic triterpenoids from the endophytic fungus *Fusarium* sp. *Phytochemistry Letters* **15**:125–130 DOI [10.1016/j.phytol.2015.12.010](https://doi.org/10.1016/j.phytol.2015.12.010).
- Ibrahim SR, Mohamed GA, Shaala LA, Banuls LM, Van Goietsenoven G, Kiss R, Youssef DT. 2012.** New ursane-type triterpenes from the root bark of *Calotropis procera*. *Phytochemistry Letters* **5**(3):490–495 DOI [10.1016/j.phytol.2012.04.012](https://doi.org/10.1016/j.phytol.2012.04.012).
- Inagaki M, Shibai M, Isobe R, Higuchi R. 2001.** Constituents of ophiuroidea. 1. Isolation and structure of three ganglioside molecular species from the brittle star *Ophiocoma scolopendrina*. *Chemical and Pharmaceutical Bulletin* **49**(12):1521–1525 DOI [10.1248/cpb.49.1521](https://doi.org/10.1248/cpb.49.1521).
- Jain R, Jain SC, Arora R. 1996.** A new cholestane derivative of *Abutilon bidentatum* Hochst. and its bioactivity. *Pharmazie* **51**(4):253 DOI [10.1002/chin.199637250](https://doi.org/10.1002/chin.199637250).
- Khadabadi SS, Bhajipale NS. 2010.** A review on some important medicinal plants of *Abutilon* spp. *Research Journal of Pharmaceutical, Biological and Chemical Sciences* **1**(4):718–729.
- Litaudon M, Jolly C, Le Callonec C, Cuong DD, Retailleau P, Nosjean O, Nguyen VH, Pfeiffer B, Boutin JA, Guéritte F. 2009.** Cytotoxic pentacyclic triterpenoids from *Combretum sundaicum* and *Lantana camara* inhibitors of Bcl-xL/BakBH3 domain peptide interaction. *Journal of Natural Products* **72**(7):1314–1320 DOI [10.1021/np900192r](https://doi.org/10.1021/np900192r).
- Mahato SB, Kundu AP. 1994.** ¹³C NMR spectra of pentacyclic triterpenoids—a compilation and some salient features. *Phytochemistry* **37**(6):1517–1575 DOI [10.1016/S0031-9422\(00\)89569-2](https://doi.org/10.1016/S0031-9422(00)89569-2).
- Migahid AM. 1978.** *Flora of Saudi Arabia*. Second Edition. 1. Riyadh: Riyadh University Publication, 23.
- Nascente PDS, Meinerz ARM, Faria ROD, Schuch LFD, Meireles MCA, Mello JRBD. 2009.** CLSI broth microdilution method for testing susceptibility of *Malassezia pachydermatis* to thiabendazole. *Brazilian Journal of Microbiology* **40**:222–226 DOI [10.1590/S1517-83822009000200002](https://doi.org/10.1590/S1517-83822009000200002).
- Ostrosky EA, Mizumoto MK, Lima ME, Kaneko TM, Nishikawa SO, Freitas BR. 2008.** Métodos para avaliação da atividade antimicrobiana e determinação da concentração mínima inibitória (CMI) de plantas medicinais. *Revista Brasileira de Farmacognosia* **18**:301–307 DOI [10.1590/S0102-695X2008000200026](https://doi.org/10.1590/S0102-695X2008000200026).
- Palomino JC, Martin A, Camacho M, Guerra H, Swings J, Portaels F. 2002.** Resazurin microtiter assay plate: simple and inexpensive method for detection of drug resistance in *Mycobacterium tuberculosis*. *Antimicrobial Agents and Chemotherapy* **46**(8):2720–2702 DOI [10.1128/AAC.46.8.2720-2722.2002](https://doi.org/10.1128/AAC.46.8.2720-2722.2002).

- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. 1999.** Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* **26(9–10)**:1231–1237 DOI [10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3).
- Rogers CB, Subramony G. 1988.** The structure of imberbic acid, A 1 α -hydroxy pentacyclic triterpenoid from *Combretum imberbe*. *Phytochemistry* **27(2)**:531–533 DOI [10.1016/0031-9422\(88\)83135-2](https://doi.org/10.1016/0031-9422(88)83135-2).
- Shahwar D, Ahmad N, Ullah S, Raza MA. 2010.** Antioxidant activities of the selected plants from the family Euphorbiaceae, Lauraceae, Malvaceae and Balsaminaceae. *African Journal of Biotechnology* **9(7)**:1086–1096 DOI [10.5897/AJB09.1622](https://doi.org/10.5897/AJB09.1622).
- Survase SA, Jamdhade MS, Chavan S. 2012.** Antibacterial activity of *Abutilon bidentatum* (Hochst.) leaves. *Science Research Reporter* **2(1)**:38–40.
- Uddin MN, Afrin R, Uddin MJ, Uddin MJ, Alam AH, Rahman AA, Sadik G. 2015.** *Vanda roxburghii* chloroform extract as a potential source of polyphenols with antioxidant and cholinesterase inhibitory activities: identification of a strong phenolic antioxidant. *BMC Complementary and Alternative Medicine* **15(1)**:1–9 DOI [10.1186/s12906-015-0728-y](https://doi.org/10.1186/s12906-015-0728-y).
- Wang CY, Chen YW, Hou CY. 2019.** Antioxidant and antibacterial activity of seven predominant terpenoids. *International Journal of Food Properties* **22(1)**:230–238 DOI [10.1080/10942912.2019.1582541](https://doi.org/10.1080/10942912.2019.1582541).
- Webster NS, Xavier JR, Freckelton M, Motti CA, Cobb R. 2008.** Shifts in microbial and chemical patterns within the marine sponge *Aplysina aerophoba* during a disease outbreak. *Environmental Microbiology* **10(12)**:3366–3376 DOI [10.1111/j.1462-2920.2008.01734.x](https://doi.org/10.1111/j.1462-2920.2008.01734.x).
- Yasmin S, Akram Kashmiri M, Anwar K. 2011.** Screening of aerial parts of *Abutilon bidentatum* for hepatoprotective activity in rabbits. *Journal of Medicinal Plants Research* **5(3)**:349–353.